ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 8, Number 4
December 2016

AGRICULTURAL SCIENCE AND TECHNOLOGY

2016

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Bojin Bojinov (Bulgaria)
Stoicho Metodiev (Bulgaria)
Svetlana Georgieva (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Zervas Georgios (Greece)
Ivan Varlyakov (Bulgaria)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Ramesh Kanwar (USA)
Martin Banov (Bulgaria)
Peter Cornish (Australia)

Product Quality and Safety

Marin Kabakchiev (Bulgaria)
Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions

on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate an author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to editorial review and the editors reserve the right to improve style and return the paper

for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc. and AGRIS (FAO).

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone.: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone.: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 8, Number 4
December 2016

AGRICULTURAL SCIENCE AND TECHNOLOGY

2016

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Quality of grain and flour of foreign bread wheat cultivars (*Triticum aestivum* L.) under the conditions of south Dobrudzha region

P. Chamurliyski^{1*}, N. Tsenov², I. Stoeva³, S. Doneva³, E. Penchev³

¹Department of Crop Science and Plant Protection, College Dobrich, Konstantin Preslavski University of Shumen, 9300 Dobrich, Bulgaria

²Agronom I Holding, 9300 Dobrich, Bulgaria

³Dobrudzha Agricultural Institute, 9521 General Toshevo, Bulgaria

(Manuscript received 23 March 2016; accepted for publication 28 September 2016)

Abstract. One of the main directions of the breeding programs in common winter wheat, besides increasing productivity, is developing of cultivars with excellent baking properties. An important prerequisite for this is the involvement of new gene plasma of variable origin, which is adequate to the growing conditions and the desired breeding direction. The aim of investigation is study of some main properties related to the grain quality and the baking properties of bread wheat accessions of foreign origin under the conditions of the South Dobrudzha region. Twenty-five foreign bread wheat cultivars of various origins were investigated for a three year period. Cultivars Aglika, Enola, Pryaspa and Yantur were used as standards. Some indices related to the quality of grain and flour were analyzed at the Bread Making Laboratory of Dobrudzha Agricultural Institute (DAI). The expression of the following parameters was followed: test weigh, % of protein, sedimentation, wet gluten yield, softening degree, pharinographic value, bread volume, and the quality index (QI) was calculated. The cultivars, which demonstrated high grain quality, were the Romanian Faur, Moldovan Dobropolka, American Wahoo and the Ukrainian Zmina. Averaged for the three years, highest variation was found for the index pharinographic value. On the whole, the materials with origin from Romania, Ukraine and USA were characterized with high values of the quality indices. Cultivars Faur and Zmina can be successfully included in the breeding program of DAI for development of strong wheat varieties.

Keywords: winter wheat, grain quality, baking properties

Introduction

In the contemporary world, wheat is of primary importance for human nutrition and is distributed in almost all geographic regions. The main task of the breeding programs in Bulgaria is developing of cultivars with high production potential in combination with very good quality of grain and flour (Tsenov et al., 2008; Chamurliyski et al., 2011). The long-term research work in this direction brought about high achievements (Panayotov et al., 2004) and lead to the dominance of group A cultivars (strong wheat varieties) which are competitive and which meet the demands of the dynamic market under the pressure of the western breeding (Stoeva, 2012).

The proper choice of initial parental forms is decisive for obtaining positive results. For many years, main sources of quality for Bulgarian breeding have been the Ukrainian and Russian wheats (Panayotov and Kostov, 2008; Tsenov et al., 2014). They combine high productivity, resistance to abiotic stress and excellent quality of grain and flour. Cultivar Bezostaya 1, for example, is included directly or indirectly in over 40 Bulgarian cultivars, which is an evidence for its enormous influence on the development of the breeding programs (Panayotov et al., 2005).

This has lead to partial genetic erosion and to the necessity of investigating new genetic plasma. Wheat is a crop of the micro climate which determines variable reaction of the accession under the environments of the specific locations (Tsenov et al., 2013), especially when breeding materials of foreign origin are concerned. The quality indices are highly dependent on the growing environment (Tsenov et al., 2004; Stoeva and Penchev, 2013). The

study on the working collections under the conditions of South Dobrudzha provides data on the economic traits of the accessions, their values as parental forms and the possibility of efficiently involving them as parental forms in the breeding process (Stoeva et al., 2009). All of the above justified the necessity to carry out this investigation with the aim to study some main properties related to the grain quality and the baking properties of bread wheat accessions of foreign origin under the conditions of the South Dobrudzha region.

Material and methods

The investigation involved a total of twenty-five common winter wheat varieties of foreign origin (USA, Moldova, Ukraine, Serbia and Romania) (Table 1). The accessions were grown during three successive years, 2010, 2011 and 2102, in the experimental field of Dobrudzha Agricultural Institute (DAI) – General Toshevo. Four standard cultivars were used for comparing their quality traits and baking properties – cultivars Aglika, Enola, Pryaspa and Yantar. They are national models for the respective quality groups at the Bulgarian Executive Agency of Variety Testing, Field Inspection and Seed Control.

The baking and quality properties of the varieties were analyzed at the Bread Making Quality Laboratory of DAI. The expression of the of the following indices was followed: protein content (%) determined according to the conventional method of Kjeldahl, using Keltec Auto 1030 Analyzer, test weight (TW, g) according to BS - ISO

* e-mail: plam_s@yahoo.com

Table 1. Common winter wheat cultivars used in this investigation

Serbia (Novi Sad)	Moldova (Baelti)	USA	Romania (Fundulea)
Kantata	Podoima	Yuamar	Glosa
Sonata	Livada	Harry	Boema
Tina	Dobropolka	Praire Red	Faur
Dragana		Wahoo	
Vila	Odessa (Ukraine)	Cougar	BG Chek - Quality Groups
Lilyana	Vesta	Wesley	Aglika (Group A)
	Kiria	New Sky	Enola (Group B)
	Kolumna	Jerry	Yantar (Group C)
	Zmina	Prowers	Pryaspa (Group D)

9771/2; sedimentation value (SDS, ml) according to Pumpyanski, 1971; wet gluten content (WGC, %) in 70% flour according to BDS - 13375-88; rheological properties of dough: softening degree (Sd, f.u.); valorimetric value (Val); bread volume (Lvol, ml) by pharinograph according to ICC115/1. The quality index (Qi) was calculated according to Hlaverson and Zeleny (1988). The data were processed and analyzed by applying variation, two-factor dispersion, correlation and cluster analyses using software XLstat 7.0, SPSS V.19 and Unistat 6.0.

Meteorological characterization of the investigated period (Figure 1)

The autumn of 2009 was characterized by a mean temperature higher than the norm. The high amount of rainfalls in October provided good moisture reserves. Although sowing was late, wheat emergence was uniform and the crops overwintered in good condition. The abundant rainfalls in May and June of 2010 made harvesting difficult and had negative effect on grain quality.

The hydrothermal conditions of 2011 were not particularly

favorable for wheat. The lack of snow cover and the extreme low temperatures in February lead to freezing of some of the plants. Spring was cool and long. Grain filling occurred under sufficient moisture content of soil and the temperatures were close to the long-term norm. Harvesting of the crop went under dry weather but nevertheless the investigated materials could not express their genetic potential with regard to the quality indices.

The abundant precipitation during the autumn of 2011 delayed the preparation of the area for sowing so planting was done in November, after the optimum agro technical dates. The low temperatures in November prolonged the emergence and development of the crops, and the plants entered the beginning of winter not sufficiently hardened. The long and cooler spring of 2012 additionally delayed the development of wheat and the rainfalls in May in combination with temperature contrasts had unfavorable effect on productivity. The temperatures higher than the climatic norm and the long drought in June and July influenced positively the formation of grain quality.

Figure 1. Meteorological characterization of the investigated 3-year period

Results and discussion

The obtained results, averaged for three years of the investigation are presented in Table 2. The exceeding of the standard cultivar Aglika, the national model for group A of strong wheats, at the highest level of significance should be noted. Ten of the cultivars are worth mentioning for their test weight, a maximum value of 80.3 g being measured for cultivar New Sky. All twenty-four accessions formed lower percent of protein according to Aglika, only the Romanian cultivar Faur showing a level close to the standard. Four of the investigated genotypes had higher sedimentation than it but the exceeding was not significant at $p=0.01$ %.

Nevertheless, these results had positive effect on the calculation of the quality index and the wheat cultivars Dobropolka, Zmina и Wahoo had significant differences with the standard cultivar. With regard to wet gluten yield, twelve of the varieties were clearly different from the standard, while the softening degree was

lowest in Livada and Zmina. Valorimetric value 50 was obtained only for cultivar Faur, the value of Aglika being 44. Bread volume was a trait which also did not show significant exceeding of the standard cultivar.

The variation coefficients of the investigated indices were calculated. Lowest variation was found for test weight, protein content and bread volume. High variability was registered for the indices valorimetric value, sedimentation and softening degree.

Depending on the values of the individual indices during the investigated period, years 2010 and 2012 can be considered favorable, while in 2011 a lower level of grain quality was determined on the whole. The effect of the year on the traits was significant, without exceptions (Table 3).

The effect of the cultivar was also significant, with the exception of bread volume. The results were to a large extent analogous to those reported by other authors proving the determining role of the year for the expression of the quality potential of wheat (Tsenov et

Table 2. Values of the investigated indices averaged for three years of investigation and their variation

Varieties	TW, g	Protein, %	SDS, ml	Qi	WGC, %	Sd, f.u.	Val	Lvol, ml
Vesta	75.7	9.67	31.0	3.13	14.1	147.3	28.7	576.7
Kiria	76.6	10.07	41.3	4.10	18.5	126.3	35.3	643.3
Kolumna	79.7***	10.90	48.6	4.43	23.4***	118.0	36.0	703.3
Zmina	77.3	10.37	52.3	5.0***	19.8	91.3***	39.3	621.7
Kantata	77.9	9.63	37.3	3.87	19.8	150.0	26.3	591.7
Sonata	75.5	9.50	38.3	4.04	19.3	155.7	25.0	578.3
Tina	76.2	9.73	41.0	4.20	18.7	141.7	28.0	686.7
Dragana	77.9	10.13	35.7	3.52	21.0	154.7	26.0	623.3
Vila	78.4	10.43	46.6	4.45	23.6***	123.0	22.5	675.0
Lilyana	78.2	9.93	41.3	4.16	22.6***	124.0	32.7	680.0
Yuamar	78.5***	10.80	48.8	4.43	23.1***	113.3	34.3	658.3
Harry	75.9	10.10	50.3	4.97	20.1	127.0	33.7	631.7
Praire Red	78.7***	11.13	48.3	4.34	20.6	99.3	46.3	671.7
Wahoo	77.6	9.77	50.0	5.10***	20.1	137.7	33.0	615.0
Cougar	79.1***	10.90	45.3	4.14	21.9	120.7	43.0	648.3
Wesley	78.2	10.57	48.6	4.58	23.0***	113.0	36.7	661.7
New Sky	80.3***	10.97	49.0	4.45	22.9***	116.0	39.0	648.3
Jerry	78.3	10.57	52.0	4.88	22.7***	108.0	40.7	633.3
Prowers	80.0***	11.43	52.6	4.58	24.0***	111.0	39.3	666.7
Podoima	79.0***	10.30	41.7	4.02	19.0	124.7	41.0	646.7
Livada	79.5***	10.53	51.3	4.87	22.4***	97.3***	42.0	656.7
Dobropolka	77.8	10.10	52.0	5.13***	19.5	107.7	37.0	625.0
Glosa	78.9***	11.07	47.0	4.24	22.2***	105.0	37.3	663.3
Boema	76.5	11.13	47.3	4.25	23.0***	99.6	43.3	640.0
Faur	79.2***	11.23	47.0	4.18	22.2***	107.3	50.0	656.7
Aglika	77.5	11.23	50.3	4.46	19.9	112.3	44.3	696.7
Enola	78.4	11.97	39.7	3.31	17.7	146.0	50.7	658.3
Yantar	76.0	10.93	30.7	2.76	18.0	152.7	25.0	560.0
Pryaspa	75.9	10.67	32.0	2.96	19.2	158.7	23.3	588.3
SE	1.39	0.11	1.24	0.11	0.42	3.68	1.46	6.76
VC %	1.8	5.8	15.0	14.6	10.9	16.0	21.9	5.7

*** - $p \leq 0.01$

Table 3. Variance analysis of factors „variety” and „year” in the investigated indices

Source of variation		df	Mean Square	F-ratio	P-value
TW, g	Genotype	28	5.8	5.8	0.0000
	Year	2	468.4	471.8	0.0000
	Residual	56	1.0		
Protein, %	Genotype	28	1.1	2.2	0.0051
	Year	2	21.9	43.2	0.0000
	Residual	56	0.5		
SDS, ml	Genotype	28	134.2	3.5	0.0000
	Year	2	1273.8	32.8	0.0000
	Residual	56	38.9		
Qi	Genotype	28	1.1	5.0	0.0000
	Year	2	2.4	10.7	0.0001
	Residual	56	0.2		
WGC, %	Genotype	28	15.5	3.9	0.0000
	Year	2	914.7	229.9	0.0000
	Residual	56	4.0		
Sd, f.u.	Genotype	28	1176.0	2.4	0.0000
	Year	2	32654.1	65.9	0.0000
	Residual	56	495.5		
Val	Genotype	28	173.5	1.7	0.0424
	Year	2	17391.1	172.4	0.0000
	Residual	56	100.9		
Lvol, ml	Genotype	28	3974.7	1.5	0.0924 n.s.
	Year	2	12752.0	4.9	0.0113
	Residual	56	2621.1		

al., 2004; Stoeva et al., 2006). Other sources underline the effect of the cultivar on the total variation (Trethowan et al., 2002; Penchev and Stoeva 2004). These somewhat contradictory results confirm the specific role of the external conditions and environment for the formation of the quality parameters in wheat in combination with the various genetic potential of the cultivars.

The use of correlation analysis allowed evaluating the correlations between the investigated traits (Table 4). Most of them correlated significantly with test weight. This confirmed the importance of the index for grain quality which can serve as initial basic evaluation. Similar tendency was observed for sedimentation value as well. The high significant positive correlation with wet gluten, valorimetric value and bread volume proved the influence of

sedimentation on the rheological properties of dough and the baking properties. There were significant negative correlations only of the softening degree with the other indices but they are normal due to the nature of this trait. Very strong was the correlation between the protein content and the valorimetric value. The good combination between high test weight, sedimentation and gluten was decisive for high bread volume.

The groups of cultivars formed through cluster analysis based on Euclidean distances according to the investigated indices are presented in Figure 2. The investigated accessions were divided into groups based on their similarities into four main clusters. The effect of the environmental conditions did not allow good differentiation of the four quality groups. The standards Aglika and Enola were at

Table 4. Correlation analysis of the studied parameters

	TW, g	Protein, %	SDS, ml	Qi	WGC, %	Sd, f.u.	Val
TW, g	1						
Protein, %	0.50**	1					
SDS, ml	0.54**	0.28 n.s	1				
Qi	0.36*	-0.09 n.s	0.93***	1			
WGC, %	0.66***	0.36*	0.64***	0.52**	1		
Sd, f.u.	-0.53**	-0.40*	-0.85***	-0.71***	-0.57**	1	
Val	0.53**	0.67***	0.57**	0.34 n.s.	0.22 n.s.	-0.67***	1
Lvol, ml	0.60***	0.42*	0.59**	0.45**	0.56**	-0.56**	0.49**

*** - $p \leq 0.01$; ** - $p \leq 0.05$; * - $p \leq 0.01$ and ; n.s. – nonsignificant

1. Vesta; 2. Kiria; 3. Kolumna; 4. Zmina; 5. Kantata; 6. Sonata; 7. Tina; 8. Dragana; 9. Vila;
10. Lilyana 11. Yuamar; 12. Harry; 13. Praire Red; 14. Wahoo; 15. Cougar; 16. Wesley; 17. New Sky;
18. Jerry; 19. Prowers; 20. Podoima; 21. Livada; 22. Dobropolka; 23. Glosa; 24. Boema; 25. Faur;
26. Aglika; 27. Enola; 28. Yantar; 29. Pryaspa

Figure 2. Cluster analysis based on Euclidean distances according to the investigated indices

short distance and therefore groups A and B were difficult to distinguish. Similar was the situation with the other two model cultivars for groups C and D – Yantar and Pryaspa. In this case again differentiation by quality could not be made. Due to the specificities discussed above, it could be considered that the wheat cultivars which are close to Aglika and Enola and fall into their or the neighboring cluster are quality wheats and can be referred to the group of increased strength (B), while the materials with greatest similarity to cultivars Pryaspa and Yantar are with low quality and can be referred to group D of the fillers. Thus grouped, cultivars Faur and Dobropolka from Romania, Zmina from Ukraine and the cultivars with origin from USA can be pointed out as the wheat varieties with highest grain quality.

The grain quality is genetically determined and is always the result from the interaction of the cultivar with the environment. For many seasons the quality potential of a cultivar can not be entirely expressed, which makes difficult the study on the correlations existing between the individual indices. Nevertheless, the systematic investigation on the expression of quality through the main characteristics which determine it provides sufficient objective information for the values of each cultivar. In this relation the study on the winter wheat accessions of foreign origin is a powerful source of enriching the gene pool of quality wheats for the national breeding.

Conclusion

The quality indices, the rheological properties and the baking properties are mainly influenced by the growing conditions but

simultaneously the genetic potential of the genotype is also expressed. The test weight and the sedimentation value were in positive correlation with most of the investigated traits; this makes them suitable for evaluation of grain and flour quality. The investigated accessions can be referred to quality group B of wheats with increased strength, some cultivars from Romania, Ukraine and USA exceeding the standards by the indices sedimentation, wet gluten content and valorimetric value. The Romanian cultivar Faur and the Ukrainian cultivar Zmina were the accessions with highest quality which have to be involved in breeding for quality.

Acknowledgments

Thanks are due to the management of DAI, General Toshevo for the opportunity provided to perform the trials discussed in this paper.

References

- Chamurliyski P, Tsenov N and Stoeva I**, 2011. Productivity and quality of modern Bulgarian bread wheat varieties (*Triticum aestivum* L.). Field Crop Studies, 7, 233-241.
- Hlaverson J and Zeleny L**, 1988. Criteria of wheat quality. In: Pomeranz, Y. (ed.), Wheat: Chemistry and Technology, American Association of Cereal Chemist. St. Paul, MN, USA, 1, 15-43.
- Panayotov I, Todorov I, Stoeva I and Ivanova I**, 2004. High quality wheat cultivars created in Bulgaria during the period 1994 - 2004 -

achievements and perspectives. *Field Crop Studies*, 1, 13-19.

Panayotov I, Kostov K and Todorov I, 2005. Impact of Bezostaya 1 in wheat breeding in Bulgaria. In: "Bezostaya 1 - 50 years of triumph", International Conference, Krasnodar – KNIISKH, p. 130-137 (Ru).

Panayotov I and Kostov K, 2007. Combining the Bulgarian and Ukrainian breeding for improving quality and productivity in wheat. International Conference "Plant gene pool – the basis of modern Agriculture", 13-14 June Plovdiv, 371-374.

Penchev E and Stoeva I, 2004. Evaluation of ecological plasticity of a group of common winter wheat varieties. *Field Crop Studies*, 1, 30-33.

Poumpyanskiy AY, 1971. Technological properties of soft wheat, L: 22 (Ru).

Stoeva I, Tsenov N and Penchev E, 2006. Environmental impact on the quality of bread wheat varieties. *Field Crop Studies*, 3, 7-17.

Stoeva I, Chamurliyski P and Tsenov N, 2009. Investigation on Bulgarian and foreign common winter wheat varieties and lines with a view of their using in breeding for productivity and quality, *FCS*, 5, 253-260.

Stoeva I, 2012. Technological evaluation of new common winter wheat lines developed at Dobrudzha Agricultural Institute – General

Toshevo. *Agricultural Science and Technology*, 4, 10-14.

Stoeva I and Penchev E, 2013. Effect of the environment on the quality of flour from common winter wheat cultivars. *Agricultural Science and Technology*, 5, 230-233.

Trethowan R, van Ginkel M and Rajaram S, 2002. Progress in breeding wheat for yield and adaptation in global drought affected environments. *Crop Science*, 42, 1441-1446.

Tsenov N, Kostov K, Gubatov T and Peeva V, 2004. Study on the genotype x environment interaction in winter wheat varieties. I. Grain quality. *Field Crop Studies*, 1, 20-29.

Tsenov N, Kostov K, Todorov I, Panayotov I, Stoeva I, Atanasova D, Mankovski I and Chamurliyski P, 2009. Problems, achievements and prospects in breeding for grain productivity of winter wheat. *Field Crop Studies*, 5, 261-273.

Tsenov N, Atanasova D, Chamurliyski P and Stoeva I, 2013. Influence of extreme environmental changes on grain quality of winter common wheat (*Triticum aestivum* L.). *Bulgarian Journal of Agricultural Sciences*, 19, 685-690.

Tsenov N, Atanasova D, Stoeva I and Tsenova E, 2014. Study of opportunities for effective use of varieties from Ukraine for creating early winter wheat lines II. Grain quality. *Agricultural Science and Technology*, 6, 156-161.

Review

- Triticale – past and future** 271
St. Dobрева

Genetics and Breeding

- Productivity performance of bread winter wheat genotypes with local and foreign origin** 276
G. Raykov, P. Chamurliyski, S. Doneva, E. Penchev, N. Tsenov
- Production properties of flax (*Linum usitatissimum* L.) cultivated in Strumica region, Republic of Macedonia** 280
P. Vuckov, M. Ilievski, D. Spasova, L. Mihajlov, N. Markova-Ruzdić
- Quality of grain and flour of foreign bread wheat cultivars (*Triticum aestivum* L.) under the conditions of south Dobrudzha region** 283
P. Chamurliyski, N. Tsenov, I. Stoeva, S. Doneva, E. Penchev
- Breeding programme for developing new sweet cherry cultivars in the Fruit Growing Institute, Plovdiv, Bulgaria** 289
S. Malchev, A. Zhivondov
- Heterosis manifestations by survival and larval duration of F1 *Bombyx mori* L. hybrids** 292
R. Guncheva, M. Panayotov, P. Tsenov, Y. Dimitrova

Nutrition and Physiology

- Reproductive performance of breeding rabbits fed by graded levels of cassava (*Manihot esculenta*) leaf meal** 297
T. Ahemen, I.I. Bitto, O.I.A. Oluremi
- Clinical toxicological investigations on acute carbofuran intoxication in quails (*Coturnix coturnix*)** 302
R. Binev, I. Valchev, R. Mihaylov, Y. Nikolov

Production Systems

- Efficacy and timing of some new products against pear psylla (*Cacopsylla pyri* L.) (Hemiptera: Psyllidae): II. Spirodiclofen** 306
V. Arnaudov
- Biochemical and chemical investigations of pikeperch fingerlings (*Sander Lucioperca* L.) after wintering** 310
A. Ivanova, R. Atanasova

Effect of fluorescence on the technological characteristics of cocoons at different cooking temperatures	313
---	------------

M. Panayotov

Comparative analysis of plane geometric parameters of various types of cow milking parlors	318
---	------------

D. Georgiev

Agriculture and Environment

<i>Panthaleus major</i> /Duges/ of cereals in Bulgaria	323
---	------------

V. Maneva, D. Atanasova

Selectivity and stability of herbicides and their tank mixtures for the seed yield of sunflower (<i>Helianthus Annuus</i> L.)	326
--	------------

G. Delchev, T. Barakova

Effect of green manure cover crops on tomato greenhouse production	332
---	------------

I. Tringovska, V. Yankova, D. Markova

Reclamation of lands disturbed by mining activities in Bulgaria	339
--	------------

I. Kirilov, M. Banov

Product Quality and Safety

Fish production and meat quality traits in rainbow trout (<i>Oncorhynchus mykiss</i>) farmed in different production systems	346
---	------------

St. Stoyanova, Y. Staykov, G. Zelqzkov, I. Sirakov, G. Nikolov

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors
The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings: Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows. IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 8, Number 4
December 2016

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz