ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 8, Number 4
December 2016

AGRICULTURAL SCIENCE AND TECHNOLOGY

2016

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Bojin Bojinov (Bulgaria)
Stoicho Metodiev (Bulgaria)
Svetlana Georgieva (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Zervas Georgios (Greece)
Ivan Varlyakov (Bulgaria)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Ramesh Kanwar (USA)
Martin Banov (Bulgaria)
Peter Cornish (Australia)

Product Quality and Safety

Marin Kabakchiev (Bulgaria)
Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions

on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate an author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to editorial review and the editors reserve the right to improve style and return the paper

for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc. and AGRIS (FAO).

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone.: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone.: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 8, Number 4
December 2016

AGRICULTURAL SCIENCE AND TECHNOLOGY

2016

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Nutrition and Physiology

Reproductive performance of breeding rabbits fed by graded levels of cassava (*Manihot esculenta*) leaf meal

T. Ahemen^{1*}, I.I. Bitto¹, O.I.A. Oluremi²

¹Department of Animal Breeding and Physiology, University of Agriculture, Makurdi, Nigeria

²Department of Animal Nutrition, University of Agriculture, Makurdi, Nigeria

(Manuscript received 12 March 2016; accepted for publication 21 September 2016)

Abstract. Twenty breeding does and 4 rabbit bucks aged 28 weeks, weighing between 1622 to 1746g, were used to investigate the reproductive performance of breeding does fed by graded levels of cassava leaf meal (CLM). The rabbits were randomly assigned to four dietary treatments viz: T_1 (control), T_2 , T_3 and T_4 with CLM inclusion levels of 0%, 10%, 20%, and 30%, respectively, after balancing for live weight. Each rabbit was individually caged and served as a replicate. The feeding trial lasted for 17 weeks. The results showed that does on diet T_2 (100%) achieved the highest conception rate, followed by T_3 (75%) and T_4 (75%) and the least in T_1 (66.67%). Average daily weight change and change in weight during pregnancy were the highest ($p < 0.05$) in T_4 . Litter size at birth was significantly ($p < 0.05$) lower in T_4 (2.67), while litter size at weaning was significantly ($p < 0.05$) higher in T_3 (3.67). Average birth and weaning weights were significantly ($p < 0.05$) higher in treatment T_4 . Percentage of still birth was higher in T_2 (60%) while pre-weaning mortality was the highest in T_1 (5). It was concluded that the inclusion of up to 30% of CLM in rabbit diet may support reproductive performance in female rabbits. It is recommended that further studies on the feeding potential of cassava leaf meal at higher levels of inclusion on reproductive performance be investigated.

Keywords: Cassava leaf meal, rabbit does and bucks, reproductive performance

Introduction

Nigeria like many other developing countries in the world has a protein deficiency gap, especially that of high quality animal protein. Ajayi et al. (2007) reported that rabbit production has enormous potential in alleviating the problems of animal protein supply in a developing economy like Nigeria. The most advantageous attribute of rabbits is their high reproductive potential (Lebas et al., 1997). This is as a result of their short gestation length, early sexual maturity, high prolificacy and ability to re-breed shortly after parturition, all leading to a short generation interval (Effiong and Wogar, 2007). In spite of the above advantages, studies on the productive system showed that among other factors, feeding is one of the major limiting factors in achieving maximum performance in rabbits (Iheukwumere et al., 2005). Asuquo (1993) observed that poor nutrition will delay sexual maturity in rabbits resulting in low financial returns for the farmer. Lebas (1993) and Effiong and Wogar (2007) also observed that increased feed and nutrient levels have been advocated for breeding rabbits as a means of increasing litter size, adequate maintenance of pregnancy and subsequent milk let down by the does.

Reports from literature indicate that various levels of cassava leaf meal can be incorporated in the diets of rabbits without adverse effect on growth performance. Fomunyan and Meffega (1984) fed weaner rabbits with diets constituting up to 75% cassava leaf meal without any adverse effect on growth performance. Adegbola and

Okonkwo (2002) fed rabbits with varying levels of cassava leaf meal up to 60% and obtained good growth performance. But little information is available on the effect of incorporation of cassava leaf meal in the diets on the reproductive performance of rabbits. This study was designed to evaluate the effect of cassava leaf meal on reproductive performance of breeding does.

Materials and methods

Location

The study was conducted at the Rabbitry unit of the Livestock Research Farm, University of Agriculture, Makurdi, Nigeria. Makurdi is located at Latitude 7° 14' North and Longitude 8° 21' East and lies within the Southern guinea savannah region of Nigeria. The area is warm with temperature range of 24.20 to 36.33°C and high temperature is experienced between late February and April. The rainfall is between 508 and 1016mm (TAC, 2009).

Source of Cassava leaves and preparation of experimental diets

The test ingredient, cassava leaves (bitter variety) were obtained from the local farms in and around Makurdi after cassava roots had been harvested. The leaves were sun-dried single layered on a concrete platform for 3 – 4 days to attain less than 10%

* e-mail: drtahemen@gmail.com

Table 1: Gross composition of experimental diets

Ingredients	Experimental diets			
	T ₁	T ₂	T ₃	T ₄
Maize	43.78	38.53	33.32	29.38
Full fat soybean meal	28.17	23.37	18.63	14.50
Cassava leaf meal	0.00	10.00	20.00	30.0
Brewers dried grain	10.00	10.00	10.00	10.0
Rice offal	15.00	15.00	15.00	15.0
Mineral/Vitamin premix*	0.25	0.25	0.25	0.25
Bone meal	2.50	2.50	2.50	2.50
Salt	0.30	0.30	0.30	0.30
Total	100	100	100	100
Calculated values:				
Crude protein (%)	18.20	18.23	18.09	18.35
Crude fibre (%)	8.71	9.93	11.15	12.39
Hydrogencyanide (mg/kg)**	0.0	5.39	10.77	16.16
DE (kcal/kg)	3067.88	2980.03	2891.03	2803.20

*0.25kg of Mineral/Vitamin Premix contained the following: Vitamin A 1,800IU, Vitamin D 250IU, Vitamin E 8,000IU, Vitamin K 750mg, B1 750mg, B2 1000 mg, B6 800mg, B12 25mg, Folic 300mg, Niacin 5000mg, Pantothenate 3000mg, Biotin 25mg, Cholin 160g, Thyroxin 300mg, Copper 0.4g, Iron 4g, Manganese 5.5g, Iodine 0.2g, Zinc 5g, Cobalt 0.15g, Selenium 0.15g. **Calculations were based on determined HCN content of CLM (53.87mg/kg). T₁=0% Cassava leaf meal, T₂ =10% Cassava leaf meal T₃=20% Cassava leaf meal T₄=30` % Cassava leaf meal

moisture. The leaves were thereafter milled using hammer mill to obtain the cassava leaf meal (CLM) which was then mixed with other ingredients to produce the experimental diets. The sun dried leaves were included at 0, 10, 20 and 30% levels in experimental diets T₁, T₂, T₃ and T₄, respectively (Table 1). The diets were formulated to meet the 18% crude protein requirement for pregnant/lactating rabbits. Other ingredients used were maize, full fat soya beans, brewers dried grain, rice offal, bone meal and mineral/vitamin premix.

Experimental animals and management

Twenty healthy rabbit does and 4 rabbit bucks of mixed breeds (California x New Zealand white x American Chinchilla) were used for the study. Their individual weights varied between 1622-1746g. The rabbits were allowed a preliminary period of two-weeks before the feeding trial commenced. During this period they were fed a common diet. The rabbits were treated with ivermectin subcutaneously at a dosage of 0.1ml per rabbit against ecto and endo-parasites. After the adjustment period, the experimental animals were weighed and assigned randomly to 4 groups of 5 rabbits, and each rabbit served as a replicate. The rabbits were housed individually in hutches and managed intensively. The experimental diets and clean drinking water were supplied to the rabbits *ad libitum* throughout the feeding period which lasted for 17 weeks. Close observation was made during experimental period for any disease problem.

At the 7th week of the experimental period, does were taken to the intact bucks for mating. Mating was repeated if the first mating was not successful. After successful mating, they were returned to their hutches. Initial weights of the does were taken after successful mating and were weighed subsequently every week throughout the trial. Does were palpated for the presence of developing foetus by gentle compression of the walls of the abdomen towards the middle line with the right hand under the abdomen and in front of the pelvis after properly restraining the rabbits with the left hand. The

abdominal palpation was done at 14 days post-coitus. Non-pregnant does were re-mated also at 14th day post-coitus. On the 26th day of mating open top wooden nest boxes were introduced into the hutches of each of the pregnant does for kindling. After kindling the litter size was taken by counting the number of kittens per doe. To measure kits birth weight, individual kit was carefully transferred with gloved hand to adjusted (zero) sensitive weighing scale and weight read off-from the scale. Litter size and weight of kits were monitored at birth and at weekly intervals up to weaning at four weeks after birth.

Data collection

Initial weight, weight at mating, change in weight during pregnancy and feed intake during pregnancy were recorded. Number of does mated, number of pregnant does at day 14, conception rate (%), individual birth weight (g), weaning weight, average (g) litter size at birth, average litter size at weaning, average litter weight at birth (g), average litter weight at weaning (g), pre-weaning litter mortality (%) and percentage of still birth were recorded.

Statistical Analysis

Data obtained were subjected to analyses of variance and correlation using GenStat (Release 4.24) statistical package (GenStat, 2005). Significant differences between treatment means were separated using Duncan's Multiple Range Test of the same statistical package.

Results

The result of weight changes of rabbit does during pregnancy is presented in Table 2. Conception rate ranged from 66.67% in diet T₁

Table 2. Performance of pregnant female rabbits fed varying dietary levels of cassava leaf meal (CLM)

Parameters	Experimental diets			
	T ₁	T ₂	T ₃	T ₄
Live weight at mating (g)	1933.0±32.80	1870.0±27.50	1900.0±26.50	1767.0±21.90
Conception rate (%)	66.67	100.00	75.00	75.00
Live weight at 28th day of pregnancy (g)	1966.70±30.0	1880.0±27.20	1983.30±10.40	1916.70±11.70
Change in weight in pregnancy (g)	33.33±4.70 ^c	10.00± 5.11 ^d	83.00±2.11 ^b	150.00±8.30 ^a
Av. daily weight change (g)	1.19±0.06 ^c	0.43±0.03 ^d	2.96±0.10 ^b	5.36±1.0 ^a
Av. daily feed intake (g)	75.40±5.10	67.30±4.40	79.50±7.50	74.20±6.90
Gestation period (days)	32.0	30.67	31.0	31.0

^{a,b,c,d}Means in the same row with different superscripts are significantly different (P<0.05), Av.=Average
T₁=0% Cassava leaf meal, T₂=10% Cassava leaf meal, T₃=20% Cassava leaf meal, T₄=30% Cassava leaf meal

Table 3. Reproductive performance of does fed varying dietary levels of cassava leaf meal (CLM)

Parameters	Experimental diets			
	T ₁	T ₂	T ₃	T ₄
Av. litter size at birth	4.00±1.00 ^a	3.5±0.50 ^a	4.67±0.67 ^a	2.67±0.3 ^b
Av. litter size at weaning	1.50±0.50 ^b	2.50±0.50 ^b	3.67±0.33 ^a	2.33±0.33 ^b
Av. litter weight at birth (g)	191±20.00	158.70±10.70	188.00±44.20	149.70±19.13
Av. litter weight at weaning (g)	560.0±21.0 ^b	885.0±26.50 ^{ab}	1103.0±16.0 ^a	926.0±15.20 ^b
Percentage of still birth (%)	33	60	0	0
Pre-weaning mortality	5	2	2	1

^{a,b}Means in the same row with different superscripts are significantly different (P<0.05), Av.=Average
T₁=0% Cassava leaf meal, T₂=10% Cassava leaf meal, T₃=20% Cassava leaf meal, T₄=30% Cassava leaf meal

to 100% in diet T₂. Weight at mating was not significantly different among treatments. Weight at 28 days of pregnancy was also not significantly affected by dietary treatments. Change in weight during pregnancy was significantly (p<0.05) higher in treatment T₄, followed by T₃ and the least in T₂. Average daily weight change followed the same trend. Average gestation length was the longest in T₁ with 32 days.

The result of post kindling traits of does fed diets with graded levels of cassava leaf meal is presented in Table 3. Litter size at birth was significantly (p<0.05) lower in treatment T₄. Average litter size at weaning and average litter weight at weaning were both significantly (p<0.05) higher in treatment T₃. Percentage of still birth was higher in T₂ with 60% and 33% in T₁. Treatments T₃ and T₄ had no still birth. Pre-weaning mortality was higher in T₁ followed by T₂ and T₃, and the

least in T₄.

The result showing progressive body weight increase of kits in the first 28 days is presented in Table 4. Mean body weight of kits in treatment T₄ was significantly higher at birth and weaning. It was observed that weight of kits at weaning on treatment T₃ was significantly (p<0.05) lower than all the other treatments.

The correlation between dietary cassava leaf meal and reproductive performance of female rabbits is presented in Table 5. A negative association existed between levels of inclusion of cassava leaf meal and litter size at birth (r=-0.43; p>0.05) but there was a positive correlation between levels of inclusion of cassava leaf meal and litter size at weaning (r=0.53; p<0.05). There was positive relationship between level of inclusion of CLM and BW (r=0.43; p<0.05), as well as weaning weight (r=0.11; p>0.05).

Table 4. Weights (g) of kits from does fed varying dietary levels of cassava leaf meal (CLM)

Parameters	Experimental diets			
	T ₁	T ₂	T ₃	T ₄
Birth weight	47.80±3.30 ^a	37.50±0.40 ^b	40.30±1.60 ^b	56.00±1.0 ^a
Weight at 1 st Week	110.00±7.60	117.90±15.10	112.27±8.0	125.3±6.40
Weight at 2 nd Week	190.0±11.70	192.0±15.60	162.30±9.20	207.90±7.10
Weight at 3 rd Week	283.30±64.0 ^a	250.00±14.60 ^{ab}	206.36±11.60 ^{bc}	218.60±8.60 ^b
Weight at 4 th Week	373.30±14.50 ^a	354.00±19.70 ^a	310.0±13.0 ^b	398±10.0 ^a

^{a,b}Means in the same row with different superscripts are significantly different (P<0.05)
T₁=0% Cassava leaf meal, T₂=10% Cassava leaf meal, T₃=20% Cassava leaf meal, T₄=30% Cassava leaf meal

Table 5. Correlation matrix of CLM in diets and reproductive performance of female rabbits

1	AALE	BW	LSB	LSW	LWB	LWW	WW
AALE	1.00						
BW	0.43	1.00					
LSB	-0.43	-0.65**	1.00				
LSW	0.53*	-0.46*	0.45*	1.00			
LWB	-0.59**	-0.33	0.89**	0.06	1.00		
LWW	0.75**	0.26	0.13	0.94**	-0.26	1.00	
WW	0.11	0.80**	-0.88**	-0.78**	-0.57*	0.54*	1.00

** Correlation is significant at 1% ($P < 0.01$), * Correlation is significant at 5% ($P < 0.05$). AALE= levels of inclusion of cassava leaves, BW=Birth weight, LSB=Litter size at birth, LSW=Litter size at weaning, LWB=Litter weight at birth, LWW=Litter weight at weaning, WW=Weaning weight.

Discussion

The conception rate ranging from 66.67 to 100% obtained in this study agrees with the 67 to 100% conception rate reported by Raharjo et al. (1986) in New Zealand White breed of rabbits. This suggests that cassava leaf meal diets provided enough nutrients for the maintenance and reproduction of the rabbits. Gestation duration ranged from 30 to 32 days. Gestation length of 30.50 ± 0.35 days to 31.50 ± 0.00 days was reported by Imasuen et al. (2001) on rabbits fed some tropical leafy forage. Gestation period of does in different treatment groups was not significantly different, though the control group had longer gestation of 32 days. Gestation length in animal species is fixed and only varies within a very narrow gap as observed in this study. Similar observation was made by Ajayi et al. (2005). Average weight changes during pregnancy were significantly ($P < 0.05$) higher in rabbits on diets T_4 and the least with rabbits on diet T_2 . Small body weight changes indicate that dietary energy and/or protein intake may not be adequate for intensive production by the does without the mobilization of body reserves. However, this may not be the reason for low average weight change observed in T_2 since the rabbits were supplied with adequate dietary protein and energy.

Litter size at birth is comparable to 3.00 – 5.60 (Imasuen et al., 2001) and within the range of 2.5 ± 0.3 – 6.51 ± 0 (Ijaiya et al., 2002). However, the values are lower than litter size of 7-8 kits reported by Lukefahr and Cheeke (1990). The lower litter size at birth in this study could be as a result of the does being in the first parity. Litter size at birth increases with successive parity (Brambella, 1972). The observed negative correlation between CLM diets and litter size at birth and, positive correlation between CLM diets and litter size at weaning were because of the higher pre-weaning mortality in the control diet (T_1). Pre-weaning mortality was higher in T_1 and lower in T_4 with absolute values of 5 and 1, respectively. The observed low mortality recorded in T_4 with the lowest litter size is in agreement with the report of Schloant (1980) who stated that most cases of pre-weaning kit mortality occurred because of inadequate milk production by does. Positive relationship between litter size at birth and weaning imply that the higher the number of kits born/litter/doe, the higher the number weaned/litter/doe. The negative correlations between birth weight and litter size at birth, weaning weight and litter size at weaning is in consonance with the reports by Rao et al. (1977). According to Inglis (1980), litter size and birth weight vary depending on the breed and size of the dam. Average litter size ranges from 5-10 kits, while birth weight ranges from 30 – 80 g and both are higher in larger and medium breeds than smaller breeds.

Factors that increase litter size include the age at which the does are served and the optimum weight at service, which is an index of breed (Hafez and Hafez, 2000). The higher Average litter size at weaning and higher average litter weight at weaning observed in T_3 agree with the report of Abdelli-Larbi et al. (2014) who reported higher average litter weight at weaning from does with larger average litter size at weaning.

Birth weight and weight of the young rabbits at consecutive four weeks were comparable to 52.1 ± 3.3 g to 55.7 ± 1.9 g, 100.2 ± 2.4 g to 122.7 ± 3.0 g, 162.8 ± 4.8 g to 198.4 ± 5.7 g, 230.7 ± 7.0 g to 282.0 ± 8.0 g and 333.0 ± 9.0 g to 402.0 ± 11.0 g for birth weight, weight at 7 days, weight at 14 days, weight at 21 days and weight at 28 days, respectively, reported by Abdelli-Larbi et al. (2014). The significant ($p < 0.05$) lower birth weight in T_2 and T_3 is in conformity with the earlier report by Hafez and Hafez (2000) who reported decrease in birth weight with increase in litter size. This worker attributed the weight advantage to competition for nutrients and the lesser inter-uterine space in cases where does carry more fetuses. Eze et al. (2009) observed that in rabbits the blood supply to each implantation site is reduced when the number of fetus increases. Positive associations were found between CLM in the diets and birth weight as well as weaning weight. This suggests that kits of does on cassava leaf meal based diets were obtaining enough nutrients in their feed to support tissue gain. Average kit weight of rabbits in T_3 was significantly ($p < 0.05$) reduced at weaning. This might be because of average larger litter size of rabbits in T_3 . This is in consonance with the findings of Eshiet et al. (1980) who observed that kits from larger litters were on the average lighter than those from the smaller litters. They attributed this to the greater competition for the dam's milk especially during the first 3 weeks of age when it constitutes the main source of nourishment.

Conclusion

The results showed that does on diet T_2 (100%) achieved the highest conception rate, followed by T_3 (75%) and T_4 (75%) and the least in T_1 (66.67%). Average change in weight during pregnancy was the highest ($p < 0.05$) in T_4 . Average birth and weaning weights were significantly ($p < 0.05$) higher in treatment T_4 . Percentage of still birth was higher in T_2 (60%), while pre-weaning mortality was the highest in T_1 (5). The results indicate that inclusion of up to 30% cassava leaf meal (CLM) in the diet of breeding rabbits support reproductive performance in female rabbits. It is recommended that

further studies on the feeding potentials of cassava leaf meal at higher levels of inclusion be investigated.

References

- Abdelli-Larbi O, Mazouzi-Hadid F, Berchiche M, Bolet G, Garreau H and Lebas F**, 2014. Pre-weaning growth performance of kits of a local Algerian rabbit population: Influence of dam coat color, parity and kindling season. *World Rabbit Science*, 22, 231-240.
- Adegbola TA and Okonkwo JC**, 2002. Nutrient intake digestibility and growth of rabbits fed varying levels of cassava leaf meal. *Nigerian Journal of Animal Production*, 29, 21-68.
- Ajayi FO, Balogun OO, Ovuru SS and Mgbere OO**, 2005. Reproductive performance of rabbits fed maize milling waste based diets. *African Journal of Biotechnology*, 4, 439-443. <http://www.academicjournals.org/AJB>
- Ajayi AF, Farinu GO, Ojebiyi OO and Olayeni TB**, 2007. Performance evaluation of male rabbits fed diets containing graded levels of blood-wild sunflower leaf meal mixture. *World Journal of Agriculture Sciences* 3, 250-255.
- Asuquo BO**, 1993. Optimum mash level in a mixed mash-forage feeding system for growing rabbits. *Animal Production Research*, 13, 99-104.
- Brambella OE**, 1972. *Grzimels Animal life. Encyclopedia*, 2, 420.
- Effiong OO and Wogar GS**, 2007. Litter performance traits of rabbits under mixed feeding regime. *Proc. 32nd Annual Conference of the Nigerian Society of Animal Production*, Calabar, March 18-21, 2007, 155-158.
- Eshiet NO, Ademosun HA and Omole TA**, 1980. The effect of feeding cassava root meal on the reproductive performance of rabbits. *Journal of Nutrition*, 110, 697-702.
- Eze CA, Nwodo CP and Ogbu EO**, 2009. Effect of hemiovariectomy on reproductive performance of chinchilla rabbits. *Nigerian Journal of Animal Production*, 36, 167-174.
- Fomunyam AT and Meffeja F**, 1984. Cassava by-products in rabbit and sheep diets. *Institute of Animal Research, Mankon Bamenda, Cameroon*.
- GenStat**, 2005. *GenStatUser's Guide (Release 4.24)*. Lawes Agricultural Trust, Rothamsted Experimental station. UK.
- Hafez ESE and Hafez B**, 2000. Reproductive cycles. In: E.S.E.Hafez and B.Hafez (eds). *Reproduction in Farm Animals*. 7th Edition. Balliere Tindal, London, Great Britain.
- Iheukwumere FC, Okoli IC, Annyanwu GA and Esonu BO**, 2005. Growth performance, haematology and serum biochemical constituents of grower rabbits fed *Microdesmispuberula*, Hook-Euphorbia ceae. *Animal Production Research Advances*, 1, 24-31.
- Ijaiya AT, Fasanya OOA and Ayanwale BA**, 2002. Reproductive performance of breeding rabbit does fed maize and fermented cassava peel meal. *Proceedings 27th Ann. Conference. Nigerian Society for Animal Production*, 14. NSAP. March 17-21, Fed. University of Technologies, Akure, Nigeria.
- Imasuen JA, Ikhatua UJ, Otoikhian CSO and Arijeniwa A**, 2001. Evaluation of growth and reproductive performance of rabbits fed some tropical leafy forages. *Tropical Journal of Animal Science*, 4, 109-116.
- Ingli JK**, 1980. *Introduction to Laboratory Animal Science and Technology*. Pergamon Press, Oxford, 3-4.
- Lebas F**, 1993. Small Rabbit production, feeding and management system. *World Animal Review*, 46, 11-17.
- Lebas F, Coundert P, De Rochambeau H and Thebault RG**, 1997. *The Rabbithusbandary, health and production*. FAO Animal Production and Health series. Rome, Italy, 21, 203.
- Lukefahr SD and Cheeke PK**, 1990. Rabbit project development strategies in subsistence farming systems. *Research applications. World Animal Review*, 68, 26-35.
- Raharjo TC, Cheeke PR, Patton NM and Supriyati K**, 1986. Evaluation of tropical forages and by – product feeds for rabbit production. 1. Nutrients. digestibility and effect of heat treatment. *Journal of Applied Rabbits Research*, 9, 6-11.
- Rao DR, Sunki GR, Johnson WM and Chen CP**, 1977. Post-natal growth of the New Zealand White Rabbit. *Journal of Animal Science*, 44, 1021-1025.
- Schloant W**, 1980. Trends in the development of rabbit production and related technical problems. *Animal Research and Development (GDR)*, 11, 110-116.
- TAC**, 2009. *Makurdi Weather Elements Records, MakurdiMetereologicalStation*. Nigeria Air Force, Tactical Air Command, Makurdi, Nigeria.

Review

- Triticale – past and future** 271
St. Dobрева

Genetics and Breeding

- Productivity performance of bread winter wheat genotypes with local and foreign origin** 276
G. Raykov, P. Chamurliyski, S. Doneva, E. Penchev, N. Tsenov
- Production properties of flax (*Linum usitatissimum* L.) cultivated in Strumica region, Republic of Macedonia** 280
P. Vuckov, M. Ilievski, D. Spasova, L. Mihajlov, N. Markova-Ruzdić
- Quality of grain and flour of foreign bread wheat cultivars (*Triticum aestivum* L.) under the conditions of south Dobrudzha region** 283
P. Chamurliyski, N. Tsenov, I. Stoeva, S. Doneva, E. Penchev
- Breeding programme for developing new sweet cherry cultivars in the Fruit Growing Institute, Plovdiv, Bulgaria** 289
S. Malchev, A. Zhivondov
- Heterosis manifestations by survival and larval duration of F1 *Bombyx mori* L. hybrids** 292
R. Guncheva, M. Panayotov, P. Tsenov, Y. Dimitrova

Nutrition and Physiology

- Reproductive performance of breeding rabbits fed by graded levels of cassava (*Manihot esculenta*) leaf meal** 297
T. Ahemen, I.I. Bitto, O.I.A. Oluremi
- Clinical toxicological investigations on acute carbofuran intoxication in quails (*Coturnix coturnix*)** 302
R. Binev, I. Valchev, R. Mihaylov, Y. Nikolov

Production Systems

- Efficacy and timing of some new products against pear psylla (*Cacopsylla pyri* L.) (Hemiptera: Psyllidae): II. Spirodiclofen** 306
V. Arnaudov
- Biochemical and chemical investigations of pikeperch fingerlings (*Sander Lucioperca* L.) after wintering** 310
A. Ivanova, R. Atanasova

Effect of fluorescence on the technological characteristics of cocoons at different cooking temperatures	313
---	------------

M. Panayotov

Comparative analysis of plane geometric parameters of various types of cow milking parlors	318
---	------------

D. Georgiev

Agriculture and Environment

<i>Panthaleus major</i> /Duges/ of cereals in Bulgaria	323
---	------------

V. Maneva, D. Atanasova

Selectivity and stability of herbicides and their tank mixtures for the seed yield of sunflower (<i>Helianthus Annuus</i> L.)	326
--	------------

G. Delchev, T. Barakova

Effect of green manure cover crops on tomato greenhouse production	332
---	------------

I. Tringovska, V. Yankova, D. Markova

Reclamation of lands disturbed by mining activities in Bulgaria	339
--	------------

I. Kirilov, M. Banov

Product Quality and Safety

Fish production and meat quality traits in rainbow trout (<i>Oncorhynchus mykiss</i>) farmed in different production systems	346
---	------------

St. Stoyanova, Y. Staykov, G. Zelqzkov, I. Sirakov, G. Nikolov

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors

The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings:

Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows. IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 8, Number 4
December 2016

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz