ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 9, Number 1
March 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate on author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc. and AGRIS (FAO).

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No DNP 05-21/20.12.2016, financed from Fund 'Scientific Research' grant Bulgarian scientific Periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 9, Number 1
March 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Agro-ecological assessment of manure from different farm animals by content of biogenic elements

D. Dermendzhieva^{1*}, G. Kostadinova¹, G. Petkov¹, D. Dimov¹, T. Dinev², T. Penev¹, Tch. Miteva¹, J. Mitev¹

¹Department of Applied Ecology and Animal Hygiene, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria

²Department of Biochemistry, Microbiology and Physics, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria

(Manuscript received 12 November 2016; accepted for publication 23 January 2017)

Abstract. The aim of the present study was to investigate and to make agro-ecological assessment of fresh (excreted up to 24 h) and stored (for 6 months) manure from different farm animals (dairy cows, pigs, laying hens and turkeys) by biogenic elements content (N, P, K) for a period of one year. Samples of fresh and stored manure were collected and analyzed monthly for total N, total P and total K content. It was found that: a) biogenic elements in fresh and stored manure decreased in the following gradation by average values for the studies period: turkeys – total N (13.9 - 9.16 g/kg DM), total P (9.94 - 8.67 g/kg DM) and total K (5.02 - 3.74 g/kg DM); laying hens – total N (13.6-9.95 g/kg DM), total P (9.27- 7.94 g/kg DM) and total K (5.56 - 4.60 g/kg DM); pigs – total N (8.30 -6.77 g/kg DM), total P (6.08-4.65 g/kg DM) and total K (3.56 - 1.61 g/kg DM); dairy cows – total N (5.62 -4.76 g/kg DM), total P (1.72 - 1.48 g/kg DM) and total K (2.99 - 1.72 g/kg DM); b) the change in nutrients content of different types of manures by seasons both in fresh and in stored manure are divergent; c) coefficients of variation values determined surveyed manures by content of nutrients from minor to significant variable ($C_v = 9.30-37.5\%$); d) during the storage of the tested manures nutrient losses are the greatest for potassium (17.3-54.8%), followed by nitrogen (15.3-34.1%) and phosphorus (12.8-23.5%); e) the average losses of biogenic elements in manure by kind of animal species are in varying degrees: dairy cows (total N 15.3%, total P 14.0%, total K 42.5%); pigs (total N 18.4%, total P 23.5%, total K 54.8%); laying hens (total N 26.8%, total P 14.3%, total K 17.3%) and turkeys (total N 34.1%, total P 12.8%, total K 25.5%); f) the ratio between nutrients (N:P:K), both in fresh and in stored manure from different animal species vary in relatively wide limits both by seasons (12.6%N:18.1%P:56.3%K – 41.5%N:16.0%P:39.7%K) and by average of the studied period (15.3%N:14.0%P:42.5%K – 34.1%N:12.8%P:25.5%K).

Keywords: dairy cows, pigs, laying hens, turkeys, fresh and storage manure, biogenic elements, quality assessment

Introduction

Worldwide the needs of N, P and K, for plant food and forage production increase annually. According to FAO (2015) the demand for fertilizer nutrients ($N+P_2O_5+K_2O$) in the world up to 2018 can reach 200 500 000 t, with growth of 1.8% per year (N 119 418 000 t, P_2O_5 46 648 000 t and K_2O 34 456 000 t). Europe's share in global consumption of total fertilizer nutrients is about 12.5% of the total amount. The share of the region in nitrogen fertilizer consumption is 13.5%, phosphate 9.0% and potash 14.0%. A significant portion of these quantities has been provided by farm animal manure, which is an inevitable byproduct in the livestock production systems.

The main trend in animal manure use today, and also in the foreseeable future, is its application as a high quality organic fertilizer to improve soil fertility (Katzarov et al., 2003; Fonseca, 2011; MMPKR, 2014). The organic manure is an excellent plant nutrient source when used properly. Its composition is not constant and depends on many factors - species, age and health condition of the animals, quality and feed composition of the ration (especially protein and fiber content), level of animal metabolism and diet digestibility, housing and addition of bedding, method of cleaning and consistence of manure – solid or liquid, manner (in manure storages, on the ground, lagoons, etc.) and period of storage, environmental conditions, etc. (Petrov et al., 1983; Faassen and Dijk, 1987; Velthof et al., 2000; Katzarov and Pavlov, 2002;

Zaharinov et al., 2013). Therefore, when analyzing the quality of a particular type of manure the specific conditions that determine the formation of its physical properties and chemical composition should also be taken into consideration (Reijs et al., 2003; Wang and Gaston, 2014).

The benefits of livestock manure usage in crop production are improvements of the physical properties of soil and the provision of N, P, K and other biogenic elements (FAO, 2003). The application of livestock manure increases soil organic matter content, and thus leads to improved water infiltration and water holding capacity as well as an increased cation exchange capacity (Hoffmann, 2002; Goss et al., 2013). Zhang and Schroder (2014) added that manure in soil enhances retention of nutrients, reduces wind and water erosion, and promotes growth of beneficial organisms.

Organic manure contains 13 of the most important plant nutrients - N, P, K, Ca, Mg, S, Fe, Mn, Zn, Cu, B, Mo and Cl, making it a valuable fertilizer. Within the group of macronutrients, which are needed for plant growth in larger amounts, the primary nutrients are N, P and K. Nitrogen is the basic element for plant growth as it forms amino acids and proteins; phosphorus plays a key role to transfer energy because it is essential for photosynthesis and other chemical-physiological processes in the plant; potassium has many functions – it activates more than 60 enzymes (FAO, 2000; Chastain et al., 2002). Organic manure is more cost effective in comparison with the use of inorganic fertilizers (Zakikhani et al., 2016).

Improper treatment and storage of manure can make it lose a lot

* e-mail: d.dermendzhieva@abv.bg

of valuable ingredients for improving soil fertility. One of the keys to a successful manure management system is good nutrient management, which takes into account nutrients in the animal diet, nutrients excreted by the animal, lost in storage, transportation and land application with the final use of nutrients by the crops (Svoboda, 2003; Roberts, 2009). According to Cederberg et al. (2012) livestock absorb between 15-40% of N and P ingested in their diet, and consequently the majority of these nutrients in feed end in the manure. The losses of nutrients during storage of manure depend on many factors - type and consistence of manure, manner and duration of its storage, climatic conditions, etc. Hellmuth and Hochmuth (2015) pay attention to the fact that N is highly mobile, can quickly change the form, and can be lost rapidly from the manure during its storage and soil application. Clanton (1993) estimated that: nutrient losses from storage by volatilization, regardless of the storage system (stacks, bunkers, earthen or other pits) and kind of manure (dairy cows, swine, layers, turkeys) can reach 35% for N, while for P and K they are negligible; N losses between excretion and land application are 15-30% for solid and 15-80% for liquid manure; nutrient losses during land application vary depending on moisture content of the manure, application method, manure and soil pH, temperature and wind action. Another source (MMPKR, 2014) reported that dairy cows and poultry manure losses during accumulation and storage 25% of organic substance, while for pig manure they are 15%.

Significant problem that can arise from improper use of organic manure as a source for soil fertilization is environmental pollution (HSUS, 2008; Bywater, 2011; Cederberg et al., 2012; Mikkelsen et al. 2014; Scotti et al., 2015). Agricultural practices, based on large application of fertilizers, including farm manure, reduce the ability of agro-ecosystems to provide goods and services. This incorrect agricultural management can change species composition or reduce biodiversity in ecosystems thus affecting natural abilities of ecosystems (Loreau et al., 2001; Tillman et al., 2002). In this direction of great importance are the sanitary and hygienic properties of produced manure (absence of pathogens and odour) as they directly affect the air, surface and groundwater pollution, as well as agrochemical indices such as nutrients contents, absence of weed seeds and homogeneous structure. Environmentally friendly are some practices to reduce the quantity of N and P in animal excretion by their diets. The study of Clanton (1993) showed that properly balanced diet can reduce the amount of N and P in animal excretion by 50 and 60%, respectively. Report of Capper (2011) revealed that for the period 1977 – 2007, in the beef industry in the USA N excretion decreased by 12.3%, and P excretion by 10.3% due to implementation of some good farm management practices – improving the production systems for housing of animals, more efficient usage of energy and minerals in the diet, water reuse, etc.

A more accurate way to balance the crop fertility program is to test the manure before land application (Ap Dewi, 1994; Katzarov, 2005). Massie et al. (1993) noted that when N is used as a limiting nutrient, more P and K will be applied than crop needs and in the future their levels in soil profile and groundwater will increase.

Generally, despite numerous studies to characterize and use organic manure as fertilizer on farmland, topical questions remain about the content of biogenic elements in manure from different farm animal species, losses of nutrients during storage of manure and the ratio of the nutrients (N:P:K) in manure (fresh and storage). All this largely determines the agro-ecological quality of farm manure. This investigation was carried out to provide more data in these aspects and to search for answers of some of the above-mentioned issues.

Material and methods

Study area

The study was conducted over a period of one year - from January to December, during the first decade of 21st century in 3 livestock farms, situated in the south-central part of Bulgaria. The characteristics of the investigated farms are as follows:

- Dairy farm: Capacity – 63 cattle, including 32 cows, 8 heifers and 23 calves. Animal nutrition – with summer and winter ration; consisting of the diets – summer: green alfalfa and meadow grass (on the pasture), winter: lucerne hay, barley straw, corn silage; year-round in the composition of the ration was included beer mash; the concentrate mixture consisted of barley and wheat in the ratio 60:40%. Manure management – manure (mixture of solid manure with straw and urine) from the main production building and the building for calves was removed daily and transported to a surface earthen platform (manure storage), where it was stacked on the ground to the height of 1.20 - 1.40 m and left to mature for 6 months; then it was used to fertilize farmland.

- Pig farm: Capacity – 390 sows with average daily number of reared pigs of different categories 4552. Production buildings - 5: 1 building for 300 mating and pregnant sows and 22 boars, 1 building for 90 lactating sows with piglets prior to weaning and 450 weaned pigs; 1 building for 1200 growing pigs and 250 finishing pigs; 2 buildings for fattening pigs – for 940 and 1300 animals, respectively. Animal nutrition – it was done with concentrate mixtures according to the requirements of the relevant category pigs, the main ingredients of the rations were corn, barley, wheat, soybean and sunflower meal. Watering – it was done by automatic drinkers. Manure management – manure (mixture of solid manure, urine and technological water - slurry) in the premises was collected in channels under the pens of the animals and after filling up to 8-10 cm to the upper edge of the channel, it was pushed with additional water to the two earthen lagoons, which were filled consistently for a period of 6 months. During this time, the manure in the full lagoon matured and after that it was used to fertilize farmland.

- Poultry farm: The farm was mixed - for laying hens and turkeys, with a total capacity of 13.350 birds (11.250 laying hens and 2.100 turkeys). The object of the study was a production building for laying hens (9.50/60.0 m, 570 m²), with capacity for 2.000 birds (1.800 hens and 200 roosters) and similar production building for turkeys with capacity of 300 birds (240 female turkeys and 60 turkey cocks). Feeding of the birds during the controlled period was performed with dry forage mixtures containing nitrogen - 28.60 g/kg Dry Matter (DM), phosphorus - 8.34 g/kg DM and potassium - 7.34 g/kg DM. After completion of the production cycle and removal of laying hens and turkeys from the premises, the accumulated feces and deep litter-loose housing (mixture of feces and straw/litter) were matured indoor for 2 months and afterwards were pushed on the ground platform (manure heap), located close to the relevant production building (for laying hens and turkeys). The poultry deep litter was stored on platforms for 4 months, and then used to fertilize.

Sampling and sample preparation

Samples of fresh (excreted up to 24 h) and stored manure (for 6 months) from different animal species were collected monthly from 3-4 different places in the premises, and from 3-4 places of the manure heap/lagoons, at an amount of 1.5 kg for both types of manure from each farm. After mixing with a spatula, from the amount collected for analysis was taken 300 – 400 g of average sample in chemically clean glass containers. The total number of the collected

manure samples was 96 (48 from dairy cows and pigs and 48 from laying hens and turkeys). Each sample was analyzed in triplicate.

Manure nutrients and methods for their analyzes

Fresh and stored dairy cows, pigs, laying hens and turkey manure was analyzed for content of: total nitrogen - N (g/kg DM) – by the Kjeldahl method (Bulgarian State Standard /BSS/ ISO 11261); total phosphorus - P (g/kg DM) - spectrophotometrically by vanadate – molybdate method (BSS ISO 11263); total potassium - K (g/kg DM) - by hydrochloric acid solution according to the method of flame spectrometry (BSS ISO 11047).

Analyses of the tested indicators of manure were carried out at the Research Laboratory of the Faculty of Agriculture at Trakia University - Stara Zagora.

Agro-ecological assessment of the analyzed manures by nutrients

It was made by the method of comparative analysis as follows:

- on the base of the results for total N, total P and total K content in fresh and stored manure from different farm animal species;

- based on losses of nutrients (total N, total P and total K) during the storage of different types of manure;

- on the base of the values for the ratio between total N, total P and total K content (N:P:K) in fresh and stored manure from different animal species.

Statistical analysis

Statistical analysis was carried out by ANOVA, the software used was Statistica 6.1 (Statistica for Windows, StatSoft. Inc., Tulsa, OK, USA, 1984-2002).

Results and discussion

Total N, total P and total K content in fresh and stored manure from different farm animals. The data for these parameters are presented in Table 1.

Total nitrogen content: Fresh manure. The total N content varies widely from 5.32 g/kg DM in dairy cows manure to 18.4 g/kg DM in turkeys manure (the difference between the maximum and minimum

Table 1. Average (Cx), minimum (Cmin) and maximum (Cmax) content of total N, total P and total K in manure of different farm animals (g/kg DM)

Season	n	Species of animals	Manure					
			Fresh			Stored		
			N*	P*	K*	N*	P*	K*
Winter	n=3	dairy cows	5.32±0.61	1.55±0.18	2.16±0.11 ^c	4.45±0.53	1.37±0.29	1.50±0.13 ^c
		pigs	8.13±0.38 ^a	5.29±0.35	3.43±0.23 ^c	6.88±0.25 ^a	4.58±0.23	1.30±0.08 ^c
	n=3	laying hens	16.5±1.23	9.20±0.48 ^a	5.50±0.45	10.6±0.91	7.40±0.33 ^a	3.60±0.28
		turkeys	18.4 ±0.72 ^c	10.3±0.61	4.43±0.32	9.76 ±0.63 ^c	9.70±0.78	3.53±0.31
Spring	n=3	dairy cows	5.60±0.26 ^a	1.71±0.12 ^c	4.67±0.16 ^b	5.00±0.21 ^a	1.49±0.11 ^c	2.23±0.19 ^b
		pigs	8.43±0.33	5.52±0.35 ^a	3.73±0.31 ^b	7.37±0.46	4.52±0.23 ^a	1.63±0.20 ^b
	n=3	laying hens	10.9±0.41 ^a	8.81±0.44	5.23±0.31	9.53±0.28 ^a	7.50±0.37	4.83±0.22
		turkeys	17.1±1.02 ^b	9.90±0.84	5.90±0.77	10.0±0.63 ^b	8.32±0.81	3.56±0.25
Summer	n=3	dairy cows	5.73±0.14	1.86±0.11 ^c	2.53±0.09 ^c	4.57±0.52	1.47±0.13 ^c	1.73±0.08 ^c
		pigs	8.77±0.34 ^c	7.77±0.42 ^b	3.73±0.31 ^c	6.88±0.18 ^c	4.88±0.29 ^b	2.00±0.11 ^c
	n=3	laying hens	10.5±0.86	8.76±0.81	6.10±0.54	9.09±0.95	7.75±0.48	5.01±0.36
		turkeys	10.6±1.01	9.79±0.88	6.10±0.71	8.25±0.42	8.40±0.43	4.60±0.40
Autumn	n=3	dairy cows	5.83±0.52	1.77±0.17	2.60±0.12 ^c	5.03±0.37	1.59±0.24	1.43±0.08 ^c
		pigs	7.87±0.58	5.73±0.54	3.33±0.20 ^c	6.15±0.44	4.63±0.62	1.51±0.14 ^c
	n=3	laying hens	17.3 ±1.13 ^a	10.3±1.03	5.40±0.53	12.9 ±0.90 ^a	9.10±0.79	4.97±0.51
		turkeys	11.6±0.83	9.75±0.79	3.63±0.21	9.50±0.86	8.26±0.57	3.27±0.34
Average for the period (x ± Sx)	n=12	dairy cows	5.62±0.19 ^b	1.72±0.05 ^c	2.99±0.31 ^b	4.76±0.20 ^b	1.48±0.05 ^c	1.72±0.15 ^b
		Cv, %**	11.3	9.30	34.6	13.9	11.0	28.9
		pigs	8.30±0.30 ^a	6.08±0.37 b	3.56±0.15 ^c	6.77±0.29 a	4.65±0.17 ^b	1.61±0.11 ^c
		Cv, %	11.8	20.1	14.2	14.2	12.3	22.1
	n=12	laying hens	13.6±1.15 ^c	9.27±0.47	5.56±0.53	9.95±1.78 ^c	7.94±0.13	4.60±0.43
		Cv, %	29.8	23.7	17.3	16.0	25.0	35.2
		turkey	13.9±1.10 ^c	9.94±0.32 ^a	5.02±0.27 ^b	9.16±1.40 ^c	8.67±0.21 a	3.74±0.82 ^b
		Cv, %	27.1	24.4	28.6	16.4	24.5	37.5

* g/kg DM (Dry matter);

** Coefficient of variation (Cv), %;

*** Differences between values by rows are significant at P<0.05 – aa, P<0.01– bb, P<0.001 – cc

value is 3.45 times), both of which values were recorded in the winter (Table 1). The average values for the period were the lowest in dairy cows manure (5.62 ± 0.19 g/kg DM) and increased in the following sequence in the other type of manure: pig manure (by 1.47 times compared to dairy manure), laying hens manure (2.42 times higher, respectively) and turkey manure (2.47 times higher, respectively). By seasons, the values of total N in manure from different animals showed a different trend of change. The values increase by seasons as follows: dairy manure - winter $\rightarrow$ spring $\rightarrow$ summer $\rightarrow$ autumn; pig manure - autumn $\rightarrow$ winter $\rightarrow$ spring $\rightarrow$ summer; laying hens manure - summer $\rightarrow$ spring $\rightarrow$ winter $\rightarrow$ autumn; turkey manure - summer $\rightarrow$ autumn $\rightarrow$ spring $\rightarrow$ winter. This divergence of the results confirms once again that many different factors influence the formation of total nitrogen content in fresh manure - animal species, quantity and composition of the ration, environmental conditions, season, etc.

The coefficient of variation characterizes this parameter as slightly variable for dairy and pig manure with average values for the period of study $C_v = 11.3\%$ and $C_v = 11.8\%$, respectively, and moderately variable for laying hens ($C_v = 29.8\%$) and turkeys ($C_v = 27.1\%$) manure. These results give reason to conclude that factors and environmental conditions that affect the total N content in fresh manure of dairy and pigs are more constant throughout the different seasons and significantly more variable in hens and turkeys manure.

Stored manure. The total N content in stored manure also varies widely as in fresh manure - from 4.45 g/kg DM in dairy cows manure during the winter to 12.9 g/kg DM in laying hens manure during the autumn, but within narrower range (2.90 times is the difference between the maximum and minimum value). Average concentration of total N for the controlled period increases in the following sequence for manure from different animals: dairy cows (4.76 ± 0.20 g/kg DM), pig (1.42 times higher compared to dairy manure), turkey (1.92 g/kg DM times higher, respectively) and laying hens (2.09 times, respectively).

By seasons, the total N content in stored manure from different animals increased as follows: dairy cows - winter $\rightarrow$ summer $\rightarrow$ spring $\rightarrow$ autumn; pigs - autumn $\rightarrow$ winter/summer $\rightarrow$ spring; laying hens - summer $\rightarrow$ spring $\rightarrow$ winter $\rightarrow$ autumn; turkeys - summer $\rightarrow$ autumn $\rightarrow$ winter $\rightarrow$ spring. These seasonal changes in the levels of total N are fully reflected for fresh and stored manure from laying hens, while for the other types of manure there were some disorganizations by the seasons.

The coefficient of variation reveals a fluctuation of the parameter in a narrower range for all kinds of stored manure ($C_v =$ from 13.9% for dairy manure to 16.4% for turkeys manure) compared with the fresh manure ($C_v = 11.3-29.8\%$). Obviously, during the storage the environmental factors influence in a similar way the total N content of stored manure from different animal species, regardless of its initial concentration in the substrate.

When comparing the results obtained for total N content in fresh and stored manure from dairy cows, pigs, laying hens and turkeys, with those of other authors the following picture was revealed. The results for dairy cows manure are: higher than these, reported by Goranov et al. (1978) - 3.40 g/kg DM and those by Clanton (1993) - 3.90 g/kg DM; a little lower than the data by Webb and Archer (1994) - 6.00 g/kg DM; partially coincide with the results by Petrov et al. (1983) 2.10 - 5.00 g/kg DM; in the range of values obtained by Ap Dewi (1994) - 4.00 - 9.00 g/kg DM (for fresh and stored manure) and Christozov et al. (1998) for an average decayed manure - 5.0 g/kg DM.

The results for total N content in manure from pigs are in the range of the results established by Petrov et al. (1983) - 4.20-7.80 g/kg DM, Ensminger and Parker (1984) - 4.00 - 9.00 g/kg DM, Clanton (1993) - 7.50 g/kg DM and Ap Dewi (1994) - 5.00-6.00 g/kg DM for a mixture of pig excrement and urine in natural ratio, but they are higher than the results published by Goranov et al. (1978) - 4.50 g/kg DM, Christozov et al. (1998) - 3.00-6.00 g/kg DM and Katzarov (2005) - 5.50 g/kg DM for fresh manure and 2.60 g/kg DM for slurry from anaerobic lagoons.

The data for total N content in manure from laying hens fully corresponds with the results reported by Petrov et al. (1983) for laying hens manure reared in industrial production systems - 8.90-12.4 g/kg DM, by Tisdale et al. (1985) - 4-21 g/kg DM and by Kaytazov (2006) - 10.0-18.0 g/kg DM. The results for turkeys are similar to those published by Kaytazov (2006) for growing turkey poulters manure - 12.0-25.0 g/kg DM.

Total phosphorus content: Fresh manure. The total P content varies in considerably larger scope than the content of total N - from 1.55 g/kg DM in dairy manure in the winter to 10.3 g/kg DM in laying hens manure in the autumn and in turkeys manure in the winter (the difference between these two values is 6.64 times), (Table 1). The same tendency of change is observed in the average values of total P for the period of investigation by type of manure as well as the total N content. The average values are the lowest in dairy manure (1.72 ± 0.05 g/kg DM) and increase successively in pig manure (3.53 times higher compared to dairy manure), laying hens manure (5.39 times higher, respectively) and turkey manure (5.78 times higher, respectively).

By seasons, the values of total P in manure from dairy cows and pigs change in a similar manner and increase as follows: winter $\rightarrow$ spring $\rightarrow$ autumn $\rightarrow$ summer. The changes in total P content for manure of laying hens and turkeys by seasons are not the same. They increase in the following sequence: laying hens - summer $\rightarrow$ spring $\rightarrow$ winter $\rightarrow$ autumn; turkeys - autumn $\rightarrow$ summer $\rightarrow$ spring $\rightarrow$ winter. Although there is no one-way trend of change in total P content in different types of manure by seasons, it is obvious that seasonal factors have well pronounced influence on the processes involved in the formation of one or another level of phosphorus in the manure.

The coefficient of variation characterizes this parameter as slightly variable for dairy fresh and stored manure, and for stored pig manure ($C_v = 9.30\% - 12.3\%$), and much more variable for fresh pig, laying hens and turkey manure, and for stored laying hens and turkey manure ($C_v = 20.1-25.0\%$). These results somewhat explain the seasonal changes in total P content in fresh and stored manure.

Stored manure. The total P content in stored manure varies from 1.37 g/kg DM in dairy manure to 9.70 g/kg DM in turkey manure, both concentrations are determined in winter (Table 1). The difference between the maximum and minimum value is 7.08 times, i.e. more than in fresh manure. The average concentration of total P in different types of manure for the period of study increases in the same order as for total N content: dairy manure (1.48 ± 0.05 g/kg DM), followed by pig, laying hens and turkey manure, where the average values are higher compared to cattle manure 3.14, 5.36 and 5.86 times, respectively.

The seasonal changes in the values of total P content are divergent for manure from different animals and increase as follows: dairy cows - winter $\rightarrow$ summer $\rightarrow$ spring $\rightarrow$ autumn; pigs - spring $\rightarrow$ winter $\rightarrow$ autumn $\rightarrow$ summer; laying hens - winter $\rightarrow$ spring $\rightarrow$ summer $\rightarrow$ autumn; turkeys - autumn $\rightarrow$ spring $\rightarrow$ summer $\rightarrow$

winter. It should be emphasized that the differences for the total P content in manure by different animal species are not significant and they are not statistically proven. This gives grounds to assume that the established ranking of the results by seasons may have a random character.

By coefficient of variation values the tested manures are allocated in two groups – manures with low fluctuation of the parameter (dairy and pig manure, $C_v = 11.0-12.3\%$) and manures with a moderate fluctuation (laying hens and turkeys manure, $C_v = 24.5-25.0\%$).

The average seasonal values for total P in dairy manure are: in the range of the data reported by Christozov et al. (1998) – 1.00-3.00 g/kg DM and by Ap Dewi (1994) – 1.00-8.00 g/kg DM, bringing it closer to the lower boundary; very close to the results of Goranov et al. (1978) and Clanton (1993) – 1.60 g/kg DM; considerably lower than the results of Petrov et al. (1983), found for the manure of cattle reared in intensive production systems (4.60-7.10 g/kg DM) and the results, published by Pagliari (2014) – 2.8-18.3 g/kg DM.

The findings for pig manure are: within the ranges specified by Petrov et al. (1983) – 0.60-7.20 g/kg DM, by Ap Dewi (1994) – 1.0-6.0 g/kg DM for pig feces and urine, mixed in the natural ratio and by Pagliari (2014) – 3.9-48.7 g/kg DM; close to the data of Clanton (1993) – 5.70 g/kg DM, Katzarov et al. (2003) – 5.90 g/kg DM for slurry stored in anaerobic lagoons and Ensminger and Parker (1984) – 3.10-5.80 g/kg DM; lower than the results published by Goranov et al. (1978) – 1.90 g/kg DM and Christozov et al. (1998) – 2.0-4.0 g/kg DM. The results for laying hens manure confirm the data reported by Petrov et al. (1983) 5.40-15.9 g/kg DM, by Ap Dewi (1994) – 3-15 g/kg DM, and by Kaytazov (2006) – 8.0-12.0 g/kg DM. What distinguishes them is that our results fluctuate within much narrower ranges than those of the quoted authors. The established values for total P in turkey manure are a little lower than the lower limit values reported by Kaytazov (2006) for growing turkey poult's manure (10.0-14.0 g/kg DM) and by Pagliari (2014) for turkey manure – 10.1-28.2 g/kg DM.

Total potassium content: Fresh manure. The levels of total K vary between 2.16 g/kg DM in dairy cows manure during the winter and 6.10 g/kg DM in laying hens and turkey manure during the summer (Table 1). The difference between the maximum and minimum value for total K content is 2.82 times. These ranges of variation are significantly narrower in comparison with the ranges of variation in total N (3.45 times) and total P (6.64 times) content in fresh manure. These results indirectly support the fact that K in fresh manure is a more stable element in comparison with N and P. The average values of total K for the monitored period reveal a similar trend of alteration in manures from the studied animal species as of total N and total P content. The lowest average value is found for dairy manure (2.99 ± 0.31 g/kg DM) followed by pig, turkey and laying hens manure, where values increase 1.19, 1.68 and 1.86 times, respectively, compared to dairy manure. The only difference is in the ranking of the last two types of manure that are with exchanged positions. Bearing in mind that the differences between the values for total K in turkey and laying hens manure are minor, they can be ignored.

By seasons, the values of total K in manure from different animals increase as follows: dairy cows - winter → summer → autumn → spring; pigs - autumn → winter → summer/spring; laying hens - spring → autumn → winter → summer; turkeys - autumn → winter → spring → summer. The coefficient of variation reveals the biggest variation of the parameter for dairy manure ($C_v = 34.6\%$)

followed by that of turkeys ($C_v = 28.6\%$), laying hens ($C_v = 17.3\%$) and pigs manure ($C_v = 14.2\%$). The results clearly emphasize the role of animal species on the formation of total K content in manure.

Stored manure. The total K content in stored manure varies from 1.43 g/kg DM in dairy manure in autumn to 5.01 g/kg DM in laying hens manure in summer (Table 1). The difference between the maximum and minimum value is greater than in fresh manure – 3.50 times compared to 2.82 times, respectively. The average concentration of total K is the lowest in pig manure (1.61 ± 0.11 g/kg DM) followed by dairy (1.07 times higher compared to pig manure), turkey (2.32 times higher, respectively) and laying hens (2.85 times higher, respectively) manure.

The seasonal changes in the values of total K content are divergent in manure from different animal species and increase as follows: dairy cows - autumn → winter → summer → spring; pigs - winter → autumn → spring → summer; laying hens - winter → spring → autumn → summer; turkeys - autumn → winter → spring → summer. The coefficient of variation characterized this indicator as a significant variable with values between $C_v = 28.9\%$ (dairy manure) and $C_v = 37.5\%$ (turkey manure).

The results obtained for total K content in dairy manure are: within the range reported by Petrov et al. (1983) – 1.20-2.30 g/kg DM; higher than the results found by Penn and McGrath (2014) – 1.45 g/kg DM; lower than the data published by Goranov et al. (1978) – 4.00 g/kg DM, by Clanton (1993) – 3.10 g/kg DM, by Webb and Archer (1994) – 7.00 g/kg DM and by Christozov et al. (1998) – 3.0-6.0 g/kg DM.

With regards to the pig manure we have matching results for the fresh pig manure with those of Petrov et al. (1983) – 2.40-7.60 g/kg DM and Ensminger and Parker (1984) – 2.70-4.50 g/kg. The results for fresh pig manure are lower in comparison to those reported by Penn and McGrath (2014) – 4.91 g/kg DM. The results for stored pig manure are similar to the data by Katzarov (2005) – 1.90 g/kg DM and lower than the values found by Goranov et al. (1978) and Christozov et al. (1998) – 6.00 g/kg DM, by Clanton (1993) – 5.90 g/kg DM and Ap Dewi (1994) – 4.00 g/kg DM.

The results for total K in manure of laying hens are similar to the results published by Petrov et al. (1983) – 3.10-6.30 g/kg DM and by Kaytazov (2006) – 5.00-7.00 g/kg DM, and higher compared to the results of Penn and McGrath (2014) – 3.17 g/kg DM.

Losses of nutrients (total N, total P and total K) during the storage of manure. It is well known that during storage farm manure loses some of its biogenic elements as a result of an ongoing aerobic (on the surface of the manure) and anaerobic (inside the manure) process of organic compounds mineralization. Losses of total N, total P and total K content in stored manure for 6 months compared to that in fresh manure from different farm animals (in %) are presented in Table 2. The results show that the losses of biogenic elements are in different range as to the seasons and by type of manure.

The amount of total N in stored manure decreases compared to that in the fresh manure as follows: dairy cows manure - from 10.7% in spring to 20.2% in summer (the differences are statistically significant for spring at $P < 0.05$); pig manure - from 12.6% in spring to 21.9% in autumn (the differences are statistically significant for winter at $P < 0.05$ and for summer at $P < 0.001$); laying hens manure - from 12.6% in spring to 35.8% in winter (the differences are statistically significant for spring and autumn at $P < 0.05$); turkey manure - from 18.1% in autumn to 47.0% in winter (the differences are statistically significant for winter and spring at $P < 0.01-0.001$)

Table 2. Losses of total N, total P and total K in stored manure compared to that in fresh manure from different farm animals (in %)

Season	Species of animals	Losses, %		
		N	P	K
		In stored manure compared to fresh manure		
Winter	dairy cows	16.4	11.6	30.6
	pigs	15.4	13.4	62.1
	laying hens	35.8	19.6	34.5
	turkeys	47.0	5.8	20.3
Spring	dairy cows	10.7	12.9	52.3
	pigs	12.6	18.1	56.3
	laying hens	12.6	14.9	7.6
	turkeys	41.5	16.0	39.7
Summer	dairy cows	20.2	21.0	31.6
	pigs	21.6	37.2	46.4
	laying hens	13.4	11.5	17.9
	turkeys	22.2	14.2	24.6
Autumn	dairy cows	13.7	10.2	45.0
	pigs	21.9	19.2	54.7
	laying hens	25.4	11.7	8.0
	turkeys	18.1	15.3	10.0
Average for the period	dairy cows	15.3	14.0	42.5
	pigs	18.4	23.5	54.8
	laying hens	26.8	14.4	17.3
	turkeys	34.1	12.8	25.5

(Table 2, Figure 1). The average losses of N for the investigated period were the smallest in dairy manure (15.3%) followed by pig (18.4%), laying hens (26.8%) and turkey manure (34.1%). The differences between average values for the studied period are statistically significant for all types of manure at $P < 0.05-0.001$.

By losses of nitrogen the studied manures can be divided into two groups - with relatively low losses of total N (dairy and pig manures) and with relatively high losses of total N (laying hens and turkey manures). This suggested that the main factors which influenced N losses during the storage of manure are the type of manure and manner of its storage. Obviously storage of dairy cows manure on heaps on the ground and of pig manure in earth lagoons

Figure 1. Seasonal losses of total nitrogen in the manure of different animal species, g/kg DM

creates more favorable conditions to reduce N losses compared with the storage of laying hens and turkey manure and litter on the ground near the premises. It can be assumed that in our case the additional moistening (by precipitations) of the stored poultry manure under open air conditions is also a factor leading to greater reduction of total N, as the high humidity supports the microbial processes related to the degradation of nitrogen compounds.

The established losses of total N in dairy manure are in the range of the results obtained by Clanton (1993) in the same manner of manure storage (15-35%) and of those, reported by Christozov et al. (1998) which provide much greater variation of N losses (6-49%). Regarding pig manure, Clanton (1993) determines much larger losses of N in manure storage in lagoons (70-80%) than in our case, where the average losses of total N for the study period are 18.4%. The losses of total N in poultry manure are close to the data of Chastain et al. (2002) for storage of poultry manure in open manure heaps - 21% loss of $\text{NH}_4\text{-N}$ and 30% loss of total N. According to Nahm (2003), the majority of nitrogen (around 60-70%), which is excreted in manure is in the form of uric acid and urea, which are converted into ammonia (creating air pollution), nitrite and nitrate (penetrating the soil and the groundwater).

The losses of total P in storage manures by seasons vary in the following ranges: dairy manure - between 10.2% in autumn and 21.0% in summer (the differences are statistically significant for spring and summer at $P < 0.001$); pig manure - between 13.4% in winter and 37.2% in summer (the differences are statistically significant for spring and summer at $P < 0.05-0.01$); laying hens manure - between 11.5% in summer and 19.6% in winter (the differences are statistically significant for winter at $P < 0.05$); turkey manure - between 5.8% in winter and 16.0% in spring (the differences are not statistically significant) (Table 2, Figure 2). Average for the period losses were the smallest in turkey manure (12.8%) followed by dairy cows (14.0%), laying hens (14.4%) and pig manure (23.5%). The differences are statistically significant for dairy, pigs and turkey manure at $P < 0.05-0.001$. On this indicator the tested manures could also be divided into two groups - one group with very close losses of total N (dairy, laying hens and turkey manure), the other - with significantly higher losses of total N - pig manure. In this case, the factor with the greatest impact on losses of total K in manure is the manner of its storage. Dairy cows, laying hens and turkey manures are stored outdoors on the ground, while pig manure is stored in lagoons. This gives reason to assume that the environmental conditions in lagoons favor to a greater extent degradation of potassium in stored pig manure.

The results for total K losses in stored manures indicate that this parameter varies much more widely than the losses of total N and

Figure 2. Seasonal losses of total phosphorus in the manure of different animal species, g/kg DM

Figure 3. Seasonal losses of total potassium in the manure of different animal species, g/kg DM

total P, and are in the range as follows: dairy cows manure - from 30.6% in winter to 52.3% in spring (the differences are statistically significant for all seasons at $P < 0.01-0.001$); pig manure - from 46.4% in summer to 62.1% in winter (the differences are statistically significant for all seasons at $P < 0.01-0.001$); laying hens manure - from 7.6% in spring to 34.5% in winter (the differences are not statistically significant); turkey manure - from 10.0% in autumn to 39.7% in spring (the differences are not statistically significant) (Table 2, Figure 3). On average for the period of study the losses of total K were the smallest in laying hens manure (17.3%) followed by turkey (25.5%), dairy cows (42.5%) and pig manure (54.8%). The differences are statistically significant for dairy, pig and turkey

manure at $P < 0.01-0.001$.

Divergent trends were established for the losses of nutrients during the storage of the tested manures. For dairy cows manure the biggest losses are of total K, followed by total N and total P ($K \rightarrow N \rightarrow P$); for pig manure – $K \rightarrow P \rightarrow N$; for laying hens and turkey manure – $N \rightarrow K \rightarrow P$, respectively.

Ratio between total N, total P and total K content (N:P:K) in fresh and stored manure. The key to avoiding environmental problems associated with manure application is to apply manure based on crop nutrient requirements and implementing available best management practice in the farm. In this regard it is important to assess the tested manures based on the ratio of biogenic elements (Table 3).

It is noteworthy that the ratio between the nutrients, both in fresh and in stored manure from different animal species varies in relatively wide limits as to the seasons and by average of the studied period. Another fact that needs to be noted is that on this criterion the tested manures can be distributed in two groups. In one group can be included dairy cows manure, where ratios between nutrients are 54.4% N: 16.7% P: 28.9% K for fresh manure and 59.8% N: 18.6% P: 21.6% K for stored manure, the other group includes pig, laying hen and turkey manure, where these ratios are between 46.3% N: 33.9% P: 19.8% K for fresh pig manure and 48.2% N: 34.4% P: 17.4% K for fresh turkey manure, and between 42.5% N: 40.2% P: 17.3% K for stored turkey manure and 52.0% N: 35.7% P: 17.4% K for stored pig manure. In the second group of manures N and P content is lower, and the P content is higher as compared with the manure of the first

Table 3. Ratio between total N, total P and total K content (N:P:K) in fresh and stored manure from different farm animals (in %)

Season	Species of animals	Manure					
		Fresh			Stored		
		N : P : K, %			N : P : K, %		
Winter	dairy cows	58.9	17.2	23.9	60.8	18.7	20.5
	pigs	48.2	31.4	20.4	53.9	35.9	10.2
	laying hens	52.9	29.5	17.6	49.1	34.3	16.7
	turkeys	55.5	31.1	13.4	42.4	42.2	15.3
Spring	dairy cows	46.7	14.3	38.9	57.3	17.1	25.6
	pigs	47.7	31.2	21.1	54.5	33.4	12.1
	laying hens	43.7	35.3	21.0	43.6	34.3	22.1
	turkeys	52.0	30.1	17.9	45.7	38.0	16.3
Summer	dairy cows	56.6	18.4	25.0	58.8	18.9	22.3
	pigs	43.3	38.3	18.4	50.0	35.5	14.5
	laying hens	41.4	34.5	24.1	41.6	35.5	22.9
	turkeys	40.0	37.0	23.0	38.8	39.5	21.6
Autumn	dairy cows	57.1	17.4	25.4	62.5	19.8	17.8
	pigs	46.5	33.8	19.7	50.0	37.7	12.3
	laying hens	52.4	31.2	16.4	47.8	33.7	18.4
	turkeys	46.4	39.0	14.5	45.2	39.3	15.5
Average for the period	dairy cows	54.4	16.7	28.9	59.8	18.6	21.6
	pigs	46.3	33.9	19.8	52.0	35.7	17.4
	laying hens	47.8	32.7	19.5	44.2	35.3	20.5
	turkeys	48.2	34.4	17.4	42.5	40.2	17.3

group. These differences are highlighted in fresh as well as stored manures.

FAO (2000) identified as suitable for agriculture mineral and organic fertilizers in which the ratio between nutrients (N:P:K) is in the range as follows: 22:22:11, 14:35:14, 19:19:19, 12:24:12, 15:15:15, 11:22:22 and 10:26:26. The ratios between nutrients in all tested manures (fresh and stored) are included in all of the above ratios but in many cases the values for N, P and K exceed them, often considerably. A similar picture is observed for the other ratio – N:P recommended by FAO (2000) for fertilizers (16:20, 28:28 and 18:46). Therefore, before the use of organic fertilizers, it is necessary to determine the content of nutrients in them and in the soil which will be fertilized. The quantities of nutrients necessary for cultivation of the crops concerned should also be calculated. Only in this way excessive importation of nutrients in the soil will be avoided, which will affect the expected yields and will cause pollution of soil and groundwater. For example, land application of animal manures to meet crops N needs can lead to an accumulation of P in soil because the N:P ratio in animal manures (about 2:1 by literature source, in our study it is between 1.4:1 and 3.3:1 for fresh manure, and between 1.1:1 and 3.2:1 for stored manure) is less than the N:P ratio of about 8:1 taken up by more crops and grasses (Sharpley et al., 2003). Thus, repeated land application of manure based on plant N needs results in excessive P concentration in the soil and may saturate the soil's capacity to retain phosphorus.

Conclusion

On the basis of agro-ecological assessment of manure from different farm animals by content of biogenic elements (N, P, K) it was found that: a) biogenic elements in fresh and stored manure decreased in the following gradation by average values for the studied period: turkeys – total N (13.9 - 9.16 g/kg DM), total P (9.94 - 8.67 g/kg DM) and total K (5.02 - 3.74 g/kg DM); laying hens – total N (13.6-9.95 g/kg DM), total P (9.27- 7.94 g/kg DM) and total K (5.56 - 4.60 g/kg DM); pigs – total N (8.30 -6.77 g/kg DM), total P (6.08-4.65 g/kg DM) and total K (3.56 - 1.61 g/kg DM); dairy cows – total N (5.62 -4.76 g/kg DM), total P (1.72 - 1.48 g/kg DM) and total K (2.99 - 1.72 g/kg DM); b) the change in nutrients content of different types of manures by seasons both in fresh and in stored manure are divergent; c) coefficients of variation values determined the surveyed manures by content of nutrients from minor to significant variable ($C_v = 9.30-37.5\%$); d) during the storage of the tested manures nutrient losses are the greatest for potassium (17.3-54.8%), followed by nitrogen (15.3-34.1%) and phosphorus (12.8-23.5%); e) the average losses of biogenic elements in manure by animal species are in varying degrees: dairy cows (total N 15.3%; total P 14.0%; total K 42.5%); pigs (total N 18.4%; total P 23.5%; total K 54.8%); laying hens (total N 26.8%; total P 14.3%; total K 17.3%) and turkeys (total N 34.1%; total P 12.8%; total K 25.5%); f) the ratio between nutrients (N:P:K), both in fresh and in stored manure from different animal species vary in relatively wide limits as to the seasons (12.6%N:18.1%P:56.3%K – 41.5%N:16.0%P:39.7%K) and by average for the studied period (15.3%N:14.0%P:42.5%K – 34.1%N:12.8%P:25.5%K).

References

- Ap Dewi I, 1994. The Use of Animal Waste as a Crop Fertilizer. In: Pollution in Livestock Production Systems (Eds. Ap. Dewi and R.F.E. Axford), Cab International, Wallingford, UK, 309-332.
- Burkholder J, Libra B, Weyer P, Heathcote S, Kolpin D, Thorne PS and Wichman M, 2007. Impacts of waste from concentrated animal feeding operations on water quality. *Environmental Health Perspectives*, 115, 308-12.
- Bywater A, 2011. A review of Anaerobic Digestion Plants of UK Farms – Barriers, Benefits and Case Studies, Royal Agricultural Society of England, 13-30.
- Cederberg C, Berglund M, Gustavsson J and Wallman M, 2012. Environmental impacts from livestock production with different animal welfare potentials – a literature review, SIK report 844, Göteborg, Sweden, 19-39.
- Capper JL, 2011. The environmental impact of beef production in the United States: 1977 compared with 2007. *Journal of Animal Science*, 89, 4249-4261.
- Chastain JP, Camberto JJ and Skewes P, 2002. Poultry Manure Production and Nutrient Content. South Carolina Confined Animal Manure Managers Certification Program – Poultry. Clemson Extension. Clemson University, Clemson, SC, 66-84.
- Christozov A, Carov S, Angelova R, Pavlov D and Veleva V, 1998. Soil Science and Agriculture (Ed. A. Christozov), Stara Zagora, 26-57, 80-85 (Bg).
- Clanton CJ, 1993. Nutrient characteristics of manure, In: Proceeding of Livestock management conference (Eds. J. Shurson, G. Wagner and J. Rhoads), January 19-20, St. Paul, Minnesota, USA, pp. 73-75.
- Ensminger ME and Parker RO, 1984. Manure. In: Swine Science, 5th Edition, The interstate Printers & Publishers, Inc., Danville, Illinois, USA, 294-299.
- Faassen H and van Dijk H, 1987. Manure as a source of nitrogen and phosphorus in soils. In: Animal Manure on Grassland and Fodder Crops: Fertilizer or Waste? (Eds. H. van der Meer et al.), Martinus Nijhoff Publishers, Wageningen, 45-68.
- Fonseca MB, 2011. Supply of and access to key nutrients NPK for fertilizers for feeding the world in 2050. Report. Universidad Politécnica de Madrid, Spain, 7-33.
- Food and Agriculture Organization (FAO), 2000. Fertilizers and their use. International fertilizer industry association, 4th edition, Published by FAO and IFA, Rome, 3-28.
- Food and Agriculture Organization (FAO), 2003. Fertilizer use by crop in Poland. Published by FAO, Rome, 23-35.
- Food and agriculture organization (FAO), 2015. World fertilizer trends and outlook to 2018, Rome, 8-20.
- Goranov I, Gudev R, Garanicheva K and Dossev N, 1978. Cleaning, storage and use of manure in livestock farms. Zemizdat Publishing house, Sofia, 5-91 (Bg).
- Goss MJ, Tubeileh A, Goorahoo D, 2013. A review of the use of organic amendments and the risk of human health. *Advances in Agronomy*, 120, 275-239.
- Hellmuth R and Hochmuth G, 2015. Managing Nitrogen Inputs and Outputs on a Dairy Farm. Livestock & Forages, Cooperative extension, University of Florida, USA, 1-4.
- Hoffmann I, 2002. Crop-livestock interactions and soil fertility management in Northwest Nigeria, In: Proceeding of the First Virtual Global Conference on Organic Beef Cattle Production, 02 September-15 October, pp. 1-11.
- Humane Society of the United States (HSUS), 2008. The Impact of Industrialized Animal Agriculture on the Environment, USA, 1-8.
- Katarov C and Pavlov D, 2002. Use of manure from pig farms in agriculture, KOTAPublishing house, Stara Zagora, pp. 24-30 (Bg).

- Katzarov V, Pavlov D and Stoykov A**, 2003. Opportunities to use manure from pig farms in agriculture. *Ecology and Future*, 1, 86-90 (Bg).
- Katzarov V**, 2005. Nutrient balance in manure from pig farms when used in agriculture. *Ecology and Future*, 4, 69-74 (Bg).
- Kaytazov G**, 2006. The problem with poultry manure and can we reap the benefits of its solution. *Poultry-farming*, 4, 10-14 (Bg).
- Loreau M, Naeem S, Inchausti P, Bengtsson J, Grime GP, Hector A, Hooper DU, Huston MA, Raffaelli D, Schmid B, Timan D and Wardle DA**, 2001. Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science*, 294, 804-808.
- Manure Management Plan for Kaliningrad Region (MMPKR)**, 2014. BASE Project - Implementation of the Baltic Sea Action Plan in Russia. Baltic Marine Environment Protection Commission HELCOM, 6-31.
- Massie L, Converse J and Holmes B**, 1993. A system approach to waste management. In: *Proceeding of Livestock management conference* (Eds. J. Shurson, G. Wagner and J. Rhoads), January 19-20, St. Paul, Minnesota, USA, 56-63.
- Mikkelsen MH, Albreksten R and Gyldencoerne S**, 2014. Danish emission inventories for agriculture. *Scientific reports from Danish Center for Environment and Energy*, 108, 36-45.
- Nahm KH**, 2003. Evaluation of the nitrogen content in poultry manure. *World's Poultry Science Journal*, 59, 77-88.
- Nguyen TLT, Hermansen JE and Mogensen L**, 2010. Environmental consequences of different beef production systems in the EU. *Journal of Cleaner Production*, 18, 756-76.
- Pagliari PH**, 2014. Variety and Solubility of phosphorus forms in animal manure and their effects on soil test phosphorus. In: *Applied manure and nutrient chemistry for sustainable agriculture and environment* (Eds.: Zhongqi H and Zhang H). Springer Dordrecht, Heidelberg, New York, London, 141-150.
- Penn CJ and McGrath DM**, 2014. Chemistry and application industrial by-products to animal manure for reducing phosphorus losses to surface water. In: *Applied manure and nutrient chemistry for sustainable agriculture and environment* (Eds.: Zhongqi H and Zhang H). Springer Dordrecht, Heidelberg, New York, London, 210-226.
- Petrov P, Bozhilov A, Vankov I, Mladenov I, Petrov G, Marinova S and Bildirev N**, 1983. Liquid manure - treatment and utilization in agriculture (Ed. P. Petrov). Zemizdat Publishing house, Sofia, 282-298 (Bg).
- Reijs JW, Meijer WH, Bakker EJ and Lantinga EA**, 2003. Explorative research into quality of slurry manure from dairy farms with different feeding strategies. *NJAS - Wageningen Journal of Life Sciences*, 51, 67-89.
- Roberts TL**, 2009. The Role of Fertilizer in Growing the World's Food. *Better Crops*, 93, 12-15.
- Sharpley AN, Weld JL, Beegle KPJA, Gburek WJ, Moore PA Jr, Mullins J**, 2003. Development of phosphorus indices for nutrient management planning strategies in the USA. *Journal of Soil and Water Conservation*, 58, 137-152.
- Scotti R, Bonanomi G, Scelza R, Zoina A and Rao MA**, 2015. Organic amendments as sustainable tool to recovery fertility in intensive agricultural systems. *Journal of Soil Science and Plant Nutrition*, 15, 333-352.
- Svoboda IF**, 2003. Anaerobic digestion, storage, oligolysis, lime, heat and aerobic treatment of livestock manures. Final report, FEC Services Ltd., Scotland, 22-52.
- Tazisong IA, Senwo Z N and Zhongqi He**, 2015. Phosphatase Hydrolysis of Organic Phosphorus Compounds. *Advances in Enzyme Research*, 3, 39-51.
- Tilman D, Cassman KG, Matson PA, Naylor R and Polasky S**, 2002. Agricultural sustainability and intensive production practices. *Nature*, 418, 671-677.
- Tisdale SL, Nelson WL and Beaton JD**, 1985. Soil Fertility and Fertilizers, 4th edition, Macmillan, New York, USA, 58-76.
- Velthof GI, Bannink A, Oenema O, van Der Meer HG and Spoelstra SF**, 2000. Relationships between animal nutrition and manure quality: A literature review on C, N, P and S compounds. Alterra, Wageningen, p. 44.
- Voermans JAM**, 1993. Waste management practices in Europe. In: *Proceeding of Livestock management conference* (Eds. J. Shurson, G. Wagner and J. Rhoads), January 19-20, St. Paul, Minnesota, USA, 8-13.
- Wang JJ and Gaston LA**, 2014. Nutrient chemistry of manure and manure-impacted soils as influenced by application of bauxite residue. In: *Applied manure and nutrient chemistry for sustainable agriculture and environment* (Eds.: Zhongqi H and Zhang H). Springer Dordrecht, Heidelberg, New York, London, 239-266.
- Webb J and Archer J**, 1994. Pollution of soils and water sources by wastes from livestock production systems. In: *Pollution in Livestock Production Systems* (Eds. Ap. Dewi and R.F.E. Axford), Cab International, Wallingford, UK, 189-204.
- Zaharinov B, Shindarska Z, Marinova – Garvanska S, Kirov V and Baykov B**, 2013. Agroecologic valuation of organic waste in different technologies of storage. In: *Proceedings of the 10th International Symposium Modern Trends in Livestock Production* (ed. Z. Pavlovski), October 2-4, Editor Institute for Animal Husbandry, Belgrade, Serbia, pp. 580-589.
- Zhang H and Schroder J**, 2014. Animal manure production and utilization in the USA. In: *Applied manure and nutrient chemistry for sustainable agriculture and environment* (Eds.: Zhongqi H and Zhang H). Springer Dordrecht, Heidelberg, New York, London, 2-22.
- Zakikhani K, Kashani A and Paknejad F**, 2016. Effect of nitrogen level, green and animal manure on the growth attribute of corn crop (*Zea mays* L.). *Journal of Experimental Biology and Agricultural Science*, 4, 225-231.

Review

- Antimicrobial activity of *Lactobacillus acidophilus* against pathogenic and food spoilage microorganisms: A review** 3
T. Dinev, G. Beev, S. Denev, D. Dermendzhieva, M. Tzanova, E. Valkova

Genetics and Breeding

- Heterosis and degrees of dominance of grain yield and grain yield elements in maize hybrids in different groups of ripeness** 10
M. Ilchovska
- Use of recurrent selection of early flowering in late maize synthetic population. Results of second cycle of breeding.** 16
N. Petrovska, V. Valkova
- Productivity and adaptability of new genotypes field pea (*Pisum sativum* L.) cultivated under environmental condition of Southern Romania** 19
R. Sturzu, A. M. Ene, Cr. Melucă, J. M. Cojocar
- Nitrogen uptake and expense in durum wheat depending on genotype and nitrogen fertilization** 26
G. Panayotova, M. Almaliev, S. Kostadinova

Nutrition and Physiology

- Haematological investigations upon acute intoxication with carbofuran in dogs** 35
R. Binev, I. Valchev, R. Russenov, Y. Nikolov

Production Systems

- Phytosanitary status and yield of kamut (*Triticum turgidum polonicum* L.) grown in organic and biodynamic farming** 42
V. Maneva, D. Atanasova, T. Nedelcheva
- Hot-water treatment of gladiolus cormels for control of corm-borne fungal diseases** 45
S. Bistrichanov, T. Vatchev, Z. Avramov
- Productivity of common wheat (*Triticum aestivum* L.) grown after various predecessors and nitrogen fertilization rates** 48
M. Gerdzhikova

Agriculture and Environment

- Agro-ecological assessment of manure from different farm animals by content of biogenic elements** 53
D. Dermendzhieva, G. Kostadinova, G. Petkov, D. Dimov, T. Dinev, T. Penev, Ch. Miteva, J. Mitev

Screening of cucurbitaceous rootstocks against root-knot nematodes (*Meloidogyne* spp.) and soilborne pathogens (*Fusarium* spp. and *Pythium* spp.) **62**

V. Yankova , D. Markova, N. Velkov, S. Masheva

Animal hygiene assessment of microclimate in semi open free-stall barns for dairy cows **67**

D. Dimov, Ch. Miteva, I. Marinov, Zh. Gergovska, T. Penev, A. Enchev

Product Quality and Safety

Accumulation of astaxanthin and canthaxanthin in muscle tissues of Rainbow trout (*Oncorhynchus mykiss* W.) fed with xanthophyll supplemented feed **77**

M. Tzanova

Chemical composition and technological characteristics of wines from red grape varieties, selected in Bulgaria **83**

V. Haygarov, T. Yoncheva, Z. Nakov, M. Ivanov, D. Dimitrov

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors
The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings: Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows. IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 9, Number 1
March 2017

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz