ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 9, Number 2
June 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate on author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc. and AGRIS (FAO).

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No DNP 05-21/20.12.2016, financed from Fund 'Scientific Research' grant Bulgarian scientific Periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 9, Number 2
June 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Review

Alternatives for optimisation of rumen fermentation in ruminants

T. Slavov*

*Department of Morphology, Physiology and Nutrition, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria

(Manuscript received 12 March 2017; accepted for publication 26 May 2017)

Abstract. *The proper knowledge on the variety of events occurring in the rumen makes possible their optimisation with respect to the complete feed conversion and increasing the productive performance of ruminants. The inclusion of various dietary additives (supplements), biologically active substances, nutritional antibiotics, probiotics, enzymatic preparations, plant extracts etc. has an effect on the intensity and specific pathway of fermentation, and thus, on the general digestion and systemic metabolism. The optimisation of rumen digestion is a method with substantial potential for improving the efficiency of ruminant husbandry, increasing of quality of their produce and health maintenance.*

Keywords: nutrients, degradation, digestive processes, modification

Introduction

The digestion physiology of ruminants is determined by anatomo-physiological features of forestomachs' structure and function. Intensive microbial processes of degradation and synthesis occur in forestomachs along with mechanical changes in feed. The entire digestive tract of ruminants is inhabited by microorganisms but a true symbiosis with the host is exhibited only by the microflora colonising the forestomachs (Santra and Karim, 2003; Paul et al., 2010).

The rumen environment provides natural conditions for the development of obligate and facultative anaerobes. There is constant inflow of nutrients and fluids in the rumen. Under the influence of rumen contractions, feed matter is continuously mixed and undergoes mechanical changes. Apart these processes, the undegraded and non-utilised feed ingredients, as well as a substantial biomass of microbial and infusorian cells moves to the more distal compartments of the digestive tract – the rumen, the reticulum and the abomasum. The osmotic pressure in the rumen environment is constant. Its temperature varies within the optimum range for microbial life processes: 38-40 °C, rumen pH is about 6-7. This parameter also varies within a narrow range due to the neutralising effect of alkaline constituents of the saliva. The absorption of volatile fatty acids (VFA) and ammonia resulting from protein degradation, also contributes to the slight variations in rumen pH.

The feed nutrients undergo a number of chemical, biochemical and physical alterations in the rumen of ruminants where they are converted into simpler compounds. Multiple research studies have investigated the role of rumen ciliates on rumen digestion depending on the age and morphological development of forestomachs in small ruminants (Rusev et al., 1973; Petkov, 1982), feeding of rations with various structure and type, supplemented with enzymatic preparations (Sivkova, 2007), the supplementation of diets with biotechnological products (Radev, 2000), rumen defaunation

etc. The effects of defaunation on the digestion in forestomachs and the health was investigated in small ruminants by Enev (1994). The end products of nutrients' degradation are utilised by microorganisms for production of carbohydrates, proteins and other organic compounds, which are subsequently degraded by digestive juices in the abomasum and small intestine.

The aim of the present study is to review the available literature on existing options for optimisation of fermentation in the rumen of ruminants.

Degradation of the main groups of nutrients in forestomachs

Degradation of carbohydrates

Under the influence of extracellular enzymes of rumen bacteria and infusorians, carbohydrates undergo complex changes to simpler compounds, most notably monosaccharides.

Carbohydrates comprise 40-80% of the dry matter of plant feeds. They are the main source of energy for herbivores. Carbohydrates are divided into water-soluble – mono, di- and polysaccharides including compounds as cellulose, hemicellulose, starch and glycopolysaccharides (pectins), fructans etc. Each of these compounds experiences complex changes, mainly to simple sugars, in the forestomachs of ruminants under the influence of rumen microbial enzymes. The simple sugars are converted into VFA, alcohols and gases (carbon dioxide, hydrogen, methane). At the same time, the end products of carbohydrate degradation are used by bacteria and infusorians for synthesis and deposition of complex carbohydrates and other compounds.

The degradation of polysaccharides by rumen microflora occurs on several stages. First, simple sugars with a short carbon chain are extracellularly produced. Then, intracellularly, the sugars are further converted by hydrolysis or phosphorylation to monosaccharides, and through pyruvate and phosphoenolpyruvate glycolysis – to VFA, methane etc. (Rusev et al., 1984). The

* e-mail: todor_slavov@abv.bg

degradation of crude fibres from cellular walls allows the utilisation of the content of plant cells.

It is presumed that the mechanism of cellulose degradation in the ruminant forestomachs consists in production of extracellular cellulose by cellulolytic bacteria. Holotrichic ciliates are able to attach to substrates containing soluble carbohydrates through chemotaxis. Thus, they utilise the monosaccharides glucose, fructose, galactose, some soluble oligomers and polysaccharides containing one or several of these sugars (Orpin and Letcher, 1984). The extent of carbohydrate fermentation is influenced by the concentration and the type of available substrate, and the factors determining the rumen ecosystem environment.

Cellulose and hemicellulose are among the most important ingredients of crude fibre. They are the principal components of plant feeds - about 40-50%. Crude fibre makes up the cellular walls and thus their degradation allows for easier utilisation of cell content. The mechanism of cellulose degradation in the forestomachs of ruminants comprises production of extracellular cellulase of cellulolytic bacteria, which gradually degrades cellulose. The enzyme is β 1-4 glucosidase. Its activity results in production of cellobiose, which, via phosphorylation, is converted to glucose. The glucose in rumen content is degraded very rapidly and its level is usually low. The degree of *in vitro* uptake of glucose by *Dasytricha ruminantium* is approximately 20 $\mu\text{mol/mg}$ cell protein, while the uptake of glucose by *Isotricha* spp. is twice higher (Yarlett et al., 1981).

The digestibility of cellulose in the rumen depends on the number of factors, the most important among which are the type of feeds, the size and activity of rumen bacterial and ciliate populations. In general, cellulose digestibility ranges from 30 to 80%. As a result from the growth and metabolic activity of rumen microflora, polysaccharides and oligosaccharides in plant feeds are converted to monosaccharides - mainly glucose. Then, monosaccharides undergo intracellular degradation to many simpler products. Most important for the energy metabolism of ruminants are the volatile fatty acids (VFA). The largest formed VFA amounts are those of the acetic (C_2), propionic (C_3), butyric (C_4) and valeric (C_5) acids. The amounts of the isobutyric (C_{4i}) and isovaleric (C_{5i}) acids are lower. Apart from them, some higher fatty acids, alcohols, carbon dioxide, methane, hydrogen etc. are produced. These end products are produced by a chain of reactions with primary roles of pyruvate and nicotinamide adenine dinucleotide (NAD). The amounts of fatty acids produced in the forestomachs of ruminants depends on several factors - diet type, feed ratios, preliminary technological processing (Sivkova, 2007). Other factors are the age of animals, the size and activity of rumen microbial populations (Enev, 1994), supplementation with nutritional preparations: bioproducts (Radev, 1999), exogenous enzyme preparations (Sivkova, 2007) and other factors. Some rumen content parameters, e.g. rumen pH, osmotic pressure, carbohydrate level, also have an effect on the total VFA amount (Sutton, 1981).

Degradation of proteins

Protein metabolism in forestomachs of ruminants consists in two primary interrelated processes. The first is dietary protein degradation by extracellular microbial enzymes. The second is microbial protein synthesis by the rumen microflora in the course of their replication in the rumen environment. In the forestomachs of ruminants, especially in the rumen, both dietary and endogenous proteins undergo complex microbiological and chemical changes. Proteins are degraded to peptides, amino acids and ammonia. VFA,

CO_2 etc. are also produced. The extent of feed protein degradation depends on the rate of hydrolysis, duration of the stay and feed mass evacuation rate to the more distal digestive compartments. The rumen content pH has a significant influence on proteolytic activity (Bach et al., 2005). According to Jagos (1984), when rumen fluid pH is about 6.5, ammonia concentration is optimum from the point of view of fermentation. In the rumen, feed proteins are attacked by bacteria and ciliates. The ciliates are able to ingest bacteria and due to the consequent proteolysis, ammonia concentrations in the rumen fluid increase. Ciliates are not capable to utilise efficiently the degraded protein. Only half of bacterial nitrogen is contained in infusorian cells. The rest of the nitrogen returns back into the rumen wall under the form of amino acids which are further hydrolysed to ammonia (Konechnii, 1985).

It is established that protein hydrolysis in the rumen occurs in two stages. Initially, feed proteins are degraded to form relatively large protein molecules of not less than 2 amino acids. Polypeptides are then degraded to amino acids and ultimately, to ammonia (Wallace et al., 1990).

The extent of proteolysis in the forestomachs depends on ration structure, protein solubility, the amount of available nitrogen and energy for microorganisms etc. One part of nitrogen needs of rumen microflora could be met by the liver-rumen urea recycling. Protein metabolism results from the metabolic activity of rumen microflora. Protein structure is a key factor for the activity of microbial proteases. The proteolytic activity in the rumen varies with the type of the fed diet (Siddons and Paradine, 1982; Hvelplund, 1991). Bacterial activity is determined by the type and solubility of proteins (Bach et al., 2005; Calsamiglia et al., 2010). The options and ways for modification of rumen protein metabolism, in the view of Wallace (1991), are not adequately understood.

Ammonia is the primary metabolite of proteins and non-protein nitrogen compounds, product of rumen microflora activity (Broderik et al., 1991). Every type of feed is characterised with specific level and rate of ammonia production. The ammonia concentration in the rumen content depends not only on dietary protein quality, but also on the extent of their solubility. The protein that is not fully degraded in forestomachs, could be absorbed in small intestine as free amino acids and peptides. They could be either used directly by the ruminant organism or undergoes deamination to urea in the liver. According to Bohnert et al. (2002a), when ammonia level in forestomachs is low, the nitrogen recycling through the gut increased while nitrogen elimination with urine is reduced.

Degradation of fats

Feeding traditional rations results in hydrolysis of over 85% of dietary fat (Doreau et al., 1997). According to Beam et al. (2000), a number of factors influence the extent of hydrolysis. Increased dietary fat in feeds reduces the hydrolysis extent. Low pH and the supplementation with ionophore antibiotics which suppress the activity and growth of bacteria and ciliates, also reduce the level of fat hydrolysis (Enev, 1994; Slavov, 2013). The breakdown of double bonds of unsaturated fatty acids and their saturation with hydrogen is another substantial change in dietary fat occurring in the forestomachs of ruminants. This process is known as fat biohydrogenation and occurs with the participation of ruminal microflora (Lundy et al., 2004). A prerequisite for biohydrogenation is the preliminary hydrolysis of fats. The biohydrogenation occurs only when fatty acids have an available free carboxylic group. That is why, biohydrogenation of esterified and non-esterified fatty acids is different (Harfoot and Hazlewood, 1997).

The extent of fat biohydrogenation in forestomachs depends on their form in feeds, diet formulation and other factors (Noble, 1984). The extent of hydrogenation of linoleic and linolenic acids in most rations ranged within 70-95% and up to 85-100% (Beam et al., 2000). When ruminants are fed high-concentrate rations, the high biohydrogenation level decline resulting in inhibition of lipolysis by the low pH (Van Soest and Demeyer, 1996b). Hydrogenation is also adversely influenced when rations contain unprotected lipids. Main intermediate compounds of biohydrogenation are trans-11 18:1 (vaccenic acid). It is produced by linoleic and linolenic acids and from cis-9, trans-11 conjugated linoleic acid formed during the biohydrogenation of linolenic acid. These intermediate metabolites are in a ratio of 3:1, but in the rumen, conjugated linoleic acid is only a transient metabolite (Lock and Bauman, 2003).

Options for modification of digestive processes in the forestomachs of ruminants

According to Demeyer et al. (1981) rumen metabolism is characterised with relatively stable fermentation model, characterised with the following main parameters:

- total VFA production;
- molar ratio of the different VFA,
- ratio between fermented organic matter and methane production;
- level of anaerobic catabolic processes;
- size of bacterial and ciliate populations;
- produced amount of microbial and ciliate protein.

The modification of rumen fermentation could be realised through maximisation or minimisation of biochemical events in forestomachs. The intervention is successful when the processes are discussed and managed using an integrative approach, as they are a constant and essential part of systemic metabolism in ruminants' body. Every intervention could lead to specific long-term consequences and a chain of interrelated effects (Harison and Mc Allan, 1988).

This aspect of digestion in the forestomachs of ruminants is flexible and suitable for management and optimisation through:

- diet structure
- biotechnological products and probiotics
- enzymatic preparations
- dietary protein
- dietary fat

Effect of diet structure

The composition and structure of the diet are essential for the enzymatic activity and development of microbial populations in rumen content. The complex interpopulational and inter-species interactions established among different bacterial genera, among bacteria, ciliates and fungi, are strongly influenced by the type and amount of dietary carbohydrates. Feeding rations composed by voluminous feeds and roughages, cellulolytic bacteria are predominant, while selenomonads, streptococci and lactobacilli dominated over the other microbial species in high-concentrated diets (Dehority and Orpin, 1988). The extent of utilisation of low-quality voluminous feeds and roughages depends on the cellulolytic activity of rumen bacteria (Bohnert et al., 2002b). When ruminants are fed a high-concentrate diet, the proportion of propionates formed in the rumen increases while the ciliate counts are reduced (Warner, 1966). The reduction in pH, observed in this type of feeding, has an adverse effect on the development and activity of ciliate colonies. It is believed that ciliates have a stronger positive effect on organic

matter digestion when a diet made mainly of roughages is supplemented with starch. Cereal feeds contain between 40 and 60% starch. According to Jouany et al. (2000) The general amylolytic activity in the rumen increases when added starch comprises more than 40% of dry matter.

The maximum productive performance of animals in modern intensive livestock production systems is achieved by feeding high-energy diets. Feeding diets with high cereal percentage results in pH reduction. A subsequent effect is alteration in microbial populations – decrease in cellulolytic bacterial counts and increase in amylolytic microbial counts. As a result, the level of rumen fermentation is high, posing a risk of acidosis (Jouany et al., 2000).

Effect of bioproducts and probiotics

By the end of the last century, production performance in ruminants was promoted with the so-called nutritional antibiotics. These comprise antibiotics and other chemical products of pharmaceutical and chemical industry. Ionophore antibiotics have a specific effect on rumen populations. They change the ratio of rumen microbial populations and indirectly, influence the systemic metabolism of ruminants. Applied at a subtherapeutic dose, they maintain the equilibrium in the digestive tract in favour of non-pathogenic microflora (Rowe et al., 1983). By the end of the 1990s, the use of ionophore antibiotics in EC countries was banned. It was established that their effect was cumulative. Antibiotics were accumulated in the body of animals and reach people through the consumer chain. It was also confirmed that they induced antimicrobial drug resistance. By reason of all these facts, alternatives were sought. It was demonstrated that efficient substitutes of ionophore antibiotics are biotechnological products, probiotics, prebiotics, synbiotics, preparations on the basis of live non-pathogenic microorganisms – yeasts, bacteria and fungi/ These microorganisms were shown to colonise successively the forestomachs and the other gut compartments of ruminants and through their metabolic activity, have a positive effect on their metabolism and health. In polygastric animals (experiments with cows, sheep and lambs), the most numerous microbial rumen populations comprised bacteria degrading complex carbohydrates - *Fibrobacter*, *Ruminococcus*, *Butyrivibrio* and *Bacteroides* along with primary groups as *Prevotella*, *Selenomonas*, *Streptococcus*, *Lactobacillus* and *Megasphaera*. Some anaerobic fungi and ciliates are also found in the rumen, as well as a large variety of methanogenic bacteria (Mackie et al., 2000).

According to Williams and Newbold (1990) products on the basis of *Saccharomyces cerevisiae* and *Aspergillus oryzae* could be successfully used for stimulation of rumen microflora activity without negative effect on the catabolic ruminal events. Ruminal microorganisms were confirmed to be beneficial for the meat and milk production of ruminants. The inclusion of yeasts as dietary supplement had a substantial effect on rumen fermentation as at moderate doses, they influenced positively the vitality of ruminal microflora. The yeasts are unicellular fungi. Their cellular wall is composed of 30% glucans, 30% mannan, 7% protein, 10% fat and other constituents. The cellular wall contains also different enzymes, the most important among which is invertase. As a source of energy, the yeasts utilise sugars, organic acids and higher alcohols. Nitrogen is obtained from both organic and inorganic sources. The yeast-mediated breakdown of sugars is oxidative. They have a 78.64% general digestibility and contain 13.119 MJ/kg metabolisable energy. The *in vitro* yeast protein digestibility attains 78%, and metabolisable energy is estimated at 13.15 MJ/kg,

(Krishna et al., 1991).

Cole et al. (1992) investigated the effect of dietary yeast supplementation in lambs and calves at a dose of 0.75% on transportation stress and the subsequent recovery. It was found out that the presence of yeast colonies in the digestive tract did not have a considerable effect on their health, but influenced the metabolism of nitrogen, iron and zinc. The inclusion of yeasts together with lactobacilli (*Lactobacillus celobiosus* CCM-400) at a dose of 0.5 g resulted in increased number of colonies of pectinolytic, amylolytic, xylanolytic and enteric bacteria in sheep forestomachs. At the same time, rumen pH increased without changes in VFA levels (Zonrcova et al., 1992). According to Dawson et al. (1990) the metabolic activity of *Saccharomyces cerevisiae* in the rumen of ruminants is low. They do not take directly part in the digestion, and the effect of their application is due to their ability to stimulate the development of the obligate rumen microflora.

One of the most popular preparations at a global scale is the probiotic Yea-Sacc 1026. The probiotic is a natural bioactive feed supplement and one of the leading products of the manufacturer. The inclusion of Yea-Sacc 1026 in the ration of dairy cows lead to increased rumen pH, reduction of lactic acid and ammonia in the rumen content. The level and direction of fermentation processes depend on the type of diet that is supplemented with Yea-Sacc 1026 and the proportion of concentrate in it. The addition of 10 g of the probiotic per capita daily has resulted in increased total amount of produced milk, increased amount of milk protein and fat (Gombos et al., 1995). The experiments of Williams (1989) investigated the effect of Yea-Sacc 1026, supplemented to the ration of calves and dairy cows. According to the results, the addition of Yea-Sacc 1026 increased milk protein and altered the rumen metabolism of supplemented animals. The protein in duodenal chyme and the total cellulolytic bacterial counts in the rumen increased. According to the author, the effects of the probiotic on carbohydrate and mineral metabolism and microbial protein production are still disputable.

Effect of enzyme preparations

The rationale of including synthetic enzymes in rations is to assist in the digestion of nutrients and their more efficient utilisation. The global animal feed production industry has responded to this trend with production of preparations with mono- or multienzymatic activity. Commercial products intended for use in ruminants are a complex of enzymes, most commonly with fibrolytic (cellulase, xylanase) and at a lesser extent, with proteolytic (protease, peptidase) primary activity. The goal for the supplementation of diets with these products is the more efficient breakdown of the main structural polysaccharides in roughages.

Multienzymatic preparations for ruminants' diets are produced with several microbial species with proven effect and metabolic activity. Most commonly, these are the 4 bacterial species *Bacillus subtilis*, *Lactobacillus acidophilus*, *Lactobacillus plantarum* and *Streptococcus faecium* and the fungi *Aspergillus oryzae*, *Trichoderma reesei* and *Saccharomyces cerevisiae* (Muirhead, 1996). The efficiency of enzymatic preparations varies depending on the growth substrate and environmental conditions (Lee et al., 2004). So far, a considerable number of experiments are reported in the world literature. They allow concluding that the mechanism of action of enzyme preparations was mostly dependent on the structure of rations and other physiological and technological factors (Hatfield, 1993; McAllister et al., 2000). The addition of enzymes to the rations entails structural changes which make it more susceptible to breakdown by the rumen microflora (Nsereko et al.,

1999). The mechanism of action of enzyme preparations is not yet clear including whether the effect of enzyme supplementation is realised before feed intake, in the rumen or there are post-ruminal effects.

The changes occurring after inclusion of exogenous enzymes to ruminant diets are not resulting from a single effect, but are rather a combination of a pre- and post-feeding mechanisms (Colombatto et al., 2003a; Sarnklong et al., 2010). McAllister et al. (2001) discussed that each of these pathways was important and that the enzyme-provoked degradation of feed ingredients before its consumption probably entailed respective consecutive events in the rumen and the gut. The pre-consumption effect of exogenous enzymes was probably due to release of soluble carbohydrates and complexes, or removal of structural barriers impeding the microbial breakdown of substrates in forestomachs. In the rumen, exogenous enzymes could act directly on nutrients or could stimulate indirectly the digestive activity through a synergic effect on available microflora. Supplemented enzymes remained active also in the more distal alimentary tract compartments (McAllister et al., 2001), where they could induce post-ruminal breakdown of fibres or indirect improvement of nutrients absorption through reduction of intestinal chyme viscosity. The way exogenous enzymes influence the rumen fermentation and consecutive effects of nutrients' metabolism in ruminants is extremely complex and unpredictable, that is why today, the research in this field is continuously evolving.

Effect of dietary protein

The extent of proteolytic activity of ruminal microflora varies within a wide range. It depends on numerous factors as the ration's composition, the size of proteolytic bacterial colonies, the complex interpopulation interactions with other species, the presence and dietary level of easily digestive protein etc. In most cases, bacterial proteolytic enzymes are intracellular. They are locked inside the cellular walls and are released after bacterial cell breakdown. An insignificant part of enzymes (10-30%) is released in the environment inhabited by microorganisms. Ivan et al. (2004) concluded that the inclusion of sunflower seed in high-concentrate diets reduced rumen fauna and as a result, dietary protein was preserved and feed digestion – was improved. In cows, Bargo et al. (2001) found out that the increased level of sunflower meal in the diet increased the amount of ruminally undegraded protein. Having studied the effect of different dietary nitrogen sources, Ahmed and Abdalla (2005) did not observe any changes in pH and VFA after feeding ruminants with rations containing sunflower meal.

The studies on the possibilities for influencing protein degradation in the rumen date back to the middle of the 20th century (Mills et al., 1944; Belasco, 1956). The authors have studied the effect of carbohydrates and fermentation type on the possibility of utilisation of non-protein nitrogen (urea) as a protein source in the rations of cattle. Later research has investigated the effect of non-protein nitrogen compounds after adding carbohydrates for improvement of ammonia utilisation through management of rumen microbial fermentation (Chamberlain et al., 1985; Rooke et al., 1987; Oh et al., 1999). In 1966, Virtanen established an important fact that has given a new focus in studies on rumen digestion – ruminants were shown to be able to convert non-protein nitrogen into milk protein.

As non-protein nitrogen compounds are relatively inexpensive compared to some feeds as meals, they are a suitable alternative to the natural plant protein. From non-protein nitrogen compounds, urea is most commonly used in diet formulation for ruminants. Urea

is easily dissolved in water and rapidly hydrolysed to ammonia and CO₂ in the rumen. The rapid release of ammonia from urea in the rumen, poor dietary nitrogen utilisation from high-urea rations and the potential risk from intoxication are the factors limiting the use of urea in ruminant nutrition (Bohnert et al., 2002b). Apart urea, other alternatives of dietary plant proteins are ammonium sulfate, ammonium phosphate, ammonium lactate, biuret etc. Biuret is not water-soluble and its degradation rate is slower than that of urea. The hydrolysis of biuret to ammonia is lower than that of urea (Currier et al., 2004c). In order to increase the amount of synthesised protein through guided energy protein rumen fermentation, the provision of supplemental energy for a specific ration is a prerequisite. The second purpose should be the provision of a definite amount of protein needed for the total amount of energy that is expected to be released in the rumen over 24 hours. Bacterial protein is produced under optimum conditions. For efficient utilisation of urea nitrogen, the diet should contain not less than 1 kg easily fermentable carbohydrates (e.g. starch) per 100 g urea. One of the important problems of using synthetic nitrogen compounds in ruminant nutrition is the creation of conditions for a relatively slow hydrolysis of proteins in forestomachs. The dynamic equilibrium between ammonia release and ammonia nitrogen utilisation of rumen bacteria should be maintained within the optimal range.

Effect of dietary fat

The effect of fat supplementation on fermentation is largely dependent on the type of added fat and its dietary level (Onetti et al., 2001). The inclusion of 4% fat in the rations of lactating cows according to the authors had no effect on rumen pH and total VFA amount. The acetic/propionic acid ratio, ammonia and protozoa counts in the rumen were significantly lower. In experiments with sheep, Cwiek and Borowiec (2005) established no positive effect from dietary fat supplementation on rumen fermentation and nutrients' digestibility. The use of sunflower seed as additive to high-concentrate rations for ruminants reduces rumen microflora and as a result, decreased the need for protein supplementation and increased digestibility of feeds (Ivan et al., 2004).

Salehpour and Amanlou (2007) have investigated the effect of whole cotton seed as a source of crude fibre in the diets of lactating cows. The supplement, according to the authors, had no effect on ruminal and faecal pH of experimental animals. The adverse effect on rumen fermentation of fat supplemented to ruminant diets was mainly attributed to the unsaturated oleic acid (C18:1) and partly on palmitic acid (16:0). The importance of the degree of unsaturation of fatty acids and the length of their carbon chain is confirmed by the studies of Jenkins (1987).

In vitro experiments performed by Zhang et al. (2008) have demonstrated the effect of C18 fatty acids – according to the authors, they exerted an insignificant effect on pH, Ammonia nitrogen amount and the total volatile fatty acids amount. The addition of these unsaturated fatty acids inhibited the growth of some cellulolytic bacteria and fungi, including *F. succinogenes* and *R. flavefaciens*. The effect of unsaturated C18 fatty acids comprised inhibition of methanogenesis, probably related to its direct effect on rumen microflora, which is associated to methane formation.

Conclusion

The digestion in the forestomachs of ruminants are an

important stage of their systemic metabolism. The dietary carbohydrate, protein and fat breakdown begins in the complex stomach to yield intermediate and end products as VFA, carbon dioxide, methane, amino acids, ammonia, nitrogen, glycerol, fatty acids, gases etc. which are included in metabolic pathways of nutrients and energy metabolism. The forestomachs environment is inhabited by multiple and various microbial colonies – ciliates, bacteria, fungi, yeasts etc. whose living activity is relevant for ruminant digestion in general. The presence of microflora is a classic example for symbiosis in nature. The extracellular enzymes produced by microorganisms start to degrade the complex carbohydrates from feeds, dietary protein and fat. At the same time, during their development and replication, the microflora produces microbial protein and fat of different composition and quality. The intricate interpopulation interactions both within and among the microbial species, their growth, replication, metabolic activity and end products of their life activity have been investigated for many decades. The knowledge of the complex physiological, biochemical and physical events occurring in the forestomachs of ruminants enables their guiding and modification with respect to the more efficient breakdown of nutrients and their optimum utilisation by ruminants. The advances in microbiology, physiology, animal nutrition and biotechnologies permit the future quest for new and modern means of digestion manipulation aimed at more balanced nutrition and improved performance of ruminants.

References

- Enev E**, 1994. Effect of rumen defaunation on digestion and productivity in sheep. Thesis for PhD, Sara Zagora (Bg).
- Petkov A**, 1982. Digestion and digestive behaviour in lambs. Thesis for PhD, Sara Zagora (Bg).
- Radev V**, 1999. Effect of yeast biotechnological products on rumen fermentation in sheep. Thesis for DSc, Sara Zagora (Bg).
- Radev V**, 2000. Effect of Yea-Sacc 1026 on the development and metabolic activity of rumen ciliates in sheep. *Animal Sciences (Sofia)*, XXXVII, pp. 34-38 (Bg).
- Rusev V, Tosev and Petkov A**, 1984. Physiology of digestion in ruminants. Zemizdat, Sofia (Bg).
- Rusev V, Lazarov V and Petkov A**, 1973. Physiological events in the complex stomach of ruminants and their productivity. *Proceedings of reports*, Sofia (Bg).
- Sivkova K**, 2007. Digestive events in ruminants depending on the composition and structure of their rations. Thesis for DSc, Sara Zagora (Bg).
- Slavov T**, 2013 Effect of dietary lipids supplementation on digestion, metabolism and productivity of small ruminants. Thesis for PhD, Sara Zagora (Bg).
- Konechnii Y**, 1985. Investigation of proteases produced by rumen bacteria. *Scientific Works of the Faculty of Zootechnics, HIZVM, Stara Zagora*, 297-300.
- Bach A, Calsamiglia S and Stern MD**, 2005. Nitrogen Metabolism in the rumen. *Journal of Dairy Science*, 88, 9-21.
- Bargo FD, Rearte H, Santini FJ and Muller LD**, 2001. Ruminal digestion by dairy cows grazing winter oats pasture supplemented with different levels and sources of protein. *Journal of Dairy Science*, 84, 2260-2272.
- Beam TM, Jenkins TC, Moate PJ, Kohn RA and Palmquist DL**, 2000. Effects of amount and source of fat on the rates of lipolysis and biohydrogenation of fatty acids in ruminal contents. *Journal of Dairy*

Science, 83, 2564-2573.

Belasco IJ, 1956. The role of carbohydrates in urea utilization cellulose digestion and fatty acid formation. *Journal Animal Science*, 15, 496-508.

Bohnert DW, Schauer CS, Falck SJ and DelCurto T, 2002a. Influence of rumen protein degradability and supplementation frequency on steers consuming low-quality forage: II. Ruminal fermentation characteristics. *Animal Science*, 80, 2978-2988.

Bohnert DW, Schauer CS and Del Curto T, 2002b. Influence of rumen protein degradability and supplementation frequency on performance and nitrogen use in ruminants consuming low-quality forage: Cow performance and efficiency of nitrogen use in wethers. *Journal Animal Science*, 80, 1629-1637.

Calsamiglia S, Ferret A, Reynolds CK, Kristensen NB and Vuuren van AM, 2010. Strategies for optimizing nitrogen use by ruminants. *The Animal Consortium*, 4, 1184-1196.

Chamberlain DG, Thomas PC, Wilson W, Newbold CJ and MacDonald JC, 1985. The effect of carbohydrate supplements on ruminal concentrations of ammonia in animal given diets of grass silage. *Journal of Agricultural Science*, 104, 31-340.

Cole NA, Purdu CW and Hutcheson DP, 1992. Influence of yeast culture on feeder calves and lambs. *Journal Animal Science*, 70, 1682-1690, 19 ref.

Colombatto D, Bhat MK and Morgavi DP, 2003a. Influence of fibrolytic enzymes on the hydrolysis and fermentation of pure cellulose and xylan by the mixed ruminal microorganisms in vitro. *Journal Animal Science*, 81, 1040-1050.

Currier TA, Bohnert DW, Falck SJ, Schauer CS and Barle SJ, 2004c. Daily and alternate day supplementation of urea or biuret to ruminants consuming low-quality forage: III. Effects on ruminal fermentation characteristics in steers. *Journal Animal Science*, 82, 1528-1535.

Cwik A and Borowiec F, 2005. Effect of dietary fat source on nutrients digestibility and rumen fermentation in sheep. *Polish Academy of Sciences*, 14, 239-242.

Dawson KA, Numan KE and Boling JA, 1990. Effect of microbial supplement containing yeast and microbaci on roughage-feed ruminal, microbial activities. *Journal Animal Science*, 68, 3392-3398.

Dehority BA and Orpin CG, 1988. Development of and natural fluctuations in rumen microbial population. In: *The Rumen Microbial Ecosystem* (Ed. P. N. Hobson). Elsevier Applied Science, London, UK, 196-245.

Demeyer DI, Van-nevel CJ and Aerts J, 1981. The role of protozoa in rumen fiber digestion. In K. H. Ferranti, O. Theander (Editors). *Proceeding OECD/COST Workshop on Improved utilization of lignocellulosic materials for animal feeds*, Brounsc Hweig, pp. 69-71.

Doreau M and Chilliard Y, 1997. Digestion and metabolism of dietary fat in farm animals. *British Journal of Nutrition*, 78, 15-35.

Harfoot GC and Hazlewood GP, 1997. Lipid metabolism in the rumen. In: P.N. Hobson, C.S. Stewart (Eds.). *The rumen microbial ecosystem*. Chapman & Hall Publishing house, London, 382-426.

Hatfield RD, 1993. Cell wall polysaccharide interactions and degradability. Phenolic constituents of plant cell walls and wall biodegradability. In: Jung, H.G., Buxton, D. R., Hatfield, R. D. and Ralph, J. (eds) *Forage Cell Wall Structure and Digestibility*. American Society of Agronomy, Crop Science Society of America, Soil Science Society of America, Madison, Wisconsin, pp. 285-313.

Hvelplund T, 1991. Volatile fatty acids and protein production in the rumen. In: "Rumen Microbial Metabolism and Ruminant Digestion". Ed. J. P. Jouany, INRA, Paris, 131-150.

Ivan M, Mir PS, Mir Z, Entz T, He ML and McAllister TA, 2004.

Effect of dietary sunflower seeds on rumen protozoa and growth of lambs. *British Journal of Nutrition*, 92, 303-310.

Jagos JP, 1984. Poruchu metabolism a system preventive diagnostiky scout, 1, vud. Brno.

Jenkins TC, 1987. Effect of fats and fatty acid combinations on ruminal fermentation in semi – continuous In Vitro cultures. *Journal Animal Science*, 64, 1526-1532.

Jouany JP, Michalet-Doreau B and Doreau M, 2000. Manipulation of the rumen ecosystem to support high performance beef cattle. *Asian-Australasian Journal of Animal Sciences*, 13, 96-114.

Krishna JA, Mandal B, Palival VK and Yadav KR, 1991. Fermented yeasts as a feed for adult animals. *Indian Veterinari Journal*, 68, 1126-1129.

Lee SS, Choi NJ, Ahn BH, Moon YH, Kim CH and Ha JK, 2004. In vitro stimulation of rumen microbial fermentation by a rumen anaerobic fungal culture. *Animal Feed Science & Technology*, 115, 215.

Lock AL and Bauman DE, 2003. Dairy products and milk fatty acids as functional food components. *Proceedings Cornell Nutrition Conference*.

Lundy FP, Block E, Bridges WC, Bertrand JA and Jenkins TC, 2004. Ruminal bio hydrogenation in Holstein cows fed soybean fatty acids as amides or calcium salts. *Journal of Dairy Science*, 87, 1038-1046.

Mackie RI, Aminov RI, White BA and McSweeney CS, 2000. Molecular ecology and diversity in gut microbial ecosystems. P.B. Cronjé (Ed.), *Ruminant Physiology: Digestion, Metabolism, Growth and Reproduction*, CAB International, London, UK, pp. 61-77

McAllan AB, Cockburn JE, Williams AP and Smith RH, 1988. The degradation of different protein supplements in the rumen of steers and the effects of these supplements of carbohydrate digestion. *British Journal of Nutrition*, 68, 669-682.

McAllister TA, Stanford K, Bae HD, Treacher RJ, Baah J, Shelford JA and Cheng KJ, 2000. Effect of a surfactant and exogenous enzymes on digestibility, growth performance and carcass traits of lambs. *Canadian Journal of Animal Science*, 80, 35-44.

Mills RC, Lardinous CC, Rupel IW and Hart EB, 1944. Utilization urea and growth of ruminal heifer calves with corn molasses or cane molasses as the only readily available carbohydrate in ration. *Journal of Dairy Science*, 27, 571-578.

Muirhead S, 1996. *Direct Fed Microbial, Enzyme and Forage Additive Compendium*, 3rd edn. The Miller Publishing Company, Minnetonka, Minnesota, pp.391.

Noble RC, 1984. *Proceedings of the 37th Nottingham Easter School*, pp. 185-200.

Nsereko VL, Wallace RJ, Rode LM, Beauchemin KA, McAllister TA, McKain N, Morgavi DP, Wang Y, Iwassa AD and Yang WZ, 1999. Effects of enzymic and non-enzymic components of a fungal enzyme preparation on digestion of grass silage by rumen microorganisms in vitro. *Journal of Dairy Science*, 82, 123.

Oh YG, Kim KH, Chong JJ and Chamberlain DG, 1999. The effect of the form nitrogen in the diet on ruminal fermentation in the yield of microbial protein in sheep consuming diet of grass silage supplementation with starch or sucrose. *Animal Feed Science & Technology*, 78, 227-237.

Onetti SG, Shaver RD, McGuire MA and Grummer RR, 2001. Effect of type and level of dietary fat on rumen fermentation and performance of dairy cows fed corn silage – based diets. *Journal of Dairy Science*, 84, 2751-2759.

Orpin CG and Letcher AJ, 1984. Effect of absence of ciliate

protozoa on rumen fluid volume from rate and bacterial population in sheep. *Animal Feed Science & Technology*, 10, 145-153.

Paul V, Guertler P and Wiedemann S, 2010. Degradation of Cry1Ab protein from genetically modified maize (MON810) in relation to total dietary feed proteins in dairy cow digestion. *Transgenic Research*, 19, 683-689.

Rooke JA, Lee NH and Armstrong DG, 1987. The effect of intraruminal infusion of urea, casein, glucose syrup and mixture of casein and glucose transport syrup on nitrogen digestion in the rumen of cattle receiving grass-silage diets. *British Journal of Nutrition*, 57, 89-98.

Rowe R, Davies A and Broome A, 1983. Effect of the novel ionophore ICI139603 on nitrogen digestion in sheep. In „Protein metabolism and nutrition“. BDS M. Arnal, R. Rion and D. Boonin. 2 INRA.

Salehpour M and Amanlou H, 2007. Evaluation of whole cottonseed as effective fiber source in early lactating Holstein cows. *Pakistan Journal of Nutrition*, 6, 294-298.

Santra A and Karim A, 2003. Rumen manipulation to improve animal productivity. *Journal Animal Science*, 16, 748-763.

Sarnklong C, Cone WJ, Pellikaan W and Hendriks HW, 2010. Utilization of Rice Straw and Different Treatments to Improve. *Asian-Australasian Journal of Animal Sciences*, 23, 5, 680 – 692. Its Feed Value for Ruminants: A Review

Sutton JD, 1981. Concentrate feeding and milk composition. In *resent Advances in Anim. Nutrition*. Ed. W. Haresing. Butterworths, London, pp. 35-48.

Wallace RJ, 1991. Rumen protolysis and its control. In: *Rumen Microbial Metabolism and Ruminant Digestion*. Ed. J.P. Jouany, INRA, Paris, 131-150.

Wallace RJ and McKain N, 1990. A comparison of methods for determining the concentration of extracellular peptides in rumen fluid of sheep. *Journal of Agricultural Science, Camb*, 114, 101-105.

Warner ACI, 1966. Periodic change in the concentrations of microorganisms in the rumen of sheep fed a limited ration every three hours. *Journal of General microbiology*, 45, 237-242.

Williams PEV, 1989. Understanding the biochemical mode of action of the yeast culture Yea-Sacc 1026. *Indian dairyman*, 41, 7p, 353-366.

Williams PEV and Newbold C, 1990. Rumen probiotic: The effect of novel microorganisms on rumen fermentation and ruminant productivity. *Butterworths, London*, pp. 211-227.

Yarlett NC, Hann AC, Leoyd D and Williams A, 1981. Hydrogenosomes in the rumen protozoon *Dasytricha ruminantium*. *Schuberg Biochemical Journal*, 200, 365-372.

Zhang CM, Guo YQ, Yuan ZP, Wu YM, Wang JK, Liu JX and Zhu WY, 2008. Effect of octadeca carbon fatty acids on microbial fermentation, methanogenesis and microbial flora in vitro. *Animal Feed Science and Technology*, 141, 5 – 6. populations. *Journal of Dairy Science*, 71, 2967-2975, 26 ref.

Zonrcova ZR and Nemcova V kmet, 1992. Effect of yeast cels and lactobacilli on rumen fermentation in sheep. *Xvocisna-Vyroba*, 37, 771-776, 19-6 ref.

Review

- Alternatives for optimisation of rumen fermentation in ruminants** 91
T. Slavov

Genetics and Breeding

- Characterization of a new winter malting barley cultivar Ahil** 98
B. Dyulgerova, D. Vulchev, T. Popova
- Evaluation of high yielding mutants of *Hordeum vulgare* cultivar Izgrev** 103
B. Dyulgerova, N. Dyulgerov

Nutrition and Physiology

- In vitro gas production of different feeds and feed ingredients at ruminants** 105
E. Videv, J. Krastanov, S. Laleva, T. Angelova, M. Oblakova, N. Oblakov, D. Yordanova, V. Karabashev
- Evaluation of chemical composition of raw and processed tropical sickle pod (*Senna obtusifolia*) seed meal** 110
Augustine C., Kwari I.D., Igwebuike J.U., Adamu S.B.
- Effect of urea-fortified all concentrate corncob diets on serum biochemical and hematological indices of West African dwarf goats** 114
U. M. Kolo, A. A. Adeloye, M. B. Yousuf

Production Systems

- Analysis of the technological dairy cows traffic "to and from" herringbone milking parlors** 119
K. Peychev, D. Georgiev, V. Dimova, V. Georgieva
- Effect of pre-sowing soil tillage for wheat on the crop structure and the yield components under the conditions of slightly leached chernozem soil in Dobrudzha region** 124
P. Yankov, M. Drumeva
- Study on the process of unloading grain harvesters at the end of the field** 129
G. Tihanov

Agriculture and Environment

- Modeling and simulation of fuzzy logic controller for optimization of the greenhouse microclimate management** 132
Didi Faouzi, N. Bibi-Triki, B. Draoui, A. Abene

Floristic diversity of 'Chinarite' protected area – Rodopi municipality, Bulgaria L. Dospatliev, M. Lacheva	140
Heavy metal pools in urban soils from city parks of Sofia, Bulgaria V. G. Kachova, I. D. Atanassova	144
Ecological characteristics of reclaimed areas in Pernik mines region, Bulgaria I. Kirilov, M. Banov	151
Reclamation of soil excavated from construction and mine searching areas in Turkey F. Apaydin	160
 Product Quality and Safety	
Concentration of sulfur-containing amino acids at turkey broilers during and after muscle dystrophy, fed with deficient feed supplemented with oxidised fat K. Stoyanchev	164
Exopolysaccharide influence's on acid gel formation K. Yoanidu, P. Boyanova, P. Panayotov	167
Carcass characteristics and technological properties of <i>Musculus Longissimus Lumborum</i> at lambs from the Bulgarian dairy synthetic population and its F1 crosses with meat breeds N. Ivanov, T. Angelova, S. Laleva S. Ribarski, D. Miteva, D. Yordanova, V. Karabashev, I. Penchev	171

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors
The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings: Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows. IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 9, Number 2
June 2017

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz