ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 9, Number 2
June 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate on author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc. and AGRIS (FAO).

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No DNP 05-21/20.12.2016, financed from Fund 'Scientific Research' grant Bulgarian scientific Periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 9, Number 2
June 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Ecological characteristics of reclaimed areas in Pernik mines region, Bulgaria

I. Kirilov*, M. Banov

Nikola Poushkarov Institute of Soil Science, Agrotechnologies and Plant Protection, Sofia, Bulgaria

(Manuscript received 5 August 2016; accepted for publication 29 April 2017)

Abstract. A referential literary analysis of the reclaimed area around coal mines near the town of Pernik, Bulgaria, is made in this paper. Affected areas (damaged, altered, etc.) around mining companies in the country are a serious environmental problem, including the territory of coal mines of Pernik which is a problem for the country. The area affected by the mining industry around Pernik occupies a territory of about 30 000 ha. As a result of the natural factors and the anthropogenic impact this region is characterized by a rather complicated soil situation. There are markedly expressed erosion and land-sliding processes on the slopes as well as conditions for surface over-humidification, swamping and salinization in some land lowering and on separate sections of the terraces of the rivers. The situation has acquired an exceptionally technogenic character after the development of the mines, the enormous pits and heaps and the construction of the industrial pads and equipment. The reclamation of environment in industrial areas is closely connected above all with the reclamation of disturbed areas and materials and can be performed in different ways according to their properties and economic effect. In connection with the extremely unfavorable conditions of a significant proportion of disturbed technogenic areas one of the basic methods is planting forests of production character or for recreational purpose. The inevitable degradation of the landscape can be corrected mainly by technical and biological reclamation of the disturbed terrains. A large part of the green belt of the town of Pernik was constructed on former heaps – mainly by afforestation. These activities have solved the problem of the separation of the town from the mining sites by a noise-preventing and dust-stopping embankment especially when the prevailing northwestern and northeastern winds are considered. Our main goal is to analyze and suggest the appropriate methods for biological reclamation of territories degraded by mining activities in Pernik coal mines region. That all proves the relevance and necessity of fundamental research and experimental work to identify opportunities for its targeted guidance and acceleration of the process. It has been established that almost all lands provided for agricultural uses are not cultivated and enter the fund of deserted lands. The basic reason for that fact is the lack of projects for biological reclamation and the insufficient justification of technical reclamation.

Keywords: Pernik mines region, coal mining, disturbed areas, biological reclamation, plants

Introduction

Modern intensive industrial development to a constantly growing degree reinforces the anthropogenic impact on the various components of the natural environment - terrain, soil, water, air, vegetation (Mitkov et al., 2013). One of the main problems in this regard is the total destruction of natural surface and vegetation by the influence of the mining industry, which leads to abrupt

disturbance of the ecological balance in these areas and the transformation of large territories into "industrial deserts" (Zheleva, 2010). This issue is important and topical for our country. Banov (2005) reported that the greatest disturbances and contamination of environment in Bulgaria are due to the mining industry (Figures 1 and 2).

In aggregate data so far over 30 000 ha have been disturbed with a clear trend to progressive increase of the phenomenon

Figure 1. An open pit for coal mining – Pernik Mines (Banov, 2005)

* e-mail: stingra@abv.bg

Figure 2. An open pit for coal mining – Pernik Mines (Banov, 2005)

(Zheleva, 2010). In this regard, reclamation and optimization of these technogenic landscapes is becoming more and more important with a certain complex environmental, social and economic impact (Kuzev, 2002). Depending on the composition and properties of the substrates and the specific economic, environmental and social impact reclamation can be performed in different ways - biological, hydro-technical, residential and industrial development (Regulation No. 26, 1996). In recent decades, in Mini Pernik only technical reclamation has been used (Hristova et al., 2011). The main goal of this paper is to analyze and suggest the appropriate methods for biological reclamation of degraded by mining activities territories in "Pernik Mines" PLC region.

Research sites in "Pernik Mines" PLC region

The location of "Pernik Mines" PLC areas was conditioned by the natural development of mining in the Pernik mining basin (Arsov, 1997). To a great extent, the town planning of Pernik is a function of the occurrence and liquidation of mining pits and sites from the infrastructure of Pernik Mines. They are related to the regulations of coal mining. In different periods of time there have been the following mines:

- Mines for underground extraction (old mines): "T. Nenkov", "Beli briag", "Tsaleva Krusha", "Popov Dol", ("Brigadir"), "Bela voda" ("Tolbuhin", "Bogdanov Dol"), "D. Blagoev-1", "D. Blagoev-2", "Sveta Anna" ("G. Dimitrov"), "Hristo Botev", "Gladno Pole", "Humni Dol", "Ivan Garvanov", "Kladnitsa".
- Mines for open pit mining: "Kutsian" ("7th of September"), "Republika", "Al. Milenov".
- Mining sites: one in "Bela Voda mine", one in "Sveta Anna mine" and one in "Teva mine".

The harmful impact of the underground mining activities is connected with rock movement (sinking and destruction) on the surface, destruction of buildings and facilities and/or swamping of the sunken terrain. Generally the urban development of the town is outside the dangerous zones. And where the neighborhoods were built prior to the exploitation of the deposits, expropriation and compensation with houses in safe areas is necessary. The sunken land has been reclaimed and handed over to other organizations for use in accordance with applicable regulations.

As far as open mines are concerned there is just one that is still functioning – "Republika" mine. "Kutsian" mine was closed in 1958 and "Al. Milenov" mine was closed in 1981. At the moment part of the former pit of "Kutsian" mine is used as sludge storage for "Republika Thermal Power Plant". The remaining part has been reclaimed and handed over to other organizations. The pit of the former "Al. Malinov" open mine is used for a railway dump of the "Republika" mine. Its bottom has turned into a lake as part of the water catchment area.

The impact of the open mining activities is expressed mainly in the following consequences:

- Disturbance of the landscape and the soil layer;
- Seismicity during the performance of explosion activities;
- Dusting of the air and noise from the heavy mechanization.

Conditions of the above sites

The scale of the transformations of the natural landscapes of the four sites from Pernik Mines – "Sveta Anna mine", "Teva mine", "Bela Voda mine" and "Republika mine" is extremely large as the share of the technogenic (anthropogenic) landscapes constantly increases (Arsov, 1997). The most essential changes in the existing landscape structure occur alongside the open mining

developments. In them there already are new, totally different from the ones before combinations of landscape components. The disturbance of the natural landscapes is basically performed at the expense of agricultural land and partially at the expense of the forest fund, the settlement fund and infrastructure facilities (Donov, 1976).

It is necessary to add that in the natural landscape, outside the territories for open mining, there are already deformities of the terrain and landslides as a result of the underground mining. The deteriorated general technogenic landscape structure of the open mines is further worsened by the compacted technical infrastructure – facilities for transportation of the coal and rock waste, road network, railway lines, power lines, etc. The total transformation of the natural landscapes has led to a completely destroyed plant component and thorough alteration of the soil-ecological conditions. Therefore, at first there must be development of a single concept of the structure of the territory and the landscape within the range of the open mining activities of Pernik Mines – a concept that will be based on the "Method of Landscape-Ecological Planning of Territories". A separate project for the planning of landscape, based on the already developed concept must be implemented for every open mine and its adjacent dump heaps. Planting must be a priority in the transformation of landscape by reclamation.

In conclusion, we consider that an environmental risk exists due to the contaminated mining water, which flows into the gullies and therefrom into the Struma River. There is a necessity for establishment of a monitoring system for control of the mining water and design of the adequate water treatment facilities must be developed based on the data obtained.

Total area of the territory of the site

The terrains which have been acquired and conceded in the more than 100-year-old history of Pernik Mines at the moment amount to 13226 decares, including 2808 decares occupied by buildings and facilities. Besides the terrains of Mini Pernik PLC quoted above, there are also other terrains as follows:

- For the Central Management Department – 338 decares;
- For the area of the former Kutsian mine (7th September) – 100 decares;
- Rudartsi village – 182 decares for a recreation base.

Geographical location, contemporary relief and soil differences

The Municipality of Pernik is located in the Balkan sub-area of the Mediterranean soil area, the subtropical xerophyte-forest European soil sector of Luvisols, Leptosols and Vertisols. In the periphery of the valley large proluvial tails have been formed. According to the scheme of soil-geographic zoning of Bulgaria, the land of Pernik falls into the Kraishte-Gornostrumski soil region from the Southern Bulgarian xerothermal soil zone. The town of Pernik and its adjacent area falls into the Znepolski and Vitosha region according to the floristic zoning (Zheleva, 2002). As a result of the various and specific geo-morphological and hydro-climatic conditions, relief and plants on this comparatively small territory soil differences have formed, which approximately have the following distribution in the area:

- Vertisols are found on lower and better levelled to slightly hilly and rounded parts of the terrain – mostly in the area of Bela voda mine, Sveta Anna mine and partially Teva mine;
- Luvisols are found on higher and at the same time more distinctly outlined relief forms – in the northern part of the Bela voda mine and Teva mine and mainly in the area of Republika mine;
- Fluvisols have been formed on the lowering and terraces of the rivers Struma, Buchinska, Moshinska, Diva reka, Teva and

Tsarnezhka.

On the whole land and forest fund of the town of Pernik industrial embankments from the opening of the above quoted mines are scattered. The lands around them are highly anthropogenic. Parts of the embankments have been biologically reclaimed and have been included into the forest fund. A smaller part of the land has been reclaimed for agricultural use and still other part has entered the Habitation fund.

A large part of the lands in the area of Pernik are occupied by anthropogenic soils and industrial embankments. The survey of the soil diversity in the area shows that very little of the land is characterized by natural and non-degraded soils. The soils in the area of the open pit mining have been degraded to a great extent. Therefore, their recultivation after the activities of reclamation is a long process (Arsov, 1997). A plan for the selection of suitable plant types for a period of 5 years is recommended for the part of biological reclamation of each and every reclamation project. All soils around the town of Pernik are environmentally polluted to a different degree as a result of the industrial activities emitting harmful substances into the atmosphere, which is favored by the continuous thermal inversions typical for the town. Therefore, a large group of amelioration activities are needed, aiming at protection and increase of soil fertility (Penkov, 1985).

In the area around Pernik there are contaminated soils as well as soils degraded by technogenic sites. These are mainly the foundation pits of the open mining, the heaps of both open and underground mining, the mine sites, the roads, railway tracks, sewage systems, power lines, etc. As a result of these multilateral industrial activities (construction, pits, heaps, etc.) the soils in these territories have been almost irrevocably destroyed. The heaps described in this paper (7th September mine and Karamanitsa mine) are part of such degraded terrains – (Zheleva, 2002). As far as their morphological structure is concerned, the heaps have no sequence in the overlay of various materials and layers (Donov, 1978). Because of unselective depositing heaps have a heterogeneous composition with prevailing geological materials, typical of the excavated over-coal and inter-coal layers. These are mainly marls (at a different stage of weathering), a lot more slightly altered, limestone, sandstone, flint kernels, etc. The participation of clayey shale is essential and it is mixed with coal to a different degree (from 0 to over 30 %), thus forming the so called coaly shale as well as cinder. At separate places of the heaps, where the coal is on the surface or close to it, there might be oxidation and self-ignition, caused by the influence of the atmospheric conditions. In such cases burning hearths are formed which emit sulfur gases. The soil erosion in the heaps must also be prevented.

The basic contamination of the soils by heavy metals is caused mainly by Stomana PCL, the electric furnaces of Blagoi Popov Factory and Republika Thermal Power Plant. In the piled rock masses of Mini Pernik no heavy metals above the admissible norms have been found. Besides, the opportunities for reclamation of the degraded lands are few because there is no humus soil covering. For the greater part of them these lands can be reclaimed for the needs of forestry, for landscape architecture or construction – as far as the stability of the terrains allow that (Gencheva, 1984).

The deteriorated ecological conditions are a prerequisite for better landscape architecture of the town of Pernik and for restoration of the healthy and sustainable urban eco-system by using the degraded terrains for planting. At the same time part of the degraded terrains can be used for other urban planning or industrial needs without disturbing the natural soils of the suburban eco-system.

Climatic and meteorological data of the studied area

The territory under survey occupies part of the Tran-Pernik sub-area of the Kraishte-Vitosha area. It has rounded hilly surface. The altitude is from 600 to 900 m, with an average altitude of 730 m. The highest peaks in the location are Kitka and Usoitsa – 854 m, and the lowest elevation is at Batanovtsi. Biala Voda Mine is located to the west of Pernik and exploits the raw materials from the northwestern slopes of Golo Bardo. According to the climatic zoning the area falls into the moderate continental climatic sub-area – the high fields of Western Middle Bulgaria (Sabev and Stanev, 1963). The terrain has a hilly relief and is situated in a valley of a comparatively high altitude. Winter is too cold here with minimal degrees from -2 to -3.5 °C in January on average. The number of days with minimal temperature of < 10°C for the three winter months is between 15 and 20. About 50-55 days of the winter the average temperature around the clock is below 0°C. A characteristic feature of the area is the large frequency of fog – about 15-20 days for the three winter months. Winter precipitation is the lowest – about 80-120 mm, mainly snow. Around 80 days of the year are characterized by snow cover. Spring is comparatively cool. Spring precipitation is higher than that of winter - 147 - 175 mm on average. Summer is also rather cool.

The yearly precipitation has continental character, the maximum being in June and the minimum – in February. The morphographic conditions influence the regime of winds, which are light with frequent calms. The climatic elements: precipitation, temperature, evaporation and winds have an impact on the value of the run-off and regime of the surface and underground water. Having in mind all these characteristics, it is important to clarify the most essential meteorological elements. The data has been taken from the meteorological station of Pernik, which has an altitude of 700 m.

Plants

Data of previous research (Vasilev and Andreev, 1983) and present observation of the floristic composition of the grass vegetation having naturally occupied the terrain – vegetation of different origin and age, shows that three stages can be distinguished - weed - ruderal, wheat - ephemeral and rhubarb - leguminous. Sharp boundaries among the different stages are not available – neither in time, nor in space.

On the heap of underground mining Maksim Taban the total projective cover of the surface reaches 80-90 %, which already corresponds to a comparatively advanced stage of restoration of the grass vegetation – in 1978 it was 70-80%. The kinds of grass that are found there are *Anthemis coluta* L., *Bromus commutatus*, *Avena fatua*, *Senecio vulgaris* L., *Poa anna* L., *Pestuca pratensis*, *Amaranthus albus* L., *Daucus carota* L., etc. (Zheleva, 2002). In different parts of the heap of Maksim taban, which has a mound-shaped form with rather steep slopes there are the three types of plant vegetation. The prevailing type is the wheat – ephemeral one but in certain parts there are also the wheat-ruderal and the rhubarb-leguminous stages.

The aims of forest reclamation in the region are putting a stop to the erosion of heaps and the pollution of air by dust and gases from the self-igniting remains of coal. In the beginning, until 1990 the forestry activities were performed without any leveling of the terrains. At present the heaps are transformed into terrains which are more convenient to work on and favor the growing of the saplings. It must be noted down that in the aspect of afforestation or biological reclamation the main role is played by forestry. Pernik Mines provide the terrains in an unsuitable or difficult for assimilation state. The first attempts for landscaping of the heaps of Pernik mines began in the

1920s in the heaps of Kalkas Mine. They were mostly amateur and had mainly sanitation-hygienic purposes. However, in forestry afforestation is made not only for ecological but also for economic appropriateness. Therefore, an evaluation of tree production must be made assessing the sustainability of the tree types as well as their wood-producing opportunities.

Prokopiev (1962) put on a scientific basis the attempts for landscaping of the heaps by afforestation of part of the heap of Temelko Nenkov Mine. He used types of trees which in his opinion were characterized by higher smoke resistance. *Qercus rubra* L., *Fraxinus americana* L., *Robinia pseudoacacia* L., *Eleagnus angustifolia* L., *Laburnum vulgare* Berchtold et Presl., *Amorpha fruticosa* L., *Cotinus coggygia* Scop., *Lonicera tatarica* L., *Tamarix tetrandra* Pall and *Evonymus europaea* L. Later *Tilia cordata* Mill., *Tilia argentea* Desf., *Fraxinus excelsior* L., *Aesculus hippocastanum* L., *Populus pyramidalis* Roz., *Betula alba* L., *Cydonia vulgaris* Pers., *Pirus communis* L., *Prunus divaricata* Ledeb., *Acer pseudoplatanus* L., *Acer negundo* L., *Acer dasycarpum* Ehrh., *Acer platanoides* L., *Pinus nigra* Arn., *Picea excelsa* Wall., *Biota orientalis* Endl., *Juniperus virginiana* L. and other types of bushes were imported. In fact drought-resistant and heat-resistant kinds with powerful root systems must basically be used. The observation of the author of the resistance of the above mentioned types showed that the most resistant ones under the conditions of drought and heat are *Platanus*, *Qercus rubra* L., *Fraxinus excelsior* L., *Quercus robur* and *Accacia*.

The most sensitive kinds are *Acer negundo*, *Tilia cordata* Mill., *Aesculus hippocastanum* L., *Betula alba* L., *Acer dasycarpum* Ehrh., *Fraxinus americana* L. *Tilia argentea* Desf. is in medium position. *Pinus nigra* Arn., *Juniperus virginiana* L. and *Biota orientalis* Endl. are in good condition, but *Picea abies* is in very bad condition and is not to be recommended. The nature of afforestation must be appropriate to the purpose of landscaping. The use of favourable special functions of trees is most important and the logging function is secondary in importance. Out of all tree species *Pinus nigra* Arn. has the largest participation by area – 60.4%, followed by *Robinia pseudoacacia* – 7.1%. *Betula alba* L. and *Pinus sylvestris* are almost equal by area – 6%, followed by *Populus pyramidalis* Roz., *Quercus rubra* and *Fraxinus excelsior*.

Because of the heavy disturbance of the relief, the geological base and soils, the reclamation of the degraded terrains with valuable vegetation will be difficult and the types of trees that can be used in reclamation must be the same as the above mentioned. The selection of plant species is compliant with the environmental conditions of altitude, exposure, precipitation, acidity/alkalinity and partial salinization of the substrates with a lot of rock content. Forestry reclamation of the area aims at preventing erosion of the heaps and pollution of the air by dust and gases from the self-igniting remains of coal.

A clear and categorical evaluation of the impact of vegetation cannot be made, because it is created artificially with the aim of reducing the negative impact of the open pit mining and more generally of all mining activities on the ecological circumstances in the region. The positive role of vegetation in fortifying the open heaps and the reducing dust pollution are doubtless. Because of the immediate proximity of the heaps to the town of Pernik the importance of vegetation is mostly for landscaping and creating a better urban environment.

A prognosis of the tree supply by separate species for a 20-years period of time is made on the basis of taxation data (Zheleva, 2002). The results are displayed in Table 1.

Table 1. Prognosis of tree growth and tree supply by species around the year 2016

№	Tree Species	Sum of tree growth around 2016	Supply around the year 2016	
		in m ³	m ³	%
1	<i>Pinus sylvestris</i>	2 495	5 015	12.2
2	<i>Pinus nigra</i> Arn.,	14 570	24 275	59.1
3	<i>Qercus rubra</i> L.	350	525	1.3
4	<i>Quercus cerris</i>	130	270	0.7
5	<i>Ulmus</i>	45	75	0.2
6	<i>Betula</i>	1 820	2 730	6.6
7	<i>Acacia</i>	1 600	2 830	6.9
8	<i>Aesculus</i>	10	25	0.1
9	<i>Acer</i>	80	205	0.5
10	<i>Tilia platyphyllos</i>	25	40	0.1
11	<i>Fraxinus americana</i>	85	130	0.3
12	<i>Fraxinus excelsior</i>	860	1 710	4.2
13	<i>Populus Bachelieri</i>	1 970	3 005	7.3
14	I - 214	10	15	-
15	<i>Populus regenerata</i>	150	225	0.5
Total amount		24 200	41 075	100.0

Landscape of the disturbed territories, measures for its restoration

The greatest risk for disturbance of environment in the area is the total alteration of landscape. As a result, the conditions for development of the flora and fauna in the region are completely disturbed, too. The hydro-geological and hydrological conditions have also been disturbed. The destroyed natural landscapes have been replaced by anthropogenic ones, which are aesthetically unacceptable being in proximity to the urban area. A profound development of a concept for landscaping of the whole region is necessary, considering the urban building plan for the suburbs of the town. As a result of this concept it is necessary to develop particular project solutions for landscape patterns (Zheleva, 2002)

The following prognosis principles for the impact and structure of landscape can be formulated:

1. Conservation and optimization of the existing natural landscapes, which are located outside the limits of the mines and the heaps in the outlined boundaries of the mine properties.
2. From the point of view of landscape in relation with ecology and aesthetics it is appropriate to fully preserve the agrarian, forest, water and waterside landscapes.
3. The already existing technogenic landscapes and those that will appear after the future open pit mining in Mini Pernik must be restored by engineering-technical and biological reclamation. The inner diversity of the cultural landscape must correspond to the conditions of its sustainability to the ecological and aesthetic requirements.
4. The reclamation of the territories in the mines where the mining activity has ceased must be accelerated. Deserted places such as heaps, technical equipment and infrastructure elements must not remain in the newly created cultural landscapes.
5. In the restructuring of landscape by reclamation priority must be given to planting, having in mind its special functions in the landscape and also the creation of pasture landscapes with good-

quality grass composition.

6. It is necessary to develop a single concept for the design of the territory and the landscape within the range of Pernik Mines.

Results and recommendations based on research done over the years

The inevitable degradation of the landscape can be corrected mainly by technical and biological reclamation of the disturbed terrains. A large part of the green belt of the town of Pernik was constructed on former heaps – mainly by afforestation with *Accacia*, *Betula Populus*, bushes, *Abies* and *Pinus* (Prokopiev, 1962). These activities have solved the problem of the separation of the town from the mining sites by a noise-preventing and dust-stopping embankment especially when the prevailing northwestern and northeastern winds are considered.

The issues of the harmful seismic effect are solved mainly by developing specialized projects for drilling-explosive activities of such parameters that can neutralize the seismic effect. The mining waters in underground and open pit mining are previously deposited and clarified in artificially formed water catchments. After that they are pumped out and gathered in natural water receivers such as gullies and rivers which flow into the Struma River.

It has been established that almost all lands provided for agricultural use are not cultivated and enter the fund of deserted lands. The basic reason for that fact is the lack of projects for biological reclamation and insufficient justification of technical reclamation (Zheleva, 1979). The submitted projects for technical reclamation and the condition of the already reclaimed terrains give the impression of inefficient use or not having been used at all. The results and recommendations based on the implemented scientific research and the carried out vegetation and field experiments require growing of agricultural crops directly on the heap substrates without covering them with humus. The conclusions by Lichev et al. (1973), as a result of their research are that marls, which have stayed on the surface for 2-3 years, are biologically appropriate for growing basic field and fodder crops. For instance, the yields of alfalfa grown on such marls without fertilizing and watering are the same as the yields of alfalfa grown under the same conditions on Vertisols. The results are also good with *Onobrychis sativa*, ryegrass, wheat and maize but only in certain combinations of mineral fertilization.

As a result of the natural factors and the anthropogenic impact the region of Pernik is characterized by a rather complicated soil situation. There are markedly expressed erosion and land-sliding processes on the slopes. There are conditions for surface over-humidification, swamping and salinization in some land lowerings and on separate sections of the terraces of the rivers. The situation has acquired an exceptionally technogenic character after the development of the mines, the enormous pits and heaps and the construction of the industrial pads and equipment. The most important is the fact that the natural hills, slopes, flat terrains, vales and low valleys have been wiped out and replaced by newly constructed and absolutely different chaotically dispersed negative (pits) and positive (heaps) relief forms.

The geological base in these terrains is also highly degraded (Lichev and Treykiashki, 1973). The geological layers at depth of 80 to 100 m have been excavated and at the same time they have been heaped on the surface without consideration of their natural stratigraphic order.

In the general context of the above stated conclusions the influence of Mini Pernik is displayed by alterations of the relief, dust and partial aerosol pollution of the atmosphere, destruction of

vegetation, contamination and destruction of the soil cover and other components of the environment.

Quality characteristics of the reclaimed land from the region of Pernik Mines - Survey of research

The first research and practical activity in the area of biological reclamation of the technogenic landscapes in the developed industrial countries began at the end of the 19th century with the creation of forest areas (Gencheva and Zheleva, 1997). In Bulgaria the first surveys and afforestation were made in the 1950s in the region of Pernik (Gushevilo, 1977). Later two directions of biological reclamation were differentiated – forestry biological and agro-biological which have clearly defined specifics in both theoretical and experimental research and in practical activities. The theoretical concept is aimed at clarifying two main problems - the specifics of technogenic neo-relief and the opportunities for sustainable and ecologically balanced biocenoses (ecosystems), which defines the two main stages of reclamation:

1. Technical, which is connected with stabilization and correction of the neo-relief and its integration into the adjacent landscape.

2. Biological - characterization and assessment of the properties and fertility of substrates forming technogenic landscapes, determination of the type of vegetation, the application of appropriate technologies and amelioration of substrates (Ivanov et al., 2007).

At appropriate stages particular methods in the field of soil science, horticulture, forestry, agrochemistry, dendrometry, etc. were applied. The basic requirements for technical reclamation include preparation of disturbed areas for commercial use and its environmental inclusion in the adjacent landscape - creating rational forms of relief with a favorable structure of the heaps, planning their surface, leveling and normalization of the slopes (greater than 18-24 degrees), removal of the effects of subsidence, performing ameliorative activities; creating a road network; regulation of the hydrological regime (creation of drainage ditches, runoff regulation, moving riverbeds, etc.); creating the necessary engineering facilities (Miltchev, 2008). The link is the biological approach for restoration of the disturbed areas and to some extent the methodical approach in fundamental research (Gencheva et al., 1986).

The main problems of forestry biological reclamation are related to clarifying the complex relationships in the system "climatic factors - edaphic factors - biocenotic factors". The clarification of these interactions is the basis for sustainable forest ecosystems (Gencheva et al., 1995). Initially it is necessary to study the natural environmental conditions, the composition and properties of the geological materials on which to experiment and determine the type composition, options for melioration of substrates, methods and means of cultivation, afforestation and crop growing (Petrova, 1994). Subsequently the established sample sites and massive afforestation enable clarification of the reverse process - the effects of different dendrocenoses on the direction, the nature and intensity of soil formation processes under particular climatic conditions (Gencheva et al., 1986).

Until now studies both at home and abroad, are mostly focused on issues related to the opportunities of using different geological substrates as plant protection for forest trees most commonly as uniform plantations or in a group mosaic mixture as well as on the behaviour of the experimental tree types (Lichev et al., 1973). The properties of bulk materials in the major coal-producing areas in the country have been characterized. Research shows certain

differentiation on different sites - disturbed by mining of coal, ores of nonferrous and rare metals, construction and inert materials, etc. (Gencheva et al., 1982). In the first years after the construction and beginning of reclamation activities heaps were loose and porous, but after 8-10 years, their density increased. Therefore, plants which do not tolerate heavy soils deteriorate their growth (for example *P. Nigra* and *P. silvestris*).

Heaps from underground coal mining contain amounts of coal particles that change the properties of bulk materials due to the high temperature processes that develop in them - continuous burning and smoldering (Haigh and Gentcheva, 2002). Heaps which were formed as a tailing pond from processing and briquette factories in energy complexes have very bad forestry properties. Substrates in them are very toxic to the plants, due to the large amounts of salts and in some cases acidic compounds therein. All substrates have a low amount of key nutrients such as nitrogen and phosphorus but potassium has usually satisfactory or very good amounts in stock (Zheleva et al., 1995). To conduct reclamation on these heaps, especially in the lower altitudes the limiting factor turns out to be water - both underground water and air moisture is insufficient during the summer months. In the last 8-10 years drought has started from the beginning of April and has ended in late September. The plant type on these heaps complies with the forest zoning of Bulgaria, the climatic and edaphic factors of the area. The research findings require the use of xerothermic types typical of the forestry zone of the area, especially at the beginning of reclamation and also types which develop a strong root system early after afforestation or have extensive adaptive properties and ecological plasticity (Petrova, 1991). In this direction there are fewer studies and developments on land disturbed by mining and processing of ores of nonferrous metals and radioactive elements where problems of reclamation are connected not only with stone coverage, mechanical and granulometric composition, density and water regime but they are also in connection with the complex chemical composition and properties, the acidic regime, sorption properties of the substrates, the presence of elements and compounds that induce high acidity or alkalinity, etc. (Ivanov, 2007).

Key features of Waste Coal in Pernik coal province and the analysis of the geological map - Pernik map sheet (Figure 3). This map shows that the surveyed sites in the Pernik basin fall in areas

Figure 3. Geological map of Bulgaria with Pernik region (in the marked circle on the map) (Committee on geology, Enterprise geophysical surveys and geological mapping, 1991)

built of rocks and are distinguished by a number of specific physical properties such as water absorption and degradation, poor ductility and primarily light colored (white, tan, gray, greenish-gray, reddish (Kostadinova, 2016).

The study of the negative effects and the opportunities to overcome them has posed a major problem in research and practical activities in this direction. It is in connection with the standing problem of selecting the appropriate plant types to consolidate the steep heap slopes and creating economically valuable and sustainable ecosystems in these disturbed areas (Haigh, 2007).

Although forestry biological reclamation is a new direction in ecological restoration research attempts have been made to create technology for reclamation of areas disturbed by different mining activities. The emphasis is on ameliorative measures (fertilization, liming, sideration and cultivation care), schemes and density of afforestation, etc. There are certain results regarding the influence of mineral fertilization on forestry biological reclamation, but at this time of recession in our economy there is danger of underestimating the need of this activity and research on these issues. Technologically there is less research related to the opportunities of using mechanization in reclamation works (Zheleva, 2010). Changes in the nature of economy, inflation and difficult financing conditions of risky research at this stage direct our attention to methods with low investment. Under conditions of sloping terrains mechanization is specific, so research should be aimed at mechanizing operations of biological stabilization of reclaimed land, exploration of the opportunities of hand-held power tools in a spoil heap, exploration of the possibilities for remote introduction of chemical solutions hydraulically to stabilize the steeply sloping surfaces and mineral fertilizers, grass mixtures, etc., exploration of the possibilities for using existing or constructing new machines for slope correction, processing terraces on sloping land, fertilizing, afforestation, etc.

Research for clarifying the reverse process - the effects of different dendrocenoses on the direction, nature and intensity of soil formation, has begun only in the last 10 - 15 years (Filcheva et al., 2000). This is logical, since the processes of weathering and soil formation are very conservative and require longer period of influence of technogenic phytocenoses to establish some changes in geological substrates. A very important issue in methodical respect in these studies is to identify the most informative indicators affecting the degree and direction of soil formation processes - the amount and properties of diluvium, changes in acidity, organic matter, qualitative composition of humus, destruction processes and biochemical transformation of organic residues, etc. (Ivanov et al., 2007)

Specific properties of bulk materials due to reclamation activities in the region of Pernik town

The current studies have so far been focused on:

- determining the key diagnostic indicators for qualitative assessment of geological substrates in connection with forestry biological reclamation;
- optimizing the type composition of biocenoses on reclamation areas in order to create sustainable ecosystems;
- development of technological instructions corresponding to the specifics of the areas for reclamation, with special attention paid to melioration activities and mechanization;
- expanding research on the specific nature of the processes of weathering and soil formation with a view to their targeted management.

Underlying soil survey, biometric research, the assessment of soil fertility and carried out field trials give rise to certain conclusions

about the reclamation of selected areas - heaps of open pit mining of coal in the city of Pernik (heap of Al. Milenov mine) (Ivanov, 2007). Heaps of coal mining, according to the way of mining are from galleries (underground) and open pits. Tailings ponds or tailing heaps received from the enrichment of coal emerge as a third category. Open pit mining turns out to be more economical so heaps of this type of mining are preferred as object of research (Shishkov, 1985). Soil forming materials have been irregularly layered in profile, which gives respective impact on the physical and chemical properties at depth. Selected soil sections in established crops have visibly marked differences in morphological terms. Soil materials in some layers have quite a lot of skeleton, mainly coaly shale, marls and slates in Pernik. The volume weight of samples does not differentiate a lot from that of natural soil. The results show that the total porosity is relatively good especially in the surface layers and newer heaps. In depth, however, it decreases as a result of subsidence.

Water properties of the tested materials are expressed by the basic hydrological constants (in weight percentage of water to that of the total sample). The results show that the materials of heaps in Pernik have comparatively high percentage of capacity of active moisture (CAM) (Kostova, 2013). In general, materials have relatively favorable water properties which do not differ substantially from the water properties of natural soils. Chemical analyses show neutral or alkaline soil pH, low contents of humus and low availability of basic nutrients (which does not apply to Potassium) (Hristov, 2015). The data exposed so far give us grounds to determine that forest plant properties of the tested materials are not good.

The forest plant properties of soils can be improved by performing a complex of activities causing a rise of the indicators of soil fertility (Bogdanov et al., 1994). Terrain experiments conducted on heaps were based on the results of soil research. The aim was to determine the most appropriate tree types on heaps of open coal mining, the most appropriate agricultural machines for their growth and mainly the most appropriate rates of fertilization and ratio of introduced fertilizers. Mineral fertilization was performed according to specific schemes appropriate for the technical possibilities of the terrain (mainly size and compactness of areas). Field trials were conducted with pure and mixed cultures of red oak, silver lime, plain and American ash tree, silver birch, black pine and sycamore. The results show that the applied fertilizers caused the greatest reaction by acacia, plain ash and birch, in which the absolute values of all biometric parameters were the greatest at the higher rates of fertilization. Sycamore and lime occupied average position. Generally they also had a positive reaction to mineral fertilization but the change of indicators of fertilized trees was slighter. As far as the studied parameters are concerned, the weight and volume of the roots change to the greatest extent in the control figures, followed by the height of seedlings and the diameter of the root collar. This is favorable because the bigger root system creates a better opportunity for feeding and growth of tree types. Full fertilization has a significant positive effect on the black pine and lime, nitrogen - on ash (more precisely ash suffers from lack of nitrogen) and phosphorus and potassium fertilizers have a favorable impact on the black alder. Mineral fertilization also affects in one way or another poplar planted as cuttings in the terrain. Poplars prove a significant effect of mineral fertilization.

Initial results of mineral fertilization show that it can and must be a serious ameliorative measure for acceleration of the soil forming process, improvement of soil fertility and better growth and sustainability of planted seedlings. Mineral fertilizers should be applied by strict differentiation into various plant types and in order to

expand the range of vegetation used in biological reclamation we must always consider the climatic conditions, the fertility of soil materials and the biology of tree and bush types (Zheleva and Kostova, 1986).

In the geological structure of the coal bearing stratum in Pernik coal mining basin there are mostly marls, a large amount of clay, shale and sandstone. In our climatic conditions these geological materials weather relatively fast and homogeneous loose mass appears on the surface after 2-4 years. In connection with the technology of filling the relief of the heaps is either conical with high and steep slopes or ctenoid. For the same reasons and because of the differences in composition and thickness of different under-coal layers, the profile of industrial heaps is built by different in composition and power layers in the rotation of which there is no regularity (Gushevnikov and Pascalev, 1977). The skeletal fraction has also a different participation and it also depends on the geological products and weathering materials derived from them. Heaps of Pernik sometimes contain over 50% of skeleton, which is due to the larger fractions of shale and marl. The heap material from Pernik contains a larger quantity of sand fractions. For these reasons reclaimed soils on these heaps will be formed more quickly and will be characterized by a light mechanical composition. The density of the soils in the area forms a typical porosity ranging between very good and unsatisfactory. However, it constantly changes because of the subsidence of heaps.

Properties and fertility of heap materials are determined by specific studies of the morphological features, the mechanical composition, the physical properties, water and temperature regime, chemical composition and properties and microbial activity of substrates. In a wetted condition they are cut and spread, being difficult to process, and when they are dry large crevices are easily formed. In water logging soil particles are not water proof, they are easily blurred and show a lack of organic matter (Ivanov, 2007).

The physical properties are reflected on water properties of heaps in the above mentioned areas (Kostova, 2013). Data show that productive water supplies, especially at shallow depth are too low in some periods during the vegetation season. Because of its clay composition the heap materials retain more moisture, but much of it is dead stock. For these reasons and because of lack of organic substance heaps do not have high capacity of active moisture. Changes in productive water supply during the summer months in Pernik are not so sharply defined although there are quite low values here, too. Studies of the chemical composition and properties of heap materials from the two terrains show that they differ significantly (Filcheva, 2007). The only similarity between them is the neutral soil reaction of the soil solution. Total nitrogen content also varies in different ranges from 0.0126% to 0.060% in the heaps in Pernik. In most cases, however, absorbable forms of nitrogen are few, which is confirmed by the high values of the C:N ratio (mostly above 20). These results show that the rate of decomposition is low. Almost all substrates are very poorly stocked with absorbable phosphorus and well to very well stocked with absorbable potassium. The reaction of the soil solution is neutral to slightly alkaline. Only layers that contain more coal and cinder or black clays have more active and hydrolitical acidity.

The type composition of vegetation was determined according to the particular climatic conditions, the properties of the substrates, and also on the basis of the composition and condition of natural and cultural vegetation. Based on the complex ecological and biological evaluation of the determined appropriate plant species to create lawns of Montagu type, anti-erosion grassing and park grass arrays were recommended (Banov and Hristov, 1997). The main species

used by practical teams for afforestation of the new heaps in Pernik coal basin are black pine and acacia, which turn out to be suitable pioneer vegetation for these habitats because of their ecological and biological characteristics. In some cases other deciduous plants such as plain ash tree, birch and lilac bush are used. State and development of all species with the exception of the American ash tree are good, indicating that they can be successfully used in similar conditions. Fertilization combined with proper technology to create and maintain forest crops is a virtually possible activity for successful forest biological reclamation. Fertilization has positive impact on the fertility of substrates. In most cases, the amount of the imported nutrients increases, but it does not lead to accumulation and creating a deficit mode, which requires regular fertilizing in 3-4 years, thus establishing a rapid increase in organic matter (Banov et al., 1994).

On the heap of an open development in Pernik the best effect is obtained with white acacia, white birch and American ash tree, followed by the hybrid sycamore (*Platanus Acerifolia*) and silver lime at the rate of $N_{100}:P_{1500}:K_{50}$. Experiments continue with plain ash tree, chokeberry, large-fruited hawthorn and dog rose. Regarding ways of fertilization better results for the tree and bush vegetation are obtained by nest or row fertilization. Area fertilization produces massive wild development of weeds. In relation to the specific nature of industrial heaps in order to make them suitable for commercial use or green fields, scientists must deal with the problem through a comprehensive approach - carefully and reasonably. This is the only way for successful forestry biological reclamation to be applied to disturbed by coal mining terrains (Banov and Pavlov, 2014).

Conclusion

The affected areas (damaged, altered, etc.) around mining companies in the country are a serious environmental problem, including the territory of Pernik town coal mines. The area around that town, affected by the mining industry occupies a territory of about 30 000 ha. As a result of the natural factors and the anthropogenic impact this region is characterized by a rather complicated soil situation. There are markedly expressed erosion and land-sliding processes on the slopes as well as conditions for surface over-humidification, swamping and salinization. The situation has acquired an exceptionally technogenic character after the development of mines, the enormous pits and heaps and the construction of the industrial pads and equipment. In general, the influence of Mini Pernik is displayed by alterations of the relief, dust and partial aerosol pollution of the atmosphere, destruction of vegetation, contamination and destruction of the soil cover and other components of the environment.

The reclamation of the environment in industrial areas is closely connected above all with the reclamation of disturbed areas and materials and can be performed in different ways according to their properties and economic effect. The inevitable degradation of the landscape can be corrected mainly by technical and biological reclamation of the disturbed terrains. A large part of the green belt of the town of Pernik was constructed on former heaps – mainly by afforestation. These activities have solved the problem of the separation of the town from the mining sites by a noise-preventing and dust-stopping embankment. The recommendations based on the implemented scientific research and the carried out vegetation and field experiments require the growing of agricultural crops directly on the heap substrates without covering them with humus.

References

- Arsov O**, 1997. Evaluation of the Impact of Mini Pernik PLC Site on Environment", Sofia.
- Banov M, Hristov B and Filcheva E**, 1994. Humus Accumulation and its Quality in Reclaimed Lands - Soil Processes and Greenhouse Effect. USDA, Soil observation Service, 140-144.
- Banov M and Hristov B**, 1997. Comparison characteristic of humus accumulation on reclaimed land built with different geological materials - Scientific Conference "50 years Institute of Soil Science in Bulgaria", Sofia 12 - 16 May 1997.
- Banov M and Pavlov P**, 2014. Research and restore land damaged during industrial activity. Yearbook of Mining and Geology University „St. Ivan Rilski“, roll II „Mining and Mineral processing“, „St. Ivan Rilski“ Publishing house, pp. 195-199 (Bg).
- Bogdanov B, Gencheva S and Zheleva E**, 1994. Basic Treatments in Forest Reclamation. Science reports from "Jubilee Conference for 125th anniversary of BAS and 65th anniversary of the IG", Sofia, pp. 193-197.
- Donov V, Gencheva S, Zheleva E, Milanov R and Delkov N**, 1976. Park Construction of Terrains Disturbed by Mining or Industrial Activities, Report on subject, developed in the Research Section of the University of Forestry, Sofia (Bg).
- Donov V, Gencheva S, Zheleva E, Delkov N, Pavlov D and Malinov R**, 1978. Reclamation of Industrial Heaps, Zemizdat Publishing house, Sofia (Bg).
- Filcheva E**, 2007. Characteristics of soils in Bulgaria in content, composition and stocks of organic matter. Grouping of soils in Bulgaria. Published in cooperation with the project "Sustainable Land Management", Sofia, p. 191 (Bg).
- Filcheva E, Noustorova M, Gencheva S and Haigh MJ**, 2000. Organic Accumulation and Microbiological Action in Surface Coal – mine Spoils, Pernik. Bulgaria Journal of Ecological Engineering, 15, 1-15.
- Gencheva S, Delkov N and Zheleva E**, 1982. Opportunities and Results of Soil Reclamation and Landscape in Some Industrial Regions. National Conference of Conservation and Reproduction of Environment, 1-5 of November, Sunny Beach, I, 211-215.
- Gencheva S and Zheleva E**, 1997. Opportunities of Forestry Biological Reclamation for Optimization of Technogenic Landscapes, Collection of the National Conference "Environment – Landscape and Ecology. Theoretical and practical aspects." UBA, National Club, "Landscape Architecture", Sofia, pp. 30-34
- Gencheva S, Zheleva E, Bogdanov B and Petrova R**, 1995. Conditions and Problems of Forestry Biological Reclamation in Bulgaria. "Jubilee Conference for 125th anniversary of BAS and 65th anniversary of the IG", Sofia, pp. 254-258.
- Gencheva S, Zheleva E and Delkov N**, 1984. Reclamation of Terrains, Disturbed by Mining and Industrial Activities, for the Purpose of Environment Conservation, Subject in The Research Section of The University of Forestry, Sofia, code: 283. Scientific Report NIS/KOPS, CINTI-82463017.
- Gencheva S, Zheleva E and Delkov N**, 1986. Biological Reclamation of Heaps from Coal Mining for the Purposes of Forestry", Volume XXX of "Landscape Design and Environmental conservation", Zemizdat Publishing house, Sofia, pp. 63-68 (Bg).
- Gencheva S, Zheleva E and Delkov N**, 1986. Specific Characteristics of the Forestry Reclamation of Heaps from Coal Mining. Collection from the National Conference on Ecological and Genetic problems, Vratsa, II, 439-442.

- Geological map of Bulgaria - Pernik map sheet**, 1991. Committee on geology, Enterprise geophysical surveys and geological mapping. Sofia (Bg).
- Gushevilov Zh**, 1977. Influence of Mineral Fertilization on Reclamation of Marl Materials from Open Mines in West Bulgaria. Thesis for PhD, MAFF, N. Poushkarov ISS, Sofia (Bg).
- Gushevilov Zh and Paskaleva K**, 1977. Composition of the Clay Materials in Marl Heaps in Pernik Coal Mining Region and Their Importance for Agricultural Reclamation. *Soil Science and Agro-chemistry*, 6, 14-22 (Bg).
- Haigh M and Gentcheva-Kostadinova S**, 2002. Ecological erosion control on coal-spoil banks: an evaluation. *Ecological Engineering*, 18, 371-377.
- Haigh M and Gentcheva-Kostadinova S**, 2007. Geomorphological Impact of Erosion Control Measures on a Steep Coal-Spoil Embankment, Pernik, Bulgaria, pp.179-183 (Bg).
- Hristov B, Shishkov T, Katchova V, Atanassova E and Atanassova I**, 2015. Basic Chemical and physico-chemical characteristics of soils and substrates in the region of Pernik coal mine basin. In: Scientific Committee editors. *Soils and Agrotechnology in a Changing World. Digital Book of Proceedings*, May 11-15, 2015. Sofia (Bulgaria): ISSAPP "Nikola Pushkarov",
- Hristova M, Tzolova C, Banov M and Ivanov P**, 2011. Pedogenesis in 2011. Technogenic soils with different reclamation profile. B: I. Atanasov et al. (Ed.). *Scientific reports - International Conference - "100 Years soil science in Bulgaria."* 16 to 20 May 2011. Sofia, part 1. ISBN 978-954-749-088-8. PublishScieSet-Eco, Sofia, pp. 186-190.
- Ivanov P**, 2007. Soil Formation Processes in Reclaimed Areas from Restored Landscapes, Disturbed by Industrial Activities in Different Ways of Use. Thesis for PhD, NCAS, N. Poushkarov ISS, Sofia (Bg).
- Ivanov P, Dimitrova A and Banov M**, 2007. Microbiological Characteristics of Lands Reclaimed after Mining Activities. *Soil Science, Agro-chemistry and Ecology Journal*, 1, 31-36 (Bg).
- Ivanov P, Filcheva E and Banov M**, 2007. "Content and Composition of Organic Substance in Reclaimed after Mining Activities Land". *Soil Science, Agro-chemistry and Ecology Journal*, 1, 27-30 (Bg).
- Ivanov P, Tzolova V, Banov M and Hristova M**, 2007. Opportunities for Improvement of the Qualities and Productive Potential of Technogenic Soils, *Soil Science, Agro-chemistry and Ecology Journal*, 1-4, 127-130.
- Kostadinova A**, 2016. Characteristics of Mining Waste from the Coal Industry and Their Environmentally Sound Management. Thesis for PhD, University of Forestry, Sofia (Bg).
- Kostova S, Ilinkin C and Petrova R**, 2013. Physico-chemical Characterization of Technogenic Soils Formed during Coal Mining in the Region of Pernik. Collection of the Vth National Student Conference "Management and sustainable use of biological resources", Vth National Student Scientific Conference "Management and sustainable use of biological resources."
- Kuzev L**, 2002. Environmental and Technological Problems in the Separation of Raw Maksim-Taban, Yearbook of Mining and Geology "St. Ivan Rilski", 44-45, II Mining and Mineral Processing, Sofia, pp. 145-149.
- Lichev S and Gushevalov Z**, 1973. Growing Agricultural Crops on the Dumps of the Coal Basin of Pernik, Lichev S., Badzhov K., Nikolova M. (editors), V Symposium, Development of methods for reclaiming landscape, disturbed by industry. 3-10-th June, Bourgas, Sunny Beach. G. Dimitrov AA, N.Poushkarov ISS – Sofia.
- Lichev S, Treykyashki P and Gushevilov Zh**, 1973. Characteristics of Geological Materials Formed in Heaps of the Mining area of Pernik. Lichev S., Badzhov K., Nikolova M. (editors), V Symposium, Development of methods for reclaiming landscape, disturbed by industry. 3-10th June, Bourgas, Sunny Beach. G. Dimitrov AA, N.Poushkarov ISS – Sofia, pp. 371-378
- Miltchev M**, 2008. Technology Manageable Fill the Flooded Mine Workings in Underground mine "St. Anna" in the Pernik basin, pp. 1-9 (Bg).
- Mitkov V**, 2013. Report Preparation of Geo-mechanical and Hydro-geological Assessments of the Most Endangered Sites on the Surface of the Suspended Underground Mining Activities at "Mini Pernik" (I), Sofia (Bg).
- Penkov M, Dzhuninski B and Kavardzhiev Y**, 1985. Amelioration of Soils of Unfavourable Properties, Zemizdat Publ. house, Sofia.
- Petrova R, Alexandrov AI, Tsankov G, Gateva R, Shipkovenski D and Petkov P**, 1994. Condition and Ameliorative Effect of Vegetation on Maxim Heap - Pernik. *Science Forest*, 4, 33-39.
- Petrova R and Gencheva S**, 1991. Soil Fertility of Various Origins in the Industrial Embankments in the Pernik coal basin. *Forestry Journal*, 3, 10-11.
- Prokopiev E**, 1962. Opportunities of Landscaping of Heaps in the Coal Basin of Pernik. *Scientific Works of the Institute of Forests*, Sofia, 10, 108-141.
- Regulation No 26**, 2002. For Reclamation of Disturbed areas, Improvement of Low Productivity Lands, Removal and Utilization of the Humus Layer, SG, issue 30/22 March 2002.
- Sabev L and Stanev S**, 1963. Climatic Zones in Bulgaria and Their Climate, Technika Publishing house (Bg).
- Shishkov G, Kehayova M and Stoykova S**, 1985. Coal and Coal Basins in Bulgaria. Sofia, "Kliment Ohridski" Publishing house, p. 163 (Bg).
- Tomova A**, 2000. Research in the Reserve of Ostritsa in Golo Bardo Mountain, University of Forestry, Sofia (Bg).
- Vassilev P and Andreev N**, 1983. Florographic Elements in the Flora of Golo Bardo Mountain. International Symposium "The Relationship of Man and the Mountain Ecosystems", Project 6 of the program MAB, Vratsa, 24-29 of October, 1983, I, Sofia.
- Zahariev B, Donovan V and Masarov S**, 1983. Forest-plantation Zoning of Bulgaria, Zemizdat Publishing house, Sofia (Bg).
- Zheleva E**, 1979. Opportunities of Reclamation of Industrial Heaps for the Needs of Landscaping, Sofia, Center of Scientific Information at the MFF.
- Zheleva E**, 2002. Assessment of the Environmental Impact of an Investment Project for Processing of Technological Waste from the Produce of Pernik Mines – the 7th September and Karamanitsa Heaps with an Enrichment Installation for Extracting of a Fuel Product, Sofia.
- Zheleva E**, 2010. Reclamation of Disturbed Areas. PSSE, Sofia, p. 411 (Bg).
- Zheleva E, Gencheva S and Heig M**, 1995. Opportunities for Joint Science Research on Problems of Forestry Biological Reclamation in Great Britain and Bulgaria. "Jubilee Conference for 125th anniversary of BAS and 65th anniversary of the IG", Sofia, pp. 189-192.
- Zheleva E and Kostova E**, 1986. Increasing the Fertility of Industrial Heaps by Mineral Fertilization. *Forestry and Forest Industry Journal*, 9, Sofia (Bg).

Review

- Alternatives for optimisation of rumen fermentation in ruminants** 91
T. Slavov

Genetics and Breeding

- Characterization of a new winter malting barley cultivar Ahil** 98
B. Dyulgerova, D. Vulchev, T. Popova
- Evaluation of high yielding mutants of *Hordeum vulgare* cultivar Izgrev** 103
B. Dyulgerova, N. Dyulgerov

Nutrition and Physiology

- In vitro gas production of different feeds and feed ingredients at ruminants** 105
E. Videv, J. Krastanov, S. Laleva, T. Angelova, M. Oblakova, N. Oblakov, D. Yordanova, V. Karabashev
- Evaluation of chemical composition of raw and processed tropical sickle pod (*Senna obtusifolia*) seed meal** 110
Augustine C., Kwari I.D., Igwebuike J.U., Adamu S.B.
- Effect of urea-fortified all concentrate corncob diets on serum biochemical and hematological indices of West African dwarf goats** 114
U. M. Kolo, A. A. Adeloye, M. B. Yousuf

Production Systems

- Analysis of the technological dairy cows traffic "to and from" herringbone milking parlors** 119
K. Peychev, D. Georgiev, V. Dimova, V. Georgieva
- Effect of pre-sowing soil tillage for wheat on the crop structure and the yield components under the conditions of slightly leached chernozem soil in Dobrudzha region** 124
P. Yankov, M. Drumeva
- Study on the process of unloading grain harvesters at the end of the field** 129
G. Tihanov

Agriculture and Environment

- Modeling and simulation of fuzzy logic controller for optimization of the greenhouse microclimate management** 132
Didi Faouzi, N. Bibi-Triki, B. Draoui, A. Abene

Floristic diversity of 'Chinarite' protected area – Rodopi municipality, Bulgaria L. Dospatliev, M. Lacheva	140
Heavy metal pools in urban soils from city parks of Sofia, Bulgaria V. G. Kachova, I. D. Atanassova	144
Ecological characteristics of reclaimed areas in Pernik mines region, Bulgaria I. Kirilov, M. Banov	151
Reclamation of soil excavated from construction and mine searching areas in Turkey F. Apaydin	160
 Product Quality and Safety	
Concentration of sulfur-containing amino acids at turkey broilers during and after muscle dystrophy, fed with deficient feed supplemented with oxidised fat K. Stoyanchev	164
Exopolysaccharide influence's on acid gel formation K. Yoanidu, P. Boyanova, P. Panayotov	167
Carcass characteristics and technological properties of <i>Musculus Longissimus Lumborum</i> at lambs from the Bulgarian dairy synthetic population and its F1 crosses with meat breeds N. Ivanov, T. Angelova, S. Laleva S. Ribarski, D. Miteva, D. Yordanova, V. Karabashev, I. Penchev	171

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors

The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings:

Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows. IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 9, Number 2
June 2017

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz