ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 9, Number 2
June 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate on author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc. and AGRIS (FAO).

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No DNP 05-21/20.12.2016, financed from Fund 'Scientific Research' grant Bulgarian scientific Periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 9, Number 2
June 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Product Quality and Safety

Concentration of sulfur-containing amino acids at turkey broilers during and after muscle dystrophy, fed with deficient feed supplemented with oxidised fat

K. Stoyanchev*

Department of Internal Diseases and Clinical Toxicology, Faculty of Veterinary Medicine, Trakia University, 6000 Stara Zagora, Bulgaria

(Manuscript received 5 April 2016; accepted for publication 7 June 2017)

Abstract. The aim of the present study was to reproduce experimentally muscular dystrophy in 50 broiler turkeys, through early nutrition with a diet deficient in vitamin E, selenium and sulfur-containing amino acids, supplemented with oxidised fat and to study blood plasma sulfur containing amino acids (methionine and cysteine). The experiments were conducted with 1 day-old broiler turkeys. By the 3rd day of life, they were divided into 40 experimental (II group) and 10 control birds (I group); the latter were fed a standard compound feed, whereas the former group received a diet deficient in sulfur-containing amino acids methionine and cysteine (reduced up to 50%), vitamin E, and Se (from 0.2 mg/kg in standard feed to 0.01 mg/kg), further supplemented with oxidized fat containing peroxides and aldehydes with peroxide number of the food 8.0 meq O₂/kg. The clinical signs of experimental muscle dystrophy in broiler turkeys appeared first by the 25th day of feeding, when the mild clinical form (II A group) and the severe clinical form (II B group) were established. The results indicated clearly that in turkey broilers suffering from muscle dystrophy, the concentrations of sulfur-containing amino acids cysteine and methionine decreased. After the treatment of turkey broilers with muscle dystrophy, and supplementation with non-deficient forage with Se, vitamin E, but also with sulfur-containing amino acids cysteine or methionine with Seled at a dosage 0.06mg/kg, the plasma levels of sulfur-containing amino acids cysteine and methionine was normalized in the mild clinical form (II A group). The birds affected by the severe clinical form of disease (II B group), which were not treated with Seled® and whose deficient feed was not corrected, could not recover and levels of sulfur-containing amino acids cysteine and methionine did not normalize.

Keywords: turkey broilers, muscular dystrophy, methionine, cysteine

Introduction

Avian myopathies are a serious concern for the welfare and productivity of poultry (Georgiev, 1979; Gabrashanski et al., 1979; Kulikova et al., 1979; Aliev et al., 1985; Avanzo et al., 2001; Todorovic et al., 2002; Mench, 2004; Redish et al., 2005; Kuttappan et al., 2012). Through genetic selection, some progress has been achieved for reduction of these diseases, but yet, an effective strategy for their eradication is lacking (Mench, 2004).

The deficiency of Se and vitamin E in the food is considered as the primary cause for myopathies (Todorovic et al., 2002; Chang et al., 2005). Reviewing the investigations of some authors, Gabrashanski et al. (1979) outlined the role of vitamin E, vitamin A and some amino acids in the pathogenesis of muscle dystrophy, whereas Helmbold et al. (1972) believed that apart for building the proteins needed for tissue growth and egg formation, these amino acids have other functions as well. It is stated that vitamin E and sulfur-containing amino acids deficiency is the cause for the development of muscle dystrophy in young chickens. For instance, creatine, that is physiologically essential for the functioning of muscle tissue, is synthesized from the amino acids arginine, methionine and glycine. At the same time, discussing the role of vitamin E in the pathogenesis of myopathies in chickens, turkeys and ducklings, the authors' strategy for control of these states includes three factors: 1) vitamin E as a natural and most effective antioxidant for prevention of encephalomalacia; 2) its special relationship with Se for prevention of exudative diathesis and

myopathies in turkeys and 3) the relation between both Se and the amino acid cysteine for prevention of nutritional muscle dystrophy.

Taking into consideration the insufficient reports about the role of some amino acids in the pathogenesis of muscle dystrophy in birds, the present study aimed to investigate the levels of the sulfur-containing amino acids methionine and cysteine as well as that of amino acid lysine in turkey broilers, suffering from and recovered from muscle dystrophy.

Material and methods

The experiments were carried out in the Experimental Base of the Department of Internal Diseases, Faculty of Veterinary Medicine, Trakia University. They were conducted on 50 one-day old broiler turkeys from the "Konverter" hybrid, Kanada. The birds were identified by wing marks. From the 1st to the 3rd day of life, all turkeys were put under the same regimen of feeding and rearing and were fed a standard compound feed.

By day 3, they were initially divided into 2 groups: the control group (n = 10) was fed a starter ration. The experimental group (n = 40) was fed starter forage for broiler turkeys deficient in vitamin E, selenium (Se) and sulfur-containing amino acids methionine and cysteine (50%) and was supplemented with oxidized fat with peroxide number of the food 8.0 meq O₂/kg. In accordance with the national prevention programme control birds were treated with Seled® at 0.06 mg/kg per os, between 6 – 10 and 16 – 20 days of age,

* e-mail: ksto@abv.bg

while experimental birds did not receive Seled®. The clinical signs of experimental muscle dystrophy in broiler turkeys appeared first by the 25th day of feeding, when the mild clinical form (II A group, n = 20) and the severe clinical form (II B group, n = 20) were established. Reproduction of muscular dystrophy and clinical signs in broiler turkeys were similar to those in broiler chickens, described in detail in a previous study of ours (Stoyanchev and Maruzova, 2017).

In all groups, blood was sampled from the wing vein after manifestation of the disease by the 25th and 28th day, prior to the treatment with Seled® and by the 31th and 34th day after the treatment with Seled® at a dose of 0.06 mg/kg per os. Blood for biochemical analyses (5 ml) was sampled from *v. subcutaneous ulnaris*. Heparin was used as anticoagulant.

By the 25th and 28th day, 7-day oral treatment with Seled® at 0.06 mg/kg was initiated in the chickens with mild muscular dystrophy (group IIA) and simultaneously, the deficient feed was replaced with the standard one. In experimental group IIB, the birds were not treated with Seled® and the deficient feed was nonreplaced with the standard one.

The blood plasma concentrations of sulfur-containing amino acids were assayed with an automated amino acid analyzer T 339 M MICROTECHNA PRAHA in the Research Laboratory of the Faculty of Agriculture at the Trakia University.

The analysis of numerical data was done with statistical software Statistica 6.0® (Stat Soft SEM, USA).

Results and discussion

The results about the blood plasma levels of sulfur-containing amino acid cysteine in healthy and suffering from muscle dystrophy turkey-broilers prior to the treatment and after treatment are presented on Figure 1.

At diseased turkey broilers, by the 25th day of feeding cysteine decreased considerably – $0.086 \pm 0.004\%$ in group IIA and $0.078 \pm 0.005\%$ in group IIB, the difference with the respective control group ($0.118 \pm 0.003\%$) being statistically significant ($p < 0.05$). By the 28th day of disease the levels were – $0.072 \pm 0.003\%$ in group IIA and $0.072 \pm 0.006\%$ in group IIB, the difference with the respective control group being statistically significant ($p < 0.01$). It could be seen that cysteine concentrations were particularly low in diseased birds.

After treatment by the 31th and the 34th day in group IIA, blood

Figure 1. Blood plasma concentrations of the sulfur containing amino acid cysteine at turkey-broilers, either healthy or with muscular dystrophy prior to treatment and after treatment with Seled® (* $p < 0.05$, ** $p < 0.01$)

cysteine levels changed, increasing to $0.110 \pm 0.005\%$ and to $0.109 \pm 0.007\%$, respectively, but the difference with the respective control group was insignificant ($p > 0.05$). In experimental group IIB, where the birds were not treated with Seled® and the deficient compound feed was not replaced with normal, by the 31th and the 34th day, blood cysteine levels did not change – $0.070 \pm 0.006\%$ and to $0.071 \pm 0.005\%$, respectively, and were statistically significantly different ($p < 0.01$) vs controls.

The analysis of blood methionine levels prior to the treatment demonstrated a similar tendency (Figure 2). In turkeys with muscle dystrophy, by the 25th day of feeding it was highly reduced – $0.057 \pm 0.007\%$ in group IIA and $0.055 \pm 0.005\%$ in group IIB (statistically significantly lower than respective controls: $0.081 \pm 0.009\%$ at $p < 0.01$). The highest reduction occurred in the diseased poult by the 28th day of feeding – $0.051 \pm 0.004\%$ in group IIA and $0.051 \pm 0.006\%$ in IIB, also statistically significantly different from control concentrations ($p < 0.01$).

In healed turkey-broilers by the 31th and 34th day, after treatment, methionine concentrations increased in group IIA: to $0.077 \pm 0.008\%$ and $0.078 \pm 0.009\%$, respectively, although they reached control levels and the differences vs respective controls were insignificant ($p > 0.05$). In group IIB, the birds were not treated with Seled® and the deficient compound feed was not replaced with normal, so by the 31th and the 34th day, blood methionine levels did not change – $0.052 \pm 0.006\%$ and to $0.056 \pm 0.007\%$, respectively ($p < 0.01$ vs control levels).

In conclusion, the analysis of blood plasma concentrations of the sulfur containing amino acids showed a reduction in turkey-broilers suffering from muscle dystrophy. These data supported the involvement of sulfur-containing amino acids in muscle dystrophy. Our results corresponded to the opinion of Helmboldt et al. (1972) who assume that the deficiency of vitamin E and sulfur-containing amino acids in the food results in the development of nutritional muscle dystrophy in chickens. The amino acids methionine, arginine and glycine participate in creatine synthesis that is physiologically essential for the function of muscle tissue.

Our data about the role of sulfur-containing amino acids in the muscle dystrophy in turkey-broilers are also confirmed by the fact, that the treatment protocol of muscle dystrophy includes the supplementation of non-deficient forage not only with Se, vitamin E, but also with cysteine or methionine.

Explaining the pathogenesis of myopathies in chickens, turkey

Figure 2. Blood plasma concentrations of the sulfur containing amino acid methionine at turkey-broilers, either healthy or with muscular dystrophy prior to treatment and after treatment with Seled® (** $p < 0.01$)

poults and ducklings, the inadequate amount of sulfur-containing amino acids is outlined together with vitamin E deficiency (Gordon and Jordon, 1985). These deficiencies result in impaired membrane structure. The decreased blood plasma concentrations of sulfur-containing amino acids in turkey-broilers suffering from muscle dystrophy, observed by us, is supported by the views of aforementioned authors.

Conclusion

In turkey-broilers suffering from muscle dystrophy, the blood concentrations of sulfur-containing amino acids cysteine and methionine decreased. After the treatment of turkey-broilers with muscle dystrophy, and supplementation with undecified forage with Se, vitamin E, but also with sulfur-containing amino acids cysteine or methionine with 0.06 mg/kg Seled the plasma levels of sulfur-containing amino acids cysteine and methionine were normalised in birds affected with the mild clinical form (II A group). The birds suffering from the severe clinical form of disease (II B group), which were not treated with Seled® and whose deficient feed was not corrected, could not recover and levels of sulfur-containing amino acids cysteine and methionine did not become normal.

References

- Aliev A, Barei V, Bartko P, Bolda Y, Djordjak P, Gabrashanski P, Jlek J, Jagosh P, Karsai F, Koni E, Kovash G, Nedkova L, Sova Z, Shefer M and Yrzgula L**, 1985. Metabolic Diseases in domestic animals. Zemizdat Publishing house, Sofia (Bg).
- Avanzo JL, Mendonca Junior C X de and Cesar M de C**, 2002. Role of antioxidant systems in induced nutritional pancreatic atrophy in chicken. *Comparative Biochemistry and Physiology, Biochemistry & Molecular Biology*, 131, 815-823.
- Chang WP, Combs Jr, Scanes CG and Marsh JA**, 2005. The effects of dietary vitamin E and selenium deficiencies on plasma thyroid and thymic hormone concentrations in the chicken. *Developmental and Comparative Immunology* 29, 265-273.
- Gabrashanski P, Kovalski V, Rish M, Nedkova I, Daskalova A, Raacka U and Abdulaev D**, 1979. Trace elements and trace elementoses in industrial animal husbandry. Zemizdat Publ. house, Sofia-Moscow (Bg).
- Georgiev H**, 1979. Studies of some diseases in turkeys related to selenium and vitamins E deficiencies. Thesis for PhD, VVMI, Sofia, (Bg).
- Gordon KF and Jordon ET**, 1985. Poultry diseases, Bailliere Tindal, London.
- Helboldt W, Reid M and Joder H**, 1972. Disease of Poultry, The Iowa State University Press, Ames.
- Kulikova WF, Kudriavcev U and Ilushechkin UP**, 1979. Diseases of turkeys. Leningrad, Kolos Publ. house (Ru).
- Kuttappan VA, Goodgame SD, Bradley CD, Mauromoustakos A, Hargis BM, Waldroup PW and Owens CM**, 2012. Effect of different levels of dietary vitamin E (DL- α -tocopherol acetate) on the occurrence of various degrees of white striping on broiler breast fillets. *Poultry Science*, 91, 3230-5.
- Manual BUT**, 2000. The team breeds success. Warren Hail Broughth Chester CH4OEW, England, 1-48.
- Mench J**, 2004. Measuring and auditing broiler welfare. In: Lameness (ed. Weeks CA, Butterworth A.). Wallingford, CABI Publishing, p. 3-17.
- Reddish JM, Latshaw JD, St-Pierre NR, Pretzman C and Mar WM**, 2005. Myosin Heavy Chain Isoform Expression Is Not Altered in the Pectoralis Major Muscle in Selenium-Deficient, Chickens Recovering from Exudative Diathetic Myopathy. *Poultry Science*, 84, 462.
- Stoyanchev K and Maruzova V**, 2017. Reproduction of muscular dystrophy in broiler chickens through early nutrition with deficient feed supplemented with oxidised fat. *Trakia Journal of Sciences*, 15, 67-73.
- Todorovic M, Jokic Z and Davidovic V**, 2002. The influence of selenium and vitamin E in poultry nutrition / Znacaj selena i vitamina E u ishrani zivine. *Biotechnology in Animal Husbandry*, 18, 231-238.

Review

- Alternatives for optimisation of rumen fermentation in ruminants** 91
T. Slavov

Genetics and Breeding

- Characterization of a new winter malting barley cultivar Ahil** 98
B. Dyulgerova, D. Vulchev, T. Popova
- Evaluation of high yielding mutants of *Hordeum vulgare* cultivar Izgrev** 103
B. Dyulgerova, N. Dyulgerov

Nutrition and Physiology

- In vitro gas production of different feeds and feed ingredients at ruminants** 105
E. Videv, J. Krastanov, S. Laleva, T. Angelova, M. Oblakova, N. Oblakov, D. Yordanova, V. Karabashev
- Evaluation of chemical composition of raw and processed tropical sickle pod (*Senna obtusifolia*) seed meal** 110
Augustine C., Kwari I.D., Igwebuike J.U., Adamu S.B.
- Effect of urea-fortified all concentrate corncob diets on serum biochemical and hematological indices of West African dwarf goats** 114
U. M. Kolo, A. A. Adeloye, M. B. Yousuf

Production Systems

- Analysis of the technological dairy cows traffic "to and from" herringbone milking parlors** 119
K. Peychev, D. Georgiev, V. Dimova, V. Georgieva
- Effect of pre-sowing soil tillage for wheat on the crop structure and the yield components under the conditions of slightly leached chernozem soil in Dobrudzha region** 124
P. Yankov, M. Drumeva
- Study on the process of unloading grain harvesters at the end of the field** 129
G. Tihanov

Agriculture and Environment

- Modeling and simulation of fuzzy logic controller for optimization of the greenhouse microclimate management** 132
Didi Faouzi, N. Bibi-Triki, B. Draoui, A. Abene

Floristic diversity of 'Chinarite' protected area – Rodopi municipality, Bulgaria L. Dospatliev, M. Lacheva	140
Heavy metal pools in urban soils from city parks of Sofia, Bulgaria V. G. Kachova, I. D. Atanassova	144
Ecological characteristics of reclaimed areas in Pernik mines region, Bulgaria I. Kirilov, M. Banov	151
Reclamation of soil excavated from construction and mine searching areas in Turkey F. Apaydin	160
 Product Quality and Safety	
Concentration of sulfur-containing amino acids at turkey broilers during and after muscle dystrophy, fed with deficient feed supplemented with oxidised fat K. Stoyanchev	164
Exopolysaccharide influence's on acid gel formation K. Yoanidu, P. Boyanova, P. Panayotov	167
Carcass characteristics and technological properties of <i>Musculus Longissimus Lumborum</i> at lambs from the Bulgarian dairy synthetic population and its F1 crosses with meat breeds N. Ivanov, T. Angelova, S. Laleva S. Ribarski, D. Miteva, D. Yordanova, V. Karabashev, I. Penchev	171

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors

The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings:

Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows. IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 9, Number 2
June 2017

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz