ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 9, Number 3
September 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate on author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc., AGRIS (FAO) and DOAJ.

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No DNP 05-21/20.12.2016, financed from Fund 'Scientific Research' grant Bulgarian scientific Periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 9, Number 3
September 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Genetics and Breeding

Parthenogenetic responsiveness of sunflower hybrid combinations with expressed tolerance to herbicides

M. Drumeva*, P. Yankov

Department of Plant Production, Faculty of Shipbuilding, Technical University Varna, 1 Studentska, 9010 Varna, Bulgaria

(Manuscript received 6 March 2017; accepted for publication 3 July 2017)

Abstract. For accelerated development of fertility restorer lines with imidazolinone tolerance in sunflower (*Helianthus annuus* L.), the gamma-induced parthenogenesis, which combines classical breeding approaches and in vitro techniques, was applied. As initial materials six imidazolinone-tolerant hybrid combinations (F1203, F1216, F1246, F1248, F1295 and F1332), obtained from Dobrudzha Agricultural Institute, General Toshevo, were used. To initiate parthenogenetic development mixed pollen from different fertility restorer lines was applied. The mixed pollen was irradiated with a dose of 600 Gy. A total of 166 parthenogenetic immature embryos were obtained, and 84 young plants developed from them, 42 of the plants were fertile. Best parthenogenetic response, expressed as a number of embryos obtained, was observed by hybrid combination F1295, followed by hybrid combination F1332. The lowest parthenogenetic responsiveness is reported in hybrid combinations F1246 and F1248, where the total number of parthenogenetic embryos obtained was 5. In hybrid combinations F1295 and F1332, as well as in F1216, the method of gamma-induced parthenogenesis can be efficiently used for accelerated production of parental lines for heterosis breeding in sunflower. In hybrid combinations F1203, F1246 and F1248, which showed low parthenogenetic responsiveness, conventional methods for developing lines from these hybrids are recommended.

Keywords: sunflower, double haploid fertility restorer lines, embryo culture, gamma-induced parthenogenesis

Introduction

The herbicide-tolerant sunflower hybrids are generally preferred by agricultural producers in Bulgaria and this determines the main directions of contemporary sunflower breeding. Imidazolinone (IMI) and sulfonylurea (SU) tolerance to herbicides transferred to cultural sunflower from wild species is most often used (Kaya et al., 2012).

The development of parental components for the hybrid is a necessary prerequisite for the production of sunflower hybrids tolerant to herbicides. The classical breeding approach for developing lines that can be included as parental components of hybrids comprises inbreeding of plants for at least 5 – 6 years. To accelerate this process, a number of researchers have directed their attention towards non-traditional approaches for production of homozygotic lines. The development of plant biotechnologies opened new opportunities in this respect. One of the methods for accelerated development of haploid and double haploid lines is the gamma-induced parthenogenesis (Todorova et al., 1997; Gok Guler et al., 2015; Blasco et al., 2016; Wang et al., 2016). The use of this method significantly shortens the time necessary to develop parental lines for heterosis breeding in sunflower (Todorova and Ivanov, 1999, 2000). The properties of the double haploid lines obtained through this method are determined by the genetic potential of the initial material they were derived from. In spite of the unquestionable benefits from the use of biotechnology methods for acceleration of the breeding process, there are many cases in which they are influenced by the genotypic responsiveness of the investigated plant material (Sauton, 1989; Bouvier et al., 1993; Faris

et al., 1999; Todorova et al., 2004; Taşkın et al., 2013; Drumeva and Yankov, 2015). Therefore, additional investigations and certain modifications of the standard methodologies are necessary depending on the specific requirements of the genotype.

This study was aimed at investigating the genotypic responsiveness of sunflower hybrid combinations with expressed herbicide tolerance to the application of the gamma-induced parthenogenesis method for accelerated production of imidazolinones-tolerant fertility restorer lines.

Material and methods

As initial materials for parthenogenetic induction six imidazolinone-tolerant hybrid combinations (F1203, F1216, F1246, F1248, F1295 and F1332), obtained from Dobrudzha Agricultural Institute, General Toshevo, were used. For initiation of parthenogenetic embryo development in the studied hybrid combinations, pollination with gamma rays inactivated mixed pollen from fertility restorer lines was applied. The pollen collected from these lines was mixed at equal weight portions and was irradiated with dose 600 Gy, with Cs 137 as a source of ionizing radiation.

To sterilize the male gametophyte the initial plants from the hybrids were treated with water solution of gibberellic acid (GA₃) in concentration 45 mg/l. The solution was applied directly on the young button (1-1.5 cm in diameter). To avoid foreign pollination all initial female lines were isolated immediately before beginning of flowering. Only 5 plants from each of the initial female genotypes were pollinated.

* e-mail: m_drumeva@abv.bg

The isolation and cultivation of the obtained parthenogenetic embryos was done according to the method of Azpiroz et al. (1988). The formed young parthenogenetic plants were transferred to soil and stabilized at temperature 20°C/15°C day/night, relative air humidity 70% and photoperiod 16/8 h day/night for 30 days, then the day temperature was increased to 25°C ± 2°C, and the night temperature to 20°C ± 2°C.

The ploidity of the plants was determined flow-cytometrically. The plants were analyzed at growth stage first-second pair of leaves. Cell analyzer Partec CA II (Partec GmbH) was used. The samples were prepared according to the protocol of De Laat and Göhde (1987). The nuclei were stained with DAPI (4 6-diamidino-2-phenylindole) (Göhde et al., 1979).

The analysis of the histograms was done with DPAC (Partec GmbH). To evaluate the parthenogenetic response of the IMI resistant hybrid combinations involved in the experiment, one-factor dispersion analysis and cluster analyses were applied. Data were processed with Statistica, Version 7.

Results and discussion

Out of the 6 hybrid combinations used as initial material for parthenogenetic induction, a total of 166 immature embryos were obtained and 84 young plantlets developed from them. The analysis on the parthenogenetic responsiveness of the investigated hybrids revealed significant differences in their reaction (Table 1). A number of authors point out that the genotype of the initial parental forms is an important factor for the successful induction of parthenogenetic development with the aid of irradiated pollen (Sauton, 1989; Bouvier

et al., 1993; Faris et al., 1999; Todorova et al., 2004; Drumeva and Yankov, 2015). By the present study the effect of the female genotype was observed, expressed in a different response of the investigated hybrids to initiation of parthenogenetic development. Up to now there are no established parameters which can clearly indicate when satisfactory results from the application of the method can be expected and when it will not be effective, concerning the influence of the female genotype (Drumeva and Yankov, 2015). The fact that there are plant species in which parthenogenesis occurs naturally implies that there are specific genes determining the parthenogenetic development. In spite of the numerous studies up to now, it is still unclear if the parthenogenetic processes result from the action of specific genes or reflect a temporary variation in the expression of key regulatory genes usually involved in sexual reproduction (Koltunow, 1993).

In this investigation the highest number of embryos was obtained from hybrid combination F1295, and the lowest – from F1246 (Figure 1). These two hybrids outlined the parthenogenetic responsiveness of the investigated hybrids in this experiment. A good parthenogenetic induction was read in hybrid F1332. In the rest of the hybrid combinations, the frequency range of parthenogenetic induction was from 3 (formed embryos) in hybrid combination F1248 to 11 in F1216.

To obtain more details about the results from the parthenogenetic responsiveness of the hybrids involved in this investigation, the criterion of Student-t was applied (Table 2). The analysis proved that the variations in the parthenogenetic reaction of hybrid combinations F1203 and F1246 were not significant, while the variations between them and F1248 were with the lowest degree of significance. F1216, F1295 and F1332 were with high statistical

Table 1. Dispersion analysis of the parthenogenetic reaction of the investigated imidazolinone-tolerant hybrid combinations for the trait number of parthenogenetic embryos

Factors	SS	df	MS	F	Sig.
Corrected Model	23866.000 ^a	9	4773.200	2.387E3	.000
Hybrid combinations	23866.000	5	4773.200	2.387E3	.000
Error	24.000	12	2.000		
Corrected Total	23890.000	17			

Figure 1. Genotype responsiveness of the investigated hybrid combinations to parthenogenetic induction

Figure 2. Double haploid plants obtained from the investigated hybrid combinations by using the method of gamma-induced parthenogenesis

Table 2. Statistical evaluation of the variation in the parthenogenetic responsiveness of the investigated hybrids for the trait number of parthenogenetic embryos

Hybrid combinations	F1203	F1216	F1246	F1248	F1295	F1332
F1203	—	***	*	ns	***	***
F1216		—	***	***	***	***
F1246			—	ns	***	***
F1248				—	***	***
F1295					—	***
F1332						—

GD_{5%} = 2.23 *   GD_{1%} = 3.13 **   GD_{0.1%} = 4.42 ***

significance against the background of all investigated hybrid combinations.

When conditionally grouping the investigated hybrids by their parthenogenetic responsiveness, four groups were differentiated – hybrids with low parthenogenetic reaction (F1203 and F1246), hybrids with moderate expression of the parthenogenetic response (F1216), hybrids with good parthenogenetic responsiveness (F1332) and hybrids with very good parthenogenetic responsiveness (F1295). Under the conditions of this experiment, normal growth and development into young double haploid plants were observed in all parthenogenetic embryos obtained from the hybrid genotypes. The diploid status was flow-cytometrically determined in all produced plants, and 50% of the plants were fertile. Fertile plants were obtained from all investigated combinations, with the exception of F1246. The fertile plants were double haploid lines, in which the fertility restorer gene was in homozygous dominant state, while in the sterile plants it was in homozygous recessive state (Figure 2).

With regard to the number of obtained fertile plants, which were potential lines with herbicide resistance, the tendency noticed while reading the differences in the number of parthenogenetic embryos between the individual hybrid combinations remained the same to some extent, the statistical significance of the differences in some of the hybrids being with lower degree of significance. The differences between F1295 and F1332, the hybrids with the highest potential for

parthenogenetic responsiveness, were significant at P=0.05. This can be explained with the lower survival rate of the parthenogenetic embryos formed from F1295. These results did not change the initial grouping of the hybrids since the formation of the parthenogenetic embryo depended closely on the genotypic responsiveness to parthenogenetic induction, while the development of a plant from this embryo was a function of additional factors – composition of the nutrition media, conditions of cultivation, etc.

Figures 3 and 4 present the relative percent of each investigated hybrid combination, both from the total number of obtained parthenogenetic embryos, and from the total number of fertile plants with potential tolerance to herbicides. The relative percentage in the final product - the number of fertile parthenogenetic plants, decreased by 17% in hybrid F1295, compared to its share in terms of the total number of formed parthenogenetic embryos. In contrast, all other hybrid combinations, except for F1246, had an increase in the percentage of this indicator. In F1332 the relative share in the final product increased by 7%, and in F1216 - by 6%. In the other two hybrid combinations (F1203 and F1248), where a relatively low parthenogenetic induction frequency was reported, there is also an increase of their share in the final product. However, the 2% in F1203 and 3% in F1248, respectively, translated into absolute values, are not enough for efficient use of these two genotypes for the implementation of the method of gamma-induced parthenogenesis..

Figure 3. Relative percent of the investigated hybrid combinations from the total number of obtained parthenogenetic embryos

Figure 4. Relative percent of the investigated hybrid combinations from the total number of obtained parthenogenetic fertile plants

Conclusion

The efficiency of the gamma-induced parthenogenesis method depends on the genotypic responsiveness of the initial hybrid combinations. Hybrid combinations F1295 and F1332 demonstrated very good and good responsiveness to parthenogenetic induction. The highest relative percentage from the end product as a result of the application of the gamma-induced parthenogenesis method, expressed as number of fertile double haploid plants, was found in F1295 and F1332. In hybrid combinations F1295 and F1332, as well as in F1216, the method of gamma-induced parthenogenesis can be efficiently used for accelerated production of parental lines for heterosis breeding in sunflower. In hybrid combinations F1203, F1246 and F1248, which showed low parthenogenetic responsiveness, conventional methods for developing lines from these hybrids are recommended.

Acknowledgement

A part of the scientific research, the results of which are presented in this article, was conducted at Technical University, Varna as part of a research project, funded by the state budget.

References

- Azpiroz H, Vincourt P, Serieys H and Gallais A, 1988. La culture *in vitro* des embryons immatures dans l'accélération du cycle de sélection des lignées de tournesol et ses effets morpho-vegetatifs. *Helia*, 10, 35-38.
- Blasco M, Badenes, ML and del Mar Naval M, 2016. Induced parthenogenesis by gamma-irradiated pollen in loquat for haploid production. *Breeding Science*, 66, 606-612.
- Bouvier L, Zhang Y and Lespinasse Y, 1993. Two methods of haploidization in pear, *Pyrus communis* L.: greenhouse seedling selection and *in situ* parthenogenesis induced by irradiated pollen. *Theoretical and Applied Genetics*, 87, 229-232.
- De Laat A, Ghde W and Vogelzang M, 1987. Determination of ploidy of single plants and plant populations by flow cytometry. *Plant Breeding*, 99, 303-307.
- Drumeva M and Yankov P, 2015. Investigation on the parthenogenetic response of sunflower lines and hybrids. *Agricultural Science and Technology*, 7, 415-418.
- Faris N, Nikolova V and Niemirowicz-Szczytt K, 1999. The effect of gamma irradiation dose on cucumber (*Cucumis sativus* L.) haploid embryo production. *Acta Physiologiae Plantarum*, 21, 301-396.
- Gok Guler P, Sari N, Yetisir H, Yegul M, Kantoglu Y and Kunter B, 2015. The effects of genotypes and irradiation doses on haploid embryo induction and plant production in bottle gourd [*Lagenaria siceraria* (Malign) Stanley]. In V International Symposium on Cucurbits, 1151, pp. 135-142.
- Ghde W, Schumann J, Bchner Th, Otto F and Barlogie B, 1979. Pulse cytophotometry – application in tumor cell biology and clinical oncology. In Melamed M.R., Mullaney P.F. & M.L. Mendelsohn (eds.) *Flow cytometry and sorting*, John Wiley and Sons, New York, p. 599.
- Kaya Y, Jovic S and Miladinovic D, 2012. "Sunflower." *Technological Innovations in major world oil crops*, Springer New York, 1, 85-129.
- Koltunow A M, 1993. Apomixis: Embryo sacs and embryos formed without meiosis or fertilization in ovules. *The Plant Cell*, 5, 1425-1437.
- Sauton A, 1989. Haploid gynogenesis in *Cucumis sativus* induced by irradiated pollen. *Cucurbit Genetics Cooperative*, 12, 22-23.
- Taşkın H, Yücel NK, Baktemur G, Çömlekçioğlu S and Büyükalaca S, 2013. Effects of different genotypes and gamma ray doses on haploidization with irradiated pollen technique in watermelon (*Citrullus lanatus* L.). *Canadian Journal of Plant Science*, 93, 1165-1168.
- Todorova M and Ivanov P, 1999. Induced parthenogenesis in sunflower: Effect of pollen donor. *Helia*, 22, 49-56.
- Todorova M and Ivanov P, 2000. Induced parthenogenesis in sunflower: Effect of gamma irradiation doses. 15th International Sunflower Conference, Toulouse, France, June, 2000, L46-L51.
- Todorova M, Ivanov P, Nenova N and Encheva J, 2004. Effect of the female genotype on the efficiency of -induced parthenogenesis in sunflower (*Helianthus annuus* L.). *Helia*, 27, 67-74.
- Todorova M, Ivanov P, Shindrova P, Christov M and Ivanova I, 1997. Dihaploid plant production of sunflower (*Helianthus annuus* L.) through irradiated pollen-induced parthenogenesis. *Euphytica*, 97, 249-254.
- Wang SM, Lan H, Jia HH, Xie KD, Wu XM, Chen CL and Guo WW, 2016. Induction of parthenogenetic haploid plants using gamma irradiated pollens in 'Hirado Buntan'pummelo (*Citrus grandis* L. Osbeck). *Scientia Horticulturae*, 207, 233-239.

Reviews

- Problems and achievements of cotton (*Gossypium Hirsutum* L.) weeds control** 179
T. Barakova, G. Delchev
- Achievements and problems in the weed control in grain sorghum (*Sorghum Bicolor* Moench.)** 185
G. Delchev, M. Georgiev

Genetics and Breeding

- Parthenogenetic responsiveness of sunflower hybrid combinations with expressed tolerance to herbicides** 190
M. Drumeva, P. Yankov
- In vitro propagation of oil-bearing rose (*Rosa damascena* Mill.)** 194
V. Badzhelova

Nutrition and Physiology

- Variation in the chemical composition and physical characteristics of grain from winter barley varieties** 198
B. Dyulgerova, N. Dyulgerov, D. Dimova
- Haematological and serum biochemical indices of broiler chickens fed raw sickle pod (*Senna obtusifolia*) seed meal** 203
C. Augustine, I.D. Kwari, J.U. Igwebuike, S.B. Adamu
- Prey size selectivity of pikeperch (*Sander Lucioperca* L.) fed with topmouth gudgeon (*Pseudorasbora Parva* Temminck & Schlegel)** 209
M. Gevezova-Kazakova, M. Yankova, T. Hubenova, A. Zaikov, G. Rusenov
- Influence of organic nitrogen amendment, containing amino acids on the cellulase and xylanase, produced by *Trichoderma* spp. isolates** 213
D. Draganova, I. Valcheva, Y. Kuzmanova, M. Naydenov

Production Systems

- Justification of a method for determining the moment for switching on the level one signaling of filled grain harvester hoppers** 218
G. Tihanov, B. Kolev, K. Trendafilov, N. Delchev, Y. Stoyanov
- Mathematical approaches for assessment and classification of the European Union member states based on the average yield of vegetables for the period 1961-2014** 223
N. Keranova

Present status of <i>Zymoseptoria tritici</i> (<i>Mycosphaella graminicola</i> /Fuckel/ Schroter) of the wheat cultures in the Republic of Macedonia	227

I. Karov, E. Arsov

Agriculture and Environment

Influence of basic agrotechnical activities on the productivity and yield of <i>Triticum monococcum</i> L.	230

S. Stamatov, K. Uzundzhaliyeva, E. Valchinova, G. Desheva, P. Chavdarov, B. Kyosev, T. Cholakov, R. Ruseva, N. Velcheva

Avifauna abundance and diversity in Jos wildlife park, Nigeria	234

B.T. Kwaga, D. Iliya, A. Ali, D. Khobe

Ecological analysis of the flora in the 'Chinarite' protected area - Rodopi municipality, Bulgaria	240

L. Dospatliev, M. Lacheva

Product Quality and Safety

Food emulsions with amidated pectin from celery (<i>Apium graveolens</i> var. <i>rapaceum</i> D. C.) tubers	246

Iv. Petrova, N. Petkova, M. Ognyanov, Ap. Simitchiev, M. Todorova, P. Denev

Sensory and instrumental texture analysis of Bulgarian commercial pates	251

M. Tonchev, T. Atanasov, A. Todorova, Ts. Atanasova, N. Shtrankova, M. Momchilova G. Zsivanovits

Short Communication

Influence of elevated platform (wire-mesh or wooden) in the cage on domestic rabbit (<i>Oryctolagus cuniculus</i>) activity	257

S. Peeva, E. Raichev, D. Georgiev, A. Stefanov

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors
The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings: Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows, IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 9, Number 3
September 2017

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz