ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 9, Number 3
September 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate on author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc., AGRIS (FAO) and DOAJ.

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No DNP 05-21/20.12.2016, financed from Fund 'Scientific Research' grant Bulgarian scientific Periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 9, Number 3
September 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Product Quality and Safety

Food emulsions with amidated pectin from celery (*Apium graveolens* var. *rapaceum* D.C.) tubers

Iv. Petrova¹, N. Petkova^{2*}, M. Ognyanov³, Ap. Simitchiev⁴, M. Todorova², P. Denev²

¹Department of Catering and Tourism, Faculty of Economic, University of Food Technologies, 4002 Plovdiv, Bulgaria

²Department of Organic Chemistry and Inorganic Chemistry, Technological Faculty, University of Food Technologies, 4002 Plovdiv, Bulgaria

³Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Laboratory of Biologically Active Substances, 4000 Plovdiv, Bulgaria

⁴Department of Machinery for Food Industry, Technical Faculty, University of Food Technologies, 4002 Plovdiv, Bulgaria

(Manuscript received 29 May 2017; accepted for publication 22 August 2017)

Abstract. Hydrocolloids, especially polysaccharides from traditional plant sources and their derivatives possessed significant emulsifying properties. Pectin was isolated from celery tubers by accelerated "green" method for extraction based on ultrasonic irradiation. Further chemical modification of celery pectin was performed with 4 mol/L NH_3 . The amidated celery pectin was obtained with the following characteristics: the degree of esterification (DE) 31%, the degree of amidation (DA) 16%, degree of acetylation (DAc) 2% and anhydrouronic acid content (AUAC) 68%. This modified pectin was incorporated in preparation of model 30, 40 and 50% oil-in-water emulsions. The effect of amidation of celery pectin on the stability of emulsions was investigated. The results showed that amidation increased the emulsifying properties of pectic polysaccharides. It affected also the rheological characteristics of model emulsion. The current study demonstrated preparation of emulsion with low-caloric amidated pectin as proper alternative to the traditional emulsifiers.

Keywords: celery, amidated pectin, IR-FT spectra, emulsions, rheology

Introduction

The celery (*Apium graveolens* var. *rapaceum* D. C.) tubers are evaluated as a rich source of pectin (Siddiqui, 1990; Hansen et al., 2001; Petrova et al., 2014). Our previous studies demonstrated significant yield of pectin from celery – 30% dw (Petrova et al., 2014; Murdzheva et al., 2016). Celery pectic polysaccharides are characterized with low degree of acetylation around 4% in comparison of sugar beet and sunflower pectin and anhydrouronic acid content (AUAC) 75% (Hansen et al., 2001). Our earlier investigations showed that the main pectic polysaccharide fraction from celery tubers were with average degree of esterification (DE) and good gelling strength (Denev et al., 2004; Petrova et al., 2014; Murdzheva et al., 2016).

Pectic substances are hydrocolloids widely used in food industry as a gelling agent, thickener, texturizer, emulsifier and emulsion stabilizer (Rolin and Vries, 1990; Sakai et al., 1993; Ngouemazong et al., 2015). The important characteristics of commercial pectin used as food additives is 65% galacturonic acid-based content, which may be present as free acid, methyl ester, in amidated pectin or acid amide (May, 2000).

The strong correlation was found between rheological properties (emulsion stability) and DE of the pectin (Popova et al., 1993; Einhorn-Stoll et al., 1996). Moreover, the protein moiety, feruloyl and acetyl groups, play a major role in pectin emulsifying activities, while the emulsion-stabilizing properties of the polymer are controlled by the homogalacturonan domain and the neutral sugar side chains of the rhamnogalacturonan-I (Ngouemazong et

al., 2015).

Amidated pectin are low methoxyl pectin in which some of the carboxylic acid groups are amidated. The chemical modification of pectin (ammonolysis, amidation, transesterification) is relatively easy as it does not require extreme conditions and the amide bond is sufficiently resistant to hydrolysis by acids or alkali (Kratchanov et al., 1989; Mishra et al., 2008). Moreover, pectin can be modified through amidation by the partial replacement of methoxyl groups with amide groups, following a treatment of pectin with ammonia or primary aliphatic amines (generally in alcohol media) (Kratchanov et al., 1989; Racape et al., 1989; Sinitsya et al., 2000).

The degree of amidation (DA), or the percentage of carboxyl group in the amide form (typically around 17%), allows this type of pectin to tolerate more variation in calcium content and to possess a higher degree of thermoreversibility (Matia-Merino et al., 2004). The influence of amidated citrus pectin on the rheological and microstructural properties of sodium caseinate gels (2% w/v) (Matia-Merino et al., 2004) were studied. In addition, the aggregation of whey proteins with amidated pectin and the influence of this aggregation on emulsifying properties were evaluated (Kováčová et al., 2009). Low-methyl esterified amidated celery pectin was used successfully for the preparation of fermented dairy products enriched with soluble dietary fibers (Vlaseva et al., 2016). Until now the amidated celery pectin was not investigated in details for their emulsion-stabilizing properties in model oil-in water emulsions.

The aim of the current study was to modified celery pectin by amidation and then to evaluate the emulsifying potential of characterized amidated celery pectin. The replacement of high-

* e-mail: petkovanadejda@abv.bg

caloric emulsifiers (lecithin and glycerol monostearate) with modified natural dietary fiber will allow the marketing of food emulsions with properties similar to functional foods.

Material and methods

Plant material

Fresh celery (*Apium graveolens* var. *rapaceum* D. C.) tubers were purchased from the local market in Plovdiv. The plant material was washed with tap water, cleaned from impurity, peeled, coarsely ground and dried at 40°C.

All reagents were analytical grade and used without further purification. pH values were monitored by a pH meter 7110 WTW (Germany).

Isolation of pectin from celery tubers

The dried celery tubers were finely ground in laboratory homogenizer and preliminary washed with 70% ethanol acidified with 2% hydrochloric acid to obtain alcohol insoluble solids (AIS). The pectin from celery AIS was isolated by ultrasound-assisted extraction with ammonium oxalate as previously described (Petrova et al., 2014).

Amidation of celery pectin

The amidation reaction was performed in heterogeneous media. Isolated celery pectin was soaked with 120 cm³ 70% (v/v) aqueous ethanol. Then 4 mol/L NH₃ in 70% (v/v) ethanol with temperature 4°C was added to the mixture (Kratchanov et al., 1989). The amidation reaction was carried out under conventional conditions (at 4°C for 180 min) with periodically shaking. Then 5% HCl/70% C₂H₅OH was added to adjust pH to 3.0 and vacuum filtration was done. Finally, the product was washed with 70% (v/v) aqueous ethanol to the neutral pH value. Amidated celery pectin (ACP) was dried in vacuum oven.

Characterization of celery pectin and its amides

Degree of esterification, degree of amidation and anhydrouronic acid content were determined by titration methods (Food Chemical Codex, 2004). Degree of acetylation was determined by the hydroxamic acid reaction, using β -D-glucose pentaacetate as a standard (McComb and McCreedy, 1957). Protein content was analyzed by Bradford's method with bovine serum albumin as a standard (Bradford, 1976).

FT-IR spectroscopy

FT-IR spectra of celery pectin and its amide derivatives were recorded on a Nicolet FT-IR Avatar (Thermo Science, USA) spectrometer in KBr pellets in the range 4000–400 cm⁻¹ at resolution 4 cm⁻¹ with 132 scans. The pellets were prepared with 3 mg of the sample in 100 mg of KBr. The absorption was reported in wavenumbers (cm⁻¹).

Emulsions

Model oil-in-water emulsions were obtained by following several steps. Amidated celery pectin was dissolved in hot double-distilled water (40°C) with initial stirring until a transparent water-pectin aqueous phase was obtained. Commercially available sunflower oil, which is not further processed, was used as the oil phase (OP). Emulsification was performed on a laboratory homogenizer POLYTRON® Kinematica GmbH PT-G 45/80 (Switzerland) for 30 s at 523 s⁻¹ and 20°C. Emulsion stability (ES), index of oil stability (IOS) and index of water stability (IWS) were determined in accordance with Tharp (1982).

Rheological measurements

The effective viscosity and shear stress of model oil-in-water emulsions were measured at various shear rates via digital Brookfield viscometer (RV-DV II+ Pro). Effective viscosity and shear stress were calculated by specialized software Rheocalc 32, the graphical correlation between shear stress and shear rate was automatically drawn. The rheological model was selected after approximation and evaluation of the correlation coefficient - the highest value was chosen. By using the equation of the selected rheological model the effective viscosity was calculated.

Results and discussion

Characteristic of celery pectin and its amides

Pectin was successfully extracted from AIS from celery tubers using ammonium oxalate and ultrasound irradiation 45 kHz. The obtained celery pectin was characterized as average methoxylated ones and the results were summarized in Table 1. The yields coincided with our previous studies (Petrova et al., 2014; Murdzheva et al., 2016). The degree of acetylation (2%) of celery pectin was higher than citrus pectin and lower than sugar beet pectin (Leroux et al., 2004). Protein content not exceeded 2%. AUAC was relatively high 69% and coincided with these of citrus pectin (Georgiev et al., 2012). After amidation it was found that protein content, DAc and AUAC remained unchanged, while DE decreased to 31%. The degree of amidation was 16% which characterized celery pectin as low-methoxylated amidated pectin.

The FT-IR spectra of celery pectin indicated broad bands at 3402 cm⁻¹ that was due to the stretching vibrations of –OH groups. The band at 2932 cm⁻¹ was assigned to C–H stretching vibration of CH₂ groups. The bands at 1460 and 1377 cm⁻¹ were due to –CH₂ scissoring and –OH bending vibration, respectively. The signal at 1749 cm⁻¹ was typical for C=O stretching vibration of methyl esterified carbonyl groups and the band at 1630 cm⁻¹ corresponded to the absorption of the carboxylate anions. The band at 1017 cm⁻¹ suggested –CH–O–CH– stretching. The band at 920 cm⁻¹ was assigned to ρ (CH₃) methyl group in complex ester. The absorption at 888 cm⁻¹ and 760 cm⁻¹ were typical for α -D-Galp in ⁴C, conformation.

In the FT-IR spectra of amidated celery pectin (ACP) typical

Table 1. Characteristics of celery pectin and its amide

Substance	Yield, %	AUAC, %	DE, %	DAc, %	DA, %	Protein content, %
Celery pectin	30	69	49	2	-*	2
Celery amidated pectin	99	68	31	1	16	2

AUAC - anhydrouronic content, DE - degree of esterification, DAc - degree of acetylation, DA – degree of amidation, *- absence of amidation

Table 2. Stability of model emulsions with amidated pectin

Amidated pectin, %	Separate phases, %		ES, %	IOS	IWS
	Oil	Water			
Emulsions with 30% OP					
0.4	2.0	52.0	46.0	0.9	0.3
0.6	0.0	40.0	60.0	1.0	0.4
0.8	0.0	0.0	100.0	1.0	1.0
1.0	0.0	0.0	100.0	1.0	1.0
Emulsions with 40% OP					
0.4	0.0	18.0	82.0	1.0	0.7
0.6	0.0	6.3	93.7	1.0	0.9
0.8	0.0	0.0	100.0	1.0	1.0
1.0	0.0	0.0	100.0	1.0	1.0
Emulsions with 50% OP					
0.4	0.0	6.0	96.0	1.0	0.9
0.6	0.0	0.0	100.0	1.0	1.0
0.8	0.0	0.0	100.0	1.0	1.0
1.0	35.0	20.0	45.0	0.3	0.6

ES - emulsion stability, IOS - index of oil stability, IWS - index of water stability and OP – oil phase

bands for amide groups were observed. The bands at 3400 cm^{-1} typical for $\nu(\text{OH})$ overlapped with stretching vibration for $\nu(\text{NH})$ from amide groups. In addition, the bands at 1650 cm^{-1} and 1681 cm^{-1} were assigned to amide $\text{C}=\text{O}$ stretching vibration. The bands around 1629 cm^{-1} belong to mixed mode $\delta(\text{N-H})$ and $\nu(\text{C-N})$ ($\text{N}-\text{C}=\text{O}$). Other typical bands for pectin structure remained unchanged after amidation. Therefore, the FT-IR results confirmed appearance of new amide group in celery pectin spectra. The data were in accordance with some previous reports for amidation of pectin (Mishra et al., 2008; Murdzheva et al., 2016).

Emulsion stability

This work revealed the possibility to design emulsion products enriched in dietary fibers from celery. In the present study, the emulsifying and stabilizing properties of ACP were investigated. In a series of model emulsions with increasing concentration of the amidated pectin (0.6, 0.8, 0.9% of aqueous phase) and different oil phase (30, 40, 50% of total emulsion), the stability of the dispersion systems was determined by the centrifugation method.

The results revealed (Table 2) that increasing concentration of amidated celery pectin and oil phase improved emulsion stability. In the 30% oil phase of model emulsions, the stability was improved by 24% with an increasing amidated pectin content of 0.4 to 0.6%. The stability of the emulsions was further enhanced by increasing the amount of oil phase. With an increase in the oily phase from 30% to 40%, thirty-six percent improvement in emulsion stability prepared with 0.6% pectin was observed. All model emulsions with 0.8% amidated celery pectin content were stable, regardless of the amount of oil phase ($\text{ES}=100\%$). However, in the current study it was not possible to obtain long stable emulsions with 1% ACP in a 50% oil phase, as well as in our previous reports for emulsions with 0.4% celery pectin and 30% oil phase. Unlike non-amidated celery pectin (Petrova et al., 2014) at 50% increasing oily phase it is not possible to emulsify 1% of the amidated pectin. Other study also demonstrated good emulsion stability (from 62.1–79.4%) of sugar

beet pectin (0.5 %) with galacturonic acid content (60.2–77.8%) (Ma et al., 2013). But obtained with only 0.4 % amidated pectin 50 % emulsion showed the best emulsion stability (94 %) at this concentration (Table 2). Therefore, amidation of celery pectin resulted in improvement in the emulsifying and stabilizing properties of pectin by reducing the concentration of pectin used to obtain stable emulsions. In addition, it is known that the carbohydrate moiety also stabilizes by steric and viscosity effects in the aqueous phase (Leroux et al. 2003).

Rheology

Rheology describes the deformation of a fluids under the influence of stress. Rheograms are used to describe rheological properties for the fluids. They were constructed by plotting shear stress (τ) as a function of the shear rate (Figure 1).

Effective viscosity is a rheological indicator that shows internal friction and structural-dynamic equilibrium between system components. The relationship between shear stress and shear rate of the model emulsions at different concentration of amidated pectin were shown (Figure 1).

The increased concentration of amidated pectin led to the increasing effective viscosity and change in the rheological behavior of the emulsion system. Experimental data revealed that all model emulsions had non-Newtonian behavior - not linear relationship between shear stress and shear rate (Figure 2). Higher values of effective viscosity with increasing pectin concentration were due to better dispersion of the oil and structuring of the aqueous phase.

The change in the effective viscosity values with increasing shear rate allowed to determine the rheological behavior of the system (Figure 3).

The viscosity decreased with increasing shear stress (pseudo plasticity). Rheological behavior can be described by the Herschel-Bulkley model (equation 1) of a non-Newtonian fluid (Table 3). In this non-Newtonian character is more pronounced when emulsions with a high ratio of oil phase and increases with increasing

Figure 1. Rheograms for 40% oil phase at different concentration of amidated celery pectin (0.6; 0.8; 1%) at 20°C

Figure 2. Dependence of the plastic viscosity on a share rate for 40% oil phase at different concentration of amidated pectin (0.6; 0.8; 1%) at 20°C

Figure 3. Viscosity versus shear rate of oil-in-water emulsions with 0.8% amidated celery pectin and different oil phase (OP) concentrations

Table 3. Rheological characteristics of model emulsions

Model emulsion	τ_0 , Pa	K, Pa.s ⁿ	n
30 % OP v/v			
0.8% ACP	4.34	99.9	1.05
1.0% ACP	9.68	192.0	1.06
40 % OP v/v			
0.6% ACP	4.34	117	1.01
0.8% ACP	8.57	111.2	1.11
1.0% ACP	40.7	450	1.4
50 % OP v/v			
0.6% ACP	0.77	70.49	0.46
0.8% ACP	48.5	197	1.04

ACP – amidated celery pectin, OP – oil phase, τ_0 – yield stress, k – consistency index, n – flow behaviour index

amount of pectic polysaccharides in them.

$$\tau = \tau_0 + k \cdot \dot{\gamma}^n, \text{ Pa} \quad (1)$$

where: τ – shear stress, Pa; τ_0 – yield stress, Pa; k – consistency index, Pa.sⁿ; n – flow behaviour index; $\dot{\gamma}$ – shear rate, s⁻¹.

When the value of n (flow behavior index) is close to 1 the rheological behavior tends to pass from a shear thinning to a shear thickening fluid. The fluid acts as a shear thickening fluid when n is above 1 (Table 3).

The emulsifying properties of amidated celery pectin depended on many factors, especially DE, DAc, DA and protein content. According to Nguemazong et al. (2015) the protein moiety, feruloyl, and acetyl groups, play a major role in pectin emulsifying activities. Even though, the acetyl groups are more common in the pectin fraction which adheres the oil phase, their presence is not an absolute requirement with respect to the emulsifying capacity (Leroux et al., 2003). Protein (2% in our pectins) additionally acts as the anchor for the oil droplets to stabilize emulsions (Funami et al., 2011). Therefore, the better emulsifying properties of amidated

celery pectin could be explained with the presence of many functional groups and small protein content in polymer. They can interact with oil and water phase in emulsion system. Additional hydrophobic interaction could stabilize the obtained emulsions. Not at last, the type of oil phase strongly affects the stability of the prepared emulsions (Nguemazong et al., 2015).

Conclusion

The current study demonstrated the potential application of amidated celery pectin for preparation of emulsions enriched in dietary fibers with potential healthy effect. Incorporation of modified celery pectin improved emulsion stability. Moreover, the presence of modified dietary fibers in emulsion systems instead of traditional emulsifiers allowed preparation of low-calorie product with 30% sunflower oils. The replacement of natural emulsifiers containing saturated fatty acids (lecithin and glycerol monostearate) with modified natural dietary fiber (amidated celery pectin) was expected to have a potential beneficial health effect on the cardiovascular and

gastrointestinal system. Experimental data revealed that all model emulsions have non-Newtonian behavior - not linear relationship between shear stress and shear rate. Rheological properties of oil-in-water emulsions prepared with amidated celery pectin demonstrated the potential of application of this polysaccharide not only in mayonnaise but in other food emulsion products (dressings, salad dressings, etc.).

References

- Bradford M**, 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72, 248-254.
- Denev P, Kirtchev N and Panchev I**, 2004. Kharakteristika pectinovih veshtestva v othodakh pri proizvodstve sokov, II Mezhdunarodnoya Konferenciya Lesnie Biologicheski Aktivni Resursi, September 21-23, Habarovsk, Russia (Ru).
- Einhorn-Stoll D, Glasenapp N and Knuzek H**, 1996. Modified pectin in whey protein emulsions. *Food/Nahrung*, 40, 60-67.
- Funami T, Nakauma M, Ishihara S, Tanaka R, Inoue T and Phillips G**, 2011. Structural modifications of sugar beet pectin and the relationship of structure to functionality. *Food Hydrocolloids*, 25, 221-229.
- Georgiev Y, Ognyanov M, Yanakieva I, Kussovski V and Kratchanova M**, 2012. Isolation, characterization and modification of citrus pectins. *Journal of BioScience and Biotechnology*, 1, 223-233.
- Hansen K, Thuesen A and Soderberg J**, 2001. Enzyme assay for identification of pectin and pectin derivatives, based on recombinant pectate lyase. *Journal of AOAC International*, 84, 1851-1854.
- Institute of Medicine**, 2003. *Food Chemicals Codex*. Fifth edition, The National Academies Press, Washington, USA.
- Kováčová R, Synytsya A and Štětina J**, 2009. Characterisation of whey proteins-pectin interaction in relation to emulsifying properties of whey proteins. *Czech Journal of Food Sciences*, 27, S4-S8.
- Kratchanov Chr, Denev P and Kratchanova M**, 1989. Reaction of apple pectin with ammonia. *International Journal of Food Science and Technology*, 24, 261-267.
- Leroux J, Langendorff V, Schick G, Vaishnav V and Mazoyera J**, 2003. Emulsion stabilizing properties of pectin. *Food Hydrocolloids*, 17, 455-462.
- Ma S, Yu S, Zheng X, Wang X, Bao Q and Gyo M**, 2013. Extraction, characterization and spontaneous emulsifying properties of pectin from sugar beet pulp. *Carbohydrate Polymers*, 98, 750-3.
- Matia-Merino L, Lau K and Dickinson Er**, 2004. Effects of low-methoxyl amidated pectin and ionic calcium on rheology and microstructure of acid-induced sodium caseinate gels. *Food Hydrocolloids*, 18, 271-281.
- May C**, 2000. Pectin. In: *Handbook of Hydrocolloids* (eds. G.O. Phillips and P.A Williams), CRC, New York, USA, pp. 169-188.
- McComb E and McCready R**, 1957. Determination of acetyl in pectin and in acetylated carbohydrate polymers (hydroxamic acid reaction). *Analytical Chemistry*, 29, 819-821.
- Mishra R, Datt M, Pal K and Banthia A**, 2008. Preparation and characterization of amidated pectin based hydrogels for drug delivery system. *Journal of Materials Science-Materials in Medicine*, 19, 2275-2280.
- Murdzheva D, Petkova N, Vasileva I, Todorova M, Ognyanov M, Ivanov I and Denev P**, 2016. Accelerated modification of low-methoxylated celery pectin, 12th International Conference on Polysaccharides-Glycoscience, October 19-21, Prague, Czech Republic.
- Ngouemazong E, Christiaens S, Shpigelman A, Van Loey A and Hendrickx M**, 2015. The emulsifying and emulsion-stabilizing properties of pectin: a review. *Comprehensive Reviews in Food Science and Food Safety*, 14, 705-718.
- Petrova I, Petkova N, Kyobashieva K, Denev P, Simitchiev Ap, Todorova M and Dencheva N**, 2014. Isolation of pectic polysaccharides from celery (*Apium graveolens* var . *rapaceum* D.C.) and their application in food emulsions. *Turkish Journal of Agricultural and Natural Sciences*, 1, 1818-24.
- Popova M, Kratchanov Chr and Panchev I**, 1993. An opportunity for simultaneous purification and deesterification of pectic substances. *Food Hydrocolloids*, 7, 103-112.
- Racape E, Thibault J, Reitsma J and Pilnik W**, 1989. Properties of amidated pectin. II. Polyelectrolyte behavior and calcium binding of amidated pectin and amidated pectic acids. *Biopolymers*, 28, 1435-48.
- Rolin C and De Vries J**, 1990. Pectin. In: *Food Gels*, (ed. P. Harris), Elsevier, Amsterdam, The Netherlands.
- Sakai T, Sakamoto T, Hallaert J and Vandamme EJ**, 1993. Pectin, pectinase, and protopectinase: production, properties, and applications. *Advances in Applied Microbiology*, 39, 213-294.
- Siddiqui I**, 1990. Studies on vegetables. Investigation of water, oxalate, and sodium hydroxide soluble celery (*Apium graveolens*) polysaccharides: pectic polysaccharides. *Journal of Agricultural and Food Chemistry*, 38, 70-74.
- Sinitsya A, Copikova J, Prutyaynov V, Skoblya S and Machovic V**, 2000. Amidation of highly methoxylated citrus pectin with primary amines. *Carbohydrate Polymers*, 42, 359-368.
- Tharp B**, 1982. The effect of certain colloid emulsifier blends and processing procedures on emulsion stability. *Progress in Food and Nutrition Science*, 6, 209-220.
- Vlaseva R, Ivanova M, Hadjikinova M, Hadjikinov D, Dobрева V, Petkova N, Denev P and Dimitrov D**, 2016. Comparative characterization of possibilities of using low-esterified and amidated pectin in fermented dairy products. *Journal of Microbiology, Biotechnology and Food Sciences*, 6, 649-651.

Reviews

- Problems and achievements of cotton (*Gossypium Hirsutum* L.) weeds control** 179
T. Barakova, G. Delchev
- Achievements and problems in the weed control in grain sorghum (*Sorghum Bicolor* Moench.)** 185
G. Delchev, M. Georgiev

Genetics and Breeding

- Parthenogenetic responsiveness of sunflower hybrid combinations with expressed tolerance to herbicides** 190
M. Drumeva, P. Yankov
- In vitro propagation of oil-bearing rose (*Rosa damascena* Mill.)** 194
V. Badzhelova

Nutrition and Physiology

- Variation in the chemical composition and physical characteristics of grain from winter barley varieties** 198
B. Dyulgerova, N. Dyulgerov, D. Dimova
- Haematological and serum biochemical indices of broiler chickens fed raw sickle pod (*Senna obtusifolia*) seed meal** 203
C. Augustine, I.D. Kwari, J.U. Igwebuike, S.B. Adamu
- Prey size selectivity of pikeperch (*Sander Lucioperca* L.) fed with topmouth gudgeon (*Pseudorasbora Parva* Temminck & Schlegel)** 209
M. Gevezova-Kazakova, M. Yankova, T. Hubenova, A. Zaikov, G. Rusenov
- Influence of organic nitrogen amendment, containing amino acids on the cellulase and xylanase, produced by *Trichoderma* spp. isolates** 213
D. Draganova, I. Valcheva, Y. Kuzmanova, M. Naydenov

Production Systems

- Justification of a method for determining the moment for switching on the level one signaling of filled grain harvester hoppers** 218
G. Tihanov, B. Kolev, K. Trendafilov, N. Delchev, Y. Stoyanov
- Mathematical approaches for assessment and classification of the European Union member states based on the average yield of vegetables for the period 1961-2014** 223
N. Keranova

Present status of *Zymoseptoria tritici* (*Mycosphaella graminicola* /Fuckel/ Schroter) of the wheat cultures in the Republic of Macedonia 227

I. Karov, E. Arsov

Agriculture and Environment

Influence of basic agrotechnical activities on the productivity and yield of *Triticum monococcum* L. 230

S. Stamatov, K. Uzundzhaliyeva, E. Valchinova, G. Desheva, P. Chavdarov, B. Kyosev, T. Cholakov, R. Ruseva, N. Velcheva

Avifauna abundance and diversity in Jos wildlife park, Nigeria 234

B.T. Kwaga, D. Iliya, A. Ali, D. Khobe

Ecological analysis of the flora in the 'Chinarite' protected area - Rodopi municipality, Bulgaria 240

L. Dospatliev, M. Lacheva

Product Quality and Safety

Food emulsions with amidated pectin from celery (*Apium graveolens* var. *rapaceum* D. C.) tubers 246

Iv. Petrova, N. Petkova, M. Ognyanov, Ap. Simitchiev, M. Todorova, P. Denev

Sensory and instrumental texture analysis of Bulgarian commercial pates 251

M. Tonchev, T. Atanasov, A. Todorova, Ts. Atanasova, N. Shtrankova, M. Momchilova G. Zsivanovits

Short Communication

Influence of elevated platform (wire-mesh or wooden) in the cage on domestic rabbit (*Oryctolagus cuniculus*) activity 257

S. Peeva, E. Raichev, D. Georgiev, A. Stefanov

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors
The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings: Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows, IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 9, Number 3
September 2017

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz