ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 9, Number 4
December 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate on author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc., AGRIS (FAO) and DOAJ.

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No DNP 05-21/20.12.2016, financed from Fund 'Scientific Research' grant Bulgarian scientific Periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 9, Number 4
December 2017

AGRICULTURAL SCIENCE AND TECHNOLOGY

2017

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Production Systems

Lucrative status of improved dual purpose cowpea (*Vigna unguiculata* L., Walp) in Damboa, Borno State, North-Eastern Nigeria

B.H. Gabdo*

Department of Agricultural Economics and Extension, Adamawa State University, Mubi, Adamawa State, Nigeria

(Manuscript received 22 May 2017; accepted for publication 27 October 2017)

Abstract. This study divulged the cost-returns analysis between improved dual purpose cowpea (IDPC) varieties and other non-IDPC/local cowpea varieties in Azir, Damboa town, Kuboa and Sabongari villages of Damboa local council, Borno State, Nigeria. Socio-economic stratification of the villages into Low Population Low Market (LPLM) and Low Population High Market (LPHM) on the basis of human demographic profile and proximity to wholesale market were the criteria adopted for selecting the villages for this study. A total of 150 cowpea farmers responded to the questionnaire instrument; administered between December, 2006 and February, 2007. Results of the budgetary techniques used show that adopters of IDPC varieties generate \$386.01ha⁻¹ as net income as against \$248.37ha⁻¹ for the non-IDPC adopters; proportional to 55.42% rise in net income per hectare of cowpea resulting from the impact factor of IDPC adoption alone. Similarly, farmers in Sabongari village realized the highest net income of \$491.99ha⁻¹ as against Azir the lowest net income of \$247.21ha⁻¹ across the four villages owing to demand factor; interplay between human population and proximity to wholesale market; both in favour of Sabongari but none in favour of Azir. In terms of ratios, farmers in Sabongari were adjudged as more lucrative across adopters' category, socio-economic domain and villages respectively owing to higher return on sale ratio and lower operating ratio. Azir and Sabongari villages were found to exhibit potentiality for comparative cost advantage and comparative sales advantage respectively. Thus, the research concludes that an expansion path for cowpea is feasible if cultivation and sales are premised on cost and sales advantages.

Keywords: lucrative, improved dual purpose cowpea, socio-economic domain, return on sales ratio, operating ratio

Introduction

Improved planting materials, by generalization, agricultural genetics is a veritable weapon for global fight against unsustainability and food insecurity; it ensures remarkable productivity increase, income rise and augments the welfare of farming families. The cowpea industry is no exception having gone through series of quality improvements in seed over several decades. The improved dual purpose cowpea (IDPC) varieties, characterized by duality effect of both higher grain and higher fodder yields is one of the many improvements attained in the cowpea enterprise.

Nigeria, the highest world producing nation of cowpea, produces 1.69 million tonnes per annum; representing 56.3% of world production and generates annual revenue of US\$650 million (Ortiz, 2003; Langyintuo et al., 2005). On global index, cowpea area under cultivation stands at 12.5 million hectares yielding an estimated 3 million tonnes of cowpea grain annually; 64% (1.92 million tonnes) of the production share emanates from West and Central Africa (Quin, 1997; Singh et al., 1997). Substantial evidence abounds on productivity rise in Nigeria. Ortiz (1998) concur with the foregoing that cowpea production in the country rose by over 400% between 1961 and 1995; with an average rise in grain yield of 0.48tha⁻¹ (Quin, 1997) to 0.68tha⁻¹ for the year 2005 (FAO, 2007). The justification for these productivity rises is certainly not preclusive of the series of advances in cowpea research spanning many decades in Nigeria; particularly, the development of high yielding

planting materials such as the IDPC varieties among other yield enhancing factors.

The purpose of this study was to make the cost-returns analysis between improved dual purpose cowpea (*Vigna unguiculata* L., Walp) varieties and other non-IDPC/local cowpea varieties in some areas in Damboa, Borno State, Nigeria.

Material and methods

Study area and sampling method

Damboa is one of the 27 local government areas of Borno State, Nigeria sandwiched by Kaga, Konduga, Gwoza, Chibok, and Biu local governments areas of the State with human population of 231 573 (NPC, 2006) inhabiting the 6.219 km² of the area. Damboa lies in semi arid region with much variability and unpredictability in rainfall patterns that often last between May and October; which supports the cultivation of low rainfall requiring crops such as wheat, sorghum, maize and cover crops like cowpea and groundnuts. Damboa was purposively selected for this study being the only local government area in the Sudan Savanna of Borno State where IDPC varieties were released to farmers by IITA in 2004. Azir, Damboa, Kimba, Kuboa, Nzuda, Mungule and Sabongari were identified as the production centres for IDPC varieties in the local government. These centres were stratified in to 2 socio-economic domains (LPLM and LPHM) based on human population size and accessibility to wholesale market, viz:

- LPLM (Low Population-Low Market): Named Socio-economic

* e-mail: bashirgabdo@gmail.com

Domain (SEDOM) 1. It has low population (<20,000 inhabitants) and poor accessibility to whole sale market (market tension between 1 and 5). Except Sabongari, all other aforementioned villages satisfied this criterion.

- LPHM (Low Population-High Market): Named Socio-economic Domain (SEDOM) 2. It composes of low population (<20,000 inhabitants) and good access to wholesale market (market tension between 6 and 10). Of the production centres or villages, only Sabongari village that fits this criterion.

On the whole, based on the socio-economic categorization, random sampling was used to select four villages Azir, Damboa Town, Kuboa and Sabongari and a total of 150 respondents were administered with questionnaire materials based on proportionality; 25 each from Azir and Kuboa and 50 each from Damboa Town and Sabongari. Data collection lasted between December, 2006 and February, 2007.

Analytical methods

Net Return-Profit Model. The net-return profit model was used to estimate the costs and returns in the IDPC and non-IDPC varieties. Several models for estimating profitability in agriculture exist. Amaza (2000) strengthened the use of gross margin analysis; gross returns less gross variable cost as a proxy measure of profitability. Olukosi and Ogunbile (2005) emphasized the use of gross margin especially in peasant African farming where fixed input is highly negligible. Onyeagocha et al. (2010) stated that the net returns range between $-\infty \leq \Pi \leq +\infty$ and can be estimated using the two equations below:

$$GM = \sum P_i Q_i - \sum P_j X_j \quad (1)$$

$$NR = GM - TFC \quad (2)$$

where: $P_i Q_i$ = Value of Output, $P_j X_j$ = Value of Variable Input, GM = Gross margin, TFC = Total Fixed Cost and NR = Net Return.

The aforementioned equations can be condensed to give a compact notation as viz:

$$\Pi = \sum_{i=1}^n P_i Q_i - \left[\sum_{j=1}^n P_j V X_j + \sum_{j=1}^n P_j F X_j \right] \quad (3)$$

Furthermore, the model can be extended to accommodate more variables as possible:

$$\Pi = P_1 Q_1 + P_2 Q_2 + \dots + P_n Q_n - [P_1 V X_1 + P_2 V X_2 + \dots + P_n V X_n + P_1 F X_1 + P_2 F X_2 + \dots + P_n F X_n] \quad (4)$$

In this study the model was narrowed to capture specific variables as applicable in the Damboa cowpea research as below:

$$\Pi = P_1 Q_1 - [P_1 V X_1 + P_2 V X_2 + P_3 V X_3 + P_4 V X_4 + P_5 V X_5 + P_1 F X_1] \quad (5)$$

where: $P_1 Q_1$ = Unit price of a bag of cowpea multiplied by number of cowpea bags harvested, $P_1 V X_1$ = Unit price of a kg of seed multiplied by number of kg of seed planted, $P_2 V X_2$ = Unit price of a bag of fertilizer multiplied by number of bags of fertilizer applied, $P_3 V X_3$ = Unit price of a litre of insecticides multiplied by number of litres of insecticides sprayed, $P_4 V X_4$ = Unit price of a man day of labour multiplied by number of maydays of labour utilized, $P_5 V X_5$ = Unit price of other cost or miscellaneous cost multiplied by number of such costs incurred, $P_1 F X_1$ = Price paid as land tax or depreciation of land

Return on Sale Ratio. This ratio shows how large an operating

margin the enterprise has on its sales. The ratio divides the net income by the revenue and expressed as a percentage.

$$R.S = \frac{N.I}{R} \times 100\% \quad (6)$$

where: $R.S$ = Return on Sale, $N.I$ = Net Income, R = Revenue.

Operating Ratio. This ratio determines the level of managerial control on operating costs. It divides the operating expenses by the revenue and expressed as a percentage by formula:

$$O.R = \frac{O.E}{R} \times 100\% \quad (7)$$

where: $O.R$ = Operating Ratio, $O.E$ = Operating Expenses, R = Revenue.

Results and discussion

Summary of farmers' socio-economic attributes

The summary of socio-economic attributes in Table 1 indicates that majority (74%) of the farmers had between 41 and 60 years of age. The preponderance of this age category may not be unconnected with their higher demand for family upkeep than other age categories. This age category further indicates that the farmers were youthful adults who are expected to be sound in reasoning and rational in their farms' decision making, converse to the relatively old age category who were a minority. The poor participation by the young group (below 40 years) in cowpea cultivation may be inferred on their negative attitudes towards agriculture for white collar jobs and their relatively less responsibility for family upkeep. Educationally, 45% of the farmers had no formal training even those with formal education mostly at lower levels. The significance of education in adoption process as a factor that influences adoption cannot be underestimated. Amaza (2000) stated that farmers are more efficient in information uptake and more productive at higher level of education. Male farmers predominate (70%) owing to the religious and traditional beliefs of the majority of the inhabitants that women are precluded from non-domestic responsibilities for which farming is perceived as one. On marital status, about 95% of the farmers were found to be married, justifying further the position that farmers in the area engage in cowpea production as means of livelihood; to earn income and use for family upkeep. Most of the farmers surveyed had 1-5 years cultivation experience. Studies have shown that years of farming experience prepare farmers to be more acquainted with the production and market situations for the crop. Iheanacho (2000) affirmed that years spent cultivating a particular crop makes farmers more conversant with weather and climatic changes, social and economic factors and factors influencing farming in general. Thus, farmers are better-off in terms of productivity and profitability index with longer years of cultivation. The recent influx of farmers in cowpea production may be ascribed to the motivation associated with the introduction of the IDPC varieties in the area and the new entrants are likely to experience improved productivity over time. More than 80% of the farmers lack membership of any cowpea association. The dearth of farmers' organisation into cowpea association is perhaps a factor responsible for the poor credit status of the farmers. On household size, majority (63.3%) had between 6 and above as family size; considerably large and a very significant factor in promoting IDPC adoption. Most of the farmers were poor (50.7%) on the basis of ownership of small farm sizes, few small ruminants and walking a distance of at least 10km to and from farm sites daily.

Table 1. Summary of farmers' socio-economic attributes

Variable	Farmers	(%)	Variable	Farmers	(%)
Age (Years)			Household Size		
20-40	37	37	1-5	55	36.7
41-60	111	111	6-10	76	50.7
Over 60	2	2	11-15	17	11.3
Sex			16-20	2	1.3
Male	105	105	Wealth Status		
Female	45	45	Poor	76	50.7
Formal Education (Years)			Average	74	49.3
1-7	35	35	Small Ruminants		
8-10	39	39	0-10	18	12.0
15-21	8	8	11-20	67	44.7
No formal education	68	68	Over 20	65	43.3
Farming Experience			Large Ruminants		
1-10	95	95	0-5	135	90.0
11-20	25	25	6-10	15	10.0
Over 20	30	30	Group Membership		
Farmers' Production Category			Yes	29	19.3
Adopters	120	120	No	121	80.6
Non-Adopters	30	30	Farm Distance		
Marital Status			0-10 km	148	98.6
Married	142	142	Over 10 km	2	1.3
Single	5	5			
Divorced	3	3			

Source: Field Survey, 2007

Costs and Returns Analysis

Revenue is practically enhanced by size/scale of production and yield obtained among other factors. Table 2 revealed that IDPC production is profitable; the adopters of the varieties realized a higher income of US\$386.01ha⁻¹ as against US\$248.37ha⁻¹ for the non-adopters (producers of local cowpea varieties or other improved varieties devoid of the duality effect). This income differential attributable to higher grains and fodder yields equates to 55.42% increase in per hectare incomes of cowpea owing to the adoption of IDPC varieties in Damboa. Similarly, Alene et al. (2005) reported an income of US\$289.76ha⁻¹ and US\$124.68ha⁻¹ for adopters and non-adopters of IDPC respectively in their adoption study in Kano, Nigeria. It is evidently glaring that cost components (seeds, fertilizer,

insecticides, miscellaneous cost and labour) are higher in magnitude under the adopters' budget than the non-adopters' owing to the relative-marginal higher cost of IDPC seeds and the accompanying effects it exerts on other cost components.

Table 2 also shows that farmers in SEDOM 2 (Sabongari) made higher per hectare income of US\$491.99 as against US\$401.16 realized by farmers in SEDOM 1 (Azir, Damboa and Kuboa). This may be due to the obvious fact that the domain was classified as a high market access centre, which exerts high influence on market demand. The market access factor is also reflective of the market price, which was higher and led to higher revenue in SEDOM 2. This finding concords with Kristjanson et al. (2005) that socio-economic domain as a factor plays a distinctive role in the market price for the

Table 2. Costs and returns per hectare across adoption category of IDPC varieties in Damboa

Variable	Adopters Nigerian N (US\$)	%	Non-adopters Nigerian N (US\$)	%
Seed	3521.70 (29.79)	10	2662.25 (22.52)	10
Fertilizer	4226.04 (35.75)	12	3194.70 (27.02)	12
Insecticide	3169.53 (26.81)	9	2396.03 (20.27)	9
Labour	22891.05 (193.65)	65	17304.63 (146.39)	65
Miscellaneous cost	1408.68 (11.92)	4	1064.90 (9.01)	4
Total cost	35217.00 (297.92)	100	26622.50 (225.21)	100
Revenue	80847.00 (683.93)		55982.00 (473.58)	
Net income	45630.00 (386.01)		29359.50 (248.37)	

Source: Field Survey, 2007

IDPC grains during marketing. Analysis of net incomes across villages revealed that farmers in Sabongari and Azir villages realized the highest and the lowest incomes respectively. Again, higher market demand facilitated by improved market access and high

human population justifies the case of Sabongari, while dearth of wholesale market accounts for the low revenues in other villages; incomes were found to improve as distance to wholesale market diminishes.

Table 3. Costs and returns per hectare across socio-economic domains of IDPC varieties in Damboa

Variable	SEDOM 1 (LPLM) Nigerian N (US\$)	%	SEDOM 2 (LPHM) Nigerian N (US\$)	%
Seed	3099.18 (26.22)	10	3407.98 (28.83)	11
Fertilizer	3499.18 (29.60)	12	3500.00 (29.61)	11
Insecticide	2699.21 (22.83)	9	2825.99 (23.91)	9
Labour	19494.27 (164.91)	65	20409.95 (172.66)	65
Miscellaneous cost	1199.65 (10.15)	4	1256.00 (10.63)	4
Total cost	29991.18 (253.71)	100	31399.93 (265.63)	100
Revenue	77412.24 (654.87)		89557.69 (757.62)	
Net income	47421.06 (401.16)		58157.76 (491.99)	

Source: Field Survey, 2007

Table 4. Costs and returns per hectare across village level of IDPC varieties in Damboa

Variable	Azir Nigerian N (US\$)	%	Damboa Nigerian N (US\$)	%	Kuboa Nigerian N (US\$)	%	Sabongari Nigerian N (US\$)	%
Seed	2852.45 (24.13)	12	3171.66 (26.83)	10	2819.20 (23.85)	12	3407.98 (28.83)	11
Fertilizer	2844.76 (24.07)	12	3800.00 (32.15)	12	2903.03 (24.56)	12	3500.00 (29.61)	11
Insecticide	2000.00 (16.92)	8	2852.04 (24.13)	9	2177.28 (18.42)	9	2825.99 (23.91)	9
Labour	15409.13 (130.35)	65	20598.65 (174.25)	65	15724.77 (133.02)	65	20409.95 (172.66)	65
Miscellaneous cost	600.00 (5.08)	3	1267.00 (10.72)	4	567.69 (4.80)	2	1256.00 (10.63)	4
Total cost	23706.35 (200.54)	100	31689.35 (268.08)	100	24191.96 (204.65)	100	31399.93 (265.63)	100
Revenue	52928.89 (447.75)		80716.35 (682.82)		53714.70 (454.40)		89557.69 (757.62)	
Net income	29222.54 (247.21)		49027.00 (414.74)		29337.74 (248.18)		58157.76 (491.99)	

Source: Field Survey, 2007

On the whole, it is instructive to note that the higher grain and fodder yields characterizing the IDPC varieties is the rationale for the productivity rise and consequently per hectare incomes. Labour as cost component accounts for the bulk (65%) of total production cost of cowpea. However, Langyintuo et al. (2005) reported labour as accountable for over 70% of total cost of cowpea production. This drop in proportion could be explained by the somewhat remoteness of Damboa and its villages vis a vis low rates of labour. This study further elucidates that farmers in Sabongari and Azir villages made the highest income and the lowest cost of production per hectare respectively. The justification for the highest income was facilitated by the higher market demand due to the presence of an improved wholesale market in Sabongari. Azir, perhaps the most remote of the villages characterized by low labour rates which is also reflective of its least cost characteristics. Thus, Azir and Sabongari villages were found to exhibit criteria for comparative cost advantage and comparative sales advantage respectively in Damboa as entity.

Return on Sale Ratios and Operating Ratios

The return on sale ratio, is an index for expressing the magnitude of profit margin of an enterprise, while the operating ratio as an index for expressing cost of production. Further stated, that each can take a range between 0 and 1 when expressed as a fraction or range between 0 and 100% when expressed as a

percentage (Gabdo et al., 2005). Furthermore, the return on sale ratio that holds: the higher-the better (approaching the best point of 1 or 100%), while the operating ratio holds: the lower-the better (approaching the best point of 0 or 0%).

Table 5. Farmers' return on sales ratios and operating ratios for IDPC cowpea in Damboa

Variable	Operating Ratio	Return On Sale Ratios
Across Adoption Category		
Adopters	0.4356 (43.56%)	0.5644 (56.44%)
Non-Adopters	0.4755 (47.55%)	0.5245 (52.45%)
Across Socio-Economic Domain		
Sedom 1	0.3874 (38.74%)	0.6126 (61.26%)
Sedom 2	0.3506 (35.06%)	0.6494 (64.94%)
Across Villages		
Azir	0.4479 (44.79%)	0.5521 (55.21%)
Damboa	0.3926 (39.26%)	0.6074 (60.74%)
Kuboa	0.4504 (45.04%)	0.5462 (54.62%)
Sabongari	0.3506 (35.06%)	0.6494 (64.94%)

Source: Field Survey, 2007

Table 5 shows ratios across adoption category; farmers in the adopters' category had RS=56.44% and OR=43.56% and RS=52.45% and OP=47.55% for farmers in the non-adopters' category. This shows that the adopters of IDPC had a higher return on sale ratio and lower operating ratio relative to the non-IDPC farmers. This higher percentage of return on sale ratio on the part of the adopters imply a further justification that the adopters of IDPC varieties were more lucrative or profitable than the non-adopters owing to higher yield by the adopters. Similarly, the lower percentage of operating cost on the part of the adopters infers that the cost of production is lower compared to production cost for the non-adopters. Since the IDPC requires less management effort than the non-IDPC varieties explains lower cost of production of the IDPC varieties as against the non-IDPC varieties. Similarly, ratios across socio-economic domains indicates farmers in SEDOM 2 had higher lucrative status (profitability) than those in SEDOM 1 owing to higher return on sale ratio and lower operating ratio. Finally, ratios across villages also shows that farmers in Sabongari village were more lucrative than those in other villages primarily due to higher return on sale ratio and lower operating ratio.

Conclusion

The improved dual-purpose cowpea (IDPC) varieties have been found to be more yielding (both fodder and grain yields) than the non-improved dual purpose cowpea (non-IDPC) varieties with the IDPC farmers gaining 55.42% rise in the income levels over the non-IDPC farmers in Damboa. In view of the poor state of small holder farmers in Damboa, the introduction and possible adoption of IDPC varieties shall help to improve income, reduce poverty level and improve the standard of living of cowpea farmers in the area; these are possible owing to the duality effect associated with the IDPC varieties on higher grain and fodder yields. A vacuum still exists for improvement in terms of the lucrative status of the farmers; evidenced from the wide gap existing between the best points of return on sale ratio and operating ratio. Expansion path (higher production) is feasible when Azir is targeted as a cultivation point; due to its comparative cost advantage and Sabongari is targeted as selling point; due to its comparative sales advantage.

References

Alene AD, Manyong VM, Tipilda A and Singh BB, 2005. The adoption and impact of improved dual purpose cowpea varieties on rural livelihoods in Northern Nigeria. <http://www.usaid.gov/ng/downloads>.

Amaza PS, 2000. Resource Use Efficiency in Food Crop Production in Gombe State, Nigeria. Unpublished Thesis for PhD, Department of Agricultural Economics, University of Ibadan, Nigeria, p.166.

Food and Agriculture Organisation (FAO), 2007. Statistics, Rome, Italy. <http://faostat.org>

Gabdo BH, Tashikalma AK and Polycarp IM, 2005. Financial Appraisal of Private Agro-based Industries in Adamawa State: A Case Study of Afcott Processing Industry, Ngurore-Yola, Nigeria. Global Journal of Agricultural Sciences, Bacchudo Publishers, Calabar, 4, 81-84.

Iheanacho AC, 2000. Pattern and Technical Efficiency of Resource Use in Millet-based crop mixtures in Borno State of Nigeria. Research Journal of Science, 6, 79-103.

Kristjanson P, Okike I, Tarawali S, Singh BB and Manyong VM, 2005. Farmers' Perceptions of Benefits and Factors Affecting the Adoption of Improved Dual Purpose Cowpea in the Dry Savannas of Nigeria. Journal of Agricultural Economics, 32, 195-210.

Langyintuo AS, Lowenberg-DeBoar J, Faye M, Lambert D, Ibro G, Moussa B, Kergna AS, Kushwaha S, Musa S and Ntougkam G, 2005. Cowpea Demand and Supply in West and Central Africa. Development of Agricultural Economics, Krannert Building, Purdue University, West Lafayette, Indiana, USA.

National Population Commission (NPC), 2006. Population Statistics, Abuja, Nigeria.

Olukosi JO and Ogunbile AO, 2005. Introduction to agricultural production economics: Principles and applications, AGITAB Publishers Ltd., Zaria, Nigeria, p. 97.

Onyeagocha SUO, Ehirim NC, Emenyenu CA, Onyemauwa CS, Nwosu FO, Ohajianya DO, Eze CC, Anyanwu GA, Oguoma NNO and Esonu BO, 2010. Post-Avian flu profitability and resource use efficiency of broiler farmers in Akwa Ibom State, South-South, Nigeria. Agricultural Journals. Medwell publishers, 5, 37-44.

Ortiz R, 1998. Cowpea from Nigeria: A Silent Food Revolution. Outlook on Agriculture, 27, 125-128.

Ortiz R, 2003. Plant Breeding Success in African Rural Development, Technology Transfer in Developing World. Check Biotechnology.org. Microsoft Internet Explorer.

Quin FM, 1997. Introduction. Pp ix-xv. In: Singh BB, Mohan-Raj DR, Dashiell KE and Jackai, LEN (Eds.). Advances in Cowpea Research. Co-publication of international Institute of Tropical Agriculture (IITA) and Japan International Research Centre for Tropical Agricultural Sciences (JIRCAS), IITA, Ibadan, Nigeria.

Singh BB, Chambliss OL and Sharma B, 1997. Recent Advances in Cowpea Breeding. Pp 30-49. In: Singh BB, Mohan-Raj DR, Dashiell KE and Jackai LEN (Eds.). Advances in Cowpea Research. Co-publication of international Institute of Tropical Agriculture (IITA) and Japan International Research Centre for Tropical Agricultural Sciences (JIRCAS), IITA, Ibadan, Nigeria.

Genetics and Breeding

- Variation in the agronomic and morphological traits in spring barley** 263
N. Dyulgerov, B. Dyulgerova
- Study on the loss of accuracy of AC method for milk yield control in sheep** 268
D. Dimov, P. Zhelyazkova, A. Vuchkov
- Hordein polymorphism between spring barley cultivars by SDS-PAGE electrophoresis** 273
N. Neykov, S. Doneva

Nutrition and Physiology

- Comparative study of rapeseed, monofloral types and multifloral honey by some physico-chemical parameters** 277
I. Zhelyazkova, S. Lazarov
- Body condition score, nonesterified fatty acids and beta-hydroxybutyrate concentrations in goats with subclinical ketosis** 282
V. Marutsova, R. Binev

Production Systems

- Lucrative status of improved dual purpose cowpea (*Vigna unguiculata* L., Walp) in Damboa, Borno State, North-Eastern Nigeria** 286
B.H. Gabdo
- Study on the emptying time of grain harvester hoppers** 291
K. Trendafilov, N. Delchev, B. Kolev, G. Tihanov
- Length of the growing season and yield in *Triticum monococcum* L., in accordance with the growing conditions** 296
S. Stamatov, E. Valchinova, G. Desheva, K. Uzundzhaliyeva, P. Chavdarov, T. Cholakov, B. Kyosev, R. Ruseva, N. Velcheva
- Productivity of durum wheat cultivar Predel at nitrogen-phosphorous fertilization** 301
L. Plescuta
- Effect of the herbicide treatment dose on the weed infestation in common winter wheat** 306
Z. Petrova
- Evaluation of some technological properties of Caucasian ram wool** 311
D. Pamukova, G. Staykova, N. Stancheva, D. Panayotov

Agriculture and Environment

- Saved CO₂ emissions by using renewable sources for hot water yield in Bulgarian dairy farms** 315
R. Georgiev, R. Slavov, K. Peychev, D. Georgiev, S. Apostolov, J. Ellingsen, J. Tønnesen
- Inventory of the legal base for reclamation of lands disturbed by open-cast mining in Bulgaria** 320
M. Banov, V. Tzolova, I. Kirilov
- Taxonomic composition of phytoplankton in Black Sea area in front of the Cape Galata (2008-2016)** 326
D. Klisarova, D. Gerdzhikov
- Biodiversity of the macrozoobenthos in some protected marine areas along Bulgarian Black Sea coast** 336
E. Petrova, S. Stoykov
- Heavy metals in organs of gudgeon (*Gobio gobio* L.) from Vardar River, R. Macedonia** 340
R. Nastova, V. Kostov, I. Uselinovska

Product Quality and Safety

- Mathematical methods for assessment and analysis of honey yield data for Bulgaria and the European Union for the period 1961-2014** 347
N. Keranova
- Carcass traits and meat quality of different slow growing and fast growing broiler chickens** 351
M. Oblakova, N. Mincheva, P. Hristakieva, I. Ivanova, M. Lalev, Sv. Georgieva
- Role and importance of the awareness for whey in dairy sector at an international level** 358
M. Yilmaz, H. Celik, A.D. Karaman, K. Celik

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors

The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings:

Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows, IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 9, Number 4
December 2017

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz