ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 10, Number 1
March 2018

AGRICULTURAL SCIENCE AND TECHNOLOGY

2018

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Jean-François Hocquette
(France)
Bogdan Szostak (Poland)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate on author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc., AGRIS (FAO) and DOAJ.

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No. DNP 06-41/20.12.2017, financed from Fund 'Scientific Research' grant Bulgarian scientific periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 10, Number 1
March 2018

AGRICULTURAL SCIENCE AND TECHNOLOGY

2018

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Effect of dietary garlic powder and probiotics supplementation on growth performance of male Ross 308 broilers

H. Lukanov^{1*}, I. Pavlova², A. Genchev¹

¹Department of Animal Science - monogastric and other animals, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria

²Department of Pharmacology, Physiology of the Animals and Physiological Chemistry, Faculty of Veterinary Medicine, Trakia University, 6000 Stara Zagora, Bulgaria

(Manuscript received 6 December 2017; accepted for publication 24 January 2018)

Abstract. This study was conducted to evaluate the growth performance of male classic broiler chickens whose feed was supplemented with either garlic powder, probiotics or both. A total of 120 day-old male Ross 308 broiler chickens were divided into 4 groups: first group – control; second group: fed feed supplemented with *Lactobacillus* spp. probiotic combination; third group: fed feed supplemented with 0.4% garlic powder and fourth group: fed both 0.4% garlic powder and *Lactobacillus* spp. The experiment lasted until the 35th day of age. The results showed that the highest live weight at the end of the experiment was established in the group supplemented with probiotic+garlic powder (2257.51 ± 3.87 g), and the lowest – in the control group (2171.37 ± 4.71 g). The group supplemented with probiotic+garlic powder exhibited the highest average weekly weight gain during the 2nd, 3rd and 4th week of life and the highest total weight gain (2211.58 ± 3.91 g). The groups fed probiotic and probiotic+garlic powder had the most efficient feed conversion ratios (1.43 ± 0.2). During the study period, there was no substantial decline in the consumption of feed containing 0.4% garlic powder. The supplementation of feed with garlic powder+*Lactobacillus* spp., had a beneficial effect on the growth performance of broiler chickens. Our data, in line with those of other research studies, suggested that the addition of 0.4% garlic powder combined with *Lactobacillus* spp. to poultry feed increased meat production in broiler chickens.

Keywords: broiler chickens, probiotic, garlic powder, feed consumption, fructooligosaccharides (FOS), growth performance

Introduction

During the last decades, phytobiotics, probiotics and prebiotics became increasingly used in modern livestock husbandry by reason of emerging problems with nutritional antibiotics utilisation and general antimicrobial resistance (Neu, 1992). The application of probiotics in poultry farming is associated with increased resistance to bacterial and protozoal diseases, improved feed conversion and better growth performance (Gupta, 2013). Lactic acid bacteria (LAB): lactobacilli and bifidobacteria, are among the most frequently used probiotics. The role of *Lactobacillus* spp. as dietary probiotic in poultry nutrition comprises promotion of growth of useful microflora, higher activity of digestive enzymes, improved feed intake and digestibility, growth-promoting effect (Jin et al., 1998). Some negative issues are the most expensive feed containing probiotics, difficulty to guarantee a minimum probiotic content after prolonged storage, sensitivity to most probiotic bacteria to high temperatures, inconsistent effects on production performance and intestinal health in adult birds and last but not least, possible transfer of antimicrobial resistance to microbial pathogens.

Prebiotics are indigestible nutrients that stimulate the growth and activity of beneficial intestinal microflora and thus, have a positive effect on the whole body. The supplementation of prebiotics to poultry rations improves intestinal health of birds, their productive performance, improves nutrient utilization, reduces cost of produce and contributes to maintenance of a proper environment (Hajati and Rezaei, 2010). The most commonly used prebiotics are from the group of oligosaccharides: fructooligosaccharides (FOS), inulin, oligofructan, isomaltoligosaccharides, etc. Plants rich in FOS and

inulin are garlic, chicory, onion, artichoke, asparagus, bananas, wheat, tomatoes, etc. (Niness, 1999).

Garlic (*Allium sativum*) is used widely at a global scale as a spice, as well as for control of some infectious and non-infectious diseases in phyto- and traditional medicine, including reduction of blood and liver cholesterol contents. Garlic powder contains FOS with prebiotic effect (Campbell et al., 1997). Furthermore, it contains a varying amount of allicin, vitamins (vitamin C, thiamine, riboflavin and niacin), selenium and potassium (Anonymous, 2014a). The dietary supplementation of garlic powder in chickens was found to be beneficial for growth performance (Raeesi et al., 2010; Issa and Omar, 2012; Elagib et al., 2013).

The combination of probiotics with substances with prebiotic activity improves the effect of the former due to creation of more favorable environment for replication of probiotic bacteria in poultry gut. This study was aimed at evaluating the growth performance of male classic broiler chickens whose feed was supplemented with either garlic powder, probiotics or both.

Material and methods

A total of 120 male day-old Ross 308 broiler chickens were divided into 4 groups. Each comprised three subgroups with 10 sexed male chicks provided by a certified hatchery:

- First group (I) – control (Group 1 in figures and the table);
- Second group (II) – fed feed supplemented with *Lactobacillus* spp. probiotic (Group 2 in figures and the table);
- Third group (III) – fed feed supplemented with 0.4% garlic

* e-mail: drlukanov@gmail.com

powder (Group 3 in figures and the table);

- Fourth group (IV) – fed a combination of both 0.4% garlic powder and *Lactobacillus* spp. probiotic (Group 4 in figures and the table).

The garlic powder (country of origin China) contained 3310 kcal/kg metabolisable energy (ME); 16.55% crude protein (CP); 0.73% crude fat (CF); 9.0% crude fibre (CF); 0.77% lysine; 0.11% methionine; 0.37% threonine; 0.12% tryptophan; 0.08% Ca; and 0.41% P (Anonymous, 2014a).

The experiments were performed in the experimental base of the Poultry Breeding unit, Faculty of Agriculture, Trakia University – Stara Zagora, according to the requirements for humane treatment of birds and in compliance with animal hygienic norms of the production type and age category. Twelve cages were provided for individual subgroups. Throughout the trial, ambient temperature and daylight were controlled as per the recommendations of the producer of this meat-type hybrid (Anonymous, 2014b). The experiment continued until the 35th day of age.

The feeding was according to the age, and feed was produced *in situ* with uniform energy, protein, amino acid and mineral content for the four groups as follows:

- **Starter** (0-14 days of age): 12.65 MJ/kg ME, 23% CP, 1.43% lysine, 0.51% methionine, 0.94 % threonine, 1.05% Ca, 0.5 % available P and <4.00 CF;

- **Grower** (15-28 days of age): 13.20 MJ/kg ME, 21% CP, 1.24% lysine, 0.45% methionine, 0.83% threonine, 0.9% Ca, 0.45% available P and <4.00 CF;

- **Finisher** (29-35 days of age): 13.40 MJ/kg ME, 19% CP, 1.09% lysine, 0.41% methionine, 0.9% Ca, 0.45% available P and <4.00 CF.

The following traits were controlled:

- Live weight, g – on a weekly basis, by individual weighing of chickens. To this end, technical balance EU-6000CE with precision up to 0.1g was used;

- Weight gain, g – by statistical processing of individual weight control data;

- Feed intake, g – on a daily basis, by weighing the remaining feed in the morning and what was left after 24h. To this end, technical balance EU-6000CE with precision up to 0.1g was used;

- Total feed intake for the experimental period, g (0-35 days of age) - by statistical processing of daily feed intakes;

- Feed conversion ratio (FCR) - by statistical processing of feed intake and live weight data for the period of the trial.

- European Broiler Index (EBI) for the experimental period was made:

- $EBI (0-35) = \text{Average daily weight gain (g)} \times \text{survival (\%)} / 10 \times FCR$

Statistical analysis

All data were analysed by Statistica 13.0 software (Statistica for Windows; Stat – Soft, 2015). Mean (x), standard error of mean (SEM) and coefficient of variation (CV,%) values were calculated for each group. The differences were considered statistically significant at $P < 0.05$, using Student's t-test, if the data were normally distributed.

Results and discussion

The results presented on Figure 1 showed that average live weight of broilers during the last 2 weeks of the trial was close to that

indicated in the technological documentation for the Ross 308 hybrid. The weight of chicks from the experimental groups was superior to that of controls, with higher differences until 28 days of age. Nevertheless, a highly positive effect from dietary supplementation with garlic powder and the combination probiotic+garlic powder was noted. In this period, the highest live weight was that of broiler chickens from group IV, exceeding by 2.9% the recommendations of the hybrid manufacturer. After that age, the differences between experimental and control groups decreased and at 35 days of age they were between 0.8 and 4.0%. It was thought that until 35 days of age, the weight of broilers was positively influenced by garlic powder combined with LAB probiotic and less significantly when they were added independently to the feed. A comparable trend in the effects of dietary garlic powder on live weight was reported by Raeesi et al. (2010), Issa and Omar (2012), Elagib et al. (2013) and Lukanov et al. (2015). On the contrary, Choi et al. (2010) did not observe a beneficial effect of garlic powder on the growth of broiler chickens. Similar to our results, Ignatova et al. (2009) reported positive effect of the probiotic administration on the body weight, feed intake and feed conversion in broiler chickens. The results of our study were in line with the concepts for positive influence of synbiotic dietary combinations on growth performance of farm animals and birds. Our results about better growth performance in groups fed with probiotics and prebiotics were similar to those of Falaki et al. (2010) and Mookiah et al. (2014) but not to the conclusions of Mokhtari et al. (2010).

Figure 1. Average live weight of birds during the test period, g

Figure 2 shows that between 6 and 16 days of age as well as during the period 21-30 days, feed intake of broiler chickens from the experimental groups was lower as compared to the technological norms for Ross 308. Feed intake of control birds during the third and fourth week of the trial was similar to that specified by the hybrid producer. During the finisher period, a tendency to increased feed intake was noticed in chickens from all groups, exceeding the technological norms. Despite the observed between-group differences during the growth stages, the feed intake for the entire experimental 35-day period was 7% to 11.1% lower than the technological norms for Ross 308. In agreement with our previous study (Lukanov et al., 2015), no relationship between the presence of garlic powder at amounts up to 0.4% in feed and its intake by birds

Figure 2. Daily feed intake per bird during the test period, g

was established. On the contrary, Abdullah et al. (2010) reported lower feed consumption in groups supplemented with 0.5% and 1% garlic powder.

The time course of average weekly weight gain (Figure 3) in experimental and control groups compared to producer's norms showed that during the first half of the study, average weekly weight gain was significantly lower in all four groups vs Ross 308 norms and followed the same pattern. During the third week, a considerable increase was demonstrated and towards the end of the third week this trait attained the value specified by Aviagen®. At the beginning of the 4th week, this ascendant trend continues for all groups, attaining 930 g average weekly weight gain in group IV. In the middle of the 4th week, the studied trait declined sharply in the experimental groups until the end of the study without reaching levels specified in the technological norms. This compensated the substantial growth retardation during the first half of the study as seen from data on Figure 1 and Table 1. Due to the extensive variability and dynamics of this trait, no comparison with data from other research teams was possible.

According to data from Table 1 the highest live weight was that of chickens fed garlic powder and probiotic with feed compared to the other supplemented groups and to control birds ($P < 0.001$). In general, all groups exhibited lower live body weights than values specified by the hybrid producer. This could be attributed to the

Figure 3. Average weekly weight gain of birds during the test period, g

environmental influences, which in control studies were close to the ideal conditions for Ross 308. Similar results were obtained for weight gain throughout the entire experimental period, e.g. statistically significant differences among the groups. The variation of these two traits in the three supplemented groups and the controls by the end of the study was between 4.8% and 10.43% (CV), evidencing a good uniformity at the end of the trial.

Until 35 days of age, all four studied groups had superior feed conversion ratios than the technological norm ($FCR = 1.54$). The difference in the experimental groups was especially obvious. Feed conversion (kg/kg) until 35 days of age kept a stable trend towards beneficial effect of dietary garlic powder+probiotic as well as of independent 0.4% garlic powder and probiotic addition to feed. The feed conversion of controls was very similar to the technological norm. Most efficient feed conversion was noticed in the group supplemented with 0.4% garlic powder and the group that received both garlic powder and probiotic ($FCR = 1.43$) but the difference was statistically insignificant. Our results about FCR values agreed with those of Mokhtari et al. (2010), Awad et al. (2009) and Lukanov et al. (2015).

European Broiler Index is used to describe the efficiency of

Table 1. Growth performance of chickens from the four groups

Parameters	1 group			2 group			3 group			4 group			Ross 308
	n	$\bar{x} \pm \text{SEM}$	CV, %	n	$\bar{x} \pm \text{SEM}$	CV, %	n	$\bar{x} \pm \text{SEM}$	CV, %	n	$\bar{x} \pm \text{SEM}$	CV, %	
Final body weigh, g	28	2171.37 $\pm$ 4.71	10.11	27	2208.43 $\pm$ 4.80	4.8	28	2188.43 $\pm$ 4.22	9.02	28	2257.51 $\pm$ 3.87	8.14	2283
Significance					1:2***, 1:3**, 1:4***, 2:3**, 2:4***, 3:4***								
Final gain, g	28	2125.67 $\pm$ 4.76	10.33	27	2162.59 $\pm$ 4.85	10.43	28	2142.53 $\pm$ 4.26	9.21	28	2211.58 $\pm$ 3.91	8.31	2226
Significance					1:2***, 1:3**, 1:4***, 2:3**, 2:4***, 3:4***								
FCR, kg/kg	28	1.52 $\pm$ 0.23	11.05	27	1.43 $\pm$ 0.2	10.74	28	1.44 $\pm$ 0.20	9.29	28	1.43 $\pm$ 0.2	7.93	1.54
Significance					NS								
EBI	28	380.8 $\pm$ 2.0	10.1	27	397.1 $\pm$ 2.0	10.2	28	405.1 $\pm$ 1.8	9.0	28	420.8 $\pm$ 1.7	8.1	410.86
Significance					1:2,3,4***, 2:3**, 2:4***, 3:4***								

Note: with 1,2,3 and 4 are signed: 1 - control group; 2 - fed feed supplemented with *Lactobacillus* spp. probiotic; 3 - fed feed supplemented with 0.4% garlic powder; 4 - fed a combination of both 0.4% garlic powder and *Lactobacillus* spp. probiotic. EBI - European Broiler Index.

* $P < 0.05$ ** $P < 0.01$; *** $P < 0.001$; NS- Non Significant

broiler production. In our study we found that the higher index was calculated for group four fed with garlic powder + probiotic (420.8 ± 1.7) compared to the other groups ($P < 0.001$). The lowest index was calculated for the control group, without feed supplementations - 380.8 ± 2.0 . In general, all experimental groups showed a better efficiency of broiler production in comparison to the control group. Our results according to production efficiency for probiotic or symbiotic treated broilers are in agreement with those of Peric et al. (2010), Awad et al. (2008) and Khan et al. (2011). In contrast to our results, Biernasiak and Slizewska (2009) reported lower EBI in a probiotic supplemented group than an unsupplemented one.

Conclusion

The results showed that the dietary supplementation of 0.4% garlic powder and probiotics increased statistically significantly the live body weight ($P < 0.001$). The group fed both probiotic and garlic powder, exhibited higher live body weight, higher weight gain and better feed conversion vs either of the other three groups (control group, probiotic treated group and 0.4% garlic powder supplemented group). There are no substantial differences between control and experimental groups according to the daily weight gain parameter.

References

- Abdullah AY, Mahmoud KZ, Nusairat BM and Qudsieh RI, 2010. Small intestinal histology, production parameters, and meat quality as influenced by dietary supplementation of garlic (*Allium sativum*) in broiler chicks. Italian Journal of Animal Science, 9, 414-419.
- Anonymous (Aviagen, Ross 308), 2014b. Ross broiler management handbook, Aviagen, Ross, 308.
- Anonymous (USDA), 2014a. National Nutrient Database for Standard Reference Release 27. Basic Report 02020. Spices, garlic powder. Report Date: September 29, 2014 22:01 EDT.
- Awad W, Ghareeb K and Böhm J, 2008. Intestinal Structure and Function of Broiler Chickens on Diets Supplemented with a Synbiotic Containing *Enterococcus faecium* and Oligosaccharides. International Journal of Molecular Sciences, 9, 2205-2216.
- Awad WA, Ghareeb K, Abdel-Raheem S and Böhm J, 2009. Effects of dietary inclusion of probiotic and synbiotic on growth performance, organ weights, and intestinal histomorphology of broiler chickens. Poultry Science, 88, 49-56.
- Biernasiak J and Slizewska K, 2009. The effect of a new probiotic preparation on the performance and faecal microflora of broiler chickens. Veterinarni Medicina, 54, 525-531.
- Campbell JM, Bauer LL, Fahey GC, Hogarth AJCL, Wolf BW and Hunter DE, 1997. Selected fructooligosaccharides (I - kestose, nystose, and IF-fructofuranosylnystose) composition of foods and feeds. Journal of Agricultural and Food Science, 45, 3076-3082.
- Choi IH, Park WY and Kim YJ, 2010. Effects of dietary garlic powder and α -tocopherol supplementation on performance, serum cholesterol levels, and meat quality of chicken. Poultry Science, 89, 1724-1731.
- Elagib HAA, El-Amin WIA, Elamin KM and Malik HEE, 2013. Effect of dietary garlic (*Allium sativum*) supplementation as feed additive on broiler performance and blood profile. Journal of Animal Science Advances, 3, 58-64.
- Falaki M, Shamas SM, Dastar B, Zerehdaran S and Khomairim M, 2010. The Investigation of intestinal microflora and growth response of young broilers given feed supplemented with deferent levels of probiotic and prebiotic. Journal of animal and veterinary advances, 9, 2685-2690.
- Gupta AR, 2013. The benefits of probiotics in poultry production: an overview. International Journal of Livestock Research, 3, 18-22.
- Hajati H and M Rezaei, 2010. The Application of prebiotics in poultry production. International Journal of Poultry Science, 9, 298-304.
- Ignatova M, Sredkova V and Marasheva V, 2009. Effect of dietary inclusion of probiotic on chickens performance and some blood indices. Biotechnology in Animal Husbandry, 25, 1079-1085.
- Issa KJ and Omar JMA, 2012. Effect of garlic powder on performance and lipid profile of broilers. Open Journal of Animal Sciences, 2, 62-68.
- Jin LZ, Ho YW, Abdullah N and Jalaludin S, 1998. Growth performance, intestinal microbial populations, and serum cholesterol of broilers fed diets containing Lactobacillus cultures. Poultry Science, 77, 1259-1265.
- Khan SH, Yousaf B, Mian AA, Rehman A and Farooq MS, 2011. Assessing the effect of administering different probiotics in drinking water supplement on broiler performance, blood biochemistry and immune response. Journal of Applied Animal Research, 39, 418-428.
- Lukanov H, Genchev A and Ribarski S, 2015. Effect of feed supplementation with garlic power on meat productivity and meat quality traits of classic Ross 308 male hybrid chickens. Trakia Journal of Sciences, 1, 66-76.
- Mokhtari R, Yazda AR, Rezaei M and Ghorbani B, 2010. The Effects of Different Growth Promoters on Performance and Carcass Characteristics of Broiler Chickens. Journal of Animal and Veterinary Advances, 9, 2633-2639.
- Mookiah S, Sieo CC, Ramasamy K, Abdullah N and Ho YW, 2014. Effects of dietary prebiotics, probiotic and symbiotic on performance, caecal bacterial populations and caecal fermentation concentrations of broiler chickens. Journal of the Science of Food and Agriculture, 94, 341-348.
- Neu H, 1992. The Crisis in Antibiotic Resistance. Science, 257, 1064-1073.
- Niness KR, 1999. Nutritional and Health Benefits of Inulin and Oligofructose Inulin and Oligofructose: What Are They? American Society for Nutritional Sciences, 1402S-1406S.
- Peric L, Milosevic N, Zikic D, Bjedov S, Svetkovic D, Markov S, Mohnl M and Steiner T, 2010. Effects of probiotic and phytogetic products on performance, gut morphology and cecal microflora of broiler chickens. Archiv Tierzucht, 53, 350-359.
- Raeesi M, Hoeini-Aliabad SA, Roofchae A, Zare Shahneh A and Piralı S, 2010. Effect of periodically use of garlic (*Allium sativum*) power on performance and carcass characteristics in broiler chickens. World Academy of Science, Engineering and Technology, 68, 1213-1219.

Review

- Achievements and problems in the weed control in grain maize (*Zea mays* L.)** 3
G. Delchev, M. Georgiev

Genetics and Breeding

- Yield and coefficient of ecological valence of spring barley in the regions of Sadovo and Karnobat, Bulgaria** 8
N. Neykov, T. Mokreva
- Agronomic performance of mutant lines of winter two-rowed barley** 12
B. Dyulgerova, D. Valcheva, N. Dyulgerov
- Phenotypic diversity in six-rowed winter barley (*Hordeum sativum* L.) varieties** 16
N. Dyulgerov, B. Dyulgerova
- Evaluation of rye specimens in maturity stage on the base of mathematical – statistical analysis** 21
V. Kuneva, E. Valchinova, A. Stoyanova
- Evaluation of lentil cultivars and lines for resistance to *Fusarium oxysporum* f.sp. lentis** 25
M. Koleva, Y. Stanoeva, I. Kiryakov, A. Ivanova, P. Chamurlyiski
- Registration of a new sunflower hybrid - Sevar** 29
P. Peevska, M. Drumeva, G. Georgiev

Nutrition and Physiology

- The effect of novel xylanase on feeding value of diet containing cereal by-products for broilers** 34
J.M. Abdulla, S.P. Rose, V. Pirgozliev
- Effect of dietary garlic powder and probiotics supplementation on growth performance of male Ross 308 broilers** 37
H. Lukanov, I. Pavlova, A. Genchev
- Slaughter traits of Pharaoh Japanese quails** 41
A. Genchev, H. Lukanov, I. Penchev
- Blood count in dogs with mammary gland carcinoma** 44
Ts. Hristov, R. Binev

Production Systems

- Economic efficiency of fattening on different genotypes slow-growing and fast-growing broiler chickens** 48
M. Oblakova, Y. Popova, P. Hristakieva, N. Mincheva, M. Lalev

Effect of nutmeg extract supplementation on some productive traits and economic efficiency of common carp (<i>Cyprinus carpio</i> L.) cultivated in recirculation system	54

G. Zhelyazkov, S. Stoyanova, I. Sirakov, K. Velichkova, Y. Staykov

Agriculture and Environment

Influence of biomanipulation on the living communities and the water quality in the Strezhevo hydroecosystem, R. Macedonia	57

R. Nastova, V. Kostov, N. Gjorgovska, V. Levkov

Product Quality and Safety

Residue analysis of difenoconazole in apple fruits grown in Republic of Macedonia	63

V. Jankuloska, I. Karov, G. Pavlovska

Organoleptic properties of white yam (<i>Dioscorea rotundata</i> poir) as affected by autoclaving time	67

M. Ahmed, Y.B. Kiri, M.S. Abubakar

Influence of Goji berries on oxidative changes, microbiological status and chemical properties of sausages	70

A. Mitev, A. Kuzelov, E. Joshevska

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors
The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings: Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows, IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 10, Number 1
March 2018

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz