ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)
Volume 10, Number 3
September 2018

AGRICULTURAL SCIENCE AND TECHNOLOGY

2018

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria
E-mail: gpetkov@af.uni.sz.bg

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Atanas Atanasov (Bulgaria)
Svetlana Georgieva (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Stoicho Metodiev (Bulgaria)
Bojin Bojinov (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Ivan Varlyakov (Bulgaria)
George Zervas (Greece)
Vasil Pirgozliev (UK)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Jean-François Hocquette
(France)
Bogdan Szostak (Poland)

Agriculture and Environment

Martin Banov (Bulgaria)
Peter Cornish (Australia)
Vladislav Popov (Bulgaria)
Tarek Moussa (Egypt)

Product Quality and Safety

Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

There are no submission / handling / publication charges.

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate an author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to

editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: AGRIS (FAO), CABI, EBSCO-host, ROAD and DOAJ. DOI system is used for article identification

The journal is freely available without charge to the user or his/her institution. Users can read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This issue is printed with the financial support by Contract No. DNP 06-41/20.12.2017, financed from Fund 'Scientific Research' grant Bulgarian scientific periodicals.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820 (print)
ISSN 1314 - 412X (online)

Volume 10, Number 3
September 2018

AGRICULTURAL SCIENCE AND TECHNOLOGY

2018

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Effect of experimentally induced aflatoxicosis on haematological parameters and bone marrow morphology in mulard ducks

I. Valchev^{1*}, N. Groseva², D. Kanakov¹, Ts. Hristov¹, L. Lazarov¹, R. Binev¹

¹Department of Internal Non-Infectious Diseases, Faculty of Veterinary Medicine, Trakia University, 6000 Stara Zagora, Bulgaria

²Department of General and Clinical Pathology, Faculty of Veterinary Medicine, Trakia University, 6000 Stara Zagora, Bulgaria

(Manuscript received 12 June 2018; accepted for publication 20 August 2018)

Abstract. In this experiment, the toxic effects of AFB₁ on some haematological parameters were investigated in mulard ducks, namely red blood cell counts (RBC), haemoglobin content (HGB), haematocrit (HCT), platelet counts (PLT), white blood cell counts (WBC), differential white blood cell counts (WBC %), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH) and mean corpuscular haemoglobin concentration (MCHC). Experiments were conducted with 4 groups of 20 10-day-old mulard ducks each. The groups were as followed: group I – control, fed a standard compound feed according to the species and the age; group II – experimental, whose feed was supplemented with 0.5 mg/kg AFB₁; group III – experimental, supplemented with 0.8 mg/kg AFB₁, and group IV – experimental, supplemented with 0.5 mg/kg AFB₁ and 2 g/kg Mycotox NG. The duration of the experiment was 42 days. Haematological analysis performed on the 21st day of the trial showed reduction in RBC, HCT, HGB and PLT and increased total WBC in groups II and III. The percentages of the different leukocyte classes (differential leukocyte counts) demonstrated increased proportion of heterophils and lower percentages of lymphocytes. The observed changes tended to become more pronounced on the 42nd day of the experiment. There were no statistically significant changes in MCV, MCH, and MCHC, as well as in eosinophil, basophil, and monocyte percentages between control and treated groups ($p > 0.05$). The supplementation of the feed with mycosorbent – Mycotox NG (experimental group IV) reduced partly the harmful effect of AFB₁ on the studied haematological indices

Keywords: aflatoxin B₁, haematological parameters, bone marrow, mallard ducklings Mycotox NG

Introduction

Mycotoxins are compounds with various chemical structure and biological effects produced by filamentous fungi (Sudakin, 2003). Aflatoxins, ochratoxins, trichothecene mycotoxins, zearalenone and fumonisins are the most important for human health and animal productivity (Zain, 2011). Mycotoxicoses is a collective term comprising diseases in animals and humans produced by ingestion of mycotoxins (Hussein and Brasel, 2001). Exposure of farm animals to toxigenic fungi and their toxins occurs mainly by ingestion of contaminated feeds which contain corn, peanut or sorghum (Kamei and Watanabe, 2005; Tuzcu et al., 2010). Aflatoxins are secondary toxic metabolites of fungi from the genus *Aspergillus* (*A. flavus*, *A. parasiticus* and *A. niger*) (Kurtzman et al., 1987). They belong to the group of storehouse moulds. Under field conditions, before harvesting and storage of cereal crops, growth of moulds from the genus *Fusarium* (*Graminearum*, *Culmorum*, *Claviceps purpurea*) producing zearalenone, fumonisins and trichothecene mycotoxins is more common, while during storage of feeds, fungi from genera *Aspergillus* and *Penicillium* (*Aspergillus flavus*, *Aspergillus parasiticus* and *Penicillium verrucosum*) producing aflatoxins (B_1 , B_2 , G_1 , G_2) and ochratoxins occur (Hashmi et al., 2006). The temperature range within which *A. flavus* and *A. parasiticus* produce aflatoxins is from 12 to 41°C, with optimum range 25–32°C (Lillehoj, 1983), air humidity higher than 62% and feed humidity higher than 14% (Royes and Yanong, 2002). Aflatoxins are fluorescence compounds. Aflatoxin B_1 (AFB₁) and aflatoxin B_2 (AFB₂) emit blue fluorescence, while aflatoxin G_1 (AFG₁) and aflatoxin G_2 (AFG₂) emit green fluorescence after UV irradiation (Hussein and Brasel, 2001). Aflatoxins are highly toxic for men and animals and among the most potent natural poisons. Similarly to the

other mycotoxins, they are thermostable (Cheeke, 1998). The EC legislation as well as that of Chile, Canada and the USA allows no more than 10 ppb AFB₁ in poultry feeds and 20 ppb of total aflatoxin (AFB₁ + AFB₂ + AFG₁ + AFG₂) in feeds for the poultry industry (Directive 2002/32/EC; Fonseca, 2003). More than 18 different aflatoxins are identified (B_1 , B_2 , G_1 , G_2 , P, Q, M₁, M₂, B_{2a} etc.). The commonest aflatoxins with toxicological importance are produced by moulds growing on feeds under natural conditions – B_1 , B_2 , G_1 , and G_2 , and M₁ and M₂ in milk (IARC, 2002). Aflatoxin B_1 (AFB₁) is the most toxic metabolite for men, animals and hydrobionts (Kennedy et al., 1998). It is a potent natural hepatotoxin, carcinogen, teratogen and mutagen (Han et al., 2008). Terao and Ueno (1978) reported that the toxicity of AFG₂, AFB₂ and AFG₁ was equal to 10%, 20% and 50% respectively of that of AFB₁. Aflatoxins are found to contaminate naturally mainly cereal and oily crops, nuts and milk (Wu et al., 2009). On the other hand, 77% of naturally occurring aflatoxins is from the B₁ type (Wilson and Payne, 1994). Aflatoxins are a potential threat to poultry health and performance after consumption of contaminated fodders (Huff et al., 1986). From all fowl species, growing ducklings, goslings, turkey poult and chicks are more susceptible to the toxic effect of aflatoxins. The latter are mainly localised in the liver and are characterised with necrosis, haemorrhages and proliferation of biliary duct epithelium. Chronic aflatoxicosis in these bird species results in weight loss, reduced feed conversion, increased sensitivity to infections and lower egg production (Dalvi, 1986). In an experiment with broiler chickens, Al-Daraji et al. (2004) demonstrated changes in haematological and blood biochemical parameters (red and white blood cell counts, haematocrit, haemoglobin, heterophil and lymphocyte percentages, uric acid, blood glucose, cholesterol, total protein, calcium phosphate, gamma-glutamyltransferase and alkaline phosphatase

* e-mail: valtchev@abv.bg

activities).

One of the most appropriate methods for detoxication of mycotoxin-contaminated feeds is the use of non-nutritional adsorbents that are capable to bind toxins and to inhibit their absorption from the gastrointestinal tract, and thus, to minimise their toxic effects on animals and their occurrence in animal organs and animal foodstuffs (Ramos et al., 1996; Bintvihot et al., 2002; Kana et al., 2011; Oguz, 2012). A good mycosorbent should be able to restore the nutritional value of aflatoxin-contaminated feeds. The quality of mycosorbents is assessed by four principal parameters, namely: binding capacity, absorption efficiency, activation time and binding extent (Van Kessel and Hiang-Chek, 2004). The process of aflatoxin binding by mycosorbents is based on the electric polarity principle. The negative electrical charge of the mycotoxin binds to the positive charge of the mycosorbent and thus, mycotoxin are immobilised and subsequently, removed from the animal body. The effects of aluminosilicates, active charcoal and yeasts are extensively studied, and promising results have been reported (Ramos et al., 1996; Huwig et al., 2001; Mabbet, 2005; Girish and Devegowda, 2006; Oguz, 2012).

The aim of the present study was to monitor the changes in some haematological parameters in mulard ducks with experimental aflatoxicosis. Also, the purpose was to evaluate the possibility for prevention of toxic effects of AFB₁ by feed supplementation with the mycosorbent Mycotox NG.

Material and methods

Experiments were conducted with 80 female 10-day-old mulard ducks, divided into 4 groups according to the following design:

- Group I – control. It was fed a standard compound feed according to the age produced by Zoohraninvest feed factory, Stara Zagora and comprised pelleted starter, grower and finisher;
- Group II – experimental, the standard feed was supplemented with 0.5 mg/kg aflatoxin B₁;
- Group III – experimental, the standard feed was supplemented with 0.8 mg/kg aflatoxin B₁;
- Group IV – experimental, the standard feed was supplemented with 0.5 mg/kg aflatoxin B₁ and 2 g/kg Mycotox NG (Ceva Sante Animale, France).

The aflatoxin B₁ used in this experiment was produced by *Aspergillus flavus* (99% purity) purchased from Sigma-Aldrich, Germany. The feed of experimental groups of birds was grinded for better homogenisation of aflatoxin B₁. Microclimatic parameters were optimal and equal for all groups. In the beginning of the trial, air temperature in the living area of ducks was 35°C and until the 15th day it decreased at a daily rate of 1°C; by day 28 it was 20°C, and thereafter – +18°C with air humidity of 60–75% (Regulation No. 44/20.04.2006). The duration of the light day was 24h throughout the experiment. Control and experimental groups were housed in different sections, each with area of 4 m² in the same premise. The sections were bedded with clean dry wooden shavings with layer depth of 5cm. During the first week of life, the feeding width was 1cm and subsequently: 10cm. Blood samples were collected from *v. metatarsalis medialis* on experimental days 21 and 42 in sterile vacutainers with K₂EDTA (FL medical, Italy) for determination of: haemoglobin content (colorimetric test, Human Diagnostica Germany), haematocrit (microcentrifugation method, Angelov et al., 1999), red blood cell counts, platelet counts, white blood cell counts (counted in a chamber as per Angelov et al., 1999). Differential white blood cell counts (WBC, %), were determined on stained blood

smears. The experiment was approved by the Ethics and Animal Welfare Committee of the Faculty of Veterinary Medicine to the Trakia University (permit No. 42.10.10.2011).

The statistical analysis of data was done by one-way ANOVA, and the level of statistical significance – with the Tukey-Kramer test ($p < 0.05$).

Results

Blood picture

The parameters of the red blood picture (erythrocytes, haemoglobin, haematocrit, platelets) in control and experimental groups are presented in Table 1. Red blood cell counts in Groups II and III were statistically significantly lower at both sampling days (21 and 42) compared to control birds ($p < 0.001$). Their values on day 21 were 2.52 ± 0.05 T/l (17.92%) and 2.34 ± 0.05 T/l (23.78%) vs control group (3.07 ± 0.07 T/l) and by the 42nd day the changes were more pronounced with 2.31 ± 0.02 T/l (21.96%) in experimental group II and 2.10 ± 0.05 T/l (29.06%) in experimental group III (control group – 2.96 ± 0.06 T/l). Haemoglobin contents were statistically significantly lower in groups II and III than in untreated birds ($p < 0.001$): on the 21st day, control birds' haemoglobin was 137.57 ± 1.27 g/l vs 117.63 ± 1.10 g/l (14.5%) in group II and 111.53 ± 1.35 g/l (18.93%) in Group III while on day 42, the values in challenged groups were 107.76 ± 2.60 g/l (20.18%) and 96.96 ± 3.85 g/l (28.18 %) in groups II and III, respectively (control group had average haemoglobin of 135.00 ± 1.34 g/l). In ducklings treated only with AFB₁ (Groups II, III) haematocrit decreased significantly on both sampling days ($p < 0.001$). On day 21, haematocrit of Group II was $32.4 \pm 1.23\%$ (18.39 %), and of Group III: $28.9 \pm 0.79\%$ (27.21%) than in controls ($39.7 \pm 0.81\%$). On day 42, haematocrit in Group II was $29.6 \pm 1.07\%$ (22.92%) and in Group III: $24.9 \pm 0.95\%$ (35.16%) which was substantially lower than control values ($38.4 \pm 0.76\%$; $p < 0.001$). Similar changes were observed in platelet counts – on the 21st day, average platelet counts were 60.69 ± 2.99 G/l (26.96%) in Group II and 44.29 ± 2.80 G/l (46.7%) in Group III vs controls (83.09 ± 2.93 G/l, $p < 0.001$). At the next sampling day, changes were even more pronounced with average platelet counts of 55.00 ± 1.75 G/l (33.95%) and 40.56 ± 1.48 G/l (51.29%) in Groups II and III, respectively ($p < 0.001$). Control platelets at day 42 were 83.26 ± 6.38 G/l. In Group IV, the changes in these red blood cell parameters were reduced to a various extent compared to controls ($p = 0.05$ to $p = 0.01$). Red blood cell counts on days 21 and 42 were 2.80 ± 0.06 T/l (8.80%) and 2.77 ± 0.04 T/l (6.42%) respectively. Haemoglobin content was 130.85 ± 1.64 g/l (4.89%) on day 21 and 124.49 ± 2.47 g/l (7.79%) on day 42. On day 21, average haematocrit was $36.2 \pm 0.72\%$ (8.81%), and on day 42: $34.9 \pm 0.73\%$ (9.12%). Platelet counts on days 21 and 42 were 71.74 ± 2.07 G/l (13.66%) and 68.61 ± 2.23 G/l (17.6%) respectively.

Table 2 presents the data about white blood cell counts (total and differential) in untreated birds and those fed aflatoxin with the feed. Total white blood cell counts (WBC) was statistically significantly increased in Groups II and III than in controls. In Group II, WBC were 28.08 ± 0.79 G/l (37.57%) and 32.95 ± 0.98 G/l (47.16%) on days 21 and 42, respectively ($p < 0.001$). In Group III, WBC on day 21 was 31.85 ± 0.76 G/l (56.05%) whereas on day 42 – 38.25 ± 1.18 G/l (70.83%) ($p < 0.001$). The addition of mycosorbent to the contaminated feed (Group IV) partly reduced the deleterious effects of AFB₁ on total WBC counts which were 24.32 ± 1.07 G/l (19.15%) on day 21 and 27.69 ± 1.74 G/l (23.67%) on day 42 ($p < 0.05$).

Table 1. Changes in haematological parameters (red blood cell counts, haemoglobin, haematocrit, platelet counts) in mulard ducks receiving aflatoxin B₁ (AFB₁) either alone or together with Mycotox NG

Groups	Red blood cell counts, T/l		Haemoglobin, g/l		Haematocrit, %		Platelet counts, G/l	
	Day 21	Day 42	Day 21	Day 42	Day 21	Day 42	Day 21	Day 42
I	3.07± 0.07	2.96± 0.06	137.57± 1.27	135.00± 1.34	39.7± 0.81	38.4± 0.76	83.09± 2.93	83.26± 6.38
II	2.52± 0.05 ^{1c}	2.31± 0.02 ^{1c}	117.63± 1.10 ^{1c}	107.76± 2.60 ^{1c}	32.4± 1.23 ^{1c}	29.6± 1.07 ^{1c}	60.69± 2.99 ^{1c}	55.00± 1.75 ^{1c}
III	2.34± 0.05 ^{1c}	2.10± 0.05 ^{1c,2a}	111.53± 1.35 ^{1c,2a}	96.96± 3.85 ^{1c,2a}	28.9± 0.79 ^{1c,2a}	24.9± 0.95 ^{1c,2b}	44.29± 2.80 ^{1c,2c}	40.56± 1.48 ^{1c,2a}
IV	2.80± 0.06 ^{1a,2a,3c}	2.77± 0.04 ^{1a,2c,3c}	130.85± 1.64 ^{1b,2c,3c}	124.49± 2.47 ^{1a,2a,3c}	36.2± 0.72 ^{1a,2a,3c}	34.9± 0.73 ^{1a,2c,3c}	71.74± 2.07 ^{1a,2a,3c}	68.61± 2.23 ^{1a,2a,3c}

*Data are presented as mean ± SEM; n=20 in each group; P<0.05; P<0.01; P<0.001; 1 vs control group; 2 vs experimental group I; 3 vs experimental group II

The differential white blood cell counts exhibited statistically significant changes in heterophils and lymphocytes (Table 2). Heterophil percentages were very increased in Groups II and III on day 21 – 41.6±0.93% (38.66%) and 45.1±1.29% (50.33%), as well as on day 42 – 44.5±1.00% (44.62%) and 48.1±1.73% (54.16%) (p<0.001) compared to control group (30.00±1.57%; 31.2±1.81%). More pronounced changes were established on day 42. In Group IV, heterophil percentage was somewhat lower after the addition of mycosorbent to the feed (p<0.05): 36.2±1.57% on day 21 and 37.1±1.44% on day 42. The reduction vs control group was by 20.66% and 18.91% respectively. Lymphocyte percentage on day 21 was 51.00±1.47% (18.14%) in Group II and 47.2±1.34% (24.24

%) on Group III (p<0.001). On the 42nd day, respective values in Group II and III were 48.00±2.07% (21.70 %) and 45.00±1.07% (26.60%) in Group II (p<0.001). In Group IV, changes were less manifested after the addition of the mycosorbent (p<0.05) with values by day 21 of 57.1±1.15%, and by day 42 of 55.6±1.19%. The reduction was by 8.34% and 9.30%, respectively. Statistically significant changes between monocytes, eosinophils and basophils in untreated and experimental ducks were not found out (p>0.05).

There were no statistically significant changes in mean corpuscular volume (MCV), mean corpuscular haemoglobin concentration (MCHC) and mean corpuscular haemoglobin (MCH) between control and treated mulard ducks (Table 3).

Table 2. Changes in total and differential white blood cell counts (heterophil, eosinophil, lymphocyte, monocyte and basophil percentages) in mulard ducks receiving aflatoxin B₁ (AFB₁) either alone or together with Mycotox NG

Groups	White blood cell, G/l		Heterophil Le, %		Eosinophil Le, %		Lymphocyte, %		Monocyte, %		Basophil Le, %	
	Day 21	Day 42	Day 21	Day 42	Day 21	Day 42	Day 21	Day 42	Day 21	Day 42	Day 21	Day 42
I	20.41± 1.04	22.39± 1.19	30.00± 1.57	31.2± 1.81	2.70± 0.36	2.60± 0.13	62.3± 1.07	61.3± 1.07	3.60± 0.30	3.70± 0.36	1.40± 0.30	1.20± 0.29
II	28.08± 0.79 ^{1c}	32.95± 0.98 ^{1c}	41.6± 0.93 ^{1c}	44.5± 1.00 ^{1c}	2.30± 0.21	2.40± 0.33	51.00± 1.47 ^{1c}	48.00± 2.07 ^{1c}	3.80± 0.38	3.60± 0.26	1.30± 0.36	1.50± 0.34
III	31.85± 0.76 ^{1c,2a}	38.25± 1.18 ^{1c,2a}	45.1± 1.29 ^{1c}	48.1± 1.73 ^{1c}	2.50± 0.22	2.40± 0.50	47.2± 1.34 ^{1c,2a}	45.00± 1.07 ^{1c}	3.60± 0.34	3.40± 0.54	1.60± 0.45	1.10± 0.23
IV	24.32± 1.07 ^{1a,2a,3c}	27.69± 1.74 ^{1a,2a,3c}	36.2± 1.57 ^{1a,2a,3c}	37.1± 1.44 ^{1a,2b,3c}	2.40± 0.45	2.20± 0.41	57.1± 1.15 ^{1a,2a,3c}	55.6± 1.19 ^{1a,2b,3c}	3.00± 0.47	3.50± 0.34	1.30± 0.33	1.60± 0.40

*Data are presented as mean ± SEM; n=20 in each group; P<0.05; P<0.01; P<0.001; 1 vs control group; 2 vs experimental group I; 3 vs experimental group II

Discussion

Pathomorphological studies

The bone marrow of ducklings from Groups II and III exhibited hypocellularity, reduced number of haematopoietic stem cells and initial extent of fatty infiltration (Figure 1). These events were more pronounced in birds from Group III (Figure 2). In birds from Group IV, bone marrow hypocellularity was significantly less pronounced than that of Groups II and III (Figure 3). Histopathological changes in the bone marrow of control birds were not observed.

Aflatoxins, their metabolite and reactive oxygen species (ROS) produced by them impair haematopoiesis in bone marrow and lymphatic organs, where blood cells and other blood components are formed (Halliwell, 2007). Aflatoxins induce haematological disturbances (Dietert et al., 1983). They lead to haematopoietic suppression and anaemia by reduction of red blood cells, mean corpuscular volume and haemoglobin (Reddy and Waliyar, 2012).

Table 3. Changes in mean corpuscular volume (MCV), mean corpuscular haemoglobin concentration (MCHC) and mean corpuscular haemoglobin (MCH) in mulard ducks receiving aflatoxin B₁ (AFB₁) either alone or together with Mycotox NG

Groups	MCV, pg		MCHC, g/l		MCH, f/l	
	Day 21	Day 42	Day 21	Day 42	Day 21	Day 42
I	129.94± 4.80	130.26±4.06	347.62±6.59	351.95±7.74	45.05±1.57	45.80±1.21
II	128.28±4.14	128.57±4.10	367.42±13.13	369.02±17.34	47.15±1.21	46.62±1.29
III	123.53±2.90	118.89±4.40	388.56±11.69	394.21±20.31	47.75±0.78	46.36±1.96
IV	129.58±4.00	126.06±3.72	363.00±9.47	357.33±6.62	46.79±1.05	44.98±1.35

*Data are presented as mean ± SEM; n=20 in each group; P<0.05; P<0.01; P<0.001; 1 vs control group; 2 vs experimental group I; 3 vs experimental group II

Figure 1. Reduced number of haematopoietic stem cells in the bone marrow of ducklings from Group II, receiving 0.5 mg/kg aflatoxin B₁ with the feed. H/ E, bar 20 µm

Figure 2. Hypocellularity and higher extent of fatty infiltration of haematopoietic stem cells in the bone marrow of ducklings from Group III, receiving 0.8 mg/kg aflatoxin B₁ with the feed. H/ E, bar 20 µm

Figure 3. Considerably lower bone marrow hypocellularity in ducklings from Group IV, receiving 0.5 mg/kg aflatoxin B₁ and 2 g/kg Mycotox NG with the feed. H/ E, bar 20 µm

It is reported that aflatoxins cause lympho- and monocytopenia and at the same time, increase neutrophil percentage in blood (Donmez et al., 2012). In cattle, blood coagulation disorders with prolonged prothrombin time has been observed (Thrasher, 2012).

Reduced haematocrit, haemoglobin, erythrocytes and lymphocyte percentage was observed in broiler chickens with experimental aflatoxicosis, probably due to the inhibition effect of aflatoxins on haematopoietic organs (Campbell et al., 1983; Huff et al., 1986, 1988; Kubena et al., 1990; Kececi et al., 1995, 1998; Oguz et al., 2000; Sakhare et al., 2007; Mohamed and Mohamed, 2009). Similar changes were observed in this study as well. On the other hand, this decrease in red blood parameters results from inhibition of protein synthesis (Kubena et al., 1993; Abdel-Wahhab, et al., 2002) or the faster degradation of erythrocytes in the spleen (Mokif and Muiz, 2015). It was found out that aflatoxin B₁ decreases iron-binding capacity of erythrocytes (Harvey et al., 1991) and impairs the metabolism of some trace elements (Cu, Zn) involved in haematopoiesis (Abdel-Wahhab, et al., 2002). Reduced red blood cell counts and mean corpuscular volume is reported in broiler chickens fed feeds contaminated with aflatoxins (Singh et al., 1992), ochratoxins (Doerr and Huff, 1980; Aved et al., 1991), or combination of the two mycotoxins (Sawarkar et al., 2011). Similar changes in red blood cell counts and mean corpuscular volume were found out in fish fed feeds containing aflatoxins (El-Bouhy et al., 1993). The observed anaemia is secondary to the lower rate of DNA and protein synthesis and is enhanced by the deficiency of B₁₂ and folic acid (Tessari et al., 2006). Lower values of haematological parameters were probably due to protein synthesis inhibition (Kaneko et al., 1989), lower iron-binding capacity (Harvey et al., 1991) or impaired function of haematopoietic organs (Van Vleet et al., 1992; Abdel-Wahhab et al., 2002). Observed oligochromaemia could be explained with the small size of erythrocytes, impaired blood synthesis in bone marrow or lower rate of red blood cells formation (Sharma et al., 2011). Impaired haematopoiesis could be also attributed to aflatoxin-induced oxidative stress (Oven et al., 2000).

Increased white blood cell counts and heterophil percentages are presumably due to the irritating effect of aflatoxins on gastrointestinal mucosa and its inflammation (EL-Lethey and El-Zorb, 2004; Sakhare et al., 2007; Safamehr, 2008). Observed lymphocytopenia could be attributed to the toxic effects of aflatoxins on cells in the peripheral circulation or suppression of bone marrow and lymphoid organs function (Oguz et al., 2000; Mohamed and Mohamed, 2009). Increased white blood cell counts and heterophil percentages could be also explained with bone marrow changes (Abdel-Wahhab et al., 2002).

Increase in the heterophil percentages might indicate a tendency of the organism to compensate for the decrease of its resistance (Sova et al., 1991). In mice treated with T-2 toxin, Sinozuka et al. (1998) reported decreased cell density in the haematopoietic parenchyma. The changes were observed in the metaphysis and diaphysis regions. Hypocellularity was a result from reduction of myelocytes, immature granulocytes, erythroblasts and lymphocytes. Myelocytes exhibited karyorrhexis, karyopyknosis, fragmentation and neutrophils engulfed by adjacent macrophages. In megakaryocytes, nuclei with irregular size, increased number of nuclei and intracellular cells were found out. Bone marrow and spleen red pulp hypoplasia were reported in mice challenged with T-2 toxin (Smith et al., 1994).

Contrary to the present results, Aved et al. (1991), Mohiuddin et al. (1993) and Mohamed and Mohamed (2009), reported lower white blood cell counts in broiler chickens that received feed contaminated with aflatoxins and ochratoxins. Decrease blood cell counts is a reflects deleterious effects of AFB₁ on haemopoietic tissue (Mohiuddin et al., 1993) and immune systems (Oguz, 1997). Furthermore the variation in the hematological values may be due to the temperature variations, weather changes, dose levels, geographical distribution or age related factors (Mahmood et al., 2017).

The lack of statistically significant changes in eosinophils, monocytes and basophils observed in this study agreed with other reported results (Oguz et al., 2000; Mohamed and Mohamed, 2009). Similarly, the established lack of changes in mean corpuscular volume (MCV), mean corpuscular haemoglobin concentration (MCHC) and mean corpuscular haemoglobin (MCH) in mulard ducks were in line with the previous results of other authors (Oguz et al., 2000; Basmacioglu et al., 2005).

The utilisation of mycosorbents is one of the most reliable methods for reduction of the toxic effect of mycotoxins on animals (Dakovic et al., 2005). Adsorbents have a high affinity to aflatoxin binding in the animal gastrointestinal tract and thus, they decrease their absorption and hence, deleterious effect on birds (Raju and Dewegowda, 2000; Eraslan et al., 2005). Previously, the harmful effects of aflatoxin B₁ on haematological parameters was achieved by adding mycosorbents to the compound feeds of broiler chickens (esterified glucomannan, clinoptilolite) (Oguz et al., 2000; Aravind et al., 2003; Basmacioglu et al., 2005). The negative polarity of mycotoxins is attracted by the positive polarity of the toxin binder - a mechanism through which toxin are immobilized and eliminated from the organism of animals (Kana et al., 2014).

Conclusion

The supplementation of mulard ducks' feed with increasing doses of AFB₁ only (0.5 or 0.8 mg/kg feed) induced changes in haematological parameters with oligochromaemia, erythropania, reduced haematocrit, thrombocytopaenia, leukocytosis with heterophilia and lymphocytopaenia) as well as was accompanied with bone marrow hypocellularity. The supplementation of the feed with 2 g/kg mycosorbent – Mycotox NG to the feed containing 0.5 mg/kg AFB₁ reduced efficiently the harmful effect of AFB₁ on studied haematological indices and bone marrow lesions.

References

- Abdel-Wahhab MA, Nada SA and Khalil FA**, 2002. Physiological and toxicological responses in rats fed aflatoxincontaminated diet with or without sorbent materials. *Animal Feed Science and Technology*, 97, 209-219.
- Al-Daraji HJ, Al-Ani IA, Minati JK and Al-Hiti HE**, 2004. The use of different methods to suppress the effect of aflatoxicosis and its influence on certain blood traits in broiler. *Iraqi Journal of Agricultural Science*, 35, 103-112.
- Angelov A, Ibrishimov, N and Milashki S**, 1999. *Clinical Laboratory Investigations in Veterinary Medicine*. Prof. Marin Drinov Publishing House, Sofia (Bg).
- Aravind KL, Patil VS, Dewegowda G, Umakantha B and Ganpule SP**, 2003. Efficacy of esterified glucomannan to counteract mycotoxicosis in naturally contaminated feed on performance and serum biochemical and hematological parameters in broilers. *Poultry Science*, 82, 571-576.
- Aved IA, Dafella R, Yogi AF and Adam SEI**, 1991. Effect of ochratoxin A on Lohaman type chicks. *Veterinary and Human Toxicology*, 33, 357-360.
- Basmacioglu H, Oguz H, Ergul M, Çol R and Birdane YO**, 2005. Effect of dietary esterified glucomannan on performance, serum biochemistry and haematology in broilers exposed to aflatoxin. *Czech Journal of Animal Science*, 50, 1, 31-39.
- Bintvihok A, Thiengnin S, Doi K and Kumagai S**, 2002. Residues of Aflatoxins in the Liver, muscle and Eggs of Domestic Fowls. *Journal of Veterinary Medical Science*, 64, 11,
- Campbell M.L, May JD, Huff WE and Doeve JA**, 1983. Evaluation of immunity of young broiler chickens during aflatoxicosis and ochratoxicosis. *Poultry Science*, 62, 2138-2144.
- Cheeke PR**, 1998. *Natural toxicants in Feed, forages and Poisonous Plants*. 2nd ed. Interstate publishers, Danville, IL.
- Dakovic A, Tomasevic-Canovic M, Dondur V, Rottinghaus GE, Medakovic V and Zaric S**, 2005. Adsorption of mycotoxins by organozeolites. *Colloids and Surfaces B Biointerfaces*, 46, 20-25.
- Dalvi RR**, 1986. An overview of aflatoxicosis of poultry: its characteristics, prevention and reduction. *Veterinary Research Communications*, 10, 429-443.
- Dietert RR, Bloom SE, Qureshi MA and Nanna UC**, 1983. Hematological Toxicology following Embryonic Exposure to Aflatoxin-B₁. *Experimental Biology and Medicine*, 173, 4, 481-485.
- Directive 2002/32/EC** of the European parliament and of the council of 7 may 2002 on undesirable substances and products in animal feed. *Official Journal*, No. L, 140, 30, 5, 2002.
- Doerr JA and Huff RB**, 1980. Interactive effects of aflatoxin and ochratoxin A on some blood constituents in broiler chickens. *Poultry Science*, 59, 1600-1602.
- Donmez N, Dönmez HH, Keskin E and Kisadere İ**, 2012. Research Article Effects of Aflatoxin on Some Haematological Parameters and Protective Effectiveness of Esterified Glucomannan in Merino Rams. *The Scientific World Journal*, 10.1100/2012/342468, 1-4.
- El-Bouhy ZM, Ali AA and Helmy MS**, 1993. Preliminary studies on aflatoxicosis of contaminated food. *Zagazig Veterinary Journal* 21, 607-624.

- EL-Lethey H and El-Zorb H**, 2004. Effect of dimethyl diphenil bicarbohylate (DOB) on roilers during experimental aflatoxicosis. In: Proceedings of the 1st Annimal Conference, Faculty Veterinary Medicine, Moshtohor, September, 2004, 215-233.
- Eraslan G, Essiz D, Akdogan M, Sahindokuyucu F and Altintas L**, 2005. The Effects of Aflatoxin and Sodium Bentonite Combined and Alone on Some Ilood Electrolyte Levels in Broiler Chickens. Turkish Journal of Veterinary & Animal Sciences, 29, 601-605.
- Fonseca H**, 2003. Legislação sobre micotoxinas. www.micotoxinas.com.br.
- Girish CK and Devegowda G**, 2006. Efficacy of Glucomannan-containing yeast product (Mycosorb®) and hydrated sodium calcium aluminosilicate in preventing the individual and combined toxicity of aflatoxin and T-2 toxin in commercial broilers. Asian-Australian Journal of Animal Science, 19, 877-883.
- Halliwell B**, 2007. Oxidative stress and cancer: have we moved forward? Biochemistry Journal, 401, 1-11.
- Han XY, Huang QC, Li WE and Xu ZR**, 2008. Changes in growth performance, digestive enzyme activities and nutrient digestibility of cherry valley ducks in response to aflatoxin B₁ levels. Livestock Science, 119, 216-220.
- Harvey RB, Kubena LF, Phillips TD, Elissalde MH and Huff WE**, 1991. Diminution of aflatoxin toxicity to growing lambs by dietary supplementation with hydrated sodium calcium aluminosilicate. American Journal of Veterinary Research, 52, 152-156.
- Hashmi I, Pasha TN, Jabbar MA, Akram M and Hashmi AS**, 2006. Study of adsorption potential of yeast sludge against aflatoxins in broiler chicks. Journal of Animal and Plant Science, 16, 1-2.
- Huff W, Harvey RB, Kubena LF and Rottinghaus GE**, 1988. Toxic synergism between aflatoxin and T-2 toxin in broiler chicken. Poultry Science, 67, 1418-1420.
- Huff WE, Kubena LF, Harvey RB, Hagler WM, Sorenson SP, Philips TD and Giegler CR**, 1986. Individual and combined effect of aflatoxin and deoxynivalenol (DON, Vomitoxin) in broiler chicken. Poultry Science, 65, 1291-1298.
- Hussein HS and Brasel JM**, 2001. Toxicity, metabolism, and impact of mycotoxins on humans and animals. Toxicology, 167, 101-134.
- Huwig A, Freimund S, Käppeli O and Dultler H**, 2001. Mycotoxin detoxication of animal feed by different absorbents. Toxicology Letters, 122, 691-699.
- International Agency for Research on Cancer (IARC)**, 2002. Summaries and evaluations: aflatoxins. Lyon, IARC Press, 82, p.171.
- Kamei K and Watanabe A**, 2005. *Aspergillus* mycotoxin and their effect on the host. Medical Mycology Supplement, 43, 95-99.
- Kana JR, Ngoula F, Tchoffo H, Tadondjou CD, Sadjo YR, Teguia A and Gnonlonfin Gbemenou JB**, 2014. Effect of biocharcoals on hematological, serum biochemical and histological parameters in broiler chickens fed aflatoxin B₁-contaminated diets. Journal of Animal Science Advances, 4, 939-948.
- Kana JR, Teguia A, Mungfu BM and Tchoumboue J**, 2011. Growth performance and carcass characteristics of broiler chickens fed diets supplemented with graded levels of charcoal from maize cob or seed of *Canarium schweinfurthii* Engl. Tropical Animal Health and Production, 43, 51-56.
- Kaneko JJ**, 1989. Clinical Biochemistry of Domestic Animals (4th ed.). Academic Press, San Diego, CA.
- Kececi T, Demet Q and Oguz H**, 1995. Single and combination effects of dietary aflatoxin and adsorbent (Mycofix Plus) on some hematological and serum biochemical parameters of broiler chickens. Journal of Veterinary Science, 11, 95-101.
- Kececi T, Oguz H, Kurtoglu V and Demet O**, 1998. Effects of polyvinylpyrrolidone, synthetic zeolite and bentonite on serum biochemical and hematological characters of broiler chickens during aflatoxicosis. British Poultry Science, 39, 3, 452-458.
- Kennedy D, Delaney A and Koren G**, 1998. Mutagens, carcinogens and teratogens. Mutagens, Carcinogens and Teratogens. In: Goldfrank's Toxicologic Emergencies (ed. L. Goldfrank), 6th Edn. Appleton & Lange, Stanford, 262-273.
- Kubena L, Harvey RB, Phillips TD and Clement BA**, 1993. Effect of hydrated sodium calcium aluminosilicates on aflatoxicosis in broiler chicks. Poultry Science, 72, 651-657.
- Kubena LF, Harvey RB, Huff WE, Corrier DE, Phillips TD and Rottinghaus GE**, 1990. Efficacy of a hydrated sodium calcium aluminosilicate to reduce the toxicity of aflatoxin and T-2 toxin. Poultry Science, 69, 1078-1086.
- Kurtzman CP, Horn BW and Hesseltine CW**, 1987. *Aspergillus nomius*, a new aflatoxin-producing species related to *Aspergillus flavus* and *Aspergillus tamarii*. Antonie Leeuwenhoek, 53, 147-158.
- Lillehoj EB**, 1983. "Effect of environmental and cultural factors of aflatoxin contamination of developing corn kernels." In: Aflatoxin and *A. flavus* in corn (eds. V.L. Diener, R.L. Asquith and J.W. Dickens). Southern Coop Serv. Bull. 279, Craftmaster, Opelika, Ala, p. 112.
- Mabbett T**, 2005. Integrated management of mycotoxins. Poultry International, 44, 10-14.
- Mahmood S, Younus M, Aslam A and Anjum AA**, 2017. Toxicological effects of aflatoxin B₁ on growth performance, humoral immune response and blood profile of Japanese quail. The Journal of Animal & Plant Sciences, 27, 833-840.
- Mohamed AH and Mohamed MH**, 2009. Haemato-biochemical and pathological studies on aflatoxicosis and treatment of broiler chicks in Egypt. Veterinaria Italiana, 45, 323-37.
- Mohiuddin SM, Reddymv MV, Reddy MM and Ramakrishnan K**, 1986. Studies on phagocytic activity and hematological changes in aflatoxicosis in poultry. Indian Veterinary Journal, 63, 442-445.
- Mohiuddin SM, Warasi SMA and Reddy MV**, 1993. Haematological and biochemical changes in experimental ochratoxicosis in broiler chicken. Indian Veterinary Journal, 70, 613-617.
- Mokif SA and Muiz AMI**, 2015. Effect of some Medicinal Plants in Reducing Aflatoxicosis on Erythrocytes and Organ Weights in Albino Rat. European Journal of Academic Essays, 2, 98-101.
- Oguz H**, 2012. Detoxification of aflatoxin in poultry feed: a review from experimental trials. Lohmann Information, 47, 45-56.
- Oguz H, Kececi T, Birdane YO, Onder F and Kurtoglu V**, 2000. Effect of clinoptilolite on serum biochemical and haematological characters of broiler chickens during experimental aflatoxicosis. Research in Veterinary Science, 69, 89-93.
- Oguz H**, 1997. The preventive efficacy of polyvinyl pyrrolidone (PVPP) alone and its combination with the other adsorbents into broiler feeds against aflatoxicosis. Thesis for PhD, University of Selc, UK, Institute of Health Sciences, Kenya.
- Owen RW, Giacosa A, Hull WE, Spiegelhalter B and Bartsch H**, 2000. Identification of lignans as major components in the phenolic fraction of olive oil. Clinical Chemistry, 46, 976-988.
- Raju M and Devegowda G**, 2000. Influence of esterified glucomannan on performance and organ morphology, serum biochemistry and haematology in broilers exposed to individual and combined mycotoxicosis (aflatoxin, ochratoxin and T-2 toxin). British Poultry Science, 41, 640-650. (2012).

- Ramos AJ, Fink-Gremmels E and Hernández E**, 1996. Prevention of toxic effects of mycotoxins by means of nonnutritive absorbent compounds. *Journal of Food Protection*, 59, 631-641.
- Reddy SV and Waliyar F**, 2012. Properties of aflatoxin and its producing fungi. *Aflatoxins*. <http://www.icrisat.org/aflatoxin/aflatoxin.asp> (Accessed on 8th June 2012).
- Regulation No. 44/20.04.2006** for veterinary medical requirements to animal rearing facilities, Official Gazette 41 Appendix 9.
- Royes JB and Yanong RP**, 2002. Molds in fish feeds and aflatoxicosis. Copyright by the University of Florida, Institute of Agricultural Science (UF/IFAS).
- Safameher A**, 2008. Effects of clinoptilolite on performance, biochemical parameters and hepatic lesions in broiler chickens during aflatoxicosis. *Journal of Animal and Veterinary Advances*, 7, 381-388.
- Sakhare PS, Harne SD, Kalorey DR, Warke SR, Bhandarkar AG and Kurkure NV**, 2007. Effect of Toxiroak polyherbal feed supplement during induced aflatoxicosis, ochratoxicosis and combined mycotoxicoses in broilers. *Veterinarski arhiv*, 77, 129-146.
- Sawarkar AR, Sonkusale PM, Kurkure NV, Jangade CR, Maini S and Ravikanth K**, 2011. Experimental afla & ochratoxin induced mixed mycotoxicosis in broiler and its amelioration with herbomineral toxin binder toxiroak gold. *International Journal of Poultry Science*, 10, 560-566.
- Sharma V, Gupta R, Mishra N and Sharma S**, 2011. Influences of *Tinospora cordifolia* root extract supplementation on hematological and serological parameters of male mice exposed to aflatoxin B₁. *International Journal of Pharmacology*, 7, 659-663.
- Shinozuka J, Suzuki M, Noguchi, N, Sugimoto T, Uetsuka K, Nakayama H and Doi K**, 1998. T-2 toxin-induced apoptosis in hematopoietic tissues of mice. *Toxicologic Pathology*, 26, 674-681.
- Singh A, Satija KC and Mahipal SK**, 1992. Haematological and biochemical studies on broiler chicks fed aflatoxin B₁ and after its withdrawal. *Indian Journal of Poultry Science*, 27, 153-156.
- Smith BJ, Holladay SD and Blaylock BL**, 1994. Hematopoietic alterations after exposure to T-2 mycotoxin. *Toxicon*, 32, 15-1123.
- Sova Z, Pohunkova H, Reisnerova H, Slamova A and Haisl K**, 1991. Hematological and histological response to the diet containing aflatoxin B₁ and zeolite in broilers of domestic fowl. *Acta Veterinaria Brno*, 60, 31-40.
- Sudakin DL**, 2003. Trichothecenes in the environment: relevance to human health. *Toxicology Letters*, 143, 97-107.
- Terao K and Ueno Y**, 1978. Morphological and functional damage to cells and tissues. In: *Toxicology, Biochemistry and Pathology of Mycotoxins* (eds. K. Uraguchi and M. Yamazaki). Kodansha Press, Tokyo. pp. 189-238.
- Tessari ENC, Oliveira CAF, Cardoso ALSP, Ledoux D and Rottinghaus G**, 2006. Parâmetros hematológicos de frangos de corte alimentados com ração contendo aflatoxina B₁ e fumonisina B₁. *Ciência Rural*, 36, 924-929.
- Thrasher JD**, 2012. Aflatoxicosis in animals. *Aflatoxins and Health*, www.alphabiojuice.com/AFLATOXICOSIS_IN_ANIMALS.pdf.
- Tuzcu M, Sur E, Celik I, Oznurlu Y and Ciftci MK**, 2010. Effects of aflatoxin on the proportions of peripheral blood leukocyte and Alpha-Naphthyl Acetate Esterase (ANAE) positive lymphocyte in the mouse. *Kafkas University Veterinary Faculty Journal*, 16, 337-341.
- Van Kessel TFM and Hiang-Chek N**, 2004. Aflatoxin binders-how to get the best value for money. *International Poultry Production*, 12, 33-35.
- Van Vleet JF and Ferrans VJ**, 1992. Etiologic factors and pathologic alterations in selenium-vitamin E deficiency and excess in animals and humans. *Biological Trace Element Research*, 33, 1-21.
- Wilson DM and Payne GA**, 1994. Factors affecting *Aspergillus* *Flavus* group infection and aflatoxin contamination of crops. In: *The Toxicology of Aflatoxins. Human Health, Veterinary, and Agricultural Significance* (eds. D.L. Eaton and J.D. Groopman). Academic Press, San Diego, California, pp. 309-329.
- Wu Q, Jezkova A, Yuan Z, Pavlikova L and Dohnal Vand Kuca K**, 2009. Biological degradation of aflatoxins. *Drug Metabolism Reviews*, 41, 1-7.
- Zain ME**, 2011. Impact of mycotoxins on humans and animals. *Journal of Saudi Chemical Society*, 15, 129-144.

Genetics and Breeding

- Usability of metadata analysis of goat genetic resources among five countries from Africa, Asia and Europe: Metadata analysis of goat genetic** 183
M.M. Musthafa, T. Hussain, M.E. Babar, R.S. Aljumaah, M.A. Alshaikh, I. Muritala, V. Landi, A. Martinez, M. Amills, O. Dadi, J.V. Delgado, A.B.J. Aina, A.A. Onasoga, O.A. Adebambo, C. Visser, E. Van Marle-Köster, A.O. Adebambo, F.M.M.T. Marikar
- Knezha 560 – a new mid-late maize hybrid** 191
V. Valkova, N. Petrovska
- Sources of resistance in chickpea (*Cicer arietinum* L.) to ascochyta blight (*Ascochyta rabiei*)** 195
M. Koleva, Y. Stanoeva, I. Kiryakov, A. Ivanova
- Variability and grain yield potential of maize (*Zea mays* L.) genotypes under irrigated condition in central Sudan** 199
M.B. Alhussein, S.H. Suliman, A.A. Mohammed

Nutrition and Physiology

- Effect of monosodium glutamate dietary supplementation on some productive traits of common carp (*Cyprinus carpio* L.), cultivated in net cages** 204
G. Zhelyazkov
- Effect of experimentally induced aflatoxicosis on haematological parameters and bone marrow morphology in mulard ducks** 208
I. Valchev, N. Groseva, D. Kanakov, Ts. Hristov, L. Lazarov, R. Biinev
- Effect of dietary phytoextracts supplementation on the chemical composition and fatty acid profile of rainbow trout (*Oncorhynchus mykiss* W.), cultivated in recirculation system** 215
K. Georgieva, G. Zhelyazkov, Y. Staykov, D. Georgiev

Production Systems

- Yield and seed quality of some soybean (*Glycine max*. L) varieties, cultivated in Osmaniye region, Turkey** 222
F.F. Aşık, R. Yıldız
- Productivity and yield stability at late treatment of durum wheat (*Triticum durum* Desf.) with antibroadleaved herbicides.** 227
I. Influence at treatment during 1st stem node stage
Gr. Delchev, D. Delchev
- The effects of inoculation and N fertilization on soybean [*Glycine max* (L.) Merrill] seed yield and protein concentration under drought stress** 232
O. Basal, A. Szabó
- Soil structure after treatment with different operation modes of spading machine** 236
Y. Stoyanov, K. Trendafilov, N. Delchev, G. Tihanov

- Application of herbicides on common winter wheat (*Triticum aestivum* L.) at different doses and their reflection on the structural elements of spike** **241**
Z. Petrova, M. Nankova

Agriculture and Environment

- Differences in carbon forms under two land use types in Abia State, South-east Nigeria** **246**
B.N. Ndukwu, D.N. Osujieke, C.M. Ahukaemere, P.E. Imadojemu

- Theoretical analysis of the heat energy savings in wood pellets production** **253**
R. Georgiev, K. Peychev, V. Dimova, D. Georgiev

- Agricultural characteristics of sugar factory waste products** **257**
B.B. Aşık, S. Dorak

Product Quality and Safety

- Ontogenetic and diurnal variations of essential oil content of *Hypericum montbretii* Spach, cultivated in Kazdağı (Edremit/Balıkesir), Turkey** **262**
C. Paşa, E. Esendal, T. Kiliç

- Effect of *Artemisia annua* L. extract on growth performance, biochemical blood parameters and meat quality of rainbow trout (*Oncorhynchus mykiss* W.), cultivated in recirculating system** **266**
R. Koshinski

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors
The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings: Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows, IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 10, Number 3
September 2018

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz