UDC 553.982.2

Article / Статья

© PNRPU / ПНИПУ, 2020

Development of Zone Forecast Probability Models for Oil and Gas Potential in the Central Part of the Permian Uplift by Structural and Capacity Criteria**Konstantin A. Koshkin¹, Ilya A. Tatarinov²**¹Uraloil LLC (4 Sibirskaya st., Perm, 614990, Russian Federation)²NAST-M LLC (Office 1, 12a Makarenko st., Perm, 614107, Russian Federation)**Разработка вероятностных моделей зонального прогноза нефтегазоносности центральной части Пермского свода по структурно-мощностным критериям****К.А. Кошкин¹, И.А. Татарин²**¹ООО «УралОйл» (Россия, 614990, г. Пермь, ул. Сибирская, 4)²ООО «НАСТ-М» (Россия, 614107, г. Пермь, ул. Макаренко, 12а, офис 1)

Received / Получена: 14.07.2020. Accepted / Принята: 02.11.2020. Published / Опубликовано: 11.01.2021

Keywords:

oil and gas geological zoning, probabilistic and statistical criteria for oil and gas content, non-localized oil resources, Visean terrigenous oil and gas complex, structural and capacity criteria, correlation coefficient, probability, oil, prospecting and exploration work, structural wells, marking surfaces, zones demarcation, statistical analysis, seismic exploration, absolute mark.

The task to which this article is devoted has not received due attention in recent years, since the preparation of structures for deep drilling is carried out using seismic exploration. At the same time, there is a huge array of data on structured drilling, unfortunately, it has not been fully used. Thus, the goal of the study is to use data on structured drilling to solve not only structural problems for marking surfaces, but also more complex ones related to zonal oil and gas potential of territories.

The forecast of oil and gas content for marking and oil and gas horizons was carried out in three zones of oil and gas geological zoning. Using the data on these territories, studies were carried out to build probabilistic models for the zonal forecast of oil and gas content. To substantiate the joint use of data on marking surfaces and data on the tops of oil-bearing horizons, materials pertaining to 447 deep wells were studied.

Probabilistic models of zonal forecast of oil and gas content of the central part of the Permian uplift were developed according to structural and capacity criteria. The complex use of data on the absolute marks of deep and structured wells made it possible to rank the territory of the central part of the Permian uplift by the degree of zonal oil and gas content.

The Severokamskoye (0.73), Krasnokamskoye (0.67), Baklanovskoye (0.67), Polaznenskoye (0.67), Rassvetnoye (0.64) and Mezhevskoye (0.63) fields were characterized by the maximum values of PCOM. For the Kozubavskoye field, the PCOM was 0.57. The Gorskoye, Lobanovskoye, Talitskoye, Zorinskoye and Shemetinskoye fields were characterized by the minimum values of PCOM, varying in the range of 0.51–0.53. This scheme can be used when designing prospecting and exploration works in this area.

Ключевые слова:

нефтегазогеологическое районирование, вероятностно-статистические критерии нефтегазоносности, нелокализованные ресурсы нефти, визейский терригенный нефтегазоносный комплекс, структурно-мощностные критерии, коэффициент корреляции, вероятность, нефть, поисково-разведочные работы, структурные скважины, маркирующие поверхности, зоны районирования, статистический анализ, сейсморазведка, абсолютная отметка.

Задаче, которой посвящена данная статья, в последние годы не уделяется должного внимания, так как подготовку структур к глубокому бурению производят с помощью сейсморазведки. В то же время имеется огромный массив данных по структурному бурению, к сожалению, используются не полностью. Таким образом, актуальность применения данных по структурному бурению для решения не только структурных задач по маркирующим поверхностям, но и более сложных, связанных с зональной нефтегазоносностью территорий, не вызывает сомнений.

Прогноз нефтегазоносности по маркирующим и нефтегазоносным горизонтам проводился в трех зонах нефтегазогеологического районирования. С использованием данных по этим территориям выполнялись исследования для построения вероятностных моделей зонального прогноза нефтегазоносности. При обосновании совместного использования данных по маркирующим поверхностям и данных по кровлям нефтесодержащих горизонтов исследовались материалы по 447 глубоким скважинам.

Разработаны вероятностные модели зонального прогноза нефтегазоносности центральной части пермского свода по структурно-мощностным критериям. Комплексное использование данных по абсолютным отметкам глубоких и структурных скважин позволило ранжировать территорию центральной части Пермского свода по степени зональной нефтегазоносности.

Максимальными значениями $P_{\text{КОМ}}$ характеризуются Северокамское (0,73), Краснокамское (0,67), Баклановское (0,67), Полазненское (0,67), Рассветное (0,64) и Межевское (0,63) месторождения. Для Козубавского месторождения $P_{\text{КОМ}}$ равно 0,57. Горское, Лобановское, Талицкое, Зоринское и Шеметинское месторождения характеризуются минимальными значениями $P_{\text{КОМ}}$, изменяющимися в интервале 0,51–0,53. Эти данные могут быть использованы при проектировании поисково-разведочных работ.

Konstantin A. Koshkin (Author ID in Scopus: 18042091000) – Head of the Department of Geology and Licensing (tel.: +007 912 887 25 26, e-mail: koshkin@uraloil.com). The contact person for correspondence.

Ilya A. Tatarinov – Director (tel.: +007 912 059 81 99, e-mail: i_tatarinov@mail.ru).

Кошкин Константин Александрович – начальник отдела геологии и лицензирования (тел.: +007 912 887 25 26, e-mail: koshkin@uraloil.com). Контактное лицо для переписки.

Татарин Илья Анатольевич – директор (тел.: +007 912 059 81 99, e-mail: i_tatarinov@mail.ru).

Please cite this article in English as:

Koshkin K.A., Tatarinov I.A. Development of Zone Forecast probability Models for Oil and Gas Potential in the Central Part of the Permian Uplift by Structural and Capacity Criteria. *Perm Journal of Petroleum and Mining Engineering*, 2021, vol.21, no.1, pp.2-8. DOI: 10.15593/2712-8008/2021.1.1

Просьба ссылаться на эту статью в русскоязычных источниках следующим образом:

Кошкин К.А., Татарин И.А. Разработка вероятностных моделей зонального прогноза нефтегазоносности центральной части Пермского свода по структурно-мощностным критериям // Недропользование. – 2021. – Т.21, №1. – С.2–8. DOI: 10.15593/2712-8008/2021.1.1

Introduction

Probabilistic methods were used in a number of studies for the purpose of forecasting zonal and local oil and gas potential [1–17]. It is worth noting that a whole range of geological events were predicted using probabilistic models across territories that are characterized by contrasting geological structures [3–50].

This article focuses on the task that has been largely neglected in recent years as the structures are prepared for deep drilling using seismic exploration. That said, there is a plenitude of information available on the data regarding structured drilling that is regrettably overlooked.

The purpose of the study is to demonstrate that the data on structured drilling can be of use when it comes to handling both structural problems in marking surfaces and more complex ones related to zonal oil and gas potential of territories. The task was carried out based on the survey of the central part of the Permian uplift that enjoys a vast array of data on marking horizons and oil-bearing strata. The oil and gas potential was forecast pertaining to marking surfaces and oil- and gas-bearing horizons across this area of the Permian uplift within the three zones of oil and gas geological zoning (Fig. 1). Zone 1.B.4 includes the Krasnokamsky (Severokamskoye and Krasnokamskoye oil fields) and Osinsky (Belyayevskoye field) arches (Fig. 1). This zone also includes the northwest and central areas of the Permian uplift; zone 2.A.3 spans across the Gorskoye, Rassvetnoye, Baklanovskoye, Kozubayevskoye, and Lobanovskoye fields. Zone 2.A.4 features the Talitskoye, Yuzhno-Mezhevskoye, Mezhevskoye, Polaznenskoye, and Shemetinskoye fields.

The study drew on the data about these territories to develop probabilistic models of zonal forecasts for the oil and gas potential. The combined usage of data on marking surfaces and tops of oil-bearing horizons was substantiated based on materials retrieved across 447 deep wells.

Grounds for Building

Probabilistic Models to Forecast Zonal Oil and Gas Potential

The probabilistic models of zonal oil and gas potential forecasting were developed using the following absolute elevations of horizon tops and stages:

- Bobrikovskiy horizon – C_1bb ,
- terrigenous bench of Tula horizon – C_1tl_p ,
- carbonate bench of Tula horizon – C_1tl_k ,
- Bashkir stage – C_2b ,
- Vereysky horizon – C_2vr ,
- terrigenous bench of Artinsky stage – P_1ar_p ,
- carbonate bench of Artinsky stage – P_1ar_k ,
- Iremsky horizon – P_1ir .

The first round of the statistical analysis centered around calculating the values of correlation coefficients r across all absolute elevations under study (Table 1).

The r values demonstrate that there are two groups that show significantly high correlation ratios within. The first group of high correlation ratios is found between the marking surfaces. There are significant direct correlation ratios (P_1ar_p , P_1ar_k , P_1ir). The second group with high correlation ratios can be seen across all oil- and gas-bearing horizons under study (C_1bb , C_1tl_p , C_1tl_k , C_2b , C_2vr). It is important to point out that the correlation ratios are low between the oil- and gas-bearing horizons and marking surfaces, statistically insignificant, and range from 0.01 to –0.33 (see Table 1). It means that using absolute elevations of marking surfaces for zonal oil and gas potential forecasting proves highly problematic across the entire territory of the central area of the Permian uplift. Therefore, the zonal forecasting models that draw on data from marking

Fig. 1. Diagram of oil and gas geological zoning

horizons were developed as differentiated across the areas of oil and gas geological zoning.

The statistical models of zonal oil and gas potential forecasting were developed based exclusively on the data pertaining to the following absolute elevations:

- Vereysky horizon – C_2vr ,
- terrigenous bench of Artinsky stage – P_1ar_p ,
- carbonate bench of Artinsky stage – P_1ar_k ,
- Iremsky horizon – P_1ir .

It is predefined by two reasons: firstly, the Vereysky horizon in section is located in the closest proximity to marking surfaces; secondly, all oil- and gas-bearing horizons (C_1bb , C_1tl_p , C_1tl_k , C_2b , C_2vr) show positive statistically significant correlation ratios (r in the range from 0.94 to 0.99). Combining data on these horizons is not reasonable as the results are nearly identical. In addition, the study considered layer thickness between marking surfaces and the Vereysky horizon. The suggested technique of zonal oil and gas potential forecasting can be illustrated using data across the wells that are located within zone 1.B.4. Some statistical characteristics were defined for this zone (Table 2) using the data on the wells located within (Class 1) and outside the reservoir limits (Class 2). The values of layer elevations that mark the top of the Iremsky and Vereysky horizons as well as the thickness of section between them – m ($vr-ir$) are given in Table 2 as pertaining to the defined classes.

This shows that the absolute elevations of the top of the Iremsky horizon within the reservoir limits are much higher than those outside the limits. The average value $H(ir)$ for Class 1 is 147.2 m higher than the average value for Class 2. To estimate how the average values vary in terms of $H(ir)$, we shall find the value of t criterion using the following formula:

$$t_p = \frac{|X_1 - X_2|}{\sqrt{\frac{1}{n_1} + \frac{1}{n_2} \left(\frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2} \right)}}$$

where X_1 , X_2 stand for the average values of $H(ir)$ for Classes 1 and 2 correspondingly; and S_1^2 , S_2^2 are mean square deviations.

The difference in average values is deemed statistically significant if $t_p > t_r$. Values t_r are determined depending on

Table 1

Correlation matrix

Parameter	$H(ir)$	$H(ar_i)$	$H(ar_o)$	$H(vr)$	$H(b)$	$H(tl_o)$	$H(tl_i)$	$H(bb)$
$H(ir)$	1.00	0.98	0.98	-0.30	-0.33	-0.13	-0.10	-0.16
$H(ar_i)$		1.00		-0.18	-0.21	-0.02	0.01	-0.05
$H(ar_o)$			1.00	-0.18	-0.21	-0.02	0.01	-0.05
$H(vr)$				1.00	0.99	0.95	0.95	0.94
$H(b)$					1.00	0.95	0.94	0.94
$H(tl_o)$						1.00	0.99	0.98
$H(tl_i)$							1.00	0.98
$H(bb)$								1.00

Note: Significant correlation coefficients are highlighted.

Table 2

Some statistical characteristics for Zone 1.B.4

Parameter	Average	St. dev.	Min.	Max.
Parameter			$H(ir)$, m	
Class 1	-7.1	55.9	-193.6	105.2
Class 2	-154.3	89.4	-336.1	54.0
Parameter			$H(vr)$, m	
Class 1	-822.9	23.5	-906.6	-789.1
Class 2	-901.6	44.8	-1007.6	-818.2
Parameter			$m(vr-ir)$, m	
Class 1	815.8	55.0	709.0	938.0
Class 2	747.3	69.3	660.0	907.0

Table 3

 Distribution of values $H(ir)$

Item class	Variability range $H(ir)$							
	-350...-300	-300...-250	-250...-200	-200...-150	-150...-100	-100...-50	-50...0	0...50
Areas within the reservoir limits, $n = 70$				0.042		0.085	0.471	0.271
Areas outside the reservoir limits, $n = 49$	0.043	0.022	0.285	0.326	0.040	0.163	0.061	0.040

the amount of compared data and level of significance ($p = 0.05$). Indicator $H(ir)$ is characterized by $t_p = 11.286$, $p = 0.000000$. It shows that the average values for $H(ir)$ are statistically different.

In order to estimate the probability of finding the oil and gas potential for this zone using $H(ir)$, an individual forecasting model was built using a technique that is described quite in detail in the study [1]. Let us take a brief look at the technique of building individual probabilistic models using $H(ir)$ as an example. This technique was applied to study the distribution density across both classes. The first case considers data pertaining to values of $H(ir)$. Within the territories of oil fields is Class 1, $n_1 = 70$, the second case draws on the data across the areas outside the reservoir limits, which is Class 2, $n_2 = 49$. Then the cumulative value (set) of $H(ir)$ should be divided into units belonging to Class 1. Following the applied technique, the first step of building a probabilistic model includes histograms that are drawn for Classes 1 and 2 based on the $H(ir)$ data. The optimal values of the $H(ir)$ range are found using the Sturges' rule:

$$\Delta X = \frac{X_{\max} - X_{\min}}{1 + 3.32 \cdot \lg N},$$

where $X_{\max}$ is the maximum value of the parameter, $X_{\min}$ is the minimum value of the parameter, N is the amount of data.

Frequencies are determined for each range:

$$P(X) = \frac{N_k}{N_g},$$

where $P(X)$ is a frequency in the k -th range for Group $K_{\text{BMT}}^{\text{MO}}$ ($q = 1$), $q = 2$ corresponds to Group $K_{\text{BMT}}^{\text{MO}}$, N_k stands for the number of cases that contain indicator $P(X)$ within

the k range, and N_q is the volume of samples for Classes 1 and 2. The ratios of the share of units that fell within different variability ranges of $H(ir)$ were processed using the interval analysis shown in Table 3.

It is clearly seen that, as it relates to oil- and gas-bearing areas, higher $H(ir)$ entails more frequent occurrence of values. It is worth noting that oil- and gas-bearing areas within the range of -350...-200 m do not show any values, whereas outside the oil- and gas-bearing reservoir limits the sum frequency of their detection stands at 0.341. The areas outside the oil- and gas horizon starting from the range of -200...-150 m demonstrate less frequent occurrence of values. The modal range for oil- and gas-bearing areas spans across -50...0 m, and the areas outside the oil- and gas-bearing limits range within -200...-150 m. While in the range of -350...-100 m the ratio of the first class occurrence frequency to the second class is 0.055, in the range of -100...150 m this ratio is much higher and stands at 3.373. It all shows that $H(ir)$ values are distributed noticeably differently across the classes under study. The quantitative difference in distribution of values was based on Pearson's chi-squared test χ^2 following the formula

$$\chi^2 = N_1 N_2 \sum_{i=1}^e \frac{1}{M_1 + M_2} \left(\frac{M_1}{N_1} - \frac{M_2}{N_2} \right)^2,$$

where N_1 , N_2 are the number of $H(ir)$ values in Classes 1 and 2, respectively; M_1 , M_2 are the number of values that fall within the set range in both classes correspondingly; e is the number of intervals.

The values of χ^2 as per parameter $H(ir)$ equal to 85.061 at $p = 0.000000$. It indicates that distribution densities are statistically different. Further, each range gets specified in terms of probabilities of affiliation with oil-

 Fig. 2. Dependence: *a* – $P(ir)$ on $H(ir)$; *b* – $P(vr)$ on $H(vr)$; *c* – $P(vr-ir)$ on $m(vr-ir)$

 Fig. 3. Probabilistic diagram of oil and gas potential for Zone 1.B.4 as per parameter: *a* – $P(ir)$; *b* – $P(vr)$; *c* – $P(vr-ir)$; *d* – P_{COM} for Zone 1.B.4

Table 4

Item class	Distribution of P_{COM} values							
	Variability range, P_{COM}							
	0–0.1	0.1–0.2	0.2–0.3	0.3–0.4	0.4–0.5	0.5–0.6	0.6–0.7	0.7–0.8
Areas within the reservoir limits, $n = 70$		0.050				0.100	0.450	0.316
Areas outside the reservoir limits, $n = 49$	0.061	0.244	0.387	0.081	0.122	0.081	0.020	0.017

bearing areas. The interval probabilities of affiliation with Class 1 are juxtaposed with average interval values of $H(ir)_n$. The values of $P(H(ir)_n)$ and $H(ir)_n$ are used to calculate the matching correlation coefficient r and draw up a regression equation. The resulting model is further modified on the condition that the average value of probabilities for oil-bearing areas should exceed 0.5 and those outside the reservoir limits should be below 0.5.

The linear dependence of $P(ir)$ on $H(ir)$ is expressed in the following way: $P(ir) = 0.594 + 0.0015 H(ir)$. The linear probabilistic models for other parameters are built the same way and quoted further in the article.

The visual representation of the model is given in Figure 2, *a*.

It is clear that whenever the values of $H(ir)$ oscillate from -336.1 to 105.2 m the value of $P(ir)$ rises from 0.09 to 0.75.

This model was used to calculate $P(ir)$ for wells within the zone, which were used for the probabilistic algorithm of oil and gas potential in Zone 1.B.4 (Fig. 3, *a*).

It follows that the maximum probabilities of zonal oil and gas potential are located in the northeast part of this zone. The values of $P(ir)$ for the Krasnokamskoye and Severokamskoye fields found within the zone stand at 0.59 and 0.74, respectively.

The top of the Vereysky horizon, as can be seen in Table 2, lies much higher within the oil- and gas-bearing limits than outside them. The model based on $H(vr)$ can be expressed the following way: $P(vr) = 2.488 + 0.0023 H(vr)$. The visual representation of the model is given in Figure 2, *b*.

It shows that values of $H(vr)$ that change from -1007 to -789 m increase the value of $P(vr)$ from 0.17 to 0.67.

This model was the basis for the probabilistic algorithm of oil and gas potential for Zone 1.B.4 (Fig. 3, *b*).

The maximum probabilities are clearly seen to be located in the northeast part of this zone.

As shown in Table 2, the average value for the layer thickness between the Vereysky and Irensky horizons within the reservoir limits is higher (815 m) than that outside them (747 m). The probability of oil and gas potential in this zone can be expressed in the following way: $P(vr-ir) = 0.345 + 0.0002 m(vr-ir)$. The visual representation of this dependence is shown in Figure 2, *c*.

Therefore, whenever values $m(vr-ir)$ go up from 660 to 938 m, it increases value $P(vr-ir)$ insignificantly from 0.47 to 0.53. This indicator is typically less informative than the structural ones.

This model was the basis for the probabilistic algorithm of oil and gas potential in Zone 1.B.4 (Fig. 3, *c*).

The data presented in Figure 3 *c* demonstrate that higher values of probability ($P(vr-ir) > 0.5$) are characteristic not only of the Severokamskoye and Krasnokamskoye fields but also of the territory to the south of them.

Development of a Comprehensive Model of Zonal Oil and Gas Potential Forecasting

The analysis of the resulting models suggests that they cover the oil and gas potential of this zone but the variability ranges of the probability values differ considerably. To apply the aforementioned parameters comprehensively, we shall use a composite indicator that is calculated using the following formula:

$$P_{\text{COM}} = \frac{\prod_{j=1}^m P(W_1|X_j)}{\prod_{j=1}^m P(W_1|X_j) + \prod_{j=1}^m (1 - P(W_1|X_j))}$$

where $P(W_1/X_j)$ stands for respective probabilities: $P(ir)$, $P(vr)$ and $P(vr-ir)$.

The values of P_{COM} were set as a basis for the probabilistic algorithm of oil and gas potential for Zone 1.B.4 (Fig. 3, *d*). The P_{COM} values are distributed depending on the oil and gas potential of the territories and can be seen in Table 4.

The results in the table show that oil- and gas-bearing areas in 95 % of cases demonstrate $P_{\text{COM}} > 0.5$, whereas areas outside the reservoir limits are characterized by values below 0.5. The average value of P_{COM} in oil- and gas-bearing areas stands at 0.679 ± 0.117 , while outside the reservoir limits it is 0.309 ± 0.185 , as per parameter t they differ statistically ($t = 13.368$; $p = 0.000000$). The value of parameter χ^2 as per indicator P_{COM} stands at 103.947 at $p = 0.000000$. It means that distribution densities of P_{COM} are statistically different.

It is also obvious that higher values of probability ($P_{\text{COM}} > 0.5$) are clearly characteristic of the Severokamskoye and Krasnokamskoye oil fields, and the former shows a higher probability (0.84) than the latter (0.73). The rest of the area in Zone 1.B.4. shows little promise in terms of exploring for oil and gas deposits, as the structural parameters suggest.

References

- Galkin V.I., Rastegaev A.V., Galkin S.V. Veroyatnostno-statisticheskaya otsenka neftegazonosnosti lokal'nykh struktur [Probabilistic-statistical assessment of oil and gas content of local structures]. Ekaterinburg: Ural'skoe otdelenie Rossiiskoi akademii nauk, 2001, 277 p.
- Kozlova I.A., Galkin V.I., Vantseva I.V. K otsenke perspektiv neftegazonosnosti Solikamskoi depressii s pomoshch'yu geologo-geokhimicheskikh kharakteristik neftegazomaterinskikh porod [Evaluation of petroleum potential of Solikamsk depression based on geological and geochemical characteristics of oil and gas source rocks]. *Neftpromyslovoye delo*, 2010, no. 7, pp. 20-23.
- Krivoshchekov S.N., Galkin V.I., Nosov M.A. Otsenka nelokalizovannykh resursov nefli territorii Permskogo kraia pri pomoshchi sistemy elementarnykh uchastkov [Evaluation of non-localized oil resources in Perm region by a system of elementary sections]. *Neftianoe khoziaistvo*, 2014, no. 6, pp. 9-11.
- Krivoshchekov S.N., Kozlova I.A., Sannikov I.V. Otsenka perspektiv neftegazonosnosti zapadnoi chasti Solikamskoi depressii na osnove geokhimicheskikh i geodinamicheskikh dannykh [Estimate of the petroleum potential of the western Solikamsk depression based on geochemical and geodynamic data]. *Neftianoe khoziaistvo*, 2014, no. 6, pp. 12-15.
- Galkin V.I., Kozlova I.A., Melkishev O.A., Shadrina M.A. Geokhimicheskie pokazateli ROV porod kak kriterii otsenki perspektiv neftegazonosnosti [Geochemical indicators of dispersed organic matter (DOM) of rocks as criteria of hydrocarbon potential evaluation]. *Neftpromyslovoye delo*, 2013, no. 9, pp. 28-31.
- Kozlova I.A., Melkishev O.A. Prognosticheskaya otsenka raspredeleniya nelokalizovannykh resursov nefli v devonskom terrigenom komplekse na territorii Permskogo kraia [Predictive estimation of non-localized oil resources distribution in the Devonian terrigenous complex in Perm region]. *Geologiya, geofizika i razrabotka neflianykh i gazovykh mestorozhdenii*, 2017, no. 2, pp. 4-8.
- Galkin V.I., Kozlova I.A. Razrabotka veroyatnostno-statisticheskikh regional'no-zonal'nykh modelei prognoza neftegazonosnosti po dannykh geokhimicheskikh issledovaniy verkhnedevonskikh karbonatnykh otlozhenii [Development of probabilistic-statistical regional-zoning models of oil and gas potential prediction based on the data of geochemical studies of the Upper Devonian carbonate deposits]. *Geologiya, geofizika i razrabotka neflianykh i gazovykh mestorozhdenii*, 2016, no. 6, pp. 40-45.
- Galkin V.I., Kozlova I.A., Krivoshchekov S.N., Nosov M.A., Koltirina N.S. Otsenka perspektiv neftegazonosnosti iuga Permskogo kraia po organo-geokhimicheskim dannykh [Estimation of petroleum potential prospects in the south of Perm territory on the basis of organic-geochemical data]. *Neftpromyslovoye delo*, 2015, no. 7, pp. 32-35.
- Galkin V.I., Kozlova I.A., Krivoshchekov S.N., Nosov M.A. Reshenie regional'nykh zadach prognozirovaniya neftenosnosti po dannykh geologo-geokhimicheskogo analiza rasseiannogo organicheskogo veshchestva porod domaniyevskogo tipa [Solutions to regional problems of forecasting oil bearing according to geological and geochemical analysis of dispersed organic matter of Domanic type rocks]. *Neftianoe khoziaistvo*, 2015, no. 1, pp. 21-23.
- Galkin V.I., Kozlova I.A., Krivoshchekov S.N., Melkishev O.A. K obosnovaniyu postroeniya modelei zonal'nogo prognoza neftegazonosnosti dlia nizhnne-sredneveizeiskogo kompleksa Permskogo kraia [On the justification of the construction of models for oil and gas potential area forecast Visean deposits of Perm region]. *Neftianoe khoziaistvo*, 2015, no. 8, pp. 32-35.
- Galkin V.I., Zhukov Iu.A., Shishkin M.A. Primenenie veroyatnostnykh modelei dlia lokal'nogo prognoza neftegazonosnosti [Application of probabilistic models for local forecast of oil and gas content]. Ekaterinburg: Ural'skoe otdelenie Rossiiskoi akademii nauk, 1990, 108 p.
- Galkin V.I., Shaikhutdinov A.N. O vozmozhnosti prognoza neftegazonosnosti iurskikh otlozhenii veroyatnostno-statisticheskimi metodami (na primere territorii deiatel'nosti TPP "Kogalymneftegaz") [About possibility to forecast the oil-and-gas content of Jurassic sediments based on probable and statistical methods (case study of the territorial industrial enterprise "Kogalymneftegaz")]. *Geologiya, geofizika i razrabotka neflianykh i gazovykh mestorozhdenii*. Moscow: VNIIOENG, 2009, no. 6, pp. 11-14.
- Galkin V.I., Shaikhutdinov A.N. Postroenie statisticheskikh modelei dlia prognoza debitora nefli po verkhneiurskim otlozheniyam Kogalymnskogo regiona [Development of statistical models for predicting the oil flow rates by example Jurassic deposits of Kogalym region territory]. *Neftianoe khoziaistvo*, 2010, no. 1, pp. 52-54.
- Galkin V.I., Krivoshchekov S.N. Postroenie matritsy elementarnykh iacheek pri prognoze neftegazonosnosti veroyatnostno-statisticheskimi metodami na territorii Permskogo kraia [Construction of a matrix of elementary cells in predicting oil and gas content by probabilistic and statistical methods in the Perm Territory]. *Geologiya, geofizika i razrabotka neflianykh i gazovykh mestorozhdenii*. Moscow: VNIIOENG, 2008, no. 8, pp. 20-23.

Fig. 4. Probabilistic algorithm of oil and gas potential based on indicator P_{COM} for the central part of Permian uplift

Conclusion

The discussed technique was used to build individual and comprehensive models of other areas in the central part of the Permian uplift. The models specified the values of P_{COM} across all wells, which were further used for the probabilistic algorithm of zonal oil and gas potential for the central part of the Permian uplift (Fig. 4).

The approach suggests that combined probabilities cover oil- and gas-bearing limits in all fields under study. The maximum values of P_{COM} are shown by the Severokamskoye (0.73), Krasnokamskoye (0.67), Baklanovskoye (0.67), Polaznenskoye (0.67), Rassvetnoye (0.64), and Mezhevskoye (0.63) fields. The Kozubayevskoye field showed P_{COM} of 0.57. The Gorskoye, Lobanovskoye, Talitskoye, Zorinskoye, and Shemetinskoye fields are characterized by minimum P_{COM} values that vary within the range of 0.51–0.53.

As a result, it can be concluded that the comprehensive application of data on absolute elevations of deep and structural wells facilitates the process of ranging the area of the central Permian uplift by order of zonal oil and gas potentials. This approach can be used in designing surveys and explorations in the territory under study.

15. Galkin V.I., Krivoshechekov S.N. Obosnovanie napravlenii poiskov mestorozhdenii nefiti i gaza v Permskom krae [Justification of the directions of prospecting for oil and gas deposits in the Perm Territory]. *Nauchnye issledovaniia i innovatsii*. Perm', 2009, vol. 3, no. 4, pp. 3-7.
16. Galkin V.I., Rastegaev A.V., Kozlova I.A., Vantseva I.V., Krivoshechekov S.N., Voevodkin V.L. Prognoznaiia otsenka neftegazonosnosti struktur na territorii Solikamskoi depressii [Probable estimation of oil content of structures in territory of Solikamsk depression]. *Neftpromyslovoe delo*. Moscow: VNIIOENG, 2010, no. 7, pp. 4-7.
17. Belokon' T.V., Galkin V.I., Kozlova I.A., Pashkova S.E. Dodevonskie otlozheniia Permskogo Prikam'ia kak odno iz perspektivnykh napravlenii geologo-razvedochnykh rabot [Pre-Devonian deposits of the Perm Kama region as one of the promising areas of geological exploration]. *Geologiya, geofizika i razrabotka nefitnykh i gazovykh mestorozhdenii*. Moscow: VNIIOENG, 2005, no. 9, pp. 24-28.
18. Putilov I.S. Razrabotka tekhnologii kompleksnogo izucheniia geologicheskogo stroeniia i razmeshcheniia mestorozhdenii nefiti i gaza [Development of technologies for a comprehensive study of the geological structure and location of oil and gas fields]. Perm: Permskii natsional'nyi issledovatel'skii politekhnicheskii universitet, 2014, 285 p.
19. Galkin V.I., Kozlova I.A., Krivoshechekov S.N., Piatunina E.V., Pestova S.N. O vozmozhnosti prognozirovaniia neftegazonosnosti famenskikh otlozhenii s pomoshch'iu postroeniia veroiatnostno-statisticheskikh modelei [On the possibility of predicting the oil and gas content of the Famennian deposits using the construction of probabilistic and statistical models]. *Geologiya, geofizika i razrabotka nefitnykh i gazovykh mestorozhdenii*. Moscow: VNIIOENG, 2007, no. 10, pp. 22-27.
20. Galkin V.I., Solov'ev S.I. Raionirovanie territorii Permskogo kraia po stepeni perspektivnosti priobreteniia nefitnykh uchastkov nedr [Classification of Perm kraj areas according to prospectivity for oil fields acquisition]. *Vestnik Permskogo natsional'nogo issledovatel'skogo politekhnicheskogo universiteta. Geologiya. Neftgazovoe i gornoe delo*, 2015, no. 16, pp. 14-24.
21. Sosnin N.E. Razrabotka statisticheskikh modelei dlia prognoza neftegazonosnosti (na primere terrigenykh devonskikh otlozhenii Severo-Tatarskogo svoda) [Development of statistical models for predicting oil-and-gas content (on the example of terrigenous devonian sediments of North Tatar arch)]. *Vestnik Permskogo natsional'nogo issledovatel'skogo politekhnicheskogo universiteta. Geologiya. Neftgazovoe i gornoe delo*, 2012, no. 5, pp. 16-25.
22. Galkin V.I., Sosnin N.E. Razrabotka geologo-matematicheskikh modelei dlia prognoza neftegazonosnosti slozhnopoastroennykh struktur v devonskikh terrigenykh otlozheniakh [Geological development of mathematical models for the prediction of oil and gas complex-built structures in the Devonian clastic sediments]. *Neftianoe khoziaistvo*, 2013, no. 4, pp. 28-31.
23. Dement'ev L.F. Matematicheskie metody i EVM v neftegazovoi geologii [Mathematical methods and computers in oil and gas geology]. Moscow: Nedra, 1987, 264 p.
24. Davydenko A.Iu. Veroiatnostno-statisticheskie metody v geologo-geofizicheskikh prilozheniakh [Probabilistic and statistical methods in geological and geophysical applications]. Irkutsk, 2007, 29 p.
25. Mikhalevich I.M. Primenenie matematicheskikh metodov pri analize geologicheskoi informatsii (s ispol'zovaniem kompiuternykh tekhnologii) [Application of mathematical methods in the analysis of geological information (using computer technology)]. Irkutsk, 2006, 115 p.
26. Andreiko S.S. Razrabotka matematicheskoi modeli metoda prognozirovaniia gazodinamicheskikh iavlenii po geologicheskim dannym dlia uslovii Verkhnekamskogo mestorozhdeniia kaliinykh solei [Development of mathematical model of gas-dynamic phenomena forecasting method according to geological data in conditions of Verkhnekamskoe potash salt deposit]. *Vestnik Permskogo natsional'nogo issledovatel'skogo politekhnicheskogo universiteta. Geologiya. Neftgazovoe i gornoe delo*, 2016, no. 21, pp. 345-353. DOI: 10.15593/2224-9923/2016.21.6
27. Devis Dzh. Statistika i analiz geologicheskikh dannykh [Geological data statistics and analysis]. Moscow: Mir, 1977, 353 p.
28. Pomorskiy Iu.L. Metody statisticheskogo analiza eksperimental'nykh dannykh [Methods for statistical analysis of experimental data]. Leningrad, 1960, 174 p.
29. Cherepanov S.S. Kompleksnoe izuchenie treshchinovosti karbonatnykh zalezhei metodom Uorrena - Ruta s ispol'zovaniem dannykh seismofatsial'nogo analiza (na primere turnefamenskoi zalezhi Ozernogo mestorozhdeniia) [Integrated research of carbonate reservoir fracturing by Warren - Root method using seismic facies analysis (evidence from tournaisian-famennian deposit of Ozernoe field)]. *Vestnik Permskogo natsional'nogo issledovatel'skogo politekhnicheskogo universiteta. Geologiya. Neftgazovoe i gornoe delo*, 2015, no. 14, pp. 6-12. DOI: 10.15593/2224-9923/2015.14.1
30. Galkin V.I., Ponomareva I.N., Cherepanov S.S. Razrabotka metodiki otsenki vozmozhnostei vydeleniia tipov kollektorov po dannym krivykh vosstanovleniia davleniia po geologo-promyslovym kharakteristikam plasta [Development of the methodology for evaluation of possibilities to determine reservoir types based on pressure build-up curves, geological and reservoir properties of the formation (case study of famen deposits of Ozernoe field)]. *Vestnik Permskogo natsional'nogo issledovatel'skogo politekhnicheskogo universiteta. Geologiya. Neftgazovoe i gornoe delo*, 2015, no. 17, pp. 32-40.
31. Cherepanov S.S., Martiushev D.A., Ponomareva I.N. Otsenka filtratsionno-emkostnykh svoistv treshchinovatykh karbonatnykh kollektorov mestorozhdenii Predural'skogo kraevogo progiba [Evaluation of filtration-capacitive properties of fractured carbonate reservoir of Predural'skogo edge deflection]. *Neftianoe khoziaistvo*, 2013, no. 3, pp. 62-65.
32. Galkin V.I., Kunitskikh A.A. Statisticheskoe modelirovanie rasshiriaushchegosia tamponazhnogo sostava [Statistical modelling of expanding cement slurry]. *Vestnik Permskogo natsional'nogo issledovatel'skogo politekhnicheskogo universiteta. Geologiya. Neftgazovoe i gornoe delo*, 2017, vol. 16, no. 3, pp. 215-244. DOI: 10.15593/2224-9923/2017.3.2
33. Galkin V.I., Ponomareva I.N., Repina V.A. Issledovanie protsessa nefteizvlecheniia v kollektorakh razlichnogo tipa pustotnosti s ispol'zovaniem mnogomernogo statisticheskogo analiza [Study of oil recovery from reservoirs of different void types with use of multidimensional statistical analysis]. *Vestnik Permskogo natsional'nogo issledovatel'skogo politekhnicheskogo universiteta. Geologiya. Neftgazovoe i gornoe delo*, 2016, no. 19, pp. 145-154. DOI: 10.15593/2224-9923/2016.19.5
34. Krivoshechekov S.N., Galkin V.I. Postroenie matritsy elementarnykh iacheek pri prognoze neftegazonosnosti veroiatnostno-statisticheskimi metodami na territorii Permskogo kraia [Construction of a matrix of elementary cells in predicting oil and gas content by probabilistic and statistical methods in the Perm Territory]. *Geologiya, geofizika i razrabotka nefitnykh i gazovykh mestorozhdenii*. Moscow: VNIIOENG, 2008, no. 8, pp. 20-23.
35. Ivanov S.A., Rastegaev A.V., Galkin V.I. Analiz rezul'tatov primeneniia GRP (na primere Povkhovskogo mestorozhdeniia nefiti) [Analysis of results of applying formation hydraulic fracturing in Povkhovsky oil field]. *Neftpromyslovoe delo*. Moscow: VNIIOENG, 2010, no. 7, pp. 54-58.
36. Krivoshechekov S.N., Galkin V.I., Volkova A.S. Razrabotka veroiatnostno-statisticheskoi metodiki prognoza neftegazonosnosti struktur [Development of a probabilistic-statistical method for predicting the oil and gas content of structures]. *Neftpromyslovoe delo*. Moscow: VNIIOENG, 2010, no. 7, pp. 28-31.
37. Houze O., Viturat D., Fjaere O.S. Dinamic data analysis. Paris: Kappa Engineering, 2008, 694 p.
38. Van Golf-Racht T.D. Fundamentals of fractured reservoir engineering. *Elsevier scientific publishing company*. Amsterdam - Oxford - New York, 1982, 709 p.
39. Horne R.N. Modern well test analysis: A computer Aided Approach. 2nd ed. Palo Alto: Petroway Inc, 2006, 257 p.
40. Johnson N.L., Leone F.C. Statistics and experimental design. New York - London - Sydney - Toronto, 1977, 606 p.
41. Montgomery D.C., Peck E.A. Introduction to liner regression analysis. New York: John Wiley & Sons, 1982, 504 p.
42. Darling T. Well Logging and Formation Evaluation. Gardners Books, 2010, 336 p.
43. Watson G.S. Statistic on spheres. New York: John Wiley and Sons, Inc., 1983, 238 p.
44. Yarus J.M. Stochastic modeling and geostatistics. *AAPG*. Tulsa, Oklahoma, 1994, 231 p.
45. Koshkin K.A., Melkisev O.A. Use of derivatives to assess preservation of hydrocarbon deposits. *IOP Conf. Series: Journal of Physics: Conference Series*, 2018, vol. 1015, 032092 p. DOI: 10.1088/1742-6596/1015/3/032092
46. Zhuoheng Ch., Osadetz K.G. Geological risk mapping and prospect evaluation using multivariate and Bayesian statistical methods, western Sverdrup Basin of Canada. *AAPG Bulletin*, 2006, vol. 90, no. 6, pp. 859-872. DOI: 10.1306/01160605050
47. Ahlbrandt T.S., Charpentier R.R., Klett T.R., Schmoker J.W., Schenk C.J., Ulmshiek G.F. Global resource estimates from total petroleum systems. *AAPG Memoir*, 2005, no. 86, pp. 1-334. DOI: 10.1306/M861061
48. Pang-Ning Tan, Michael Steinbach, Vipin Kumar. Introduction to data mining. Boston: Pearson Addison Wesley, 2005, 769 p.
49. Warren J.E., Root P.J. The behavior of naturally fractured reservoirs. *Soc. Petrol. Eng. J.*, 1963, vol. 3, iss. 3, pp. 245-255. DOI: 10.2118/426-PA
50. Tiab D. Modern core analysis. Vol. 1. Theory, core laboratories. Houston, Texas, 1993, 200 p.

Список литературы

1. Галкин В.И., Растегаев А.В., Галкин С.В. Вероятностно-статистическая оценка нефтегазонасности локальных структур. – Екатеринбург: УрО РАН, 2001. – 277 с.
2. Козлова И.А., Галкин В.И., Ванцева И.В. К оценке перспектив нефтегазонасности Соликамской депрессии с помощью геолого-геохимических характеристик нефтегазоматеринских пород // Нефтепромысловое дело. – 2010. – № 7. – С. 20–23.
3. Кривошеков С.Н., Галкин В.И., Носов М.А. Оценка не локализованных ресурсов нефти территории Пермского края при помощи системы элементарных участков // Нефтяное хозяйство. – 2014. – № 6. – С. 9–11.
4. Кривошеков С.Н., Козлова И.А., Санников И.В. Оценка перспектив нефтегазонасности западной части Соликамской депрессии на основе геохимических и геодинамических данных // Нефтяное хозяйство. – 2014. – № 6. – С. 12–15.
5. Геохимические показатели РОВ пород как критерии оценки перспектив нефтегазонасности / В.И. Галкин, И.А. Козлова, О.А. Мелкишев, М.А. Шадрин // Нефтепромысловое дело. – 2013. – № 9. – С. 28–31.
6. Козлова И.А., Мелкишев О.А. Прогнозная оценка распределения не локализованных ресурсов нефти в девонском терригенном комплексе на территории Пермского края // Геология, геофизика и разработка нефтяных и газовых месторождений. – 2017. – № 2. – С. 4–8.
7. Галкин В.И., Козлова И.А. Разработка вероятностно-статистических регионально-зональных моделей прогноза нефтегазонасности по данным геохимических исследований верхнедевонских карбонатных отложений // Геология, геофизика и разработка нефтяных и газовых месторождений. – 2016. – № 6. – С. 40–45.
8. Оценка перспектив нефтегазонасности юга Пермского края по органо-геохимическим данным / В.И. Галкин, И.А. Козлова, С.Н. Кривошеков, М.А. Носов, Н.С. Колтырина // Нефтепромысловое дело. – 2015. – № 7. – С. 32–35.
9. Решение региональных задач прогнозирования нефтеносности по данным геолого-геохимического анализа рассеянного органического вещества пород доломитового типа / В.И. Галкин, И.А. Козлова, С.Н. Кривошеков, М.А. Носов // Нефтяное хозяйство. – 2015. – № 1. – С. 21–23.
10. К обоснованию построения моделей зонального прогноза нефтегазонасности для нижне-среднедевизейского комплекса Пермского края / В.И. Галкин, И.А. Козлова, С.Н. Кривошеков, О.А. Мелкишев // Нефтяное хозяйство. – 2015. – № 8. – С. 32–35.
11. Галкин В.И., Жуков Ю.А., Шишкин М.А. Применение вероятностных моделей для локального прогноза нефтегазонасности. – Екатеринбург: УрО РАН, 1990. – 108 с.
12. Галкин В.И., Шахмутинов А.Н. О возможности прогноза нефтегазонасности юрских отложений вероятностно-статистическими методами (на примере территории деятельности ТПП «Голымынефтегаз») // Геология, геофизика и разработка нефтяных и газовых месторождений. – М.: Изд-во ВНИИОЭНГ, 2009. – № 6. – С. 11–14.

13. Галкин В.И., Шайхутдинов А.Н. Построение статистических моделей для прогноза дебитов нефти по верхнеюрским отложениям Когалымского региона // Нефтяное хозяйство. – 2010. – №1. – С. 52–54.
14. Галкин В.И., Кривошеков С.Н. Построение матрицы элементарных ячеек при прогнозе нефтегазоносности вероятностно-статистическими методами на территории Пермского края // Геология, геофизика и разработка нефтяных и газовых месторождений. – М.: Изд-во ВНИИОЭНГ, 2008. – № 8. – С. 20–23.
15. Галкин В.И., Кривошеков С.Н. Обоснование направлений поисков месторождений нефти и газа в Пермском крае // Научные исследования и инновации. – Пермь, 2009. – Т. 3, № 4. – С. 3–7.
16. Прогнозная оценка нефтегазоносности структур на территории Соликамской депрессии / В.И. Галкин, А.В. Растегаев, И.А. Козлова, И.В. Ванцева, С.Н. Кривошеков, В.Л. Воеводкин // Нефтепромысловое дело. – М.: Изд-во ВНИИОЭНГ, 2010. – № 7. – С. 4–7.
17. Додевонские отложения Пермского Прикамья как одно из перспективных направлений геолого-разведочных работ / Т.В. Белоконов, В.И. Галкин, И.А. Козлова, С.Е. Пашкова // Геология, геофизика и разработка нефтяных и газовых месторождений. – М.: Изд-во ВНИИОЭНГ, 2005. – № 9. – С. 24–28.
18. Путилов И.С. Разработка технологий комплексного изучения геологического строения и размещения месторождений нефти и газа. – Пермь: Изд-во Перм. нац. исслед. политехн. ун-та, 2014. – 285 с.
19. О возможности прогнозирования нефтегазоносности фаменских отложений с помощью построения вероятностно-статистических моделей / В.И. Галкин, И.А. Козлова, С.Н. Кривошеков, Е.В. Пятунина, С.Н. Пестова // Геология, геофизика и разработка нефтяных и газовых месторождений. – М.: Изд-во ВНИИОЭНГ, 2007. – № 10. – С. 22–27.
20. Галкин В.И., Соловьев С.И. Районирование территории Пермского края по степени перспективности приобретения нефтяных участков недр // Вестник Пермского национального исследовательского политехнического университета. Геология. Нефтегазовое и горное дело. – 2015. – № 16. – С. 14–24.
21. Соснин Н.Е. Разработка статистических моделей для прогноза нефтегазоносности (на примере терригенных девонских отложений Северо-Татарского свода) // Вестник Пермского национального исследовательского политехнического университета. Геология. Нефтегазовое и горное дело. – 2012. – № 5. – С. 16–25.
22. Галкин В.И., Соснин Н.Е. Разработка геолого-математических моделей для прогноза нефтегазоносности сложнопостроенных структур в девонских терригенных отложениях // Нефтяное хозяйство. – 2013. – № 4. – С. 28–31.
23. Дементьев Л.Ф. Математические методы и ЭВМ в нефтегазовой геологии. – М.: Недра, 1987. – 264 с.
24. Давыденко А.Ю. Вероятностно-статистические методы в геолого-геофизических приложениях. – Иркутск, 2007. – 29 с.
25. Михалевиц И.М. Применение математических методов при анализе геологической информации (с использованием компьютерных технологий). – Иркутск, 2006. – 115 с.
26. Андрейко С.С. Разработка математической модели метода прогнозирования газодинамических явлений по геологическим данным для условий Верхнекамского месторождения калийных солей // Вестник Пермского национального исследовательского политехнического университета. Геология. Нефтегазовое и горное дело. – 2016. – № 21. – С. 345–353. DOI: 10.15593/2224-9923/2016.21.6
27. Девис Дж. Статистика и анализ геологических данных. – М.: Мир, 1977. – 353 с.
28. Поморский Ю.Л. Методы статистического анализа экспериментальных данных: монография. – Л., 1960. – 174 с.
29. Черепанов С.С. Комплексное изучение трещиноватости карбонатных залежей методом Уоррена – Рута с использованием данных сейсмофациального анализа (на примере турнефамской залежи Озерного месторождения) // Вестник Пермского национального исследовательского политехнического университета. Геология. Нефтегазовое и горное дело. – 2015. – № 14. – С. 6–12. DOI: 10.15593/2224-9923/2015.14.1
30. Галкин В.И., Пономарева И.Н., Черепанов С.С. Разработка методики оценки возможностей выделения типов коллекторов по данным кривых восстановления давления по геолого-промысловым характеристикам пласта // Вестник Пермского национального исследовательского политехнического университета. Геология. Нефтегазовое и горное дело. – 2015. – № 17. – С. 32–40.
31. Черепанов С.С., Мартюшев Д.А., Пономарева И.Н. Оценка фильтрационно-емкостных свойств трещиноватых карбонатных коллекторов месторождений Предуральяского краевого прогиба // Нефтяное хозяйство. – 2013. – № 3. – С. 62–65.
32. Галкин В.И., Куницких В.И. Статистическое моделирование расширяющегося тампонажного состава // Вестник Пермского национального исследовательского политехнического университета. Геология. Нефтегазовое и горное дело. – 2017. – Т. 16, № 3. – С. 215–244. DOI: 10.15593/2224-9923/2017.3.2
33. Галкин В.И., Пономарева И.Н., Репина В.А. Исследование процесса нефтеизвлечения в коллекторах различного типа пустотности с использованием многомерного статистического анализа // Вестник Пермского национального исследовательского политехнического университета. Геология. Нефтегазовое и горное дело. – 2016. – № 19. – С. 145–154. DOI: 10.15593/2224-9923/2016.19.5
34. Кривошеков С.Н., Галкин В.И. Построение матрицы элементарных ячеек при прогнозе нефтегазоносности вероятностно-статистическими методами на территории Пермского края // Геология, геофизика и разработка нефтяных и газовых месторождений. – М.: Изд-во ВНИИОЭНГ, 2008. – № 8. – С. 20–23.
35. Иванов С.А., Растегаев А.В., Галкин В.И. Анализ результатов применения ГРП (на примере Повховского месторождения нефти) // Нефтепромысловое дело. – М.: Изд-во ВНИИОЭНГ, 2010. – № 7. – С. 54–58.
36. Кривошеков С.Н., Галкин В.И., Волкова А.С. Разработка вероятностно-статистической методики прогноза нефтегазоносности структур // Нефтепромысловое дело. – М.: Изд-во ВНИИОЭНГ, 2010. – № 7. – С. 28–31.
37. Houze O., Viturat D., Fjaere O.S. Dinamie data analysis. – Paris: Kappa Engineering, 2008. – 694 p.
38. Van Golf-Racht T.D. Fundamentals of fractured reservoir engineering // Elsevier scientific publishing company. – Amsterdam – Oxford – New York, 1982. – 709 p.
39. Horne R.N. Modern well test analysis: A computer Aided Approach. – 2nd ed. – Palo Alto: PetrowayInc, 2006. – 257 p.
40. Johnson N.L., Leone F.C. Statistics and experimental design. – New York – London – Sydney – Toronto, 1977. – 606 p.
41. Montgomery D.C., Peck E.A., Introduction to liner regression analysis. – New York: John Wiley & Sons, 1982. – 504 p.
42. Darling T. Well Logging and Formation Evalution. – Gardners Books, 2010. – 336 p.
43. Watson G.S. Statistic on spheres. – New York: John Wiley and Sons, Inc., 1983. – 238 p.
44. Yarus J.M. Stochastic modeling and geostatistics // AAPG. – Tulsa, Oklahoma, 1994. – 231 p.
45. Koshkin K.A., Melkisev O.A. Use of derivatives to assess preservation of hydrocarbon deposits // IOP Conf. Series: Journal of Physics: Conference Series. – 2018. – Vol. 1015. – P. 032092. DOI:10.1088/1742-6596/1015/3/032092
46. Zhuoheng Ch., Osadetz K.G. Geological risk mapping and prospect evaluation using multivariate and Bayesian statistical methods, western Sverdrup Basin of Canada // AAPG Bulletin. – 2006. – Vol. 90, № 6. – P. 859–872. DOI: 10.1306/01160605050
47. Global resource estimates from total petroleum systems / T.S. Ahlbrandt, R.R. Charpentier, T.R. Klett, J.W. Schmoker, C.J. Schenk, G.F. Ulmshiek // AAPG Memoir. – 2005. – № 86. – P. 1–334. DOI: 10.1306/M861061
48. Introduction to data mining / Pang-Ning Tan, Michael Steinbach, Vipin Kumar. – Boston: Pearson Addison Wesley, 2005. – 769 p.
49. Warren J.E., Root P.J. The behavior of naturally fractured reservoirs // Soc. Petrol. Eng. J. – 1963. – Vol3, is. 3. – P. 245–255. DOI: 10.2118/426-PA
50. Tiab D. Modern core analysis. Vol. 1. Theory, core laboratories. – Houston, Texas, 1993. – 200 p.