LOW ABUNDANCE OF THE WHISKERED BAT *MYOTIS MYSTACINUS* (KUHL, 1817) IN POLAND – CONSEQUENCE OF COMPETITION WITH *PIPISTRELLE* BATS?

KUREK, K.¹ – TOLKACZ, K.² – MYŚLAJEK, R. W.^{3*}

¹*Association for Nature „Wolf”
Twardorzeczka 229, 34-324 Lipowa, Poland*

²*University of Warsaw, Faculty of Biology, Institute of Zoology, Department of Parasitology
Miecznikowa 1, 02-096 Warszawa, Poland*

³*University of Warsaw, Faculty of Biology, Institute of Genetics and Biotechnology
Pawińskiego 5a, 02-106 Warszawa, Poland*

**Corresponding author
e-mail: robert.myslajek@gmail.com*

(Received 19th Mar 2017; accepted 5th Jul 2017)

Abstract. Interspecific competition is an important factor influencing the distribution of animals, including bats. In this study we tested if the presence of pipistrelles *Pipistrellus* sp. affects the distribution of the whiskered bat *Myotis mystacinus* using information about the species structure of bat communities obtained by capturing with mist nets in whole Poland. Abundance of the whiskered bat is low (on average 3.1%), and it only occurs frequently in the western part of the Carpathian Mountains (24.5%). The negative correlation between the abundance of the whiskered bat and pipistrelles (especially *P. nathusii* and *P. pipistrellus*) in bat communities across Poland seems to support the hypothesis that its occurrence is shaped by competitive interactions.

Keywords: bat distribution, species range, mountains, sympatric bats, species coexistence

Introduction

The whiskered bat *Myotis mystacinus* (Kuhl, 1817) has an extensive range in Europe that covers almost the entire continent (Dietz et al., 2009; Benda et al., 2016; Sachanowicz et al., 2016). However, there is no certainty about the eastern boundary of its occurrence due to the difficulty in distinguishing it from the steppe whiskered bat *Myotis aurascens* Kuzyakin, 1935 (Hutson et al., 2008). In Poland, the whiskered bat is widespread throughout the country (Sachanowicz et al., 2006). However its abundance seems to be unequal, with some areas lacking the species (Sachanowicz et al., 2006a) and other regions dominated by them (Piksa, 2011).

Presence of the whiskered bat can be affected by interspecific competition with pipistrelles *Pipistrellus* sp. (Baagøe, 1973). Such processes are key factors shaping the distribution patterns of organisms, as ecologically analogous species sharing a niche cannot occur in the same area (Gause, 1934). This results in the exclusion of one of the competing species (Hardin, 1960), which influences the structure of communities. Interspecific competition is an important factor influencing the distribution of mammals (Ylönen, 1990; Letcher et al., 1994; Eccard and Ylönen, 2003; Bonesi et al., 2006); however it has rarely been considered during the assessment of bat ranges (Arlettaz et al., 2000; Delaval et al., 2005; Santos et al., 2014; Zeus et al., 2017).

In this paper, we attempted to validate the predictions of Baagøe (1973) by comparing the distribution of the whiskered bat with that of pipistrelle species in

Poland. Our null hypothesis was that the whiskered bat is more abundant in areas where pipistrells are rare or absent.

Material and Methods

To assess the current occurrence of the whiskered bat in Poland we used information on the species structure of bat communities obtained by capturing with mist nets. We analysed both data published since 1997 and our unpublished data from studies conducted since 2000. Initially, we obtained data from 42 locations distributed across the whole of Poland (Table 1), and for further analysis we divided these locations into three groups: locations where (1) whiskered bats were not caught, (2) the total contribution of the species accounted for less than 10%, and (3) locations where the whiskered bat accounted for over 10% of all caught individuals. Statistical analyses were performed in R (R Development Core Team, 2009).

Results

The whiskered bat was the most abundant bat species found in southern Poland, and it was most common in the western part of the Carpathian Mountains (Fig. 1, Table 1) where its average share in bat communities was 24.5% ($SE \pm 7.0$). In the remaining areas it was rather rare (Welch t-test, $t = -3.0315$, $df = 5.059$, $p < 0.05$), with its average share in bat communities only 3.1% ($SE \pm 0.5$).

Figure 1. Percentage of whiskered bats in catches in Poland, 1997–2012 (see Table 1 for details)

Table 1. Percentage of whiskered bat (*MYS*), *Nathusius's* pipistrelle (*PIN*), common pipistrelle *sensu stricte* (*PIP s.s.*), soprano pipistrelle (*PPY*), not distinguished common pipistrelle and soprano pipistrelle (*PIP s.l.*), all pipistrelles in total (*PIPS*), and total number of captured bats (*N*) in various parts of Poland, 1997–2012.

Area ID	MYS	PIPS	PIN	PIP s.l.	PIP s.s.	PPY	N	source
1	0	13.0	2.0	11.0	8.0	3.0	99	Ciechanowski et al., 2006a
2	0	15.1	10.6	4.5	–	–	132	Ciechanowski, 2002
2	0	16.2	9.3	6.9	–	–	216	Ciechanowski, 2003
3	0	83.9	64.5	19.4	–	–	31	Jarzembowski et al., 1997
4	0	44.5	22.6	–	0.8	21.1	133	Ciechanowski et al., 2008
5	0	15.4	–	–	–	–	87	Authors unpublished data
6	0	3.7	3.7	0	0	0	27	Sachanowicz et al., 2001
7	0	30.3	20.1	2.2	0	8.0	139	Kmiecik et al., 2010
8	0	0	0	0	0	0	146	Postawa and Gas, 2003
9	0	5.0	4.2	0.8	0.8	0	120	Ciechanowski et al., 2006b
10	0	17.6	11.9	0.5	5.2	–	194	Ciechanowski et al., 2002
11	0	6.4	6.4	0	0	0	31	Ciechanowski and Duriasz, 2005
12	0	+	+	–	+	+	?	Cichocki and Łupicki, 2006
13	1.0	3.1	2.4	0.7	0.7	0	289	Łochyński and Grzywiński, 2009
14	0	7.0	6.0	1.0	–	–	84	Kowalski et al., 2001
15	0	13.8	8.5	5.3	–	–	452	Rachwald et al., 2001
16	0	7.4	7.4	0	0	0	27	Łochyński, 2001
17	4.2	5.2	3.1	–	1.0	1.0	96	Hejduk et al., 2008
18	1.7	?	–	–	–	–	231	Hejduk et al., 2004
19	1.5	0.2	–	–	–	–	396	Lesiński et al., 2007
20	1.5	0.3	0.3	0	0	0	400	Lesiński et al. 2006
21	0.8	1.6	0	1.6	–	–	125	Sachanowicz and Krasnodębski, 2003
22	6.8	?	?	?	–	–	44	Dudek et al., 2001
23	3.8	10.0	5	5.0	–	–	131	Ignaczak et al., 2001
24	3.0	?	–	–	–	–	79	Hejduk et al., 2001
25	6.4	16.1	6.4	9.7	–	–	62	Ignaczak, 2003
26	2.3	+	0	0	+	0	2711	Piksa et al., 2011
27	1.5	0	0	0	0	0	65	Piskorski and Urban, 2003
28	0	5.9	3.4	2.5	1.7	0.8	116	Piskorski, 2008
29	0	5.1	0	5.14	5.14	0	214	Piskorski et al., 2009
30	7.3	4.8	2.4	2.4	0	2.4	82	Piskorski, 2007
31	2.5	16.6	5.1	4.9	1.2	5.4	721	Mysłajek et al., 2002, 2005, 2014
32	18.7	3.3	3.3	0	0	0	91	Authors unpublished data
33	15.9	0	0	0	0	0	1795	Mysłajek et al., 2007 and authors unpublished data
34	21.6	0	0	0	0	0	306	Mysłajek et al., 2010
35	17.9	0.6	0.6	0	0	0	526	Mysłajek et al., 2004 and authors unpublished data
36	8.1	0	0	0	0	0	365	Mysłajek et al., 2013

37	13.8	0	0	0	0	0	290	Piksa and Gubała, 2011
38	59.3	0	0	0	0	0	3464	Piksa et al., 2011
39	0.8	14.6	0	14.6	—	—	130	Paszkiewicz et al., 1998
40	1.2	0.3	0	0.3	0	0.3	348	Bator et al., 2008
41	2.2	0	0	0	0	0	899	Węgiel et al., 2004
42	3.1	1.2	0	1.2	1.2	0	159	Sachanowicz and Wower, 2013

The abundance of the whiskered bat in Poland was negatively correlated with abundance of pipistrelle species, especially *P. nathusii* and *P. pipistrellus* (Table 2).

Table 2. Spearman's correlation coefficient (*Rho*) for the percentage of whiskered bats (*Myotis mystacinus*) in combination with pipistrelles (considered collectively by each species or group of species) in the cohorts of trapped bats in Poland. The table shows the value of the test statistic (*S*), statistically significant correlation (*p*-value), and sample size (*N*).

<i>Myotis mystacinus</i> in combination with:	Rho	S	p-value	N
<i>Pipistrellus</i> sp.	- 0.589	14519.28	p<0.001	38
<i>P. nathusii</i>	- 0.551	12055.46	p<0.001	36
<i>P. pipistrellus</i> s.l.	- 0.396	9135.34	p<0.05	34
<i>P. pipistrellus</i> s.s.	-0.408	4613.76	p<0.05	27
<i>P. pygmaeus</i>	-0.263	3694.32	ns	26

Discussion

The observed pattern of whiskered bat abundances in Poland could be explained by their selection for mountainous habitats or intraspecific competition. A factor that may limit the occurrence of the whiskered bat in Poland is the wintering ecology of the species. Whiskered bats hibernate in a fairly broad spectrum of caves; however, they prefer caves located between 1,000 and 1,930 m above sea level (Piksa et al., 2013). Thus, the main wintering grounds of this species can be found at higher altitudes on mountains, where in a study by Piksa et al. (2013), bats from *M. mystacinus* complex group (species are usually impossible to distinguish during winter counts, although the community is probably dominated by the whiskered bat) accounted for 88.5%. The whiskered bat is a species that does not migrate and only undertakes relatively short flights up to a distance of 165 km (Hutterer et al., 2005). Thus, limited abilities for long-distance migration may also limit the distribution and abundance of the species to around the most suitable hibernaculas.

Nethertheless, the presence of mountains alone does not seem to be the most important factor shaping their distribution. Reproducing females were recorded in the lowlands, and the species roosts mainly in anthropogenic shelters and utilizes feeding grounds located primarily in woodlands scattered across agricultural lands (Buckley et al., 2012; Kurek, 2014), which are common in the lowlands. Furthermore, the wide range of the whiskered bat in Europe, which reaches the northern edges of the continent (Hutson et al., 2008), shows that this is a not thermophilic species restricted to the southern and central, i.e. warmer, parts of the continent.

Antagonistic interactions between species most likely explain the observed pattern of whiskered bat distribution in Poland. Gerell (1987), by analysing the density of

distribution of Brandt's and whiskered bats in northern Europe, suggested that the distribution of these two species is the result of competitive interactions with the common pipistrelle *Pipistrellus pipistrellus* (Schreber, 1774). Our analyses confirmed the constraining effects of at least two pipistrelle species (Nathusius's and common pipistrelle) on whiskered bats. In areas where whiskered bats were captured in large numbers, pipistrelles were not observed at all or were scarce. In addition, pipistrelles were found in these regions beyond the reproductive period (Piksa et al., 2011; Kurek, 2014), which suggests that they were migrating individuals. In other areas, pipistrelle bats are a stable part of the bat fauna (Sachanowicz et al., 2006). Both the whiskered bat and pipistrelles show a preference for similar foraging areas and shelters of anthropogenic origin (Dietz et al., 2009; Buckley et al., 2012; Kurek, 2014), and competition for such resources could be a major factor for the push-out of a significant part of the whiskered bat population to more hostile mountainous areas.

We are aware that the analysis of bat communities based solely on the results of mist-netting could be biased due to a number of factors, including the size and placement of nets, and the frequency at which investigators check the nets (Kunz et al., 2009). Nonetheless, we believe that the potential bias was minimized by using data from numerous studies conducted across the whole country, which includes the repeated sampling of targeted regions, utilization of a variety of netting sites, and use of variable net-sets as suggested by Larsen et al. (2007). Another source of bias may be the use of data obtained during swarming. Places where this phenomenon occurs are called 'hot spots', and during mist-netting researchers capture bats that are not necessarily resident in that given area (Fenton, 1969; Kerth et al., 2003; Bogdanowicz et al., 2012). However, taking into account only catches made in potential feeding grounds, the share of whiskered bats in the southern regions of the country is still estimated to be over 10% (Table 1), which supports our findings that the western Carpathians are the main area in Poland where whiskered bats occur.

Acknowledgements. The study was financed by the statutory budget of the Association for Nature "Wolf". We thank Tomasz Diserens for linguistic advice.

REFERENCES

- [1] Arlettaz, R., Godat, S., Meyer H. (2000): Competition for food by expanding pipistrelle bat populations (*Pipistrellus pipistrellus*) might contribute to the decline of lesser horseshoe bats (*Rhinolophus hipposideros*). – Biological Conservation 93: 55–60.
- [2] Baagøe, H. J. (1973): Taxonomy of two sibling species of bats in Scandinavia *Myotis mystacinus* and *Myotis brandtii* (Chiroptera). Videnskabelige Meddelelser Dansk Naturhistorisk Forening 136: 191–216.
- [3] Bator, A., Szkudlarek, R., Węgiel, A. (2008): Skład gatunkowy i struktura populacji nietoperzy odławianych na terenie Ciężkowicko-Rożnowskiego Parku Krajobrazowego. – Nietoperze 9: 193–202.
- [4] Benda, P., Gazaryan, S., Vallo, P. (2016): On the distribution and taxonomy of bats of the *Myotis mystacinus* morphogroup from the Caucasus region (Chiroptera: Vespertilionidae). – Turkish Journal of Zoology 40: 842–863.
- [5] Bogdanowicz, W., Piksa, K., Tereba, A. (2012): Hybridization hotspots at bat swarming sites. – PLoS ONE 7(12): e53334.
- [6] Bonesi, L., Strachan, R., Macdonald, D. W. (2006): Why are there fewer signs of mink in England? Considering multiple hypotheses. – Biological Conservation 130: 268–277.

- [7] Buckley, D. J., Lundy, M. G., Boston, E. S. M., Scott, D. D., Gager, Y., Prodöhl, P., Marnell, F., Montgomery, W.I., Teeling, E. C. (2012): The spatial ecology of the whiskered bat (*Myotis mystacinus*) at the western extreme of its range provides evidence of regional adaptation. – *Mammalian Biology* 78: 198–204.
- [8] Cichocki, J., Łupicki, D. (2006): Nietoperze (Chiroptera) okolic Cybinki (Środkowe Nadodrze). – *Nietoperze* 7: 57–65.
- [9] Ciechanowski, M. (2002): Community structure and activity of bats (Chiroptera) over different water bodies. – *Mammalian Biology* 67: 276–285.
- [10] Ciechanowski, M. (2003): Chiropterofauna Puszczy Darżlubskiej. – *Nietoperze* 4: 45–59.
- [11] Ciechanowski, M., Duriasz, J. (2005): Nietoperze (Chiroptera) Parku Krajobrazowego Wzgórz Dylewskich. – *Nietoperze* 6: 25–36.
- [12] Ciechanowski, M., Koziróg L., Duriasz J., Przesmycka A., Świątkowska A., Kisicka I., Kasprzyk K. (2002): Bat fauna of the Iława Lakeland Landscape Park (northern Poland). – *Myotis* 40: 33–45.
- [13] Ciechanowski, M., Anikowska, U., Nalewaja, A., Przesmycka, A., Biała, A. (2006a): Nietoperze (Chiroptera) Parku Krajobrazowego „Dolina Słupi”. – *Nietoperze* 7: 19–37.
- [14] Ciechanowski, M., Czablewska, A., Mączyńska, M., Narczyński, T., Przesmycka, A., Zapart, A., Jarzembowski, T., Rachwald, A. (2008): Nietoperze (Chiroptera) Parku Krajobrazowego „Mierzeja Wiślana”. – *Nietoperze* 9: 203–224.
- [15] Ciechanowski, M., Przesmycka A., Sachanowicz, K. (2006b): Nietoperze (Chiroptera) Wdzydzkiego Parku Krajobrazowego. – *Parki Narodowe i Rezerваты Przyrody* 25 (4): 85–100.
- [16] Delaval, M., Henry, M., Charles-Dominique, P. (2005): Interspecific competition and niche partitioning: example of a neotropical rainforest bat community. – *Revue d'Ecologie (Terre et Vie)* 60: 149–166
- [17] Dietz, C., von Helversen, O., Nill, D. (2009): Bats of Britain, Europe & Northwest Africa. – A&C Black Publishers Ltd, London.
- [18] Dudek, I., Szkudlarek R., Cieślak, M. (2001): Nietoperze Parku Krajobrazowego „Chełmy” – obserwacje wstępne. – *Nietoperze* 2: 141–142.
- [19] Eccard, J. A., Ylönen, H. (2003): Interspecific competition in small rodents: from populations to individuals. – *Evolutionary Ecology* 17: 423–440.
- [20] Fenton, M.B. (1969): Summer activity of *Myotis lucifugus* (Chiroptera: Vespertilionidae) at hibernacula in Ontario and Quebec. – *Canadian Journal of Zoology* 47: 597–602.
- [21] Gause, G. F. (1934): The struggle for existence. – Dover Publications Inc., Mineola, New York (reprinted 2003).
- [22] Gerell, R. (1987): Distribution of *Myotis mystacinus* and *Myotis brandtii* (Chiroptera) in Sweden. – *Zeitschrift für Säugetierkunde* 52: 338–341.
- [23] Hardin, G. (1960): The competitive exclusion principle. – *Science* 131: 1292–1297.
- [24] Hejduk, J., Stopczyński, M., Pawenta, W., Róg, M., Domański, J. (2001): Chiropterofauna Przedborskiego Parku Krajobrazowego – wstępne wyniki badań. – XV Ogólnopolska konferencja Chiropterologiczna, 9-11 listopada, Gdańsk-Sobieszewo: 36.
- [25] Hejduk, J., Górecki, M. T., Zieleniak, A., Pawenta, W., Gajęcka, K., Ruta, S. (2004): Chiropterofauna Bolimowskiego Parku Krajobrazowego. – XVIII Ogólnopolska Konferencja Chiropterologiczna, 5-7 listopada, Toruń: 33.
- [26] Hejduk, J., Róg, M., Olczyk, M., Zieleniak, A., Górecki, M., Błaszczuk, T. (2008): Nietoperze obszaru Natura 2000 „Pradolina Warszawsko-Berlińska”. – *Biuletyn Faunistyczny Polski Środkowej* 14: 31–33.
- [27] Hutson, A. M., Spitzenberger, F., Aulagnier, S., Coroiu, I. (2008): *Myotis mystacinus*. – In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>. Downloaded on 10 November 2013.
- [28] Hutterer, R. T., Ivanova, C., Meyer-Cords, C., Rodrigues, L. (2005): Bat migrations in Europe: a review of literature and analysis banding data. – *Naturschutz und Biologische Vielfalt* 28: 1–172.

- [29] Ignaczak, M. (2003): Nietoperze rezerwatu „Bukowa Góra”. – Nietoperze 4: 101–102.
- [30] Ignaczak, M., Radzicki G., Domański, J. (2001): Nietoperze Parku Krajobrazowego Międzyrzecza Warty i Widawki. – Nietoperze 2: 125–134.
- [31] Jarzembowski, T., Ostrach-Kowalska, A., Rymarzak, G. (1997): Chiropterofauna Kaszubskiego Parku Krajobrazowego. – Przegląd Przyrodniczy 8 (3): 123–127.
- [32] Kerth, G., Kiefer, A., Trappmann, C., Weishaar, M. (2003): High gene diversity at swarming sites suggest hot spots for gene flow in the endangered Bechstein's bat. – Conservation Genetics 4: 491–499.
- [33] Kmiecik, A., Kmiecik P., Grzywiński, W. (2010): Chiropterofauna Wigierskiego Parku Narodowego. – Nietoperze 11: 11–29.
- [34] Kowalski, M., Ostrach-Kowalska, A., Krasnodębski, I., Sachanowicz, K., Ignaczak, M., Rusin, A. (2001): Nietoperze Parków Krajobrazowych Górznieńsko-Lidzbarskiego i Welskiego. – Nietoperze 2: 117–124.
- [35] Kunz, T. H., Hodgkison, R., Weise, C. D. (2009): Methods of capturing and handling bats. – In: Kunz, T. H., Parsons S. (eds.). Ecological and behavioral methods for the study of bats. The Johns Hopkins University Press, Baltimore: 3–35.
- [36] Kurek, K. (2014): Ecology and genetic variability of the whiskered bat (*Myotis mystacinus*) population in Western Carpathians. Unpublished PhD Thesis. University of Warsaw, Faculty of Biology. (in Polish with English abstract)
- [37] Lesiński, G., Gulatowska, J., Kowalski, M., Fuszara, E., Fuszara, M., Wojtowicz, B. (2006): Nietoperze Wysoczyzny Płońskiej. – Nietoperze 7: 39–55.
- [38] Lesiński, G., Kowalski, M., Wojtowicz, B., Gulatowska, J., Lisowska, A. (2007): Bats on forest islands of different size in an agricultural landscape. – Folia zoologica 56: 153–161.
- [39] Letcher, A. J., Purvis, A., Nee, S., Harvey, P. H. (1994): Patterns of overlap in the geographic ranges of Palearctic and British mammals. – Journal of Animal Ecology 63: 871–879.
- [40] Łochyński, M. (2001): Chiropterofauna Żerkowsko-Czeszewskiego Parku Krajobrazowego. – Biuletyn Parku Krajobrazowego Wielkopolski 7 (9): 139–142.
- [41] Łochyński, M., Grzywiński W. (2009): Nietoperze Parku Krajobrazowego Puszcza Zielonka. – Nietoperze 10: 23–35.
- [42] Mysłajek, R. W., Henel, K., Nowak, S. (2002): Nietoperze Chiroptera rezerwatu „Łęczok” koło Raciborza. – Nietoperze 3: 271–276.
- [43] Mysłajek, R. W., Nowak, S., Kurek, K. 2004. Fauna nietoperzy Kotliny Żywieckiej. – Chrońmy Przyrodę Ojczystą 3: 78–85.
- [44] Mysłajek, R. W., Henel, K., Kurek, K., Urban, R., Nowak, S. (2005): Fauna nietoperzy Parku Krajobrazowego „Cysterskie Kompozycje Krajobrazowe Rud Wielkich”. – Scripta Rudensia 14: 5–14.
- [45] Mysłajek, R. W., Kurek, K., Szura, C., Nowak, S., Orysiak, P. (2007): Bats (Chiroptera) of the Silesian Beskid Mountains. – Fragmenta Faunistica 50 (1): 77–85.
- [46] Mysłajek, R. W., Kurek, K., Nowak, S., Orysiak, P. (2010): Nietoperze Chiroptera Żywieckiego Parku Krajobrazowego. – Nietoperze 11: 31–43.
- [47] Mysłajek, R. W., Kurek, K., Piksa, K., Szura, K., Nowak, S. (2013): Czy specjalny obszar ochrony siedlisk Natura 2000 Beskid Mały (PLH240023) w wystarczający sposób chroni nietoperze? Chrońmy Przyrodę Ojczystą 69 (3): 226–233.
- [48] Mysłajek, R. W., Kurek, K., Nowak, S., Henel, K. (2014): Stan poznania fauny nietoperzy specjalnego obszaru ochrony siedlisk „Stawy Łęczok” PLH240010. – Nietoperze 13: 3–9.
- [49] Paszkiewicz, R., Szkudlarek, R., Węgiel, A., Węgiel, J., Węgiel, W. (1998): Materiały do chiropterofauny Pienin – letnie stanowiska nietoperzy. – Pieniny – Przyroda i Człowiek 6: 31–46.

- [50] Piksa, K., Gubała, W. (2011): Fauna nietoperzy (Chiroptera) rojących się przy otworze jaskini Dymiąca Piwnica (Babiogórski Park Narodowy). – *Chrońmy Przyrodę Ojczystą* 67 (2): 128–136.
- [51] Piksa, K., Bogdanowicz, W., Tereba, A. (2011): Swarming of bats at different elevations in the Carpathian Mountains. – *Acta Chiropterologica* 13: 113–122.
- [52] Piksa, K., Nowak, J., Żmihorski, M., Bogdanowicz, W. (2013): Nonlinear distribution pattern of hibernating bats in caves along an elevational gradient in mountain (Carpathians, Southern Poland). – *PLoS ONE* 8(7): e68066.
- [53] Piskorski, M. (2007): Fauna nietoperzy Parku Krajobrazowego Lasy Janowskie. – *Nietoperze* 7: 3–11.
- [54] Piskorski, M. (2008): Fauna nietoperzy Lasów Sobiborskich. – *Nietoperze* 9: 3–17.
- [55] Piskorski, M., Urban, M. (2003): Nietoperze Południoworostoczańskiego Parku Krajobrazowego. – *Nietoperze* 4: 21–25.
- [56] Piskorski, M., Gwardjan, M., Kowalski, M., Wojtowicz, B., Urban, M., Bochen, R. (2009): Fauna nietoperzy Parku Krajobrazowego Lasy Strzeleckie. – *Nietoperze* 10: 15–22.
- [57] Postawa, T., Gas, A. (2003): Fauna nietoperzy Wigierskiego Parku Narodowego (Północno-wschodnia Polska). – *Studia Chiropterologica* 3–4: 31–42.
- [58] R Development Core Team. (2009): R: A language and environment for statistical computing. – R Foundation for Statistical Computing, Vienna, Austria.
- [59] Rachwald, A., Boratyński P., Nowakowski, W. K. 2001. Species composition and activity of bats flying over rivers in the Białowieża Primeval Forest. – *Acta Theriologica* 46: 235–242.
- [60] Sachanowicz, K., Krasnodębski, I. (2003): Skład gatunkowy i antropogeniczne kryjówki nietoperzy w Lasach Łukowskich. – *Nietoperze* 4: 27–38.
- [61] Sachanowicz, K., Wower, A. (2013): Assemblage structure and use of anthropogenic roosts by bats in the Eastern Carpathians: Case study in the Bieszczady National Park (SE Poland). – *Italian Journal of Zoology* 80: 139–148.
- [62] Sachanowicz, K., Marzec, M., Ciechanowski, M., Rachwald, A. (2001): Nietoperze Puszczy Rominckiej. – *Nietoperze* 2: 109–115.
- [63] Sachanowicz, K., Ciechanowski, M., Piksa, K. (2006): Distribution patterns, species richness and status of bats in Poland. – *Vespertilio* 9–10: 151–173.
- [64] Sachanowicz, K., Ciechanowski, M., Rachwald, A., Piskorski, M. (2016): Overview of bat species reported in Albania with the first country records for eight species. – *Journal of Natural History* 50: 513–521.
- [65] Santos, H., Juste, J., Ibanez, C., Palmeirim, J.M., Godinho, R., Amorim, F., Alves, P., Costa, H., de Paz, O., Pérez-Suarez, G., Martínez-Alos, S., Jones, G., Rebelo, H. (2014): Influences of ecology and biogeography on shaping the distributions of cryptic species: three bat tales in Iberia. – *Biological Journal of the Linnean Society* 112: 150–162.
- [66] Węgiel, A., Szkudlarek, R., Gottfried, T. (2004): Skład gatunkowy, aktywność i struktura populacji nietoperzy odławianych latem przy otworach jaskiń w Beskidach. – *Nietoperze* 5: 95–105.
- [67] Ylönen, H. (1990): Spatial avoidance between the bank vole *Clethrionomys glareolus* and the harvest mouse *Micromys minutus*: an experimental study. – *Annales Zoologici Fennici* 27: 313–320.
- [68] Zeus, V.M., Puechmaille, S.J., Kerth, G. (2017): Conspecific and heterospecific social groups affect each other's resource use: a study on roost sharing among bat colonies. – *Animal Behaviour* 123: 329–338.