Case Report

*Corresponding author

Giovanni Cacciamani, MD

Department of Urology

University of Verona

Polo Chirurgico "Confortini" - 5th Floor

Piazzale S. Stefani 1 - 37100

Verona, Italy

Tel. +39 340/2491750

Fax. +39 045/8127715

E-mail: giovanni.cacciamani@gmail.com

Volume 2 : Issue 2

Article Ref. #: 1000EMOJ2127

Article History

Received: July 15th, 2016

Accepted: July 29th, 2016

Published: August 2nd, 2016

Citation

Cacciamani G, Siracusano S, Cerruto MA, et al. Acute flank pain, flank mass and hypovolemic shock due to kidney cysts rupture: An emblematic case of wunderlich syndrome. *Emerg Med Open J.* 2016; 2(2): 36-38. doi: [10.17140/EMOJ-2-127](http://dx.doi.org/10.17140/EMOJ-2-127)

Copyright

©2016 Cacciamani G. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Acute Flank Pain, Flank Mass and Hypovolemic Shock due to Kidney Cysts Rupture: An Emblematic Case of Wunderlich Syndrome

Giovanni Cacciamani, MD*; Salvatore Siracusano, MD; Maria Angela Cerruto, MD; Silvia Bassi, MD; Alessandro Tafuri, MD; Vincenzo de Marco, MD; Antonio Benito Porcaro, MD; Walter Artibani, MD

Department of Urology, University of Verona, Verona, Italy

ABSTRACT

Spontaneous perinephric hemorrhage (SPH) due to kidney cysts rupture is uncommon complication but when it occurs can cause an emergency situation that may endanger the patient's life. Wunderlich first described the condition of spontaneous, non-traumatic renal bleeding confined to the subcapsular and perirenal spaces. Herein, we report a case of Wunderlich syndrome (WS) due to spontaneous rupture of a simple kidney cyst. In our case, the spontaneous retroperitoneal hemorrhage (SRH) was associated to a probable giant haemorrhagic cyst that is uncommon since cyst rupture occurs into caliceal system rather than the perinephric space. Wunderlich syndrome after spontaneous, non-traumatic kidney cyst rupture is a rare entity. A multimodality approach is necessary to have a definitive diagnosis in order to choose the best treatment possible for the patient.

KEYWORDS: Lenk's triad; Wunderlich syndrome; Renal cysts; Retroperitoneal hematoma; Spontaneous Retroperitoneal Hematoma (SRH).

INTRODUCTION

Kidney cysts (KC) include both benign and malignant etiology and can occur spontaneously or be caused by acquired cystic kidney disease, such as polycystic kidney disease (PKD). KC are masses that are predominantly composed of spaces filled with clear or straw-colored fluid.¹ Spontaneous perinephric hemorrhage due to KC rupture is uncommon complication but when it occurs can cause an emergency situation that may endanger the patient's life.

In 1856 Carl Wunderlich first described the condition of spontaneous, non-traumatic renal bleeding confined to the subcapsular and perirenal spaces² that may present with a classic triad of symptoms known as "Lenk's triad": acute flank or abdominal pain, flank mass and hypovolemic shock. Herein, we report a case of Wunderlich syndrome (WS) due to spontaneous rupture of a simple KC.

CASE REPORT

A 75-year-old man presented to the emergency department with left flank pain that lasted for 5-6 hours. His medical history included hypertension, mitral regurgitation, aspirin therapy, implantation of cardiac pacemaker for an atrioventricular block, infrarenal aortic aneurysm repair, chronic renal failure (serum creatinine 6 mg/L, creatinine clearance sec., Cockcroft-Gault 10 ml/min). He had no history of previous abdominal trauma. Physical examination indicated moderate tenderness across the whole abdomen, which was more evident on the left side. He had a severe hypotension, pulse rate of 92 beats/min, respiratory rate of 24 breaths/min and body temperature of 36.8 °C.

Figures 1A and 1B: Hemoperitoneum and a retroperitoneal hematoma on the left side displacing the spleen, the pancreatic tail and intestinal structures, extended to the pelvic cavity.

A complete blood count revealed a hemoglobin (Hgb) of 6 g/dL and a severe thrombocytopenia.

A computed tomography (CT) scan was performed. It showed a retroperitoneal hematoma on left side and left hemoperitoneum extended to the pelvic cavity (Figure 1A and 1B).

The patient underwent an angiography scan showing a hemorrhagic spreading from the left kidney. Due to the rapid worsening of clinical conditions the patient underwent an explorative laparotomy. A large retroperitoneal hematoma extending from the renal region to the pelvis was found and a left nephrectomy was done.

During this surgical manoeuvre an accidental injury of the tail of the pancreas occurred with a modest parenchymal laceration that was immediately sutured. Five hours later, a second explorative laparotomy due to persistent abdominal bleeding and hemodynamic instability was done.

Twenty-four hours later a CT scan angiography showed a modest hemorrhagic spreading from a minor branch of the superior mesenteric artery that was unsuccessfully embolized. Nevertheless, the patient showed a hemodynamic stability which was maintained as such in the following days. A haemodialysis therapy started in the subsequent days.

Histopathological examination described an atrophic kidney end-stage-type with cortico—medullary atrophy, multiple giant cystic lesions without specific coating.

DISCUSSION

The WS is a potentially life threatening condition. In a meta-analysis on spontaneous perirenal hemorrhage, the most common etiology are benign or malignant neoplasms (61%) with angiomyolipoma being predominant (29.1%) followed closely by renal cell carcinoma (26.1%) and vascular disease (17%) with polyarteritis nodosa occurring most frequently (12%).³ This emergency condition may occur with acute flank pain, flank mass and hypovolemic shock in 83%, 19% and 11% respectively.³

In our case, the spontaneous retroperitoneal hemorrhage was associated to a probable giant haemorrhagic cyst that is uncommon since cyst rupture occurs into caliceal system rather than the perinephric space. In this context large KC, hemorrhage and infection and a coexisting urinary tract obstruction are predisposing factors for spontaneous cyst rupture.⁴

In literature sparse cases of WS due to the non-traumatic rupture of a simple KC were reported. In particular Puerta et al⁵ described a case of perirenal hematoma due to rupture of a KC in a 50 years old female occurred with pain, a mass in left renal fossa and hypovolemia, who needed surgical drainage of the hematoma without nephrectomy. Finally, Lin et al⁶ described another case of spontaneous perinephric hemorrhage due a hemorrhagic renal cyst in 46-years-old women that was conservatively treated.

In our case, the presence of hemoperitoneum not allowed a conservative treatment and an explorative laparotomy was mandatory.

CONCLUSION

Wunderlich syndrome after spontaneous, non-traumatic kidney cyst rupture is a rare entity. A multimodality approach is necessary to have a definitive diagnosis in order to choose the best treatment possible for the patient.

FUNDING

The authors received no financial support for the research, authorship, and/or publication of this article.

CONFLICTS OF INTEREST

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CONSENT

The patient has provided written permission for the publication of this case details.

REFERENCES

1. Hindman NM. Cystic renal masses. *Abdom Radiol (NY)*. 2016; 41: 1020-1034. doi: [10.1007/s00261-016-0761-4](https://doi.org/10.1007/s00261-016-0761-4)
2. McDougal WS, Kursh ED, Persky L. Spontaneous rupture of the kidney with perirenal hematoma. *J Urol*. 1975; 114(2): 181-184. Web site. <http://www.ncbi.nlm.nih.gov/pubmed/1159905>. Accessed July 14, 2016
3. Zhang JQ, Fielding JR, Zou KH. Etiology of spontaneous perirenal hemorrhage: A meta-analysis. *J Urol*. 2002; 167(4): 1593-1596. Web site. <http://www.ncbi.nlm.nih.gov/pubmed/11912370>. Accessed July 14, 2016
4. Katabathina VS, Katre R, Prasad SR, Surabhi VR, Shanbhogue AK, Sunnapwar A. Wunderlich syndrome: Cross-sectional imaging review. *J Comput Assist Tomogr*. 2011; 35(4): 425-433. doi: [10.1097/RCT.0b013e3182203c5e](https://doi.org/10.1097/RCT.0b013e3182203c5e)
5. Valero PJ, Medina PM, Valpuestá FI, Sánchez GM. A simple renal cyst, the origin of Wunderlich's syndrome. *Archivos Españoles de Urología*. 1999; 52(7): 794-797. Web site. <http://www.ncbi.nlm.nih.gov/pubmed/10540773>. Accessed July 14, 2016
6. Lin YY, Chen JD, How CK, Yen DHT. Spontaneous perinephric hemorrhage from a hemorrhagic renal cyst. *Internal Medicine*. 2010; 49(19): 2189-2190. doi: [10.2169/internalmedicine.49.4021](https://doi.org/10.2169/internalmedicine.49.4021)