Editorial

*Corresponding author

Alessandro Arrigo, MD

Department of Biomedical Sciences and
Morphological and Functional Imaging
University of Messina
Via Consolare Valeria
1 Messina, 98125, Italy
Tel. +39 0902212939
E-mail: alessandro.arrigo@hotmail.com

Volume 1 : Issue 1

Article Ref. #: 1000ROJ1e001

Article History

Received: November 5th, 2015

Accepted: November 6th, 2015

Published: November 9th, 2015

Citation

Arrigo A, Mormina E, Calamuneri A.
The pre-surgical planning of brain
neoplasms: from diffusion tensor im-
aging to more advanced approaches.
Radiol Open J. 2015; 1(1): e1-e5.
doi: [10.17140/ROJ-1-e001](https://doi.org/10.17140/ROJ-1-e001)

Copyright

©2015 Arrigo A. This is an open
access article distributed under the
Creative Commons Attribution 4.0
International License (CC BY 4.0),
which permits unrestricted use,
distribution, and reproduction in
any medium, provided the original
work is properly cited.

The Pre-Surgical Planning of Brain Neoplasms: From Diffusion Tensor Imaging to More Advanced Approaches

Alessandro Arrigo*, Enricomaria Mormina and Alessandro Calamuneri

Department of Biomedical Sciences and Morphological and Functional Images, University of Messina, Messina, Italy

KEYWORDS: Brain neoplasm; Tractography; Pre-surgical planning; DTI; CSD.

The pre-surgical planning of brain neoplasms is strongly contributing to change the prognosis of neoplastic patients. Indeed, it supplies more and more detailed and reliable functional and morphological information before as well as during surgery. Both invasive and non-invasive approaches are available to achieve this goal. A powerful technique for the pre-surgical planning of brain neoplasms is Diffusion Weighted Imaging (DWI) based tractography. Differently from other approaches, tractography is able to provide, non-invasively, morphological information regarding brain pathways relationship with the neoplasm, by analyzing water diffusion within white matter. This is important especially for eloquent bundles, such as Cortico-Spinal Tract (CST) and Arcuate Fasciculus (AF), whose damages have a bad impact on the patient's Quality of Life (QoL). Tractography can be performed through several diffusion signal modeling techniques, among which Diffusion Tensor Imaging (DTI) is the most known. DTI has been widely used for neurosurgery both in pre-operative and intra-operative contexts, also in combination with other functional approaches such as functional MRI (fMRI) and cortical stimulation.¹⁻⁸ This useful technique was able to reduce post-surgical deficits as well as to improve the survival of neoplastic patients, through a better delineation of maximal safe resection.⁹ Moreover, its use provided great benefits in patients with high-grade gliomas in terms of risk of death.⁹ Several studies have demonstrated that DTI suffers from many limitations regarding its reliability to correctly model diffusion signal in different conditions.^{10,11} Furthermore, tensorial models are not able to resolve different fibers geometries (i.e. crossing fibers) within the same voxel; these complex configurations have been demonstrated to characterize more than 90% of white matter voxels,¹² thus making DTI an unreliable diffusion modeling technique. In order to overcome these limitations, several other approaches were developed; in particular, High Angular Resolution Diffusion-weighted Imaging (HARDI),¹³ Q-Ball Imaging (QBI)¹⁴ and Diffusion Spectrum Imaging (DSI)¹⁵ were found to be promising techniques for resolving voxels with multiple fibers orientations. Nevertheless, tractography was further improved by an HARDI modified approach called Constrained Spherical Deconvolution (CSD); this technique does not require very long acquisition time with respect to DSI¹⁶ and it is able to improve angular resolution if compared to QBI.¹⁷ CSD-based tractography was widely used in physiological contexts as well as in pathological ones, showing high sensitivity for the detection of white matter pathways.¹⁸⁻²² Although tensorial approaches were proved to be inadequate for reliably reconstructing brain pathways²³ and histological validation of CSD-based tractography has been recently provided,²⁴ more advanced diffusion techniques are still considered not usable in clinical settings due to their too high technical requirements.²⁵ For these reasons, DTI remains, to date, the most used technique for investigating white matter bundles, also for the pre-surgical planning of brain neoplasms.²⁶⁻²⁸ In reality, as recently highlighted by Mormina and colleagues,²⁹ CSD-based tractography is feasible in clinical settings and it is able to provide very useful information during the pre-surgical evaluation of eloquent bundles in patients with high-grade gliomas. It was shown that DTI-based reconstructions can be affected also by neoplasm's effects on white matter bundles. Indeed, brain pathways might result dislocated, disrupted and/or infiltrated by neoplasm,³⁰ with simultaneous Fractional Anisotropy (FA) dec-

rements; tensorial models are not able to distinguish among these different conditions, thus causing partial reconstructions or null detection of bundles involved. Moreover, peritumoral edema surrounding high-grade gliomas is able to reduce FA values,³¹ causing, also in this case, misleading fiber tracking. Mormina, et al.²⁹ showed how CSD-based tractography is able to clearly define fiber bundles involved by high-grade gliomas. Comparative qualitative analysis between CSD-based tractography and DTI-based one in patients with high-grade gliomas is shown in Figures 1 and 2.

Figure 1: Comparative qualitative analysis of bilateral CST reconstructions in a patient with right high-grade glioma (brown volume). DTI-based tractography (A) shows a good detection of the medial CST portion in the healthy side, whereas it is less represented in the affected side. Lateral CST portions are bilaterally missing. CSD-based tractography (B) shows robust detection of the entire CST in the healthy side. The medial CST portion results poorly involved in the affected side; moreover, the lateral CST portion is still detected, although it seems to be more involved by neoplasm. While the lack of DTI detection of lateral CST might be due to DTI intrinsic limitations, different representations of medial CST in the affected side might be caused by peritumoral edema affecting DTI-based reconstruction.

Figure 2: Comparative qualitative analysis of bilateral AF reconstructions in a patient with left high-grade glioma (blue volume). DTI-based tractography (A) shows a good AF detection in the healthy side, although several portions are poorly represented, such as the anterior one. AF seems to be widely damaged by neoplasmin the affected side. CSD-based tractography (B) shows a better detection of all AF portions in the healthy side, if compared to DTI-based one. Moreover, it shows a good AF representation in the affected side; this suggests that AF is involved by neoplasm although it is not strongly damaged. Also in this case, both DTI limitations and peritumoral edema might cause differences regarding AF representations in the affected side, if compared to CSD-based one.

Mormina and colleagues²⁹ showed also how this technique allowed a reliable analysis of diffusion-based parameters of reconstructed pathways in neoplastic patients. The quantitative evaluation of white matter streamlines based on diffusion tensor parameters, such as FA and Mean Diffusivity (MD), is considered another important measurement for detecting structural alterations of white matter caused by a loss of axonal integrity. Since DTI-based tractography does not consider voxels with low FA values, the quantitative analysis might provide the erroneous information of not significant white matter alterations; the latter, together with a partial qualitative analysis of affected pathways, might thus suggest false safe resection margins around the neoplasm. Moreover, it is worthy to note that a pre-surgical planning performed by means of DTI-based tractography cannot provide any qualitative and quantitative information regarding well-known undetected white matter bundles, such as the lateral portion of CST as well as the anterior portion of AF.^{29,32} All these limitations might be avoided by using CSD-based approach. Its sensitivity for reconstructing white matter bundles has been also recently highlighted by a comparative analysis of different tractographic approaches; in detail, CSD-based tractography was able to better detect brain pathways involved by neoplasm if compared to DTI-based one.³³

This brief description showed how the use of more advanced DWI-based approaches can provide useful information before surgical treatment of patients with brain neoplasms. It was demonstrated that these techniques are able to improve effectiveness of surgical treatments as well as their outcome by allowing a better delineation of resection margins. Moreover, they may have a good impact on the quality of life of neoplastic patients by helping to preserve eloquent fiber bundles and consequently by reducing post-surgical deficits. Several authors demonstrated that advanced diffusion signal modeling algorithms, such as CSD one, can be compatible with clinical settings and they should be more largely used for pre-surgical planning of brain neoplasms.

Interesting future perspectives for brain neoplasms treatment will regard the combined use of CSD-based tractography with functional approaches, such as neuronavigated Transcranial Magnetic Stimulation (nTMS), in order to further improve detection of eloquent bundles. Indeed, a recent study has adopted nTMS for subject-specific localization of the motor area and then DTI-based tractography for CST tracking; this approach allowed a good detection of CST, although tractographic reconstructions were found affected by above described DTI limitations, such as the peritumoral edema.³⁴ The use of CSD-based tractography in a similar study protocol might provide a better subject-specific detection of CST. Finally it will be important to assess potential improvements and advantages for neurosurgeons provided by the integration of these techniques into intraoperative navigation systems.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

CONSENT

Patients has provided written consent for submission of this manuscript for publication.

REFERENCES

1. Berman JI, Berger MS, Mukherjee P, Henry RG. Diffusion-tensor imaging-guided tracking of fibers of the pyramidal tract combined with intraoperative cortical stimulation mapping in patients with gliomas. *J Neurosurg.* 2004; 101(1): 66-72.
2. Coenen VA, Krings T, Mayfrank L, et al. Three-dimensional visualization of the pyramidal tract in a neuronavigation system during brain tumor surgery: first experiences and technical note. *Neurosurgery.* 2001; 49(1): 86-92.
3. Henry RG, Berman JI, Nagarajan SS, Mukherjee P, Berger MS. Subcortical pathways serving cortical language sites: initial experience with diffusion tensor imaging fiber tracking combined with intraoperative language mapping. *Neuroimage.* 2004; 21(2): 616-622. doi: [10.1016/j.neuroimage.2003.09.047](https://doi.org/10.1016/j.neuroimage.2003.09.047)
4. Kamada K, Todo T, Masutani Y, et al. Combined use of tractography-integrated functional neuronavigation and direct fiber stimulation. *J Neurosurg.* 2005; 102(4): 664-672.
5. Nimsky C, Ganslandt O, Hastreiter P, et al. Preoperative and intraoperative diffusion tensor imaging-based fiber tracking in glioma surgery. *Neurosurgery.* 2005; 56(1): 130-137; discussion 138.
6. Nimsky C, Ganslandt O, Merhof D, Sorensen AG, Fahlbusch R. Intraoperative visualization of the pyramidal tract by diffusion-tensor-imaging-based fiber tracking. *Neuroimage.* 2006; 30(4): 1219-1229. doi: [10.1016/j.neuroimage.2005.11.001](https://doi.org/10.1016/j.neuroimage.2005.11.001)
7. Bello L, Castellano A, Fava E, et al. Intraoperative use of diffusion tensor imaging fiber tractography and subcortical mapping for resection of gliomas: technical considerations. *Neurosurg Focus.* 2010; 28(2): E6. doi: [10.3171/2009.12.FOCUS09240](https://doi.org/10.3171/2009.12.FOCUS09240)
8. González-Darder JM, González-López P, Talamantes F, et al. Multimodal navigation in the functional microsurgical resection of intrinsic brain tumors located in eloquent motor areas: role of tractography. *Neurosurg Focus.* 2010; 28(2): E5. doi: [10.3171/2009.11.FOCUS09234](https://doi.org/10.3171/2009.11.FOCUS09234)
9. Wu JS, Zhou LF, Tang WJ, et al. Clinical evaluation and follow-up outcome of diffusion tensor imaging-based functional neuro-navigation: a prospective, controlled study in patients with gliomas involving pyramidal tracts. *Neurosurgery.* 2007; 61(5): 935-948.
10. Jones DK, Cercignani M. Twenty-five pitfalls in the analysis of diffusion MRI data. *NMR Biomed.* 2010; 23(7): 803-820. doi: [10.1002/nbm.1543](https://doi.org/10.1002/nbm.1543)

11. Farquharson S, Tournier JD, Calamante F, et al. White matter fiber tractography: why we need to move beyond DTI. *J Neurosurg*. 2013; 118(6): 1367-1377. doi: [10.3171/2013.2.JNS121294](https://doi.org/10.3171/2013.2.JNS121294)
12. Jeurissen B, Leemans A, Tournier JD, Jones DK, Sijbers J. Investigating the prevalence of complex fiber configurations in white matter tissue with diffusion magnetic resonance imaging. *Hum Brain Mapp*. 2013; 34(11): 2747-2766. doi: [10.1002/hbm.22099](https://doi.org/10.1002/hbm.22099)
13. Tuch DS, Reese TG, Wiegell MR, Makris N, Belliveau JW, Wedeen VJ. High angular resolution diffusion imaging reveals intra-voxel white matter fiber heterogeneity. *Magn Reson Med*. 2002; 48(4): 577-582. doi: [10.1002/mrm.10268](https://doi.org/10.1002/mrm.10268)
14. Tuch DS. Q-ball imaging. *Magn Reson Med*. 2004; 52(6): 1358-1372. doi: [10.1002/mrm.20279](https://doi.org/10.1002/mrm.20279)
15. Wedeen VJ, Wang RP, Schmahmann JD, et al. Diffusion spectrum magnetic resonance imaging (DSI) tractography of crossing fibers. *Neuroimage*. 2008; 41(4): 1267-1277. doi: [10.1016/j.neuroimage.2008.03.036](https://doi.org/10.1016/j.neuroimage.2008.03.036)
16. Tournier JD, Calamante F, Connelly A. Robust determination of the fibre orientation distribution in diffusion MRI: non-negativity constrained super-resolved spherical deconvolution. *Neuroimage*. 2007; 35(4): 1459-1472. doi: [10.1016/j.neuroimage.2007.02.016](https://doi.org/10.1016/j.neuroimage.2007.02.016)
17. Tournier JD, Yeh CH, Calamante F, Cho KH, Connelly A, Lin CP. Resolving crossing fibres using constrained spherical deconvolution: validation using diffusion-weighted imaging phantom data. *Neuroimage*. 2008; 42(2): 617-625. doi: [10.1016/j.neuroimage.2008.05.002](https://doi.org/10.1016/j.neuroimage.2008.05.002)
18. Arrigo A, Mormina E, Anastasi GP, et al. Constrained spherical deconvolution analysis of the limbic network in human, with emphasis on a direct cerebello-limbic pathway. *Front Hum Neurosci*. 2014; 8: 987. doi: [10.3389/fnhum.2014.00987](https://doi.org/10.3389/fnhum.2014.00987)
19. Mormina E, Arrigo A, Calamuneri A, et al. Diffusion tensor imaging parameters' changes of cerebellar hemispheres in Parkinson's disease. *Neuroradiology*. 2015; 57(3): 327-334. doi: [10.1007/s00234-014-1473-5](https://doi.org/10.1007/s00234-014-1473-5)
20. Mormina E, Briguglio M, Morabito R, et al. A rare case of cerebellar agenesis: a probabilistic constrained spherical deconvolution tractographic study. *Brain Imaging Behav*. 2015. doi: [10.1007/s11682-015-9377-5](https://doi.org/10.1007/s11682-015-9377-5)
21. Auriat AM, Borich MR, Snow NJ, Wadden KP, Boyd LA. Comparing a diffusion tensor and non-tensor approach to white matter fiber tractography in chronic stroke. *Neuroimage Clin*. 2015; 7: 771-781. doi: [10.1016/j.nicl.2015.03.007](https://doi.org/10.1016/j.nicl.2015.03.007)
22. Snow NJ, Peters S, Borich MR, et al. A reliability assessment of constrained spherical deconvolution-based diffusion-weighted magnetic resonance imaging in individuals with chronic stroke. *J Neurosci Methods*. 2015; 257: 109-120. doi: [10.1016/j.jneumeth.2015.09.025](https://doi.org/10.1016/j.jneumeth.2015.09.025)
23. Kristo G, Leemans A, Raemaekers M, Rutten GJ, de Gelder B, Ramsey NF. Reliability of two clinically relevant fiber pathways reconstructed with constrained spherical deconvolution. *Magn Reson Med*. 2013; 70(6): 1544-1556. doi: [10.1002/mrm.24602](https://doi.org/10.1002/mrm.24602)
24. Azadbakht H, Parkes LM, Haroon HA, et al. Validation of high-resolution tractography against in vivo tracing in the macaque visual cortex. *Cereb Cortex*. 2015; 25(11): 4299-309. doi: [10.1093/cercor/bhu326](https://doi.org/10.1093/cercor/bhu326)
25. Lerner A, Mogensen MA, Kim PE, Shiroishi MS, Hwang DH, Law M. Clinical applications of diffusion tensor imaging. *World Neurosurg*. 2014; 82(1-2): 96-109. doi: [10.1016/j.wneu.2013.07.083](https://doi.org/10.1016/j.wneu.2013.07.083)
26. Yao Y, Ulrich NH, Guggenberger R, Alzarhany YA, Bertalanffy H, Kollias SS. Quantification of corticospinal tracts with diffusion tensor imaging in brainstem surgery: prognostic value in 14 consecutive cases at 3t magnetic resonance imaging. *World Neurosurg*. 2015; 83(6): 1006-1014. doi: [10.1016/j.wneu.2015.01.045](https://doi.org/10.1016/j.wneu.2015.01.045)
27. Radmanesh A, Zamani AA, Whalen S, Tie Y, Suarez RO, Golby AJ. Comparison of seeding methods for visualization of the corticospinal tracts using single tensor tractography. *Clin Neurol Neurosurg*. 2015; 129: 44-49. doi: [10.1016/j.clineuro.2014.11.021](https://doi.org/10.1016/j.clineuro.2014.11.021)
28. Birinyi PV, Bieser S, Reis M, et al. Impact of DTI tractography on surgical planning for resection of a pediatric pre-pontine neurenteric cyst: a case discussion and literature review. *Childs Nerv Syst*. 2015; 31(3): 457-463. doi: [10.1007/s00381-014-2587-0](https://doi.org/10.1007/s00381-014-2587-0)

29. Mormina E, Longo M, Arrigo A, et al. MRI Tractography of corticospinal tract and arcuate fasciculus in high-grade gliomas performed by constrained spherical deconvolution: qualitative and quantitative analysis. *AJNR Am J Neuroradiol.* 2015; 36(10): 1853-1858. doi: [10.3174/ajnr.A4368](https://doi.org/10.3174/ajnr.A4368)
30. Jellison BJ, Field AS, Medow J, Lazar M, Salamat MS, Alexander AL. Diffusion tensor imaging of cerebral white matter: a pictorial review of physics, fiber tract anatomy, and tumor imaging patterns. *AJNR Am J Neuroradiol.* 2004; 25(3): 356-369.
31. Lu S, Ahn D, Johnson G, Cha S. Peritumoral diffusion tensor imaging of high-grade gliomas and metastatic brain tumors. *AJNR Am J Neuroradiol.* 2003; 24(5): 937-941.
32. Li Z, Peck KK, Brennan NP, et al. Diffusion tensor tractography of the arcuate fasciculus in patients with brain tumors: comparison between deterministic and probabilistic models. *J Biomed Sci Eng.* 2013; 6(2): 192-200.
33. Küpper H, Groeschel S, Alber M, Klose U, Schuhmann MU, Wilke M. Comparison of different tractography algorithms and validation by intraoperative stimulation in a child with a brain tumor. *Neuropediatrics.* 2015; 46(1): 72-75. doi: [10.1055/s-0034-1395346](https://doi.org/10.1055/s-0034-1395346)
34. Weiss C, Tursunova I, Neuschmelting V, et al. Improved nTMS- and DTI-derived CST tractography through anatomical ROI seeding on anterior pontine level compared to internal capsule. *Neuroimage Clin.* 2015; 7: 424-437. doi: [10.1016/j.nicl.2015.01.006](https://doi.org/10.1016/j.nicl.2015.01.006)