Editorial

*Corresponding author

Ronald S. Laura, DPhil

Professor

School of Education

Faculty of Education and Arts

The University of Newcastle

Callaghan NSW 2308, Australia

Tel. +61 2 4921 5942

Fax: +61 2 4921 7887

E-mail: ron.laura@newcastle.edu.au

Volume 3 : Issue 3

Article Ref. #: 1000WHOJ3e016

Article History

Received: December 18th, 2017

Accepted: December 19th, 2017

Published: December 19th, 2017

Citation

Laura RS. From menstruation to menopause: Have we medicalised the physiology of normalcy? (Part 1). *Women Health Open J.* 2017; 3(3): e27-e29. doi: [10.17140/WHOJ-3-e016](https://doi.org/10.17140/WHOJ-3-e016)

From Menstruation to Menopause: Have We Medicalised the Physiology of Normalcy? (Part 1)

Ronald S. Laura, DPhil*

School of Education, Faculty of Education and Arts, The University of Newcastle, Callaghan NSW 2308, Australia

UNDERSTANDING THE RELATIONSHIP BETWEEN MENSTRUATION AND MENOPAUSE

Although, there are exceptions to the rule, women generally experience 'menopause' between the ages of 48 to 54, within which time the cessation of all menstrual bleeding becomes a reality. However, many women find that their bodies desist menstruating in their late thirties (premature menopause) or in their early forties.¹ The commencement of menopause is triggered when the ovaries relinquish their responsibility for the production of estrogen. 'Estrogen' is an important sex hormone that is primarily produced in the ovaries, and one of its main functions is to regulate menstrual cycles, and moderate the inception of menopause.²

Interestingly, a small portion of the female's estrogen supply is produced in the body's fat cells. It is estrogen that hormonally prepares the uterus to accommodate the fertilised egg, and this remarkable phenomenon happens when estrogen radically stimulates the thickness of the uterine lining, so as to stabilise the implanting capacity of the fertilised egg.³

Three types of estrogen exist in the female's body: the first of these is referred to as 'estradiol'. It is regarded as the most potent of the three types, and it is manufactured in the ovaries. The next of these estrogen constituents is called 'estrone', which is produced from estradiol, and the final component of estrogen is known as 'estriol'. It is also produced in the ovaries, but in much smaller quantities than either of the other two.⁴ The subtle balance of the estrogen amalgam of these three constituents helps to reduce the incidence of heart attack, slows the rate of bone loss, reduces vaginal dryness, improves skin tone, and it can regulate mood swings, in the context of both menstruation and menopause. With the onset of menopause, estrogen levels diminish markedly, but they do not drop to Zero, as some estrogen is still produced in the fat cells which continue to provide the supporting tissue around the ovaries. By utilising precursors produced by the adrenals, estrogen is also formed in the intestinal tract.⁵ It is worth contemplating that one reason why women tend to gain weight after menopause is linked with the body's capacity to confiscate even the small pools of estrogen made available by hormonal reservoirs of fat cells in the supporting tissue around the ovaries, and in the intestinal tract.

FOCUSING ON THE TRANSITION FROM MENSTRUATION TO MENOPAUSE

Early or premature menopause can happen due to any of a range of factors, including the following:

- Although rare, the body's immune system which normally fights off diseases, may mistakenly attack the ovaries and keep them from producing the female hormones which regulate menstruation.⁶ Other autoimmune diseases such as thyroid disease, rheumatoid arthritis, and even chronic fatigue syndrome may also disrupt the otherwise regular patterns of menstrual balance and harmony. For example, research has demonstrated that women with myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) suffer from persistent and extreme tiredness, headache, feelings of chronic weakness, muscle and joint pain, memory loss, insomnia, and marginalised eyesight. Women with these problems are significantly more likely

Copyright

©2017 Laura RS. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

to experience early or premature menopause.⁷

- Women who smoke are more likely to experience menopause as early as two or three years before non-smokers, depending on the length of duration which characterise the woman's smoking habit. In addition, women who have smoked for ten years or more may also be afflicted with more severe and painful menopause symptoms such as acute stomach discomfort and intense muscle cramps.⁵ Moreover, research also shows that women who are long-term smokers will be not only succumb to premature menopause, but they will die about two years earlier than those women who are not smokers.⁸
- Chemotherapy or pelvic radiation treatments for cancer are notorious for inducing premature menopause. The problem is that chemotherapy treatments tend to damage women's ovaries, thereby causing a women's menstruation periods to cease unpredictability at any time. The resultant menstrual cessation may persist for just a while, or it may shut down the menstrual system forever.⁹ Chemotherapy may also adversely affect the fertility-organs and hormonal production mechanisms associated with pregnancy, such that a woman may not be able to get pregnant again, or never be able to get pregnant a first time. Although it is to admitted that chemotherapy or radiation do not necessarily cause infertility or unanticipated menopause, the older a woman is at the time of chemotherapy or radiation, the more likely it is that she will experience acute menopause and permanent infertility.¹⁰
- More dramatic interventions such as the surgical removal of both ovaries, called a bilateral oophorectomy are likely to precipitate menopausal symptoms almost immediately. After this surgery when both the women's ovaries are discarded, her menstrual periods will literally be eliminated, and her hormonal levels will immediately be significantly diminished. In many cases intense menopausal symptoms become manifest and hot flashes will be greatly magnified, while sexual desire will dwindle and quickly be depleted, thereby necessitating the implementation of hormone therapy.¹¹
- It sometimes happens that women will confront the problem of incessant bleeding in the context of a menstrual cycle or independently. In such cases it is generally recounted that a hysterectomy is the best way to stop the bleeding by the complete removal of the uterus, though in most cases women patients are fortuitously allowed to keep their ovaries intact. This being so, it follows that personal pregnancy is no longer an option, in as much as women who have undergone a hysterectomy will no longer have periods, or the capacity to become pregnant. However, given that the ovaries have been preserved, women who have had their uterus removed experience the almost immediate symptoms of menopause. Due to the fact that their ovaries are still functional, and their ovaries seem, almost miraculously, to produce an array of the very hormones which largely govern the extent to which the symptoms of menopause will infiltrate the women's body. In addition, if a women who has had to undergo the hysterectomy is young enough, she might still have to confront the experience of menopause a couple of years earlier than would otherwise have been anticipated.¹²
- Women with a family history of chronic disease which serves to act as a catalyst for menopause are more likely to experience an early or premature menopause than are those women without disease-disposing menopausal intrusions.
- Research has revealed that women suffering medical conditions of HIV whose infectious parameters are intractable, or ineffectively contained by the medicines being prescribed, the woman involved is very likely to experience the acute symptoms of early or premature menopause. It has also been shown that women who are dealing with HIV are far more likely to have more severe and debilitating hot flashes than women who are not afflicted with the virus.¹³
- One of the more recent medical discoveries related to menopause is that a surprising number of women experience the symptoms of early and premature menopause because they have a congenital deficiency such that they are born without certain chromosomes responsible for the endocrinal and physiological management of menopause. Similarly, some women have been found to have problems with the 'menopausally-related' and designated chromosomes that signal the production of endocrinal messaging which misguidedly prompts early episodes of menopause. For example, women with the condition called Turner's syndrome are born without all or part of one X chromosome. As a consequence, their ovaries fail to get appropriately structured at birth.¹⁴ This being so, the regulation of menstrual cycles for some women, including the scheduled sequencing of menopausal episodes are being seriously disrupted.

In PART 2 of this series, I will in the next issue of WHOJ address the issues associated with the efficacy of estrogen therapy for menopausal women.

REFERENCES

1. Lizcano F, Guzmán G. Estrogen deficiency and the origin of obesity during menopause fernando lizcano and guillermo guzmán. *BioMed Research International*. 2014(2014): 757461. doi: [10.1155/2014/757461](https://doi.org/10.1155/2014/757461)
2. Positon Statement on Menopause. 28th Annual Meeting of The North American Menopause Society October 11-14, 2017, Philadelphia, PA. *Menopause*. 2017; 24(12): 1418-1461. doi: [10.1097/GME.0000000000000997](https://doi.org/10.1097/GME.0000000000000997)
3. Gold, Ellen B. The timing of the age at which natural menopause occurs. *BioMed Research International*. 2014(2014): 757461. doi: [10.1155/2014/757461](https://doi.org/10.1155/2014/757461)
4. Stuenkel CA, Davis SR, Gompel A, et al. Treatment of symptoms of the menopause: An endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2015; 100(11): 3975-4011. doi: <https://doi.org/10.1210/jc.2015-2236>
5. Grady D. Management of menopausal symptoms. *N Engl J Med*. 2006; 355: 2338-2347. doi: [10.1056/NEJMcp054015](https://doi.org/10.1056/NEJMcp054015)
6. Marsh WK, Bromberger JT, Crawford SL, et al. Lifelong estradiol exposure and risk of depressive symptoms during the transition to menopause and postmenopause. *Menopause*. 2017; 24(12): 1351-1359. doi: [10.1097/GME.0000000000000929](https://doi.org/10.1097/GME.0000000000000929)
7. Bueno-Notivol J, Calvo-Latorre J, Alonso-Ventura V, Pasupuleti V, Hernandez AV, Pérez-López FR; Health Outcomes and Systematic Analyses (HOUSAY) Project. Effect of programmed exercise on insulin sensitivity in postmenopausal women: A systematic review and meta-analysis of randomized controlled trials. *Menopause*. 2017; 24(12): 1404-1413. doi: [10.1097/GME.0000000000000936](https://doi.org/10.1097/GME.0000000000000936)
8. Pinkerton JV, Kaunitz AM, Manson JE. Vaginal estrogen in the treatment of genitourinary syndrome of menopause and risk of endometrial cancer: An assessment of recent studies provides reassurance. *Menopause*. 2017; 24(12): 1329-1332. doi: [10.1097/GME.0000000000000996](https://doi.org/10.1097/GME.0000000000000996)
9. Pinkerton JV, Bushmakin AG, Bobula J, Lavenberg J, Komm BS, Abraham L. Hot flush frequency and severity at baseline as predictors of time to transient and stable treatment success: Pooled analysis of two CE/BZA studies. *Menopause*. 2017; 24(12): 1378-1385. doi: [10.1097/GME.0000000000000918](https://doi.org/10.1097/GME.0000000000000918)
10. Saccomani S, Lui-Filho JF, Juliato CR, Gabiatti JR, Pedro AO, Costa-Paiva L. Does obesity increase the risk of hot flashes among midlife women?: A population-based study. *Menopause*. 2017; 22(10): 1065. doi: [10.1097/GME.0000000000000884](https://doi.org/10.1097/GME.0000000000000884)
11. Sturdee DW. The menopausal hot flush—Anything new? *Maturitas*. 2015; 60(1):42-49. doi: [10.1016/j.maturitas.2008.02.006](https://doi.org/10.1016/j.maturitas.2008.02.006)
12. Reding KM, Schmidt PJ, Rubinow DR. Perimenopausal depression and early menopause: Cause or consequence? *Menopause*. 2017; 24(12): 1333. doi: [10.1097/GME.0000000000001016](https://doi.org/10.1097/GME.0000000000001016)
13. Yazdkhasti M, Simbar M, Abdi F. Empowerment and coping strategies in menopause women: A review. *Iran Red Crescent Med J*. 2015; 17(3): e18944. doi: [10.5812/ircmj.18944](https://doi.org/10.5812/ircmj.18944)
14. Elavsky S, McAuley E. Personality, menopausal symptoms, and physical activity outcomes in middle-aged women. *Pers Individ Dif*. 2017; 46(2): 123-128. doi: [10.1016/j.paid.2008.09.014](https://doi.org/10.1016/j.paid.2008.09.014)