The Volta Review

Alexander Graham Bell Association
for the Deaf and Hard of Hearing

The

Research

Facilitators and Barriers to the Integration of
Orally Educated Children and Youth With Hearing
Loss Into Their Families and Communities 5

*Alice Eriks-Brophy, Ph.D.,
Andrée Durieux-Smith, Ph.D., Janet Olds, Ph.D.,
Elizabeth Fitzpatrick, Ph.D.,
Cheryll Duquette, Ph.D., and
JoAnne Whittingham, M.Sc.*

Outcomes of an Auditory-Verbal Program for
Children With Hearing Loss: A Comparative
Study With a Matched Group of Children
With Normal Hearing 37

*Dimity Dornan, A.M., Ba.Sp.Th., F.S.P.A.A.,
Cert. AVT®, Cp.Sp., Louise Hickson, B.Sp.Thy.
(Hons), M.Aud., Ph.D., Bruce Murdoch B.Sc.
(Hons), Ph.D., D.Sc., and Todd Houston, Ph.D.,
CCC-SLP, Cert. AVT®*

An Analysis of Phonological Process Use in
Young Children With Cochlear Implants 55

*Helen C. Buhler, Ph.D., Betty DeThomasis, M.S.,
Pat Chute, Ed.D., and Anne DeCora, M.S.*

EMMACOLLEY :: age 3

Implanted June 2003 (left), January 2005 (right)

I CAN

Tell Emma jokes in the car

Teach her new songs and watch her dance

Talk to her from the next room

Read her stories when she's on my lap

Listen as she talks with her friends

Feel pride in her age-appropriate language skills

Understand exactly what she's saying

Marvel at her abilities in a mainstreamed classroom

Whisper "I love you" to her

Believe she can do and be anything she wants

Introducing the
Absolute Latest
in Cochlear Implant Technology:
the **NEW PULSARci¹⁰⁰**

hear**LIFE**

When you choose a cochlear implant from MED-EL – either for yourself or a loved one – YOU CAN open up a world of possibilities. We not only offer today's most technologically advanced cochlear implant systems, but we believe we are on the verge of breakthroughs that will offer users a lifetime of more exciting tomorrows. For more information about MED-EL, contact us today.

TOLL FREE (888) MEDEL-CI (633-3524) :: V/TDD PHONE (919) 572-2222 :: www.medel.com

The Volta Review

Volume 107, Number 1
ISSN 0042-8639

Organized in 1890 to encourage the teaching of speech, speechreading and the use of residual hearing to people who are deaf or hard of hearing, the Alexander Graham Bell Association for the Deaf and Hard of Hearing (AG Bell) welcomes to its membership all who are interested in improving educational, professional and vocational opportunities for people who are deaf or hard of hearing. Affiliated with AG Bell are the Parent Section, the Deaf and Hard of Hearing Section and the International Professional Section.

3 Editor's Preface

Patricia M. Chute, Ed.D.

Research

- 5 Facilitators and Barriers to the Integration of Orally Educated Children and Youth With Hearing Loss Into Their Families and Communities

Alice Eriks-Brophy, Ph.D., Andrée Durieux-Smith, Ph.D., Janet Olds, Ph.D., Elizabeth Fitzpatrick, Ph.D., Cheryll Duquette, Ph.D., and JoAnne Whittingham, M.Sc.

- 37 Outcomes of an Auditory-Verbal Program for Children With Hearing Loss: A Comparative Study With a Matched Group of Children With Normal Hearing

Dimity Dornan, A.M., Ba.Sp.Th., F.S.P.A.A., Cert. AVT®, Cp.Sp., Louise Hickson, B.Sp.Thy. (Hons), M.Aud., Ph.D., Bruce Murdoch B.Sc. (Hons), Ph.D., D.Sc., and Todd Houston, Ph.D., CCC-SLP, Cert. AVT®

- 55 An Analysis of Phonological Process Use in Young Children With Cochlear Implants

Helen C. Buhler, Ph.D., Betty DeThomasis, M.S., Pat Chute, Ed.D., and Anne DeCora, M.S.

- 75 Information for Contributors to *The Volta Review*

Permission to Copy: The Alexander Graham Bell Association for the Deaf and Hard of Hearing, as copyright owner of this journal, allows single copies of an article to be made for personal use. This consent does not extend to posting on Web sites or other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works of any type, or for resale without the express written permission of the publisher. For more information, contact AG Bell at 3417 Volta Place, NW, Washington, DC 20007, e-mail editor@agbell.org, or call 202/337-5220 (voice) or 202/337-5221 (TTY).

Editor's Preface

Patricia M. Chute, Ed.D.

For children with hearing loss, success in the classroom is not only a result of curricular knowledge; it also encompasses a broad range of abilities from interaction with peers to clear, articulate speech. The spring 2007 issue of *The Volta Review* includes three articles in the areas of socialization, the outcomes of children enrolled in Auditory-Verbal therapy and phonological processes of children with cochlear implants. For teachers and school personnel, the information contained in each of these articles merits a close look.

Research by Eriks-Brophy and colleagues utilizes a focus-group model to identify potential facilitators and barriers to the effective integration of children with hearing loss into their families and communities. Interviews with students with hearing loss, their parents and itinerant teachers of the deaf reveal challenges that are being addressed, as well as those that still remain. Not only does the research address the needs of students with hearing loss as they negotiate their school systems, group activities and social interactions, it also discusses the roles that mothers, fathers, siblings and professionals play in the process. Clearly, as more children with hearing loss enter mainstream settings earlier, the effects of integration on these students and their families will require further attention and study.

The outcomes of Auditory-Verbal therapy as evaluated by monitoring a group of young children over four years is the focus of the first phase of a study by Dornan and colleagues. This study reviews the speech and language abilities of a group of 29 children ages 2 to 6 with severe-to-profound hearing loss as compared with the abilities of children with normal hearing of the same age. Performance across the measures assessed in this first phase indicates no significant differences between these two groups. Although the study participants comprise a small sample within a restricted age group, it is important to recognize the effects of intensive listening training on overall language development. Data on the overall progress of the children involved in this longitudinal study will be an important contribution to the field.

Finally, the phonological processes of a select group of very young children with cochlear implants are the subject of the investigation by Buhler and

colleagues. The authors address the development of phonological skills in children with cochlear implants through the use of evaluative procedures commonly used by school and early intervention speech-language pathologists (SLPs). Their results indicate that children who had greater experience using implants to hear performed better and demonstrated fewer phonological processes. These findings are particularly informative for early intervention SLPs who are providing services to children with cochlear implants because they identify tools and strategies that can be implemented readily.

The Volta Review has evolved over the past few years into a publication that addresses both current and future needs of children with hearing loss. This evolution could not have been achieved without a remarkable managing editor and editorial board. However, as with so many things that change over time, so too do the editorial personnel. Managing Editor Dawn Scarola recently stepped down from her position at *The Volta Review* to take on a new role as the editorial production director for another Washington-based nonprofit publisher. Her energy and enthusiasm will be missed sorely. Additionally, longtime associate editor Mary Ellen Nevins is moving on to become national director of the newly developed Professional Preparation for Cochlear Implants (PPCI) program at the Children's Hospital of Philadelphia. To say she will be missed is an understatement. I am sure that I speak for the readers and the editorial board of *The Volta Review* when I offer Dawn and Mary Ellen our best wishes as they each embark upon new endeavors.

Pat Chute, Ed.D.
Editor
The Volta Review

Facilitators and Barriers to the Integration of Orally Educated Children and Youth With Hearing Loss Into Their Families and Communities

Alice Eriks-Brophy, Ph.D., Andrée Durieux-Smith, Ph.D., Janet Olds, Ph.D., Elizabeth Fitzpatrick, Ph.D., Cheryll Duquette, Ph.D., and JoAnne Whittingham, M.Sc.

Family and community interactions provide important opportunities for facilitating the integration of children and youth with hearing loss, yet these environments have received little research attention. In this study, facilitators and barriers to integration associated with the social milieus of young people with hearing loss were identified. Facilitators and barriers associated with the family were related to parental roles and relationships between family members. Community facilitators and barriers were related to the inclusion of children in sports and social activities and to accommodations made by leaders and peers participating in these events. Challenges to the successful inclusion of children and youth with hearing loss into family and social environments were identified. Nevertheless, the participants retained positive attitudes toward the benefits of inclusion and the potential of individuals with hearing loss to become fully contributing members of their families, communities and society at large.

Alice Eriks-Brophy, Ph.D., is an assistant professor in the Department of Speech-Language Pathology at the University of Toronto, where she teaches courses on aural rehabilitation and articulation and phonological disorders in children. Andrée Durieux-Smith, Ph.D., is a professor of audiology and vice-dean of professorial affairs in the Faculty of Health Sciences at the University of Ottawa. Janet Olds, Ph.D., is a psychologist at the Children's Hospital of Eastern Ontario, where she works on the Cochlear Implant Team and the Behavioural Sciences and Consultation Liaison Team. Elizabeth Fitzpatrick, Ph.D., is an audiologist and a research associate at the Children's Hospital of Eastern Ontario. Cheryll Duquette, Ph.D., is a professor in the Faculty of Education at the University of Ottawa, where she teaches courses on special education and learning theories. Jo Anne Whittingham, M.Sc., is the research coordinator for the Audiology Research Lab at the Children's Hospital of Eastern Ontario.

Introduction

The integration of orally educated children with hearing loss often is examined primarily from the perspective of inclusion within educational settings. Research on the inclusion of children with hearing loss has focused primarily on pedagogical practices and educational structures for supporting them as students in regular classrooms. The advantages of mainstreaming children with hearing loss within the school setting include (1) promotion of social interaction with the hearing world, (2) availability of normal linguistic and behavioral models provided by peers, (3) a stimulating and highly oral environment with rich linguistic input, (4) increased opportunity for learning and socialization opportunities and (5) education within the least restrictive environment (Brackett, 1993; Gresham, 1982; Ladd, Munson, & Miller, 1984; Maxon & Brackett, 1992; Nolan & Tucker, 1981; Northcott, 1984, 1990; Nowell & Innes, 1997; Ross, 1990). Clearly, such advantages exist not only within the environment of the classroom but also in the family and wider community as well. However, few studies to date have examined inclusive structures and practices that might facilitate the inclusion of orally educated children and youth with hearing loss into family, social and community environments.

Those few studies that have examined issues associated with the inclusion of individuals with hearing loss in environments other than school and the classroom have focused primarily on the perceptions and experiences of parents of preschool children (see, for example, Gregory [1995] and Meadow-Orlans, Mertens & Sass-Lehrer [2003]), whereas the stories of families of older children and young adults have been given far less consideration. Interesting exceptions include research by Green (1990) and Schwartz (1990), which both reflect some of the themes that emerged from the present study.

Green (1990) described the perceptions of three parents and three young adults with hearing loss who were asked to provide written responses to a series of questions about the social and educational advantages and disadvantages of inclusion. Parent respondents in the study discussed the importance of academic integration and the types of supports they felt were required for successful integration as well as the importance of parental advocacy in the school setting. They also described the difficulties their children experienced in establishing friendships and participating in social events and activities. The young adults with hearing loss themselves expressed the sense of self-worth and confidence they had developed through attending a mainstream school. Although all three of the young adult respondents with hearing loss mentioned periods of social isolation and difficulties with peers, they all agreed that the opportunities provided to them through the inclusive model resulted in high levels of educational achievement. Furthermore, participation in extracurricular activities, such as sports and music, was em-

phasized by the students in particular as having provided them with important opportunities for social interaction with their peers.

In her discussion of psychosocial aspects of mainstreaming, Schwartz (1990) reported on the results of an informal survey distributed to an unspecified number of teachers of integrated children with hearing loss. Participants were asked to respond to questions about the skills required for mainstream success and the role of parents in developing these skills. Although the results focused primarily on teachers' perceptions of academic aspects of inclusion, the study also summarized comments from five parents of integrated children with hearing loss. These data provided some insight into parents' perceptions of the social and community inclusion experiences of their children. These parents mentioned the importance of involvement in extracurricular activities, the acceptance of the hearing loss and the importance of building self-esteem and independence as crucial qualities for successful integration of children with hearing loss. No comments from the students with hearing loss themselves were reported as part of that study.

Research examining the parenting roles of fathers versus mothers in providing intervention and language stimulation to their children is relatively limited within the literature on hearing loss, but the subject is beginning to receive some attention. This literature notes that mothers continue to be the primary participating parents in early intervention programs and home-based language stimulation activities and that they assume primary responsibility for their children's daily care (Calderon, Bargones & Sidman, 1998; Calderon & Low, 1998; Meadow-Orlans & Sass-Lehrer, 1995; Meadow-Orlans et al., 2003). The literature further shows that fathers are not perceived as disinterested in participating in such programs but, rather, as less able to organize their work schedules to accommodate early intervention appointments during business hours. Some studies indicate that mothers of children with hearing loss experience higher stress levels, less satisfaction with their spousal relationship, insufficient spousal support and inadequate time for themselves as compared to fathers (Brand & Coetzer, 1994; Meadow-Orlans, 1995). Nevertheless, young children with hearing loss who were reared in families in which fathers were present were reported to achieve higher academic, social-emotional and language outcomes than those where fathers were not present (Calderon & Low, 1998).

These findings are also true for young children who are at risk for developmental delays and are enrolled in programs such as Head Start (Cabrera, Tamis-LeMonda, Lamb & Boller, 1999), programs for children with physical disabilities (Hadadian & Merbler, 1995; Gavida-Payne & Stoneman, 1997; Turbiville, 1995), programs providing services specifically aimed at cultural minority children with disabilities (McWilliam, McGhee & Tocci, 1998) and programs providing services to children with autism and intellectual delay (Konstantareas & Homatidis, 1992).

Research Objectives

The goal of the qualitative component of a larger research study conducted by Durieux-Smith et al. (1999) was to identify facilitators and barriers to the inclusion of students with hearing loss in the domains of school, family and peers and to develop a set of recommendations for policy and practice that would facilitate the effective integration of students with hearing loss into their academic and social environments. Qualitative data were collected through a series of 10 focus group interviews consisting of guided questions related to participants' experiences with and perceptions of the integration of students with hearing loss in academic, family and community environments. Recommendations related to policies and practices that might facilitate such inclusion also were elicited.

In this article, facilitators and barriers to the inclusion of children with hearing loss into their family and community environments obtained from these focus group participants are presented. A previous article by the authors (Eriks-Brophy, Durieux-Smith, Olds, Fitzpatrick, Duquette & Whittingham, 2006) describes the facilitators and barriers to educational inclusion obtained from the same groups of participants.

Method

A self-selected group of young people with hearing loss, their parents and the itinerant teachers of the deaf who serve these students in Ottawa, Ontario, Canada, participated in focus groups examining their perceptions of integration and their recommendations for facilitating inclusion. Participation in the focus groups was voluntary. To ensure adequate opportunities for discussion, focus groups were limited to six to eight members per group. A total of 10 focus groups were conducted. Twenty-four parents of young adults with hearing loss participated in four parent focus groups. Sixteen young adults with hearing loss participated in three focus groups. Oral interpreters were provided as needed for those who requested such services. Fourteen itinerant teachers of the deaf representing two school commissions in the Ottawa region participated in three focus groups. The research was conducted at the Children's Hospital of Eastern Ontario (CHEO), a regional pediatric hospital in Ottawa that uses an Auditory-Verbal approach to speech and language intervention. In addition to mainstreaming into regular classrooms, one of the primary objectives of Auditory-Verbal therapy (AVT) is the inclusion of children with hearing loss into community and social environments.

Participants

Students With Hearing Loss

Young people were identified as potential participants if they were, or had been, patients of CHEO's audiology department. Since CHEO opened in 1974,

virtually all children in the geographic area that CHEO serves with known or suspected hearing losses have been referred to CHEO's audiology department for assessment and, when appropriate, for rehabilitation. Additional criteria for potential participants included the following:

- age 14 years or older at the time of the study,
- previous diagnosis of a hearing loss that was not medically treatable and required hearing aids,
- enrollment in the Auditory-Verbal preschool therapy program at CHEO,
- no medical evidence of a developmental or cognitive disability that would impact communication ability and
- English as the primary language of instruction in Auditory-Verbal therapy and at school.

The 16 young adults with hearing loss who participated in the focus groups had a mean pure tone average of 84.6 dB (± 19.5 dB) and were fitted with hearing aids at an average age of 22 months (range: 4–60 months). Their average age at the time of participation in the study was 19.7 years, with a range of 15–30 years. Eight participants were male and eight were female. The participants came primarily from two-parent families (87.5%) that had an average of two children per family. The primary language of the home was English for 96% of the participating families. The majority of participants had been educated in mainstream environments throughout their educational history (71%), whereas 8% had been taught in classes for children with hearing loss at some point during their elementary school years and 21% reported having received some special education services secondary to learning difficulties not associated with their hearing loss. At the time of the study, eight of the participants were enrolled in high school, four were enrolled in post-secondary educational institutions and four had completed their education and were in the work force.

Parents and Itinerant Teachers

Ten fathers and 14 mothers participated in the four parent focus groups. Two of the mothers were single parents. All parent participants were parents of the young adults with hearing loss participating in the study. Levels of education of the parent participants were relatively high, with 74% of mothers and 87% of fathers having a postsecondary level of education. The socioeconomic status of the parent participants also was relatively high, with the majority of participants belonging to middle class or upper middle class socioeconomic groups as determined by the profession of the primary family wage earner.

The 14 itinerant teachers who took part in the focus groups were all female teachers who represented two school commissions in Ottawa, Ontario. All

participants had a bachelor of education degree and had a minimum of five years of experience as itinerant teachers of children with hearing loss. Seven of the itinerant teachers were Certified Auditory-Verbal Therapists®.

Apparatus

Focus group interviews were recorded using a Sony BM-246 conference recorder. Separate channels and microphones allowed the contributions of individual speakers to be identified throughout the transcripts.

Procedures

All focus groups were conducted by the same experienced facilitator in order to ensure comparability across groups of participants. Each focus group began with an introduction to the study as a whole, an explanation of the focus group procedure and the signing of consent forms. Focus groups were held in the late afternoon or evening at CHEO for the parents and young adults and at the offices of two of the local school commissions for the itinerant teachers. The average duration of each focus group was three hours. A pilot focus group with parents whose children were still too young to participate in the study was carried out to ensure the procedure used would elicit the information required for the study. One recommendation from the pilot focus group participants was that an information package containing guided questions be provided prior to the meeting to orient participants to the topics that would be discussed during the focus groups. Many of the participants made extensive notes prior to meeting in the focus groups, which they then used during the discussions.

At the beginning of each focus group, participants were asked to respond to the question "What factors related to school/home/community/peers acted as facilitators/barriers to integration?" by independently identifying facilitators and barriers to integration in these areas on a response sheet. The focus group facilitator then went around the table, asking each participant to contribute one of the items they had listed to the group discussion. Other participants voiced their agreement or disagreement with the item mentioned and were given the opportunity to discuss their personal experiences related to the facilitator or barrier to integration under discussion. Participants were asked to contribute novel items that had not been previously mentioned as the focus group progressed. All responses were listed on chart paper as either facilitators or barriers to integration according to the domains of interest. A period of 20 minutes was reserved at the end of each focus group to obtain recommendations for policy to enhance integration in the areas of family, school, community and peers.

Analysis

All of the focus group interviews were transcribed in their entirety. Two independent coders developed a set of coding categories related to the facilitators and barriers to integration identified in the transcripts. The CHILDES database (MacWhinney & Snow, 1990), a computerized language analysis system, was used in the analysis conducted by one of the coders, and the second coder entered codes freehand directly onto the printed transcripts. The two analyses were compared, and a final list of codes based on those derived by the two independent coders was developed. This final list of codes was used to analyze the focus group data. Results from the focus groups were reported through quantitative summaries as well as through excerpts taken from the focus group interviews. These excerpts were taken verbatim from the transcripts to illustrate ideas central to the identified themes, with the exception of the removal of hesitations, repetitions, false starts and place fillers. All names were changed in order to respect anonymity; however, these names have been used consistently to allow recognition of the contributions and perspectives of individual participants.

Results

Facilitators of Family and Community Inclusion

Facilitators of family and community integration identified across the focus groups are summarized in Figure 1. The contributions of the focus groupings of students, parents and itinerant teachers to each of the identified variables are presented in Figure 2. In the sections below, themes associated with the identified facilitators and barriers to family and community inclusion are presented. Facilitators of family and social inclusion are described first, followed by barriers to such inclusion. In many cases, facilitators and barriers were described by participants as existing along a continuum such that a facilitator, if not present or inappropriately implemented, became a barrier.

Facilitators of Inclusion Related to Family

Facilitators related to the family and family inclusion were the most frequently mentioned of all categories across all groups of participants, constituting 201 of the total 497 identified facilitators, as shown in Figure 1. Parent focus-group participants tended to focus on facilitators of family and social integration related to family variables and identified 132/201 or 65.6% of the family variables. Fewer facilitators in this domain were provided by itinerant teachers (34/201 or 16.9%) and students with hearing loss (35/201 or 17.4%). Many of the family-related factors facilitating integration were associated with parental roles and the relationships between family members. Others were related to the values and characteristics that parents promoted in their

Figure 1. Facilitators of family and community integration.

children with hearing loss. The important role of parental involvement in supporting and promoting social inclusion is highlighted through the voices of the participants that serve to illustrate the various themes extracted from the data.

Personal sacrifices. Obtaining appropriate services for their young children with hearing loss was a primary consideration for all of the parents who participated in the focus groups. Many of the participating parents had made considerable sacrifices, including leaving their families and countries of origin behind, in order to ensure their child received early speech and language intervention. Three sets of parents (25% of participating parents) specifically described moving to Ottawa in order to access services for their child. The following excerpt from a mother illustrates the lengths to which some families were willing to go to obtain such services:

Ranya: Well, in our case, we had to make a big decision. Because we left our country to come and settle over here because of Rasheed. I had a sister over here and the support from her and her family was a big thing for us to make the decision and stick together. No matter

Figure 2. Contributions of focus group participants to identified facilitators.

what, we knew we had to help this coming together as a family. And with the extended family supporting you, we knew we could do this. I think that really made a big difference, family and friends. They all wanted to understand what the problem was and how best to help.

Another type of sacrifice parents and, particularly, some of the mothers, made to ensure their children with hearing loss would have optimal stimulation for language and cognitive development was to give up their jobs, careers and the associated income in order to stay home with their preschool-aged child with hearing loss. Four of the 14 mothers specifically described issues related to having left a job they enjoyed to stay at home and work with their child. The importance of making such a decision also was discussed in two of the three focus groups held with itinerant teachers. The devotion and commitment of these parents to their children is summarized in the following excerpt:

Amy: When you're doing lesson plans every day and trying to get to meetings, and trying to get more information on how to do the job better, and going for tests, and going for medical appointments, and

having left a full-time job that I really loved and making *this* my full-time job, and really wanting to do everything I could do for her . . . it was a very interesting time, a very interesting time.

Supportive networks. The presence of a supportive network of extended family and friends was described by parents and itinerant teachers in particular as a key factor in helping them deal with the pressures and stresses of raising a child with hearing loss. These supports were perceived as crucial to maintaining the motivation and encouragement required for the constant provision of input for speech and language stimulation and development. All of the parent participants (100%) and participants in two of the three itinerant teacher groups described how a network of supportive family and friends was essential to coping with feeling isolated and overwhelmed at times by the demands placed on them as a result of their child's diagnosis of hearing loss.

Participation in parent support groups such as the Ottawa chapter of VOICE for Hearing-Impaired Children or other less formalized groups provided parents with opportunities to interact with other families of children with hearing loss, discuss problems and recognize that they were not alone in the challenges they faced in bringing up their child with hearing loss. Eighteen of the 24 parent participants (75%) reported being members of VOICE. In many instances, relationships established through this parent support group grew into strong and long-lasting friendships, both for the parents as well as their children.

Dave: I think a big part, under family support, that's VOICE for Hearing Impaired Children. None of us would have met but for VOICE, and just by talking in group sessions we realized there are common interests, common concerns, "I'm not the only one." In most cases you think, "I'm the only one that's feeling this way," and you start discussing and you're going, "Oh, everybody is in the same boat." So VOICE was almost more important than direct family.

Sandra: Yes, they became family.

Maintaining a balance in the family. Efforts to maintain a balance in the family, both in terms of parenting roles and with respect to the needs of other siblings in the family, were mentioned by all of the parent participants as essential to maintaining healthy family relationships. This topic also was discussed in two of the three focus groups held with the itinerant teachers. Extended family members who understood what these families were experiencing were only one factor in providing support to the couple as well as ensuring needed respite time. It was also important that parents were able to share the daily challenges of bringing up their children. Maintaining open communication between partners and developing good coping skills were seen as vital

components of a strong parental relationship. Parents mentioned the importance of talking openly and frankly about problems, accepting their own and their partners' feelings and reactions to the stresses and demands of raising a child with hearing loss and being good listeners.

Nevertheless, seven mothers participating in two separate focus groups discussed how the bulk of activities associated with early language stimulation and intervention fell to them, while their husbands carried on with their jobs and careers in the work force. A division of parenting roles that ensured some private time for each parent was perceived as crucial to maintaining the motivation and energy necessary for the challenges of daily life, particularly in the early years. Setting aside time within the day for the parent to engage in preferred independent activities such as reading a book or the newspaper, watching a favorite television program or listening to the radio was perceived as important to the mental health of the individual parent.

Another important coping strategy mentioned by parents was the ability to maintain a sense of humor and to share laughter with other parents in facing the challenges of bringing up a child with a hearing loss. This strategy implied having a close group of understanding friends who were able to relate to what these couples were going through. Often, these friends were parents of other children with hearing loss. The important role of humor in maintaining family balance and coping with stress was discussed in two of the four parent focus groups and one of the three itinerant teacher focus groups. Parent focus groups in particular were characterized by the use of humor and a great deal of laughter in discussing both the facilitators and obstacles to integration they had encountered.

Peter: On the positive side, the getting through with friends and, really, to laugh at some of it. At times, our coping was to see the funniness and the silliness of it and talk about it. Talk about the escapades that went on and things that happened.

Parents in two of the four focus groups discussed how the demands of Auditory-Verbal therapy had the potential to take over family interactions and activities and that it was important not to lose sight of the fact that their child was a child first, and a child with hearing loss second. These parents perceived the supportive role of family as being crucial to maintaining this balance.

Mary: When my son was two, the Christmas he was two, my mother bought him a toy train and she said, "This is for fun." Because I had already learned that, I walked into a toy store, and I saw how much language I would get out of it, how much speech I could get out of it. But she bought him a locomotive that went down the back hall and the light went on and off and it went "toot-toot" and he thought it was absolutely wonderful.

Siblings. All parent participants mentioned the importance of devoting attention and time to the overall development of the child with hearing loss within the family; however, they also were very aware that the needs of other family members could and should not take second place to those of the child with hearing loss. Maintaining a balance among the needs of other siblings in particular was seen as an important and, at times, challenging priority for families. Although parents generally agreed that such a balance was important, there was also a certain consensus that other children in the family would and could not always get the amount of attention they deserved, given the time commitments associated with working with the child with hearing loss, particularly during the preschool and early school years. Maintaining a family balance became especially problematic for families who had more than one child with special needs. Sibling rivalry, attention getting, acting out and the reality of having less quality time to spend with the sibling were mentioned in all four of the parent focus groups.

Amy: The day-to-day issues I remember were like spending three hours at night preparing a lesson plan that would take an hour to implement in the highchair the next day and then spending the whole rest of the day reinforcing the language. My plan was everything in the house and every outing. I also remember that my other child suffered for it.

Ranya: I think the siblings do lose. No matter how hard you try to balance things out, you don't give them as much attention as they should be getting. And that is one of the barriers I would say.

Amy: Yes, I fully share those comments and recollections.

Dorothy: I felt exactly the same way.

Maintaining open communication with the adolescent child who has a hearing loss. A number of facilitators specifically related to parenting an older child with hearing loss also were mentioned during the focus groups. The importance of developing the whole child and not overemphasizing the hearing loss was particularly important during the early years. As the child with hearing loss grew older, these concerns were replaced by the importance of maintaining open communication and trust and fostering independence and responsibility within the family. The fact that their parents encouraged them to pursue their goals and dreams and did not impose limits on these ambitions as a result of their hearing loss was highly valued by all of the participating young people and had an important influence on how they dealt with the challenges they encountered in their lives.

Jenny: Well, my parents from the very beginning were extremely supportive . . . I've always been raised to not accept any limits and to try and go beyond everything. I'm in a gifted program in [a local high school], and my parents don't feel they need to help anymore with school work and they are just there for moral support. My mom was really important in my life. She's a very big role model for me. She works full time and so does my dad, and she just kept telling me, "Do what ever you want and if you don't like something, let people know."

Facilitators of Social Relationships and Inclusion in the Community

All participants mentioned facilitators to social networking and community inclusion, although fewer facilitators were identified for this domain compared to those associated with the family. Within the community environment, 11 of the parents described facilitators that were related primarily to the inclusion of children in sports and social activities. Additional facilitators identified by participants in all of the groupings included accommodations made by the leaders of extracurricular activities for the inclusion of children with hearing loss and the attitudes students with hearing loss had toward themselves and others, as well as peer attitudes toward individuals with hearing loss. Administrators, itinerant teachers and technology also were perceived as playing a facilitative role in social and community inclusion, as illustrated in Figures 1 and 2.

Facilitating friendships. The importance of parental involvement in initiating and facilitating friendships for their children with hearing loss, particularly in the early years, was discussed in all of the focus groups. Parents who actively promoted the development of social relationships, encouraged their child to become involved in social activities and initiated social interactions for their young children were perceived by the adult participants as having had an important positive impact on social and community inclusion. Parent participants all described having gone out of their way to make their house an open and welcome environment for other children in the neighborhood to come over and play. As one of the parents stated, "We were the house on the corner with the cookies and the drinks." Parents spoke of driving their children all over the community so they could meet up with friends and take part in social activities, as well as allowing their child to have sleepovers, parties and get-togethers regularly in their home. Although parents joked about how much time this had involved, none of them expressed any negative feelings related to such time commitments. Three sets of parents stated that they made a conscious decision to purchase a home in an area where there were many young children in order to increase the likelihood of exposing their child with

hearing loss to natural peer interactions. As their children became older, parents continued to encourage their children's friendships through opening their homes for teen activities, pool parties and other social events. However, these facilitating roles were seen as becoming substantially more difficult as their children grew older and more independent.

Enrollment in sports and social activities. Participants also commented on the importance of being enrolled in clubs, sports and other activities as an important facilitator to social integration. The importance of opportunities for social interaction with peers outside of the school setting provided by such activities was emphasized in all of the parent and itinerant teacher focus groups. Eleven parents described the activities they had encouraged their children to participate in. These parents expressed the opinion that such activities allowed their children to participate on an equal footing with their peers and were seen as an integral component in the development of self-esteem.

Janet: One way you can do that [promote social inclusion], and it's what we did with Marsha, is to expose them to all the regular activities that kids go through, like Brownies, Guides, gymnastics, swimming . . .

Frank: The other thing is that our son is very active in sports, and so he can excel in a range of things and feel good about himself without the hearing loss even being a factor.

Linda: Yes, to find a niche that gives them some credibility . . . we were really lucky because our kid was a good athlete. It just made all the difference in the world to him, being able to make friends.

Jack: Yes, I agree. It was a way in which our son could compete as an equal . . . we used sports, but we always said it didn't have to be sports. I mean, it could be art, it could be dancing, it could be anything. But it was finding out what they liked so that they could excel at that. Because once they excelled at that, it just carried over into their self-confidence and self-esteem.

Several parents took on leading roles in sports and other community activities to facilitate their child's social inclusion. Specifically, three out of 10 fathers described becoming involved in coaching the sports their child enjoyed, and two out of 14 mothers volunteered in the Brownies and Girl Guide troops in which their daughters participated. An additional positive consequence of this community involvement was the opportunities it provided to educate the general public about hearing loss and the enormous potential of children with hearing loss to participate in and contribute to society.

Peer relationships. All participants in the focus groups expressed the opinion that it was not necessary to have a large group of friends in order to feel included, but that instead, a few close friends were more than sufficient. Characteristics of good friends mentioned by the students in particular included patience, an understanding of the communication needs of the individual with hearing loss, the willingness to repeat information not initially understood, the ability to act as social and communication facilitators in difficult listening conditions and the acceptance of the hearing loss. Being able to find these qualities in a few good friends was described by all participants as being preferable to having a larger circle of less sensitive friends. As two parents stated:

Diane: In a way, a support to integration is a friend who just sees the child as normal. They don't see the disability.

Linda: Yes, that's right. I remember way back when he was young, like four or five, around there, I was speaking at a kindergarten class about hearing impairment. And there was a little girl there who said it all. Because I was talking about people who are disabled, like granny who uses a cane, and somebody else in a wheelchair, and I asked if anybody knew someone who had a disability, a problem that they needed some help with. And little Kathy just sat there, and I thought, "And she plays at our house all the time!" So I said, "Kathy, you know someone who has a disability." And she said, "Who?" And I said, "Well, Steve!" And she said, "Oh, he's just my friend." That was the only time I ever heard that, and it made my life!

Thirteen of the 16 young adults with hearing loss discussed the important role of friends in facilitating communication by acting as interpreters of missed information and helping them to follow along more easily in social situations, particularly those with difficult listening conditions. Two of the student focus groups as well as one group of itinerant teachers specifically pointed out the important role of peer relationships in keeping up with current trends and slang associated with adolescent culture.

Meg: I think friends are important for helping us kids keep up socially. It's harder for us to pick up slang words and phrases of the day . . . and it helps if we have friends to explain them to us. Friends help us learn the slang of the day. I wouldn't pick it up if I didn't have my friends to teach me.

Supportive siblings, particularly older siblings, who were able to advocate for the child with hearing loss in social and community activities and events also

were mentioned as facilitators of inclusion in all three of the focus groups held with students with hearing loss.

Communication with other individuals with hearing loss. Although all of the participants emphasized the importance of participating in academic and social activities with hearing peers, contact with other children with hearing loss was discussed as essential to building self-esteem and a strong sense of identity in all three focus groups held with the itinerant teachers, two of the three focus groups held with parents and two of the three groups of young people. Involvement in teen groups that provided opportunities to interact with other individuals with hearing loss was described as being important particularly during the adolescent years in two of the three parent focus groups and all three of the groups held with the itinerant teachers. Eleven of the 16 young people with hearing loss discussed the supportive role of the VOICE or itinerant teacher-initiated teen groups in developing friendships and providing opportunities to discuss and come to terms with their hearing loss.

Doug: On the peer side, VOICE was very important and became very critical in our daughter's case as far as meeting kids who understood each other and their circumstances.

Mary: They all looked the same at the Christmas party, remember? All running around with their FMs on.

Doug: "I'm not the only one!"

Mary: Yes, they always said that! In fact, if you had any brothers or sisters who went, they felt out of place! My older son always said, "I feel out of place," and I said, "You're only out of place one day of the year. Brian has it the other way."

Dave: As well, our daughter's primary friends became kids she met through VOICE, still today. Her best friends are kids she met at VOICE and not through school or the neighborhood.

Thirteen parents who participated in two of the parent focus groups specifically mentioned the lack of access and interaction with other children with hearing loss as one of the few negative consequences of integration into the local school. These parents spoke of the development of "magnet schools" where several children with hearing loss could be integrated, thus receiving all of the benefits of inclusion while still having the opportunity to interact with a group of individuals with hearing loss.

Characteristics of young people with hearing loss. Particular characteristics of young people with hearing loss were perceived as being facilitative of social inclusion

and were the second most frequently mentioned variable across all focus groupings, as illustrated in Figure 1. Facilitators associated with characteristics of individuals with hearing loss were distributed relatively evenly across all of the focus groups, as shown in Figure 2. These characteristics included an openness and willingness to explain and discuss their hearing loss, an assertive attitude toward their communication needs in social and community settings and the ability to maintain a sense of humor in situations in which miscommunication or misunderstandings occurred. Parents and itinerant teachers stressed the importance of encouraging the child early on to come to terms with and accept the hearing loss as a part of their identity and to promote a positive outlook. Those young people who had accepted and were comfortable with their hearing loss were perceived as having greater confidence in developing peer and social relationships. As stated in one of the focus groups held with the itinerant teachers:

Lynn: I think the basic personality of the child could be either supportive of integration or a barrier to integration. The child who has a good attitude where they want to succeed and they can problem solve and, just further to that, a personality that is likable to other children.

Recognizing and dealing with difficult situations, taking a proactive attitude, finding alternatives and developing appropriate coping strategies were discussed in particular by all groups of students with hearing loss as a means of facilitating their own inclusion into social and community activities. An example of this approach related to an employment situation is illustrated in the following excerpt:

Jennie: Part of the thing that makes you stronger is finding a way around it. Like at work, I've had a few jobs, and a big problem was using the phone, obviously. But the thing is, you just work around it, and you'll pick it and say, "Thank you for calling, please hold." That's what I would do. And then I would get somebody who could actually hear on the phone. At least you are doing something about it.

Students with outgoing personalities, good communication skills, good speech intelligibility and a willingness to use hearing technology, particularly FM systems, in social situations were viewed as being more socially successful. Enrollment in early intervention programs emphasizing the development of spoken language was perceived by all participants as having provided the basis for social and community inclusion.

Community leaders. Leaders of social activities such as hockey, swimming, soccer, football, dance, art classes, gymnastics, Girl Guides and Brownies, Boy

Scouts and other more informal community activities were perceived by all focus group participants as facilitative of integration. Behaviors and attitudes conducive to integration included a sensitivity to the needs of the individual with hearing loss, a willingness to modify activities to suit the communication needs and abilities of the student, an open attitude to the use of assistive technology as appropriate, the ability to adopt facilitative communication strategies and a willingness to consult with parents to ensure the child is included successfully in activities. Providing and facilitating opportunities for the individual with hearing loss to capitalize on their skills and talents in these areas were seen as important contributors to the development of self-confidence and self-esteem. Five of the parents became the leaders of such activities to ensure their children would be included successfully.

Itinerant teachers. Itinerant teachers contributed to social integration through their willingness to facilitate teen groups and organize activities outside of school hours where students with hearing loss were given the opportunity to socialize. Providing time during intervention to focus on the development of social skills, including peers in intervention sessions as communicative partners and playmates and providing time to allow students to discuss their social challenges during sessions were all seen as facilitative of social inclusion. The role of the itinerant teacher in contributing to peer understanding of hearing loss and sensitivity to the communication needs of students with hearing loss was discussed by all participants as fundamental to developing an awareness of hearing loss facilitative of inclusion.

Administrators. School administrators were identified in two of the four parent focus groups and two of the three groups of itinerant teachers as facilitators of social and community inclusion when they provided administrative commitment and funding for afterschool activities in which students with hearing loss were able to participate with their hearing peers. Such activities included afterschool social clubs, individual and team sports, school newsletters and year books and other activities promoted by the school administration. Particularly when such activities were facilitated by knowledgeable teachers, they were perceived as providing important opportunities for the social integration of students with hearing loss. Administrators as facilitators of integration were discussed in two of the parent and two of the itinerant teacher focus groups.

Technology. Technology, including hearing aids and FM systems in particular, contributed to community inclusion when such technology was used appropriately in social situations. Twelve of the 16 young people discussed having used an FM system at some point in their lives. A willingness to use the FM system, circulate the microphone and encourage peers to wait their turn when using the FM were particularly appreciated by the young people with hearing

Figure 3. Barriers to family and community integration.

loss in situations such as sports coaching and Guide and Scouting activities. An understanding of the limitations of hearing aids and the adoption of facilitative communication strategies in difficult listening conditions by peers and community leaders also were perceived as greatly facilitative of integration, particularly by the young people with hearing loss.

Barriers to Family, Social and Community Inclusion

A number of important barriers to inclusion in family, community and social activities also were identified by the participants. These barriers are summarized in Figure 3. The most frequently identified barriers to family and community integration again were related to variables associated with families, constituting 135/342 or 39.5% of all identified barriers. Second in order of frequency were barriers associated with characteristics of children and youth with hearing loss, which constituted 22.8% of all identified barriers. Additional barriers were distributed across the categories of peers, community leaders, administrators and professionals and physical obstacles to inclusion in the form of acoustic and technological difficulties. The

Figure 4. Contributions of focus group participants to identified barriers.

contributions of focus groupings to each of the identified variables are presented in Figure 4. Again, the majority of barriers associated with family variables were mentioned by parent participants, who identified 82/135 or 60.7% of the family barriers to inclusion. Identification of additional barriers was distributed relatively equally across all of the focus groupings with the exception of community barriers, which were mentioned rarely in the focus groups held with the itinerant teachers.

In the sections below, barriers associated with family inclusion will be presented first according to the thematic categories that were extracted from the data, followed by barriers associated with social and community inclusion.

Barriers to Family Inclusion

As was the case for the facilitators of inclusion, the majority of family-related factors inhibiting integration were associated with parental roles and

the relationships between family members. These barriers consisted of lack of supports, family stress, guilt, parental burnout and overprotectiveness, as described below.

Lack of Support. The presence of extended family in the vicinity did not guarantee that family members would understand or assist in sharing the daily challenges faced by the parents of children with hearing loss. A lack of such a supportive family network was perceived as a significant barrier by all participating parents, particularly in the early years.

Dave: I think a lot of my family members were and still are intimidated by anything that has to do with the handicap.

Mary: Oh yes, I think they were scared.

Dave: Maybe they were scared because of a lack of a program to instruct them. In our case, they were reluctant to step in. Hardly anybody stepped in.

Joan: I've had family members that have said that I was obsessed. That's their term for it. Like my brother will say, "You served Christmas dinners year after year for the whole family, but you were angry all the time." And that's because nobody was talking to my child. They didn't understand that. And the family we are close to now are the ones who took the time to speak to our daughter.

Eleven parents who participated in two separate focus groups described dealing with this lack of support or understanding by taking on all of the challenges and stresses on their own and refusing to express any need for help. Maintaining the illusion of strength and control was mentioned as an unsuccessful coping strategy, particularly by those parents who could not count on their extended family for support. As these mothers stated:

Linda: Maybe I should have voiced that need for support, too. But I had to give the façade that everything was fine, that I was in total control of everything. You know, if I had gone to my sister and said, "I need a break," I'm sure she would have been there. But I couldn't let my guard down.

Liz: It would be nice if family realized that you really do need a break every once in a while and that maybe you're not quite as comfortable leaving your hearing impaired child with the 14-year-old kid down the street or whatever. You could if your family had your child for a couple of hours.

Pam: Yes, that was a barrier, because you came across as knowing so much and being such a strong character. Like "Oh wow! I couldn't do what you're doing!" But inside you're almost crying, saying "No! Help me!" But you give this aura that everything is under control, while inside you just were screaming for a break saying, "Take him for an afternoon." But you have to appear that way, like everything is under control.

Dealing with stress, guilt and burnout. The enormous time commitments and constant involvement of parents in the development of speech and language were mentioned in all of the parent focus groups and two of the three itinerant teacher groups as potential barriers to the proper functioning of the family and factors affecting mental health. Seven mothers in particular referred to stress, exhaustion and potential burnout as factors that could impose significant barriers to family functioning, particularly during the labor-intensive preschool years. Perceptions and emotions related to becoming their child's primary speech and language teacher were described in all of the parent focus groups and in two of the three groups of itinerant teachers, especially by the mothers.

Pam: I remember CHEO saying, "Just talk to him all the time. Expose him to language." Well, that weighed pretty heavily on me. The only time I felt any relief and any sense of being my own self was when he was having his nap. Because when, like if I washed a dish I'd say, "Look! The plate is dirty! Oh! We're going to put it in the water. Oh look! There's bubbles!" And you know what? I wanted to throw that plate right through the window! And I thought, "If I don't do this, he's never going to talk." And now when I look back, I think, "So what, taking an afternoon off, I could have."

Mary: And that's after you deal with the guilt. 'Cause that guilt is, you know, there's a little baby there and you go once a week to CHEO and they give you this lesson and you go home and you do it but you think, if one day your whole heart isn't into it, you go to bed that night and you can't sleep because you're thinking, "I didn't give my 110% today on that lesson and he's never going to talk and where is he going to go to school and how is he blah blah blah." And it's not until you have time that passes that you think, "All those nights of wasted sleep, I didn't have to!"

[Others agree].

Pam: Yes, except that if I had been told, "Well, don't worry, you don't have to talk," . . . I needed that drive. But I look back now, and I

don't know how we did it sometimes, and how it really was exhausting.

Sandra: The guilt and the worry were great motivators, but it was a lot of wasted energy also.

The importance of not becoming carried away with the role of teacher at the expense of establishing and maintaining a close parent-child relationship was discussed in two of the four parent focus groups. One of the mothers expressed specific regrets with regard to this issue when looking back at her early relationship with her daughter:

Sandra: I think, too, in terms of family support, I always felt that I became Jessica's teacher as opposed to just her parent. No question. I couldn't separate the two at all from the minute we started at CHEO. And that continued right through until I'd say the end of grade ten or eleven.

Dave: You often said that you missed out being her mother. You know, that mother-daughter relationship. You were teacher and student.

Sandra: Teacher first and always. So that became a barrier.

Overprotectiveness. Overprotectiveness as a significant family obstacle to be overcome was discussed at length in all of the young-adult focus groups. Taking the bus on their own, being allowed to go out with their friends, learning to drive and having access to the family car were particularly mentioned as signs of parental overprotectiveness. As one of the student participants pointed out:

Tony: For family barriers, not exactly a barrier but . . . as you get older, like fifteen, sixteen, like hanging out. Your parents, because they were with you all the time when you were young, very supportive and everything, it's harder for them to give you your independence. They still want to know what you are doing and about your school work. Sometimes you just want to say, "Can you just get off my back!"

Parents themselves also were aware of instances of overprotectiveness on their part and recognized that they may have had different standards of independence for their children with and without hearing loss with respect to certain activities. The question of overprotectiveness was discussed in three of the four parent focus groups.

Dave: Joan wouldn't let Brian take the bus uptown when he was in grade eight or nine.

Joan: No! You let him go up one time to the Rideau Center, and I almost had a heart attack!

Dave: He knew where to go, he had to learn.

Doug: You see that's a similar problem that Mary and I had as well. I was saying, "Let her go. Let her drive the car." Mary was saying "No wait! We'll drive her." The time comes, though, when you have to stop protecting them and let them go, when you think they're ready.

Mary: But it's a hard balance. Because Doug would say, "Well don't you think it's time?" And I'd say, "Well, yes, if she could hear."

Barriers to Social and Community Inclusion

All participants spoke of situations in which they had experienced or observed a lack of support for integration in social and extracurricular activities both inside and outside of school. These barriers included negative opinions regarding the abilities and competencies of individuals with hearing loss on the part of community leaders and peers, physical barriers to participation including poor listening conditions and the limitations of existing hearing technology, communication barriers and difficult listening situations that did not permit full participation and personal characteristics inherent to individuals with hearing loss. A lack of commitment and funding to facilitate inclusion in social and sports activities within the school setting but outside of school hours contributed to the social exclusion of students with hearing loss. Students with hearing loss described having experienced periods of social isolation, loneliness and, in some cases, bullying by their peers. The middle school years encompassing grades five to seven were identified as being particularly difficult and were described by one of the young adults as a "social wasteland." The barriers described by all of the focus group participants illustrate the challenges and obstacles to the successful inclusion of children with hearing loss that continue to exist outside of the school setting.

Characteristics of individuals with hearing loss. Characteristics associated with individuals with hearing loss themselves that were perceived as having the potential to contribute to social exclusion or isolation constituted 22.8% of all identified barriers. These characteristics included reluctance of individuals to express or advocate for their own needs, shyness or a tendency toward introversion, the lack of a sense of humor and a lack of self-confidence. Parents in three of the four focus groups mentioned the reality that as their children

became older, they as parents were no longer able to initiate friendships and participate in networking to facilitate their child's participation in extracurricular activities and were perceived as meddling or interfering when they attempted to encourage their children to participate in activities or make advances towards others. On the other hand, the danger of allowing a child to become too solitary was also discussed in two of the focus groups, as illustrated by comments from one mother:

Mary: Don't let them retreat too much into their own little world, 'cause that's the easier approach for them. "Oh, I'll just stay home and read a book." I'm always after my daughter to phone people. But she always waits for other people to invite her first, and I wonder why. I'll say, "Why don't you initiate?" and she will get mad. I don't know, maybe that's the way it is with a lot of teenagers, but maybe it's not. So we're always questioning, is it the hearing impairment or is it the normal teenage stuff? And you're never quite sure which is which.

As the above interview excerpt illustrates, participating parents described particular challenges associated with dealing with their teenaged children and expressed some concerns regarding which teen behaviors were normal at this stage of development and which might be more directly attributable to the child's hearing loss. One of the parents stated that during the adolescent years, it was particularly important to "pick your battles" and not allow every difficult situation to lead to confrontation and conflict. The participating parents expressed the opinion that a significant amount of patience was required when dealing with their adolescent children.

Inappropriate communication strategies and discriminatory attitudes of community and peers. Barriers associated with negative attitudes of administrators, leaders of social activities and peers in the community toward hearing loss were perceived as having imposed significant limitations on social and community integration, despite repeated advocacy efforts on the part of parents, itinerant teachers and students. All participants discussed having had negative experiences with leaders of social activities or sports events as well as with peers who lacked knowledge of appropriate communication strategies to include the individual with hearing loss. All student participants described instances where this lack of knowledge was accompanied by an apathetic or even resistant attitude toward their inclusion. These attitudes are highlighted by an experience related by one mother:

Sandy: Well, I have to mention that Maggie came across a couple of instances that were just bizarre. She tried every children's activity in the course of ten years or so. One of the things she tried when she was six or seven was Brownies. There was a very nice pair of ladies

who were the Brownie leaders. Once we had left Maggie there with her FM, her uniform, all her doodads for the meeting, what we found was that the leaders were very open to the notion of having her there as long as it was her father or me speaking to them. But once Maggie was there alone, they made no effort at all. First of all, they told her they really didn't have time to use the FM. And of course they're volunteers, so that point was made as well. Maggie came home and told us that . . . but they wouldn't change. They just wouldn't change! In fact, I had a very friendly discussion with them, and they were absolutely recalcitrant. They would not change, and so Maggie just said to me, "Well Mom, I guess I won't be doing Brownies."

Setting limits for the individual with hearing loss in terms of participation, potential or competence was mentioned by all of the young people, in three of the four parent focus groups and two of the three itinerant teacher groups as contributing to social exclusion. These limits seem to have been informed by societal prejudices surrounding individuals with hearing loss and applied to leaders, peers and, in a few rare instances, to professionals such as itinerant teachers and guidance counselors advising the students. Luckily, in the situation described next, any negative consequences of such discriminatory perceptions were circumvented by the clear emphasis on ability as opposed to disability promoted in this student's home.

Meg: Attitude is really important, your own and your parent's attitude. Like I was saying to somebody once, I really think I would like to do this certain job, and they said, "Well, you can't do that, you are deaf!" And I made the mistake of believing that and telling myself what I wasn't going to be. And I told my mom I wanted to do this but somebody told me not to, and she said, "What are you talking about? You do whatever you want, whatever makes you happy. Because you can do it if you believe in yourself."

Physical barriers to social participation. Participants with hearing loss referred to certain specific contexts that limited their participation in social events and had the potential to impact negatively on social and community inclusion. These barriers were particularly related to technological limitations or poor acoustic listening environments. For these adolescents and young adults, an inability to use the telephone was perceived as a significant obstacle to maintaining friendships and social inclusion. At younger ages, the participants with hearing loss described having allowed parents or siblings to interact on the telephone as interpreters for them. As they became older, their desire for privacy and direct contact with their friends precluded such intermediaries. In two of the focus groups held with young people, five of the participants

mentioned that alternative technologies, most particularly email and text messaging, had become instrumental in their ability to maintain regular contact with their friends.

Certain sports and social situations were perceived to present listening conditions that were particularly difficult for many of the participants with hearing loss and tended to be avoided. These included swimming, going to bars, dances, clubs and movies. Although these activities were seen as forming part of the normal life of young people in present-day society and had a great deal of social appeal, the young people with hearing loss expressed reluctance to participate in them due to the degree of dependence in communication that they imposed. The young people with hearing loss discussed modifications to the environment that might be introduced in order to allow them greater access to these activities. These modifications included access to up-to-date, functional assistive technology to support communication in difficult listening situations. As an example, young people in two of the three focus groups mentioned the uselessness of the assistive technology provided in cinemas. This technology often requires the individuals with hearing loss to remove their hearing aids in order to use the available headsets, which do not provide sufficient amplification to allow them to follow the soundtrack of the film. Additionally, students who were able to utilize these amplification systems stated that they were often either broken or unavailable at their local cinemas.

Lack of commitment for extracurricular activities. Administrative support and commitment to opportunities for participation in social and community activities were viewed by the adult participants as fundamental to inclusion. In the words of one parent participant, in her experience, "there was absolutely no support outside of the regular classroom day for extracurricular activities, clubs or sports. Once the school day ended, that was it as far as our kids were concerned, and that's a big part of integration." The lack of such opportunities was mentioned in two of the four parent focus groups, one of the itinerant teacher groups and one of the focus groups held with young people as a serious obstacle to inclusion that placed an additional burden on parents to enroll their children in such activities outside of school, often at their own expense.

Discussion

Interactions that take place in the family and community provide numerous opportunities for facilitating the integration of children with hearing loss, yet these environments have received little attention in the literature. Facilitators and barriers to integration that fall outside of the academic domain have been the focus of little investigation to date, with the notable exceptions of research reported by Green (1990) and Schwartz (1990). The current study

used focus groups of parents, young people with hearing loss and itinerant teachers of the deaf to examine perceptions of integration and factors that served as facilitators or barriers to integration across the contexts of school, family and community. In this article, inclusion into the family and community were described using themes extracted from the focus groups and were illustrated through the voices of the participants.

Family facilitators included participation in an early intervention program such as that provided by CHEO, accessing parent support groups such as VOICE, maintaining open communication within the parental relationship and in the family, finding ways to minimize stress and having strong support from family and friends. Balancing the needs of siblings with those of the child with hearing loss was perceived as important to preventing rivalry and jealousy in the family. Family barriers included a lack of support for the couple; the need to present a façade of control to the outside world; an unwillingness to ask for help; excessive stress, guilt or burnout and overprotectiveness of the child with hearing loss. Families described having made extensive sacrifices in order that their child might obtain the best possible services at an early age, including giving up jobs and moving to Canada from another country. Dealing with the needs of adolescent children with hearing loss was perceived as particularly challenging from the perspective of the participating parents.

Although all the participants recognized the importance of social inclusion, fewer facilitators were mentioned for inclusion into social environments than for family contexts. The young adults with hearing loss all were able to relate incidents of exclusion, discrimination and social isolation, generally attributable to ignorance or apathetic attitudes toward hearing loss. All of the participants stated that there was the potential for much improvement in facilitating inclusion in the social domain. Enrollment in sports and other extracurricular activities in which children and youth could participate as equals was seen as important to the promotion of self-esteem and proper social behavior. In some cases, the parents became the leaders of social activities and the builders of social networks and friendships so that their child might be included appropriately.

These reported findings build on and add to the results reported by Green (1990) and Schwartz (1990). However, whereas other studies have emphasized the need for interpreters to facilitate inclusion in the social world (Schwartz, 1990), participants in the present study emphasized the need for sensitivity toward the individual with hearing loss and use of facilitative communication strategies as opposed to the use of intermediaries. This perspective emphasizes the importance of independence and the ability of the individual with hearing loss to participate as an equal when communication needs are accommodated. This perspective also illustrates the strength, perseverance, positive attitude and high self-esteem exemplified by the participants with hearing loss in the present study.

The importance of having a few close friends who were sensitive to the needs of the individual with hearing loss and who could serve as interpreters in difficult listening situations was mentioned by all participants. As was the case with the participants in Schwartz's (1990) research, the quality of friendships took precedence over the quantity of friends. Having contact with other individuals with hearing loss was seen as important to the development of a sense of identity as a person with a hearing loss. The lack of opportunity to interact with such individuals was perceived as one of the few negative aspects of integration into local community schools, where the student often was the only individual with hearing loss in the school.

Several of the themes that emerged from the focus groups are reflected in the essays written by the three parents and three young people with hearing loss reported by Green (1990). As is the case with the findings outlined in the present study, parent participants in Green's study also emphasized the need for parental support, high expectations regarding their children's potential and parental involvement in social as well as school activities. Contrary to the findings of the present study, however, many parents in Green's study reported that their children were never able to "fit in" with other children in social contexts and that their children had difficulty establishing friendships and making "best friends." Nevertheless, in a similar manner to the participants in this research, the students with hearing loss in Green's study spoke of experiencing rejection and social isolation both within and outside of school. The importance of participating in sports activities as a means of developing self-esteem and contributing as a social equal is a common theme across both studies, as is the continued existence of societal prejudices and ignorance regarding hearing loss as important barriers to inclusion, particularly in the community environment. As in the present study, the need to move beyond the academic setting and to facilitate inclusion in family and particularly in social contexts is reflected clearly in the statements of the six participants in Green's (1990) study.

The importance of parental involvement to facilitating integration across a variety of settings is emphasized in the findings from the 10 focus groups. This involvement includes not only advocacy and encouragement of socialization within the school environment but also extends to all of the social contexts in which these children and youth might be involved. Parents in particular spoke of the challenges they faced as facilitators of family and social integration as their children grew older and emphasized the importance of maintaining trust, open communication and the building of self esteem and independence as crucial in the adolescent years. The importance of allowing independence, "picking one's battles" and not being overprotective were seen as being particularly important during this time. The development of programs to address issues associated with hearing loss for extended family members, peers and community members also was emphasized as a potential concrete action to facilitate family and social integration.

Results of the study support current evidence for differences in the parenting roles of fathers and mothers of children with hearing loss, particularly during the labor-intensive preschool years, and provide interesting insights into the perceptions and feelings of fathers of children with hearing loss that often are not reported in the literature. The fathers who participated in the focus groups had been very involved in raising their child with hearing loss, particularly in social activities such as team sports and coaching in the later years. Comments by these fathers indicated that they had attended school meetings and parent support groups regularly and had shared in activities to promote speech and language development when their child was young. The findings indicate that fathers want to be involved in decisions and activities affecting their child's intervention and integration. Mothers, nevertheless, appear to have participated more than fathers in direct language stimulation and intervention activities, especially in the early years. This finding is similar to that reported in other research (Calderon et al., 1998; Calderon & Low, 1998; Meadow-Orlans & Sass-Lehrer, 1995; Meadow-Orlans et al., 2003) for mothers and fathers participating in an early intervention program for children with hearing loss. As was the case for findings reported by Brand & Coetzer (1994) and Meadow-Orlans (1995), the mothers who participated in the present study expressed more feelings of guilt and stress in their comments and indicated a greater risk of potential burn out than fathers.

Ideally, the principles and practices supporting inclusion in the school setting also should be extended to the child's family and wider social communities, all of which provide essential opportunities to benefit from communication and interaction in society. Thus, integration into the school is not the only environment to examine in understanding potential facilitators and barriers to inclusion. Based on the comments of participants in the current study, it appears that many challenges and obstacles to the successful inclusion of children with hearing loss exist outside of the school setting. Nevertheless, and in spite of these challenges, the participants expressed optimistic and positive attitudes toward the benefits of inclusion and the potential of individuals with hearing loss to become fully contributing members of their families, communities and wider society. It is hoped that the findings of this study and the contributions of the participants will assist in minimizing the barriers that continue to exist and will promote the inclusion of individuals with hearing loss across all domains and dimensions of society.

The findings of this study are representative of a specific sample of parents with a distinctive set of demographic characteristics and stemming from a particular geographical area. Participants in the study all had been involved in providing or receiving AVT intervention, a specific service delivery model that emphasizes the development of spoken language. Further research is needed in order to identify factors that facilitate integration related to different service delivery models, approaches to communication intervention and access to services. It is likely that socioeconomic factors and multicultural

issues also might have a significant impact on participants' perceptions and experiences of community, social and family integration.

Acknowledgments

This research was supported by grants from the Social Sciences and Humanities Research Council of Canada, Human Resources and Development Canada, and the Children's Hospital of Eastern Ontario Research Institute. We are grateful to the parents, young adults and itinerant teachers of the Ottawa-Carleton District School Board and the Ottawa-Carleton Catholic School Board who participated in this research study.

References

- Brackett, D. (1993, March). *Social/emotional and academic consequences of hearing loss*. Paper presented at the National Institutes of Health Consensus Development Conference on Early Identification of Hearing Impairment in Infants and Young Children, Washington, D.C.
- Brand, H., & Coetzer, M. (1994). Parental response to their child's hearing impairment. *Psychological Reports*, 75(3), 1363-1368.
- Cabrera, N., Tamis-LeMonda, C., Lamb, M., & Boller, K. (1999, August). Measuring father involvement in the Early Head Start Evaluation: A multidimensional conceptualization. Paper presented at the National Conference on Health Statistics, Washington, D.C. ERIC document PS029359.
- Calderon, R., Bargones, J., & Sidman, S. (1998). Characteristics of hearing families and their young deaf and hard of hearing children: Early intervention follow-up. *American Annals of the Deaf*, 143(4), 347-362.
- Calderon, R., & Low, S. (1998). Early social-emotional, language, and academic development in children with hearing loss: Families with and without fathers. *American Annals of the Deaf*, 143(3), 225-234.
- Durieux-Smith, A., Olds, J., Fitzpatrick, E., Eriks-Brophy, A., Duquette, C., Cappelli, M., & Whittingham, J. (1999). A study of factors that have facilitated the integration of adolescents with a hearing impairment: Implications for policy and practice. *Research to Action Forum Proceedings*, 68-72.
- Eriks-Brophy, A., Durieux-Smith, A., Olds, J., Fitzpatrick, E., Duquette, C., & Whittingham, J. (2006). Facilitators and barriers to the inclusion of orally educated children and youth in schools: Promoting partnerships to support inclusion. *The Volta Review*, 106, 54-88.
- Gavida-Payne, S., & Stoneman, Z. (1997). Family predictors of maternal and paternal involvement in programs for young children with disabilities. *Child Development*, 68(4), 701-717.
- Green, R. (1990). Parents and students speak. In M. Ross (Ed.), *Hearing impaired children in the mainstream* (pp. 299-313). Parkton, MD: York Press.

- Gregory, S. (1995). *Deaf children and their families*. Cambridge, UK: Cambridge University Press.
- Gresham, F. (1982). Misguided mainstreaming: The case for social skills training with handicapped children. *Exceptional Children*, 48(5), 422–432.
- Hadadian, A., & Merbler, J. (1995). Parents of infants and toddlers with special needs: Sharing views of desired services. *Infant-Toddler Intervention*, 5(2), 141–151.
- Konstantareas, M., & Homatidis, S. (1992). Mothers' and fathers' self-report of involvement with autistic, mentally delayed, and normal children. *Journal of Marriage and the Family*, 54(1), 153–165.
- Ladd, G., Munson, H., & Miller, J. (1984). Social integration of deaf adolescents in secondary-level mainstreamed programs. *Exceptional Children*, 50(5), 420–428.
- MacWhinney, B., & Snow, C. (1990). The Child Language Data Exchange System: An update. *Journal of Child Language*, 17, 457–472.
- Maxon, A., & Brackett, D. (1992). *The hearing-impaired child: Infancy through high school years*. Boston: Andover Medical Publishers.
- McWilliam, R., McGhee, M., & Tocci, L. (1998). Cultural models of African American families receiving early intervention services. Washington, DC: Office of Special Education and Rehabilitative Services. ERIC document EC306247.
- Meadow-Orlans, K. (1995). Sources of stress for mothers and fathers of deaf and hard of hearing infants. *American Annals of the Deaf*, 140(4), 352–357.
- Meadow-Orlans, K., Mertens, D., & Sass-Lehrer, M. (2003). *Parents and their deaf children: The early years*. Washington, DC: Gallaudet University Press.
- Meadow-Orlans, K. & Sass-Lehrer, M. (1995). Support services for families with children who are deaf: Challenges for professionals. *Topics in Early Childhood Special Education*, 15(3), 314–334.
- Nolan, M., & Tucker, I. (1981). *The hearing-impaired child and the family*. London: Human Horizons/Souvenir Press.
- Northcott, W. (1984). *Oral interpreting: Principles and practices*. Washington, DC: Alexander Graham Bell Association for the Deaf.
- Northcott, W. (1990). Mainstreaming: Roots and wings. In M. Ross (Ed.), *Hearing-impaired children in the mainstream* (pp. 1–25). Parkton, MD: York Publishers.
- Nowell, R., & Innes, J. (1997). Educating children who are deaf or hard of hearing: Inclusion. *ERIC Digest*, Document E557.
- Ross, M. (1990). Overview and the long view. In M. Ross (Ed.), *Hearing-impaired children in the mainstream* (pp. 315–330). Parkton, MD: York Press.
- Schwartz, S. (1990). Psycho-social aspects of mainstreaming. In M. Ross (Ed.), *Hearing-impaired children in the mainstream* (pp. 159–179). Parkton, MD: York Press.
- Turbiville, V. (1995). Fathers and family-centered early intervention. *Infants and Young Children*, 7(4), 12–19.

Outcomes of an Auditory-Verbal Program for Children With Hearing Loss: A Comparative Study With a Matched Group of Children With Normal Hearing

Dimity Dornan, A.M., Ba.Sp.Th., F.S.P.A.A., Cert. AVT®, Cp.Sp.,
Louise Hickson, B.Sp.Thy. (Hons), M.Aud., Ph.D.,
Bruce Murdoch B.Sc. (Hons), Ph.D., D.Sc., and
Todd Houston, Ph.D., CCC-SLP, Cert. AVT®

The speech and language developmental progress of children with hearing loss educated using an Auditory-Verbal approach was compared to that of a control group of children with normal hearing. The experimental group consisted of 29 children ages 2–6 years with a mean pure tone average in the better ear of 76.17 dB HL at 0.5, 1 and 2 kHz. The 29 children in the control group were matched for language age, receptive vocabulary, gender and parental education level with the children in the experimental group. A battery of standardized speech and language tests was administered to all children at the start of the study and again nine months later. Results indicated no significant differences in developmental progress between the experimental and control groups. This study is the first phase of a longitudinal study that will examine outcomes for children in an Auditory-Verbal program over four years.

Dimity Dornan, A.M., Ba.Sp.Th., F.S.P.A.A., Cert. AVT®, Cp.Sp., is managing director and founder of the Hear and Say Centre, Queensland, Australia, and a postgraduate student at the University of Queensland, Brisbane, Australia. Louise Hickson, B.Sp.Thy. (Hons), M.Aud., Ph.D., is a professor of audiology at The University of Queensland and has extensive expertise in the (re)habilitation of adults and children with hearing loss. Bruce Murdoch, B.Sc. (Hons), Ph.D., D.Sc., is a professor of speech pathology and director of the Motor Speech Research Centre at the University of Queensland in Brisbane, Australia. K. Todd Houston, Ph.D., CCC-SLP, Cert. AVT®, is an assistant professor of communicative disorders and deaf education at Utah State University and the associate director for the National Center for Hearing Assessment and Management.

Introduction

Hearing loss in children often causes delays in speech and language development (Dodd, McIntosh, & Woodhouse, 1998; Moeller, 2000), and the gap between chronological age and language age for children with significant loss often remains or widens over time (Geers, 2005; Miyamoto, Svirsky, & Robbins, 1997; McConkey, Robbins, Svirsky, & Iler-Kirk, 1997; Svirsky, 2000). Recent research and clinical outcomes indicate potential for spoken language development in children with hearing loss who use the Auditory-Verbal approach (Goldberg & Flexer, 1993, 2001; Durieux-Smith, Eriks-Brophy, Olds, Fitzpatrick, Duquette, & Whittingham, 2001; Durieux-Smith, Olds, Eriks-Brophy, Fitzpatrick, Duquette, & Capelli, 1998; Wray, Flexer, & Guthrie, 2001). This study investigated the speech and language developmental progress of 29 children who use modern hearing technology coupled with an Auditory-Verbal approach and compared their developmental progress with children with normal hearing of the same language age.

Few studies have investigated spoken language outcomes for children educated using the Auditory-Verbal approach. A recent comprehensive review of research by Eriks-Brophy (2004) concluded there was only limited evidence in favor of the Auditory-Verbal approach because of significant problems with research design. These problems included the manner of collection of research data (retrospective, anecdotal, questionnaire/expert opinion or student file evidence), selection criteria for participants and lack of a control group. In addition, only one study reviewed by Eriks-Brophy used standardized measures with normative data allowing comparison with children with normal hearing (Rhoades & Chisolm, 2000; Rhoades, 2001). The results of this study indicated that 12 months or more of global language progress occurred in a 12-month time frame for 40 children with moderate-to-profound hearing loss in an Auditory-Verbal program. The children had been in the program for one to four years. Another finding of Rhoades' research was that, on average, a group of 14 "graduates" from the Auditory-Verbal program (i.e., children who had been professionally released) was able to close the gap between chronological age and language age. Although the number of children in this study was small and a control group was not included, the results do provide positive evidence for the Auditory-Verbal approach (Eriks-Brophy, 2004).

Duncan (1999) and Duncan and Rochecouste (1999) have reported another study that did include some comparison to children with normal hearing. These authors compared the length and complexity of utterances produced by 4- and 5-year-old children with hearing loss in an Auditory-Verbal program to peers with normal hearing. Both groups were enrolled in the same integrated setting. The authors found there were fewer differences between the results for the two groups of 5-year-olds than for the two groups of 4-year-olds, indicating that the children with hearing loss may have been

moving toward narrowing the gap between chronological age and language age that typically is seen in children with hearing loss. Duncan (1999) also found that, for the same population, there were no significant differences between groups for the majority of conversational skills. However, these positive results have some limitations that must be considered, including matching for chronological age instead of language age, no matching for gender or socioeconomic status and small sample size (11 children).

Other evidence of positive outcomes for children from Auditory-Verbal programs comes from retrospective studies (Goldberg & Flexer, 1993, 2001; Durieux-Smith, Eriks-Brophy, Olds, Fitzpatrick, Duquette, & Whittingham, 2001; Durieux-Smith, Olds, Eriks-Brophy, Fitzpatrick, Duquette, & Capelli, 1998; Wray, Flexer, & Guthrie, 2001). A high percentage of children “graduating” from these programs was found to enter mainstream education and had potential for academic achievement in regular classes. Importantly, few participants in these studies received the benefits of early cochlear implantation. Several studies also report potential for literacy development for children with hearing loss educated using the Auditory-Verbal approach (Robertson & Flexer, 1993; Wray et al., 2001; Wray, Flexer, & Vaccaro, 1997). Though not providing strong empirical evidence, these studies have provided professionals with the basis for description of the possible potential of the Auditory-Verbal approach for parents of children with hearing loss. Several authors (Easterbrooks, O’Rourke, & Todd, 2000; Wu & Brown, 2004) have investigated the characteristics typical of children who succeeded in an Auditory-Verbal program and report that parental involvement and the child’s lack of other disabilities were important indicators of success.

Overall, there is some limited evidence to support positive speech and language outcomes for children with hearing loss in Auditory-Verbal programs. However, it is difficult to generalize the evidence for the potential of the Auditory-Verbal approach in developing speech and language abilities from the existing outcomes studies because of the retrospective and anecdotal nature of the research and the inclusion of a control group in only one study (Duncan, 1999; Duncan & Rochecouste, 1999). According to the Oxford Centre for Evidence-Based Medicine (May 2001), the highest rated type of clinical evidence (class 1 evidence) comes from research involving randomized controlled trials that include multiway sensitivity analysis with a minimum of 30 subjects each for the experimental and the control groups and with results indicating significant differences between groups and reliable narrow confidence intervals (Guyatt & Rennie, 2002).

Also, the experimental group would need to be enrolled at the same point in intervention and have similar points of diagnosis and amplification. It is impossible to obtain this type of evidence in the area of Auditory-Verbal outcomes research because, for ethical reasons, children cannot be randomly assigned to other levels of treatment once their parents have chosen a particular approach and because there is usually a large number of uncontrollable

variables in any chosen population (Eriks-Brophy, 2004). The second highest-ranking form of evidence is the cohort study, where the performance of an experimental group is compared to a control group. We used this approach in the present study to determine, over a nine-month period, the speech and language progress of children with hearing loss who were enrolled in an Auditory-Verbal program compared to that of children with normal hearing.

In the present study, we made the following assumptions:

- Children with hearing loss attending an Auditory-Verbal program would make significant progress over a nine-month period in language development and speech production.
- The progress made by children with hearing loss in an Auditory-Verbal program for receptive and expressive language and speech production over a nine-month period would be the same as that for a group of children with normal hearing matched for language age, receptive vocabulary, gender and socioeconomic level.

Method

The present study employed a matched-group, repeated-measures design. The two groups of children were assessed on a range of speech and language measures at the start of the study (pretest) and again nine months later (posttest).

Matching of Groups

Pairs of children were matched at the beginning of the study on the basis of language age using the Preschool Language Scale-4 (PLS-4) (Zimmerman, Steiner, & Pond, 2002) and the Clinical Evaluation of Language Fundamentals-3 (CELF-3) (Semel, Wiig, & Secord, 1995). Had chronological age been used, as it was by Duncan (1999) and Duncan and Rochecouste (1999), the children with normal hearing typically would have had a higher language level than the children with hearing loss of the same chronological age, introducing the possibility that the group of children with normal hearing might progress faster.

This study was conducted in Queensland, Australia, and, at the time, the average age of diagnosis for children with sensorineural hearing loss in Australia was over 2 years because newborn hearing screening programs were not yet in place (Wake, 2002). Thus, it was highly likely that if we had matched children by chronological age, the children with normal hearing would have had a significant language age advantage over the children with hearing loss. It is also possible that matching children for language age could have resulted in the children with hearing loss being significantly older than the children with normal hearing (Blamey et al., 2001), introducing the

potential that they might progress faster because of their superior cognitive skills. However, it was considered that the potential cognitive “advantage” afforded to the children with hearing loss who were older was likely to be offset by the delays they often experience in speech and language development.

Materials

The following standardized test measures were used in the present study.

- Preschool Language Scale-4 (PLS-4) (Zimmerman, Steiner, & Pond, 2002). The scoring ceiling used was five consecutive items incorrect. If a child obtained the highest possible score on the PLS-4, the CELF-3 was used.
- Clinical Evaluation of Language Fundamentals-3 (CELF-3) (Semel, Wiig, & Secord, 1995). Six subtests were administered only to children who achieved the top score for the PLS-4. The subtests were Sentence Structure, Word Structure, Concepts and Directions, Formulated Sentences, Word Classes and Sentence Recalling.
- Peabody Picture Vocabulary Test-3 (PPVT-3) (Dunn & Dunn, 1997). Because this test was developed in the United States, Australian alternatives for some items were used by the testers. For example, “cupboard” for “closet,” “rubbish” for “garbage,” “biscuit” for “cookie” and “jug” for “pitcher.”
- Goldman-Fristoe Test of Articulation-2 (GFTA-2) (Goldman & Fristoe, 2001). The “Sounds-in-Words” subtest was administered.

In addition to the standardized measures, a written Parent Survey developed by the researchers was administered to parents at the time of both the pre- and the posttest. This Parent Survey was designed to determine the parent’s perspective on the developmental progress related to the listening and spoken language skills of their child with hearing loss and consisted of questions that included listening behavior and comprehension of language, as well as information on the child’s emerging spoken language in social environments. A parent survey, which was intended to give qualitative information on the child’s speech and language progress, was completed by the family at home and returned.

Participants

Experimental Group

Twenty-nine children with a range of sensorineural hearing losses and amplified with either hearing aids or cochlear implants were recruited from

five regional centers of an Auditory-Verbal program in Queensland, Australia. These centers adhere to the principles of the Auditory-Verbal approach and offer a range of services including early intervention and cochlear implant programs. All children participating in this study were receiving regular audiological follow-up to ensure optimal amplification as well as weekly individual therapy in which parents were guided and coached as primary language models for their child.

Diagnostic teaching principles also were employed, and children were integrated fully into mainstream education (kindergarten at age 3 years or preschool at age 4 years) as soon as it was possible, the decision being made jointly by the parents and the clinician. Potential participants included 65 children in the early intervention program whose ages fell within the criteria for candidacy, who were accessible geographically and whose parents agreed to participate in the research. Those children included in the sample involved every child in the 2- to 6-year-old age group who fit the selection criteria because no parent refused permission for their child to participate. Selection criteria were the following:

- Pure tone average of ≥ 40 dB hearing threshold level in the better ear for four frequencies (PTA-4) at 500 Hz; 1,000 Hz; 2,000 Hz and 4,000 Hz
- Prelingual deafness (at ≤ 18 months old)
- Weekly attendance in the educational program for intensive one-to-one, parent-based Auditory-Verbal therapy for a minimum of six months
- Consistent wearing of hearing technology such as hearing aids and/or cochlear implants
- Aided hearing within the speech range or recipient of a cochlear implant
- Absence of other significant cognitive or physical disabilities reported by parents or educators
- Aged 2-6 at the first test session
- English-only language use by parents with their child

The characteristics of the children are summarized in Table 1. The participants had bilateral sensorineural hearing loss ranging from moderate to profound, with a mean PTA of 76.17 dB HL (over 3 frequencies at 0.5, 1 and 2 kHz). The pure tone average at four frequencies (0.5, 1, 2 and 4 kHz) was 74 dB HL. There is no standard measure in Australia for defining hearing loss, and clinics vary between the use of three frequencies and four frequencies. Using four frequencies is considered a better reflection of average hearing loss (Davis & Wood, 1992), and information about hearing above 2,000 Hz is very important for speech perception; to omit it from the average may underestimate the degree of impairment.

In the sample, nine children had a moderate loss, seven children had a moderate-to-severe loss and 13 children had a profound loss. All children were fitted with hearing aids and commenced intervention within three

Table 1. Experimental Group Characteristics

N	Mean Age (years)	Gender		Mean Age at Identification (months)	Mean PTA (better ear)	Type of Loss		Mean Age at CI (months)	Mean Time in AV Program (months)	Hearing Device	
		M	F			Congenital	Prelingual			HA	CI
29	3.79 (SD 1.25)	21	8	24.6	76.17 dB (SD 23.6)	27	2	27 (SD 5.8)	20	15 13	14 16

Note: PTA-4 = pure tone average at 0.5 kHz, 1 kHz, 2 kHz and 4 kHz
Abbreviations: HA, hearing aid; CI, cochlear implant; AV, Auditory-Verbal.

months of identification of hearing loss. All children with cochlear implants in this study had unilateral Cochlear Nucleus CI 24 implants¹ and used an ACE processing strategy. The average age at implantation was 27 months. All but two cochlear implant users in the study also wore a hearing aid in the contralateral ear. Both hearing devices were balanced by an Australian Hearing audiologist according to the recommendation of Ching, Psarros and Incerti (2003). Parents reported on a written survey (to be reported in a later study) that 93% of children wore their hearing aids and/or cochlear implants for 100% of their waking hours at the time of the initial assessment, and all wore them consistently at the time of the posttest nine months later.

All tests were performed in the best-aided condition. For children with cochlear implants, the child's optimally functioning MAP and speech processor program, as assessed by the child's audiologist and Auditory-Verbal therapist, was used. Both "T" and "C" levels for the child's MAP were measured behaviorally and confirmed objectively when necessary. Optimal implant performance was verified by the stability of the MAP, consistent identification by the child of the seven-sound test (i.e., the Australian adaptation of Ling's Six Sound Test, Romanik, 1990), other speech perception tests and the cochlear implant-assisted audiogram (a record of the child's cochlear implant-aided thresholds for responses at 250 Hz; 500 Hz; 1,000 Hz; 2,000 Hz and 4,000 Hz).

For the children who wore hearing aids, best-aided condition was determined by the child's audiologist and Auditory-Verbal therapist, performance on the seven-sound test, speech perception tests and the child's aided audiogram. Although the selection criteria precluded children with other significant disabilities, the experimental group included one child who had mild cerebral palsy and two children who commenced investigation for the possibility of other disorders during the course of the study and subsequently transferred to a different type of program after the nine-month test period was completed.

Control Group

Children in this group were recruited by families and staff of the Auditory-Verbal program. Selection criteria were the following:

- Hearing threshold levels within the range of 0 to 20 dB at 500 Hz; 1,000 Hz; 2,000 Hz and 4,000 Hz for both ears
- No delay in phonetic development as assessed using the GFTA-2 (Goldman & Fristoe, 2001) (Australian norms for articulation [Kilminster &

¹This is the only device available in Australia.

Laird, 1978] were used, and results within one standard deviation of the mean for age were required for inclusion.)

- Absence of significant cognitive or physical disabilities (as evidenced by case history or parent report)
- English-only language use by parents with their child

When matching the control group with the experimental group, it was difficult to achieve a complete match for each individual child for both the total language score (PLS-4 or CELF-3) and the receptive vocabulary score (PPVT-3), as the range of total language and receptive vocabulary scores was wide. However, both groups of children were matched initially for total language scores and then for receptive language. Sixty-four children with normal hearing initially were tested to ensure that matching of children with the experimental group participants was possible. The 29 children finally selected for the control group were matched with the experimental group for total language age within three months above or below their scores on the PLS-4 (Zimmerman et al., 2002) or the CELF-3 (Semel et al., 1995). They also were matched within three months above or below their scores for receptive vocabulary on the PPVT-3 (Dunn & Dunn, 1997), gender and head of household education level. All except one parent in both groups had undertaken education beyond high school, suggesting a generally high socioeconomic level in both groups. The mean age of the control group children was 2.97 years (range 14–85 months, SD = 13 months).

Procedure

Clearance for this project was sought from the medical and ethical committee of the Auditory-Verbal program and was then referred to the program board of directors, who approved the project. Ethical clearance also was obtained from the School of Health and Rehabilitation Sciences Research and Postgraduate Studies Committee, University of Queensland, Brisbane, Australia. After consent was obtained from the parents for each participant, arrangements were made to conduct the assessments. Assessments of children in the experimental group took place at the child's program center. For the control group, testing was performed either at the center, in a quiet room at the child's education setting or at the child's home.

Speech and language testing was performed by experienced, qualified speech-language pathologists. An audiologist performed all audiological assessments. Because of geographic constraints, the most convenient and available qualified staff performed the testing and, frequently, different testers assessed the children pre- and posttest. Tester reliability was not examined in the study; however, all tests were administered according to the standardized instructions in the test manuals. The language and speech tests were administered over one session if possible; however, several children required two

Table 2. Order of Presentation for Standardized Assessments

Pretest
<i>Experimental Group</i> <ul style="list-style-type: none">● PLS-4 or CELF-3● PPVT-3● GFTA-2
<i>Control Group</i> <p>Tests Used for Screening Purposes:</p> <ul style="list-style-type: none">● Pure Tone Audiometry● GFTA-2 <p>Tests Used for Matching Purposes:</p> <ul style="list-style-type: none">● PLS-4 or CELF-3● PPVT-3
Posttest
<i>Both Groups</i> <ul style="list-style-type: none">● PLS-4 or CELF-3● PPVT-3● GFTA-2

sessions because of age or attention constraints. Children were given rest breaks between assessments, and the session was discontinued if a child showed evidence of fatigue or distress. The children’s responses to the GFTA-2 were not transcribed and scored at a later date but, instead, were judged to be correct or incorrect at the time of testing.

The order of presentation of the standardized tests used is summarized in Table 2. The order of testing for the control group was different from the experimental group to (1) account first for screening and (2) establish a match with a child in the experimental group before the child was tested unnecessarily. The mean time between pretests and posttests was 9.66 months for the experimental group and 9.40 months for the control group, with the time difference being largely due to geographical access to the children with hearing loss. A Welch two-sample t test indicated that there was no significant difference between the pre- and posttest intervals for the two groups ($t = 1.095$, $df = 49.24$, $p = 0.279$).

Results

In the first instance, a preliminary analysis was carried out to ensure the validity of matching of participant groups at the pretest; that is, the matching of language age and receptive vocabulary as indicated by total language age on the PLS-4 or CELF-3 and PPVT-3 results, respectively. The experimental

group's PLS-4/CELF-3 mean age equivalent was 3.47 (SD = 1.27), and the mean for the control group was 3.39 (SD = 1.31). Between-group *t* tests showed no significant difference between these values ($t = 0.229$, $df = 56$, $p = 0.820$). Similarly, there was no significant difference between groups for the mean vocabulary age equivalents on the PPVT-3 ($t = 0.209$, $df = 56$, $p = 0.836$). The mean age equivalent for the experimental group on the PPVT-3 was 2.65 (SD = 1.25), and the mean for the control group was 2.72 (SD = 1.27).

It is significant to note that 65.5% of the children participating in this study ($n = 19$) had scores that were within the age-appropriate range for auditory comprehension on the PLS-4 at the time of the pretest. By the time of posttest nine months later, 69% ($n = 20$) were in the age-appropriate range. Similarly, for oral expression on the PLS-4, 55% ($n = 16$) of children were within the age-appropriate range at pretest and 62% ($n = 18$) fell into this range at posttest. For total language scores on the PLS-4 or CELF-3, 58.6% ($n = 17$) of children were within the age-appropriate range at pretest and 65.5% ($n = 19$) of children were within range at posttest. In addition, two children in the experimental group had scores that exceeded one standard deviation above the mean both at pre- and posttest (one child on the CELF-3 and one on the PLS-4). This result placed 65.5% ($n = 19$) of children in the normal range or above for their total language age equivalent at the time of the pretest and 72.4% ($n = 21$) at posttest.

Table 3 summarizes the age-equivalent results on all measures obtained at pre- and posttest for both groups over the nine-month test period, as well as the results of the statistical analyses. Mixed-model analysis, considered to be the most appropriate method of analyzing longitudinal data with independent observations (Brown & Prescott, 1999) was performed to check the differences between age equivalents for both groups and to investigate gain over time for both groups. Residual diagnostic plots (including Q-Q plots) were undertaken to check the assumption of normality of the residuals, necessary for this analysis. Box plots of the residuals for each of the groups indicated that the homogeneity of variance assumption was not violated.

For the PLS-4/CELF-3 total language scores, no significant interaction was found between the groups, and a significant time effect was seen between pre- and posttests for both groups ($F(1, 56) = 28.30$; $I < 0.001$). The time effect is consistent, with both groups making significant progress from pre- to posttest, and the lack of interaction is consistent with there being no difference between the rate of increase in scores for both groups. The same pattern of results was found for the auditory comprehension scores ($F(1, 56) = 176.87$; $p < 0.001$), oral expression scores ($F(1, 56) = 126.36$; $p < 0.001$) and the PPVT-3 scores ($F(1, 56) = 119.87$; $p < 0.001$). Similar calculations were performed for the GFTA-2, and, again, there was no interaction between groups; however, the time effect was significant. There was a difference between groups on this

Table 3. Summary of Changes in Age Equivalent Scores (in years) for Both Groups Over the Nine-Month Test Period and Results of Statistical Comparison of the Groups

<i>Test</i>	<i>Group</i>	<i>Pretest</i>		<i>Posttest</i>		<i>Statistical Result</i>	
		<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>t</i>	<i>p</i>
Auditory Comprehension (PLS-4/CELF-3) Group Comparison	Experimental	3.49	1.02	4.32	0.87	9.255 ^a	<0.001
	Control	3.27	1.02	4.13	1.21	9.553 ^a	<0.001
						0.211 ^a	0.834
Oral Expression (PLS-4/CELF-3) Group Comparison	Experimental	3.21	0.97	3.95	1.00	7.526 ^a	<0.001
	Control	3.20	1.18	4.03	0.89	8.372 ^a	<0.001
						0.598 ^a	0.552
Total Language (PLS-4/CELF-3) Group Comparison	Experimental	3.47	1.27	4.33	1.32	12.42 ^b	<0.001
	Control	3.39	1.31	4.20	1.38	11.56 ^b	<0.001
						-0.491 ^b	0.1625
PPVT Group Comparison	Experimental	2.65	1.25	3.80	1.70	7.684 ^b	<0.001
	Control	2.72	1.27	3.89	1.66	7.799 ^b	<0.001
						0.082 ^b	0.935
GFTA-2 Group Comparison	Experimental	2.88	1.30	3.60	1.37	6.048 ^c	<0.001
	Control	3.79	1.27	4.60	1.61	6.410 ^c	<0.001
						0.785 ^c	0.436

^a54 df

^b56 df

^c50 df

measure at pretest, with the control group having higher mean scores at the initial test session ($F_{1, 56} = 77.06$; $p < 0.001$).

Table 4 outlines the mean change in scores for both groups over the nine-month test period. The data show that the average gain was nine months or greater on all measures for the control group and on three of the measures for the experimental group (auditory comprehension, total language and PPVT-3).

Table 4. Summary of Age-Equivalent Gains in Months for Both Groups Over Nine-Month Test Period

<i>Test</i>	<i>Experimental Group Gain (Months)</i>	<i>Control Group Gain (Months)</i>
Auditory comprehension (PLS-4/CELF-3)	9.96	9.72
Oral expression (PLS-4/CELF-3)	8.88	10.32
Total language (PLS-4/CELF-3)	10.32	9.96
PPVT-3	13.80	14.04
GFTA-2	8.76	9.72

Discussion

This study examined the speech and language developmental progress of children with hearing loss educated using an Auditory-Verbal approach as compared to that of a control group of children with normal hearing. The data were collected using standardized assessments, and a control group matched for language age, receptive vocabulary, gender and head of household education level was included. Statistical analysis revealed that both groups made significant progress and that there was no significant difference between the progress made by the two groups on language and receptive vocabulary scores. The speech production results indicated a lower score at the start of the study for the children with hearing loss compared to the children with normal hearing, but the developmental progress of both groups was the same.

The results support our hypotheses that children in an Auditory-Verbal program would make significant progress in language development and speech production over a nine-month period and that this progress would be the same as for a control group of children with normal hearing. The average rate of progress of the children with hearing loss in the study ranged from 8.76 months on the GFTA-2 to 13.80 months on the PPVT-3. These findings are similar to reports by Rhoades and Chisolm (2000) and Rhoades (2001) of children in an Auditory-Verbal program. Comparisons with other rate of development studies are problematic because of the inherent heterogeneity of the population.

In addition, advances in hearing technology and cochlear implant candidacy rapidly are changing the arena of research outcomes for children with hearing loss (Geers, 2005), making it even more difficult to compare this study with earlier studies where children used less sophisticated technology. A recent study investigating development rate of language skills carried out on Australian children (Blamey et al., 2001) is the most comparable. The authors examined the rate of change in language development for a group of 40 children using hearing aids (mean PTA of 78 dB HL) and a group of 47 children with cochlear implants (mean PTA of 106 dB HL). Test measures were the PLS-4/CELF-3 and the PPVT-3. All children were enrolled in oral communication programs. The authors found that the children with cochlear implants were developing language at one-half to two-thirds the rate reported for children with normal hearing, a rate that was the same as for the aided children with a severe loss participating in the study.

The current study included 15 children with cochlear implants and 14 with hearing aids; overall, the developmental progress found was superior to that reported by Blamey et al. (2001). A possible reason for this finding is the different educational approaches in which the children were engaged. The Auditory-Verbal approach involves parents as the child's natural language teachers, sessions that are always individual and an emphasis on listening

over vision (Auditory-Verbal International, Inc., 1993). This approach is compared to an oral communication approach, in which parents are not necessarily the primary language teachers, sessions can be held with individuals or groups and the amount of emphasis on vision versus listening varies according to program. For example, the Council for Exceptional Children (Easterbrooks, 1997) reports that the auditory/oral philosophy may incorporate visual methods of teaching. Another group reports that an auditory/oral program stresses the use of speech reading and natural gestures to aid the child's communication (Beginnings, 2005).

The majority of children in the present study (72.4%) had total language scores that fell in the age-appropriate range at the posttest. We acknowledge that, as Young and Killen (2002) note, this result does not infer that the children's spoken communication skills are the same as for children with normal hearing. Rather, these scores give an indication of level of functioning on one particular scale. These results are in agreement with other studies on the Auditory-Verbal approach (Durieux-Smith et al., 1998, 2001), in which children who had graduated from an Auditory-Verbal program had communication skills that were age appropriate. The results are in marked contrast to those of Geers, Nicholas and Sedey (2003), who reported that 30% of a group of 181 children with hearing loss scored at an age-appropriate level for receptive language, and 47% were within the normal range for expressive language. All children in that study had received cochlear implants, and 89 participated in a Total Communication program while 92 used oral communication.

It is noted that both experimental and control group children in the present study exhibited significantly higher PPVT-3 scores at pre- and posttest than the general population of children with normal hearing on which the test was standardized. A possible reason for this high level of receptive vocabulary in both groups of children is the relatively high socioeconomic status of the parents (Hart & Risley, 1995; Hoff-Ginsberg, 1991). All but one parent in each group had undertaken education beyond high school. Nevertheless, the results for the children with hearing loss are promising and contrast sharply with previous studies that have reported PPVT-3 scores for children with hearing loss that are significantly lower than those of the population with normal hearing (e.g., Young & Killen, 2002; Dodd et al., 1998). One of the desired outcomes of the Auditory-Verbal approach is a high level of family involvement and increased child-directed speech (Auditory-Verbal International, Inc., 1993). In a study by Moeller (2000), positive language outcomes in children with hearing loss have been correlated with high levels of family involvement, and Hart and Risley (1995) have shown that increased child-directed speech is associated with larger vocabularies in children and faster vocabulary growth over time.

Yoshinago-Itano, Sedey, Coulter and Mehl (1998) found that children with hearing loss who are identified early and who receive intervention before age

6 months have better language skills than those children identified later. It is important to note that children in the current study did not, as a group, have the benefits of newborn hearing screening and were identified relatively late, at a mean of 24.6 months of age. As participants in an Auditory-Verbal program, however, the children with hearing loss did receive early optimal audiological management, consistently used their hearing devices and were all users of spoken communication.

Another feature of the participants in this study was that most of the children were male. The results may have been adversely affected by this large number of male children, as females are known to have a verbal advantage over males (Fenson, Pethick, Renda, Cox, Dale, & Reznick, 2000). The advantage for females also has been noted by Easterbrooks and O'Rourke (2001) when studying children in an Auditory-Verbal program.

This study addresses some of the methodological problems in existing research noted by Eriks-Brophy (2004). Several communication domains were examined, namely language reception and expression, receptive vocabulary and speech skills. The limitations of the study were that it was conducted on a sample size of 29 children over a short period of time (i.e., nine months), and the participants had the variability typical of any group of children with hearing loss. As stated earlier, the small sample size in this study is acknowledged as a limitation, and it is not appropriate to generalize the results to the population at large.

The authors plan to extend this study longitudinally and to examine additional domains such as literacy, numeracy, social and psychosocial functioning and to further investigate the variables that may influence speech and language development for children with hearing loss in an Auditory-Verbal program, such as socioeconomic status and gender. Future research also is needed to investigate the progress of children in other Auditory-Verbal programs and those using alternative educational options.

Acknowledgments

Financial support for this study has been provided by the Hear and Say Centre; School of Health and Rehabilitation Sciences, University of Queensland, Brisbane, Australia; Queensland Council of Allied Health Professionals; and the Commonwealth of Australia through the Cooperative Research Centre for Cochlear Implant and Hearing Aid Innovation (CRC Hear, Australia). The authors also wish to acknowledge the support of the staff and parents of the Hear and Say Centre, Ellen McKeering, Catherine Bow and the Bionic Ear Institute, Melody Harrison and Peter Dornan. Statistical analysis was performed by Ross Darnell of the School of Health and Rehabilitation Sciences, University of Queensland. Special thanks are due to Carol Jackson, Jane Thompson and Collette O'Brien, who have helped in manuscript preparation.

References

- Auditory-Verbal International, Inc. (1993). *AVI principles and rules of ethics*. Alexandria, VA: Author.
- Beginnings for Parents of Children Who are Deaf or Hard of Hearing, Inc. Communications Options: Reference Chart. Retrieved November 2006, http://www.beginningssvcs.com/communication_options/comm_options_chart.shtml.
- Blamey, P.J., Sarant, J., Paatsch, L., Barry, J.G., Bow, C.P., Wales, R.J., et al. (2001). Relationships among speech perception, production, language, hearing loss, and age in children with impaired hearing. *Journal of Speech, Language and Hearing Research*, 44, 264–285.
- Brown, H. & Prescott, R. (1999). *Applied Mixed Models in Medicine*. Chichester, NY: John Wiley and Sons, Ltd.
- Ching, T., Psarros, C., & Incerti, P. (2003). Management of children using cochlear implants and hearing aids. *The Volta Review*, 103, 39–57.
- Davis, A., & Wood, S. (1992). The epidemiology of childhood hearing impairment: Factors relevant to planning of services. *British Journal of Audiology*, 26(2), 77–90.
- Dodd, B., McIntosh, B., & Woodhouse, L. (1998). Early lipreading ability and speech and language development of hearing-impaired preschoolers. In R. Campbell, B. Dodd, & D. Burnham (Eds.), *Hearing by Eye II: Advances in the psychology of speechreading and auditory-visual speech* (pp. 229–242). Hove: Psychology Press.
- Duncan, J. (1999). Conversational skills of children with hearing loss and children with normal hearing in an integrated setting. *The Volta Review*, 101, 193–211.
- Duncan, J. & Rochecouste, J. (1999). Length and complexity of utterances produced by kindergarten children with impaired hearing and their hearing peers. *Australian Journal of Education of the Deaf*, 5, 63–69.
- Dunn, L.M., & Dunn, L.M. (1997). *Peabody Picture Vocabulary Test*. Circle Pines, MN: American Guidance Service.
- Durieux-Smith, A., Eriks-Brophy, A., Olds, J., Fitzpatrick, E., Duquette, C., & Whittingham, J. (2001). Facilitating the Integration of Children and Youth with Hearing Loss. *The Listener, Special Edition* (Fall 2001), 19–27.
- Durieux-Smith, A., Olds, J., Eriks-Brophy, A., Fitzpatrick, E., Duquette, C., & Capelli, M. (1998). Outcome of AVT: Results of a follow-up study. *ASHA Leader*, 3(16), 99.
- Easterbrooks, S.R. (1997). Information Center on Disabilities and Gifted Education. Educating children who are deaf or hard of hearing: Overview. Retrieved November 2006 from <http://ericec.org/digests/e549.html>.
- Easterbrooks, S.R., & O'Rourke, C.M. (2001). Gender differences in response to auditory-verbal intervention in children who are deaf or hard of hearing. *American Annals of the Deaf*, 146, 309–319.

- Easterbrooks, S.R., O'Rourke, C.M., & Todd, N.W. (2000). Child and family factors associated with deaf children's success in auditory-verbal therapy. *The American Journal of Otology*, 21, 341–344.
- Estabrooks, W. (2006). *Auditory-Verbal Therapy and Practice*. Washington, DC: Alexander Graham Bell Association for the Deaf and Hard of Hearing.
- Eriks-Brophy, A. (2004). Outcomes of Auditory-Verbal Therapy: A review of evidence and a call for action. *The Volta Review*, 104, 21–35.
- Fenson, L., Pethick, S., Renda, C., Cox, J., Dale, P., & Reznick, J. (2000). Short-form versions of the MacArthur communicative inventories. *Applied Psycholinguistics*, 21, 95–115.
- Geers, A. (2005). Spoken language in children with cochlear implants. In P.E. Spencer & M. Marschark (Eds.) *Advances in the Development of Spoken Language by Deaf Children*. New York: Oxford University Press.
- Geers, A., Nicholas, J., & Sedey, A.L. (2003). Language skills of children with early cochlear implantation. *Ear and Hearing*, 24(1), 46s–57s.
- Goldberg, D.M., & Flexer, C. (1993). Outcome survey of auditory-verbal graduates: Study of clinical efficacy. *Journal of the American Academy of Audiology*, 4, 189–200.
- Goldberg, D.M., & Flexer, C. (2001). Auditory-verbal graduates. An updated outcome survey of clinical efficacy. *Journal of American Academy of Audiology*, 12(8), 406–414.
- Goldman, R., & Fristoe, M. (2001). *Goldman-Fristoe Test of Articulation-2*. Shoreview, MN: American Guidance Service.
- Guyatt, G.H., & Rennie, D. (2002). *Users' guides to the medical literature: A manual for evidence-based clinical practice*. Chicago: AMA Press.
- Hart, B. & Risley, T.R. (1995). *Meaningful differences in the everyday experience of young American children*. Baltimore: Paul H. Brooks.
- Hoff-Ginsberg, E. (1991). Mother-child conversation in different social classes and communicative settings. *Child Development*, 62, 782–796.
- Kilminster, M.G.E., & Laird, E.M. (1978). Articulation development in children aged three to nine years. *Australian Journal of Human Communication Disorders*, 6(1), 23–30.
- McConkey Robbins, A.M., Svirsky, M. & Kirk, K.I. (1997). Children with implants can speak, but can they communicate? *Otolaryngology, Head and Neck Surgery*, 117, 155–160.
- Miyamoto, R.T., Svirsky, M.A. & Robbins, A.M. (1997). Enhancement of expressive language in prelingually deaf children with cochlear implants. *Acta Oto-Laryngologica (Stockholm)*, 117, 154–157.
- Moeller, M.P. (2000). Early intervention and language development in children who are deaf and hard of hearing. *Pediatrics*, 106(3), 43.
- Oxford Centre for Evidence-based Medicine (May, 2001). Levels of evidence and grades of recommendation. Retrieved May 2003, from http://www.cebm.net/levels_of_evidence.asp.

- Rhoades, E. (2001). Language progress with an auditory-verbal approach for young children with hearing loss. *International Pediatrics*, 16(1), 1–7.
- Rhoades, E.A., & Chisolm, T.H. (2000). Global language progress with an auditory-verbal approach. *The Volta Review*, 102, 5–25.
- Robertson, D., & Flexer, C. (1993). Reading development: A parent survey of children with hearing impairment who developed speech and hearing through the auditory-verbal method. *The Volta Review*, 95, 253–261.
- Romanik, S. (1990). *Auditory-Skills Program: For students with hearing impairment*. Sydney: New South Wales Department of Education Special Education and Focus Programmes Division.
- Semel, E., Wiig, E.H., & Secord, W.A. (1995). *CELF: Clinical Evaluation of Language Fundamentals* (3rd ed.). San Antonio: The Psychological Corporation, Harcourt Brace & Company.
- Svirsky, M.A. (2000). Language development in children with profound and prelingual hearing loss without cochlear implants. *The Annals of Otolaryngology, Rhinology and Laryngology*, 109 (12, Supplement 2), 99–102.
- Wake, M. (2002). Newborn hearing screening: Decision time for Australia. *Medical Journal of Australia*, 177, 172–173.
- Wray, D., Flexer, C., & Guthrie, B. (2001). Classroom performance of auditory-verbal students: A comparative update. Paper presented at American Speech-Language-Hearing Association National Conference. November 2000. Washington, D.C.
- Wray, D., Flexer, C., & Vaccaro, V. (1997). Classroom performance of children who are deaf or hard of hearing and who learned spoken language through the Auditory-Verbal approach: An evaluation of treatment efficacy. *The Volta Review*, 99, 107–119.
- Wu, C.D., & Brown, P.M. (2004). Parents' and teachers' expectations of Auditory-Verbal Therapy. *The Volta Review*, 104, 5–20.
- Yoshinaga-Itano, C., Sedey, A.L., Coulter, D.K., & Mehl, A.L. (1998). Language of early and later identified children with hearing loss. *Pediatrics*, 102(5), 1161–1171.
- Young, G.A., & Killen, D.H. (2002). Receptive and expressive language skills of children with five years of experience using a cochlear implant. *The Annals of Otolaryngology, Rhinology and Laryngology*, 111, 802–810.
- Zimmerman, I.L., Steiner, V.G., & Pond, R.E. (2002). *Preschool Language Scale-4*. San Antonio: The Psychological Corporation.

An Analysis of Phonological Process Use in Young Children With Cochlear Implants

Helen C. Buhler, Ph.D., Betty DeThomasis, M.S., Pat Chute, Ed.D., and Anne DeCora, M.S.

Phonological process use was investigated in five children who used Nucleus 24 cochlear implants (CIs). All participants were <3 years of age at the time of cochlear implantation and ranged from 4;2 to 4;7 years of age at onset of study. Speech samples obtained from the GFTA-2 were analyzed using the KLPA-2 to evaluate participants' phonological systems. Results indicated that four out of five participants used phonological processes for more than 40% of possible occurrences. Data suggest that participants exhibited atypical use of phonological processes relative to their chronological age but these processes were age appropriate considering participants' hearing age. Findings imply that intricate interactions exist between cochlear implantation and phonological development. Findings may be important to clinicians considering selection of assessment tools with the CI population.

Introduction

Over the past two decades, the benefits of cochlear implant use on speech perception and production have been well documented (Blamey et al., 2001a; Chin, 2003; Chin & Kaiser, 2000; Dawson et al., 1995; Kirk & Hill-Brown, 1985). The principal aim of cochlear implant (CI) use is to provide individuals with improved perception of speech and nonspeech sounds in order to enhance the overall communicative abilities of individuals with hearing loss (Chin & Pisoni, 2000).

Helen Buhler, Ph.D., is associate professor and program director of the communication disorders program at Mercy College, Dobbs Ferry, N.Y. Betty DeThomasis, M.S., is a speech-language pathologist who works with children who have hearing loss ages birth to 3 and in the school system in Phoenix, Ariz. Pat Chute, Ed.D., is division chair of the health professions and professor of the department of communication disorders at Mercy College in Dobbs Ferry, N.Y. Anne DeCora, M.S. is a speech-language pathologist working with school-aged children in the Douglas Grafflin Elementary School in Chappaqua, N.Y.

There is well-researched affirmation that CI use in children with severe-to-profound hearing loss results in an enhancement of speech perception skills, and additional research supports its benefits on speech production (Blamey et al., 2001a; Chin & Kaiser, 2000; Dawson et al., 1995; Kirk & Hill-Brown, 1985; Kirk, Miyamoto, Ying, Perdew, & Zuganelis, 2002), speech intelligibility (Blamey et al., 2001b; Dawson et al., 1995; Kirk & Hill-Brown, 1985; Osberger, Maso, & Sam, 1993; Peng, Spencer, & Tomblin, 2004), phonetic inventory development and articulation (Blamey, Barry, & Jacq, 2001; Chin & Kaiser, 2000; Dawson et al., 1995; Kirk & Hill-Brown, 1985; Serry & Blamey, 1999; Tobey, Pancamo, Staller, Brimacombe, & Beiter, 1991) and phonological development (Chin, 2002, 2003; Chin & Finnegan, 2000; Chin & Pisoni, 2000). Ultimately, research has indicated that improvement in the speech comprehension of children with CIs positively influences their ability to organize and monitor their own speech production and to develop a more sophisticated and adult-like phonological system (Tye-Murray, Spencer, & Woodworth, 1995; Waltzman & Cohen, 1998). Although there is evidence that CI use in young children may positively influence their speech perception and production (Colletti et al., 2005; Horn, Davis, Pisoni, & Miyamoto, 2005; Tomblin, Barker, Spencer Zhang, & Gantz, 2005), many children who receive CIs as early as 2 years of age experience delays in speech development. Causal factors may include the absence of auditory and linguistic input during their first two years (Chin, 2003). Researchers have found that children who receive CIs before age 2 have the potential benefit of language stimulation during the most rapid period of speech and language development (Kirk et al., 2002; Waltzman & Cohen, 1998).

Other researchers have examined speech intelligibility measures of CI recipients (Chin, Finnegan, & Chung, 2001; Chin, Tsai, & Gao, 2003; Miyamoto, Kirk, Robbins, Todd, & Riley, 1994; Peng et al., 2004; Svirskey, Robbins, Kirk, Pisoni, & Miyamoto, 2000). Tobey et al. (2003) found that the speech intelligibility of 181 children ranging in age from 8 years to 9 years with an average post-implantation age of 5.5 was 64%. Although there was variability in the participants' overall intelligibility, hearing duration was a positive factor. Similarly, Chin et al. (2003) reported that children with CIs as a group generally are less intelligible than their hearing peers; however, intelligibility increases with age and longer CI use. In a related study, Peng et al., (2004) found that three unfamiliar listeners were able to understand approximately 70% of tested sets of utterances produced by CI users with 7 years postimplant experience. Their scores were higher than previous reports. However, the authors suggest that although the increase in CI users' speech intelligibility is reflective of longer device experience, many other researchers report considerable individual variability in CI recipients' spoken language development. Contributing variability factors include age at receipt of CI, onset of deafness, type and duration of device use, motivation, type of linguistic input, amount of speech language therapy/auditory training, hearing loss

etiology, number of electrodes inserted into the cochlea, mode of communication and cognitive processing skills (Kirk et al., 2002). Tomblin et al. (2005) studied children who received CIs between ages 10 months and 40 months and found more rapid growth curves for those who received CIs as infants. Age at initial stimulation accounted for almost 15% of the variance.

According to Chin (2002), a limited number of studies have investigated specific information regarding the overall linguistic and phonological systems of children who use cochlear implants. Those researchers who have assessed the speech production of children with implants have examined the changes in percentage of correct target phonemes (Serry & Blamey, 1999), syllable structure (Chin & Pisoni, 2000) and substitution/omission patterns (Chin & Pisoni, 2000; Tye-Murray et al., 1995). The Test of Articulation Competence (Fisher & Logemann, 1971) was used to examine the articulation skills of 10 children with CIs in a study by Dawson et al. (1995). Approximately half of the participants improved their articulation skills over time. However, there were no indications as to whether their participants' improvements reflected age-appropriate skills, and their phonological skills were not assessed.

Stampe's (1969) natural theory of phonological development is based on the premise that typically developing children attempt to reproduce the adult's model or form of speech, but because of limitations in their motor and/or perceptual abilities, the productions are altered or simplified. These changes are referred to as phonological processes. Developing a more mature phonological system involves the gradual suppression or elimination of these phonological processes that do not occur in the adult language to which the child is exposed (Bankson & Bernthal, 2004; Bauman-Waengler, 2004). All children universally use phonological processes as they develop speech; however, the majority of them are typically suppressed by 5 years of age (Grunwell, 1987).

When assessing the articulation and phonological skills of CI recipients, investigators have employed various measures, including those designed for children with hearing loss, with normal hearing and with CIs. However, as the articulation and phonological development of CI recipients continue to improve, it may become more fitting to use standardized tests designed for hearing children to assess articulation and phonological competence for CI users (Chin & Kaiser, 2002). Furthermore, Chin and Kaiser assert that as additional CI recipients are mainstreamed into typical classroom settings, school-based clinicians may want to utilize ready-made and easily accessible diagnostic tools to assess articulation and phonological skills. Standardized assessments are norm referenced and allow for comparisons to typically developing children. These components often are necessary to justify the need for speech and language services in the schools. Chin and Finnegan (2000) used a phonology probe for the generative analysis of functionally misarticulated phonological systems. The probe consists of 107 words with all English

consonants except /3/ in three-word positions. The probe contains a picture-naming task, and there are between two and 19 opportunities to produce a given consonant in one of the positions to assess the use of consonant clusters. All 12 participants in their study demonstrated the ability to produce approximately 50% of the consonant clusters elicited. However, there was no indication of how different this percentage would be when the group was compared to typically developing peers.

Some researchers have examined the phonological systems of children with CIs within terms of the mature phonological sound patterns they are able to use. Serry and Blamey (1999) discussed the ability of their participants to produce stops, glides, liquids, fricatives, affricates and nasals after receiving a CI. However, when particular types of consonants were not produced correctly, the authors did not test for the presence of phonological processes to see if the errors included whole classes of sounds as opposed to sound-by-sound errors. Tye-Murray et al. (1995) examined the phonological error patterns used by their CI participants; however, their analysis was limited to errors of voicing and stopping of fricatives.

Information about the level of phonological functioning of children with CIs is essential, as it has the potential to aid in the remediation of speech production errors that are responsible for reduced speech intelligibility and overall communicative competence. The Goldman-Fristoe Test of Articulation-2 (GFTA-2) (Goldman & Fristoe, 2000) and the Khan-Lewis Phonological Assessment-2 (KLPA-2) (Khan & Lewis, 2002) are popular diagnostic tools used by school clinicians to assess the articulation and phonology skills of young children. The KLPA-2 was developed for use in conjunction with the GFTA-2 for phonological process analysis and is a standardized measurement. In a study by Chin and Kaiser (2000), the Sounds in Words subtest of the Goldman-Fristoe Test of Articulation (GFTA) (Goldman & Fristoe, 1986) was administered to 20 children to examine their articulation development. Nine of 20 participants scored below the first percentile on the GFTA when compared to children their own age with normal hearing. The percentile ranks for all the participants ranged from below the first percentile to the 33rd percentile. Relative to typically developing children with normal hearing of similar chronological age, their articulation skills were not considered age appropriate. The authors selected the GFTA because it allows for interpretation of the number of errors, standard scores and percentile ranks for children age 2 and older.

Although the GFTA-2 has been used in research with pediatric CI users to assess articulation and phonological skills, there has been little or no reported research on the outcomes of the KLPA-2. The KLPA-2 is a standardized measure that analyzes speech for up to 10 typically developing phonological processes, including reduction, place and/or manner and voicing processes. Children exhibiting a phonological delay are generally older but are expected

to acquire phonology at a developmental level comparable to younger children (Ingram, 2002). The KLPA-2 authors do not include nondevelopmental processes in their scoring system (Khan & Lewis, 2002), but they are included in the test for qualitative purposes. Nondevelopmental patterns are rarely used by typically developing children, and these patterns tend to reduce speech intelligibility more than developmental processes (Edwards, 1983; Grunwell, 1987; Khan & Lewis, 1984; Stoel-Gammon & Dunn, 1985). Adults as a whole generally are not as familiar with decoding such idiosyncratic patterns, and intelligibility is more compromised as a result (Smit, 2004). Examples of idiosyncratic processes might include initial consonant deletion, epenthesis, fricatives substituted for stops and glottal replacement (Stoel-Gammon & Dunn, 1985; Smit & Hand, 1997). A child who persists in using nondevelopmental processes for a long period of time is indicative of one who is either unaware of his unintelligibility or who is not yet capable of changing his speech productions. Thus, intervention is most likely warranted, and nondevelopmental processes generally are recommended to be targeted first (Smit, 2004). Such information can be helpful to speech-language pathologists (SLPs) who are concerned about what to target first in the intervention of CI users and for tracking clients' progress over time.

The KLPA-2 was developed for use in conjunction with the GFTA-2 for phonological process analysis and is a standardized measurement that Chin (2003) asserts is a combination of both independent and relational approaches. Stoel-Gammon and Dunn (1985) consider the child's phonological system as both an independent and relational system. Although both approaches can be used separately, they recommend that assessment of a child's system should have both methods. An independent analysis provides a description of a child's own individual system (self-contained), whereas a relational analysis provides a comparison to that of an adult standard (Dyson, 1988). The present study combines independent and relational approaches. The phonetic inventory section of the KLPA-2 is an independent analysis based on an organized list of the types of speech sounds the child produces, whether they are used correctly or not. This information can help determine which structures the child is missing compared to typically developing peers. The relational analysis consists of two types: segmental (obtained from GFTA-2 results) and phonological process evaluation (obtained from KLPA-2 results) (Smit, 2004).

The purpose of this study was to examine the overall phonological system of children who use CIs as measured by a standardized assessment, the KLPA-2. This study investigated whether children who received CIs prior to age 3 years suppressed phonological processes in a manner similar to typically developing children. The hypotheses of this study were that (1) children who use CIs will demonstrate a greater use of phonological processes as compared to the normative data of typically developing children of comparable

age with normal hearing and (2) children with longer durations of hearing as accessed through a CI will exhibit fewer phonological processes than children with shorter durations of hearing. Such information can inform school clinicians who need to make diagnostic and therapeutic decisions when working with the pediatric CI population.

Methods

Participants

Five children (three females, two males) who received the Nucleus 24 cochlear implant system and were recruited from cochlear implant facilities and schools participated in this study. Participant age at the time of data collection averaged 55 months ($SD = 2.387$). The hearing age (chronological age – age at cochlear implant activation) averaged 28 months ($SD = 8.672$). Because a small number of children participated in this study, the authors present individual factors gained from parent questionnaires that may have influenced phonological process use.

Participant 1 had his hearing loss identified at birth. He was the second child in the family to have a diagnosis of hearing loss; therefore, his parents had prior knowledge regarding amplification and early intervention services. He received amplification and began Auditory-Verbal therapy at age 4 months and continues to receive these services. In addition, he also has been enrolled in a mainstream preschool program.

Participant 2 had her hearing loss identified at age 1 year and did not begin receiving speech and audiological intervention until age 2. At the time of the study, she was receiving speech therapy daily in her auditory/oral program as well as two additional times a week at a separate location.

Participant 3's hearing loss was identified at birth, and she received amplification at age 6 months. At that time, she began receiving speech intervention and Auditory-Verbal therapy daily. She attended an auditory/oral program and a mainstream daycare from age 1 year and resides with her parents, who are both profoundly deaf.

Participant 4 was referred for further audiological assessment at birth; however, she did not receive a diagnosis of hearing loss until approximately age 10 months. She began receiving early intervention services at age 1 year, 6 months and currently receives Auditory-Verbal therapy on a daily basis at her auditory/oral school and in the home with her mother.

Participant 5 had his hearing loss identified at birth and began receiving early intervention services soon thereafter. Participants 1–4 received CIs between ages 2 years, 3 months and 2 years, 9 months, whereas participant 5 received a CI at age 12 months and was activated at 13 months. Additionally, participants 1, 2, 4 and 5 communicated primarily with spoken language and were not exposed to any formal manual language. Participant 3 communicated equally with spoken and manual (American Sign Language) languages.

Table 1. Participant Demographics

	<i>Gender</i>	<i>Chronological Age (Months)</i>	<i>Hearing Age (Months)</i>	<i>Cochlear Implant Device</i>	<i>Method of Communication</i>
Participant 1	M	56	26	Nucleus 24	Oral
Participant 2	F	51	22	Nucleus 24	Oral
Participant 3	F	56	28	Nucleus 24	Total communication
Participant 4	F	57	22	Nucleus 24	Oral
Participant 5	M	56	43	Nucleus 24	Oral
		Mean: 55	Mean: 28		

According to Yoshinago-Itano et al. (1998), children who are identified early, fitted with hearing aids and receive early intervention before age 6 months perform better than those who are fitted later. For this study, all participants used their CIs during all waking hours and, as per parent report, had no cognitive or learning disabilities (see Table 1).

Materials

The GFTA-2 (Goldman & Fristoe, 2000) was administered to the participants in one of three settings: a college speech and hearing center, a preschool or at home. Participant responses to the GFTA-2 were recorded and transcribed using the International Phonetic Alphabet (IPA, 1999). These transcriptions were used to complete the KLPA-2 (Khan & Lewis, 2002).

Procedures

All children were tested in a designated quiet area by one of two graduate students who were enrolled in the graduate program in communication disorders at Mercy College. Both students were experienced with each of the tests. Prior to the administration of the GFTA-2, each participant's basic auditory ability when using a CI was assessed with the Ling Six Sound Test (Ling, 1998). After all data were collected, each investigator independently transcribed all responses phonetically. Inter-rater agreement was between 90% and 95% for both tests. Errors were calculated and raw scores, standard scores, percentiles and test-age equivalents were determined for each participant. A phonological analysis was conducted using the KLPA-2. From this analysis, the percent of occurrence for phonological processes was determined for each participant. The 10 processes were divided into three separate categories: reduction, place and/or manner and voicing processes. Percentage of occurrence of phonological processes exhibited by each participant on the KLPA-2 was determined by dividing the number of occurrences of each

Table 2. GFTA-2 and KLPA-2 Standard Scores for Each Participant

	<i>Participant 1</i>	<i>Participant 2</i>	<i>Participant 3</i>	<i>Participant 4</i>	<i>Participant 5</i>
GFTA-2 Standard Score	79	65	73	56	100
KLPA-2 Standard Score	95	63	67	47	99

process exhibited by the participant by the total number of possible occurrences that differ for each process (Khan & Lewis, 2002). In the event that a participant used a single process more than once in a given word, the process was scored only once.

Results

Scores on GFTA-2 and the KLPA-2 were calculated using the participants’ chronological age (CA) at time of testing. Individual standard scores on the GFTA-2 and KLPA-2 for each participant are presented in Table 2. Graphic representations of these descriptive statistics are presented in Figure 1.

Figure 1 shows a scatter plot for the standard scores attained by each participant on the GFTA-2 and KLPA-2. A standard score of 100 on the GFTA-2 is average; however, four of the five participants attained standard scores below 80, more than 1.3 SD below the mean. The highest standard score of 100 (attained by participant 5) on the GFTA-2 indicates average performance. It is apparent that there is variability between participants’ GFTA-2 standard scores, indicating a difference in articulation skills.

Figure 1. Scatter plot representing GFTA-2 and KLPA-2 standard scores for each participant.

It should be noted that all GFTA-2 and KLPA-2 values were derived in relation to the performance of chronologically age-matched peers included in the standardization sample. Hearing age (HA) was not considered as a criterion for measuring articulation skills and may have contributed to the low values attained by the majority of participants. Table 1 indicates that participant 5 had the narrowest gap between CA and HA, which may have positively influenced his articulation abilities as reflected by average GFTA-2 values.

Although the average standard score on the KLPA-2 is 100, participants 2, 3 and 4 attained standard scores at least 2 SD below the mean. However, scores for participants 1 and 5 fell within the average range (95 and 99, respectively). The proximity of HA to CA for participant 5 is likely a contributing factor to his near age-appropriate phonological skills. However, participant 1, who had a 30-month gap between CA and HA, also presented with near age-appropriate phonological skills. Therefore, factors other than increased HA contributed to participant 1's phonological abilities.

Overall, there is great variability among all of the KLPA-2 values. The data reveal that two of the participants (1 and 5) attained scores that were consistently higher than those of the other three participants (2, 3 and 4). However, an overall analysis of the data also indicates that the participants on average attained scores on the GFTA-2 and the KLPA-2 that were lower than scores of the normative sample population. These results suggest that the participants of this study have less sophisticated articulation and phonological skills than those of their typically developing peers. However, if articulation and phonological skills had been examined with respect to HA rather than CA, all participants of this study may have revealed more age appropriate or average abilities.

Phonological Processes

Percentage of occurrence of phonological processes exhibited by each participant on the KLPA-2 is presented in Table 3. Of the 10 phonological processes (error patterns) analyzed by the KLPA-2, participant 1 used the fewest number of processes (six), followed by participants 3 and 5 who used eight processes and participants 2 and 4 who used nine processes. Further analysis of the data revealed that the overall percentage use of phonological processes was higher for participants 2, 3 and 4 than it was for participants 1 and 5.

Although all of the participants exhibited the use of phonological error patterns, some of these patterns did not meet the criteria for remediation. Most children vary in their use of processes; therefore, it is useful to clinicians to have a criterion to determine when a child uses a process productively. As a result, Hodsden and Paden (1991) recommend that 40% or more usage of a process warrants intervention for that process. Additionally, phonological process use may be considered appropriate if a child uses it before the typical age of suppression (the age at which 90% or fewer of children in the normative sample continued to use a process) (Smit & Hand, 1997).

Table 3. KIPA-2 Comprehensive Analysis of Developmental Phonological Processes and Percentage of Occurrence for Each Participant

	<i>Participant 1</i>	<i>Participant 2</i>	<i>Participant 3</i>	<i>Participant 4</i>	<i>Participant 5</i>
Reduction Processes					
Deletion of final consonants	2.27%	34.09%	15.91%	56.82%	6.82%
Syllable reduction	0%	7.69%	0%	7.69%	0%
Stopping	6.45%	32.26%	48.39%	19.36%	3.23%
Cluster simplification	3.85%	30.77%	42.31%	76.92%	15.39%
Liquid simplification	41.94%	54.84%	12.90%	29.03%	16.13%
Place and Manner Processes					
Velar fronting	5.26%	10.53%	15.79%	26.32%	15.79%
Palatal fronting	0%	33.33%	66.67%	0%	11.11%
Deaffrication	0%	0%	16.67%	50%	16.67%
Voicing Processes					
Initial voicing	0%	11.54%	11.54%	11.54%	3.85%
Final devoicing	18.75%	15.63%	0%	3.13%	0%

Table 4 shows each participant's CA and HA, as well as their productive phonological process use (40% or more of the time), the percentage of occurrence and the typical age of suppression. All participants, with the exception of participant 5, used at least one phonological process in excess of 40% of possible occurrences. Of these processes, only participants 3 and 4 used ones that were not developmentally appropriate for their CAs. Participant 3 (CA = 56 months, HA = 28) used the process of cluster simplification in 42% of possible occurrences. Participant 4 (CA = 57 months, HA = 22 months) used deletion of final consonants (57%), cluster simplification (77%) and deaffrication (50%). However, it should be noted that all phonological processes used by participants in this study were considered age appropriate relative to HA.

Pearson product-moment correlation coefficients were calculated to investigate possible relationships between the variables of CA, HA, standard scores, percentile ranks and age equivalents on the GFTA-2 and KLPA-2, and phonological processes used by participants. Significant correlations for the variables of this study can be seen in Table 5.

Correlation coefficients and p values showed that test measures obtained from the GFTA-2 and KLPA-2 are all highly correlated with each other. These correlations were anticipated because all standard scores, percentiles and age equivalents were obtained from the same standardized assessment tools. There are also high correlations between the GFTA-2 and the KLPA-2, which may be a result of using the same speech samples for both the GFTA-2 and the KLPA-2 analyses.

HA was highly correlated with GFTA-2 standard score ($r = 0.939$, $p < 0.018$), GFTA-2 percentile ($r = 0.972$, $p < 0.005$) and GFTA-2 age equivalency ($r = 0.975$, $p < 0.005$). The relationship between duration of hearing as accessed through a CI and performance on the GFTA-2 revealed that as the duration of CI use increased, the articulation measures attained on the GFTA-2 also increased.

There was a high negative correlation between the standard scores on the KLPA-2 and deletion of final consonant ($r = -0.919$, $p < 0.018$), cluster simplification ($r = -0.912$, $p < 0.031$) and initial voicing ($r = -0.893$, $p < 0.041$). This inverse relationship showed that the higher a participant's KLPA-2 standard scores, the lower his/her percentage of use of final consonant deletion, cluster simplification and initial voicing. Also, the KLPA-2 percentile measures are highly negatively correlated with initial voicing ($r = -0.908$, $p < 0.033$), and KLPA-2 age equivalency was negatively correlated with initial voicing ($r = -0.925$, $p < 0.024$). These inverse relationships imply that the lower the participant's percentage use of initial voicing, the higher the participant's standard scores, percentile ranks and age equivalent scores on the KLPA-2. An overall analysis of these correlations revealed a direct relationship between a decrease in phonological process use and an increase in phonological skill development.

Table 4. Participant Use of Phonological Processes (Occurring at Least 40% of the Time) Along With the Typical Age of Process Suppression

	<i>Chronological Age (yrs; mo.)</i>	<i>Hearing Age (yrs; mo.)</i>	<i>Phonological Process</i>	<i>Percentage of Use</i>	<i>Typical Age of Suppression (yrs; mo.)</i>
Participant 1	4;8 yrs. (56 mo.)	2;2 yrs. (26 mo.)	Liquid simplification	41.935%	5;0+ yrs. (Grunwell, 1987)
Participant 2	4;3 yrs. (51 mo.)	1;10 yrs. (22 mo.)	Liquid simplification	54.839%	5;0+ yrs. (Grunwell, 1987)
Participant 3	4;8 yrs. (56 mo.)	2;4 yrs. (28 mo.)	Stopping of fricatives and affricates	48.387%	2;6–6;0+ yrs. (Grunwell, 1987)
			Cluster simplification	42.308%	3;9 yrs. (Grunwell, 1987)
			Palatal fronting	66.667%	5;0 yrs. (Smit, 1993b)
Participant 4	4;9 yrs. (57 mo.)	1;10 yrs. (22 mo.)	Deletion of final consonants	56.818%	3;3 yrs. (Grunwell, 1987)
			Cluster simplification	76.923%	3;9 yrs. (Grunwell, 1987)
			Deaffrication	50%	4;0 yrs. (Lowe, 1996)

Note. Chart shows phonological process use (over 40%). Participant 5 process use was below 40% and is not included in this table.

Table 5. Pearson Product-Moment Correlation Coefficients for Variables of Significance

<i>Variable</i>	<i>Correlation (r)</i>	<i>Significance (p)</i>
KLPA-2 Standard Score		
Deletion of Final Consonants	-.919*	.027
Cluster Simplification	-.912*	.031
Initial Voicing	-.893*	.041
KLPA-2 Percentile		
Initial Voicing	-.908*	.033
KLPA-2 Age Equivalent		
Initial Voicing	-.925*	.024
GFTA-2 Standard Score		
Hearing Age	.939*	.018
GFTA-2 Percentile		
Hearing Age	.972**	.005
GFTA-2 Age Equivalent		
Hearing Age	.975**	.005

*Correlation is significant at the 0.05 level.

**Correlation is significant at the 0.01 level.

Discussion

Results of the present investigation support the hypothesis that children who use CIs demonstrate a greater use of phonological processes as compared to the normative data of typically developing children of comparable age. Results from the KLPA-2 revealed very low scores for three of five participants, indicating that they possessed less sophisticated phonological skills than their typically developing peers. Additionally, these scores affirmed that these participants presented with a moderate to severe phonological disorder. Conversely, two participants (1 and 5) attained standard scores within the average range, indicating more typical phonological development.

Further analysis of KLPA-2 results provided insight into the factors that may have contributed to the difference in phonological skills among the participants. Results of standardized phonological analysis indicated that in terms of CA, four of five participants suppressed phonological processes at a rate similar to that of their typically developing peers. Although the participants did not exhibit phonological process use in excess of 40% of possible occurrences, they did demonstrate use of multiple phonological processes at lower percentage levels (e.g., 32.26% or 26.32%). The lower percentage levels indicate an active suppression of processes, which is normal in children's development. Moreover, analysis of phonological process use in terms of participants' HA revealed that all error patterns exhibited were developmentally appropriate.

This study provides converging evidence that children with CIs use some of the same immature speech patterns as typically developing children, but they may also use atypical speech patterns. Nondevelopmental phonological error patterns are not formally analyzed nor included in the KLPA-2's standard scores and percentile ranks; however, they may be recorded during assessment. Further analysis of the KLPA-2 speech samples revealed that all five participants used nondevelopmental phonological error patterns (e.g., initial consonant deletion: /cup/→[[^]p], backing: /tub/→[k[^]b] and affrication: /tin/→[tʃɪn]). These findings support Chin's (2003) results that voiced affricates were replaced with voiceless affricates, and velar stops were missing from several children's inventories. Furthermore, findings by Chin (2002, 2003) are supported because our participants' nondevelopmental process use differed from that of their ambient language and from one another in terms of number and type of errors. Such nondevelopmental patterns have a potential to increase the negative effects on children's speech by changing syllable structure (Bauman-Waengler, 2004) and increasing unintelligibility, as listeners are unfamiliar with these types of speech patterns (Smit, 2004). Information on atypical process use can be helpful to SLPs who are concerned about what to target first in intervention.

The use of multiple phonological processes in children with CIs does not indicate definitively a need for remediation, particularly when phonological processes are not productive (less than 40% of the time for all possible occurrences) (Hodson & Paden, 1991). Although the participants in this study may continue to suppress developmental phonological processes spontaneously, periodic monitoring by clinicians would be prudent. Conversely, nondevelopmental phonological processes are not likely to be suppressed independently and therefore, therapy is recommended with these unusual processes as first targets (Edwards, 1983; Grunwell, 1987; Khan & Lewis, 1984; Smit, 2004; Stoel-Gammon & Dunn, 1985) and idiosyncratic processes as recommended targets (Smit, 2004). Such information has importance to clinical practice in that it highlights the need for clinicians to recognize patterns of speech sound errors using both standardized and less formal measures. Additionally, clinicians who are concerned with when and how to intervene with CI users may find this information useful in determining what processes to target first.

The authors of this study expected that those participants with longer duration of auditory exposure would exhibit fewer phonological processes than participants with shorter durations. Surprisingly, review of the data revealed that participant 5, (longest HA of 43 months and shortest gap between CA and HA) did not use fewer phonological processes. However, his percentage use of individual phonological processes was, on average, much lower than that of the other participants (less than 16.67%) and may have contributed positively to his overall intelligibility. Moreover, it is likely that

the phonological development of participant 5 may have been a direct result of early CI use and a longer duration of improved speech perception. This notion is supported by Kirk et al. (2002) who found that increased CI use over time led to improved speech perception and production, specifically when it occurred during the most critical and rapid periods of speech acquisition. Participant 5's CI use starting at age 13 months (age at implant activation) may have provided auditory and linguistic input during infancy, facilitating increased speech comprehension and production skills (Chin, 2003).

Although early implantation and a narrow gap between CA and HA provided a basis for the average KLPA-2 standardized scores of participant 5, these factors did not hold true for participant 1, who also attained average scores. In comparison to the other participants in the study, participant 1 did not have a narrow gap between CA and HA but had the second largest gap of 30 months. He may have presented with more typical phonological skills than the remaining three participants due to variability factors of familial, educational and environmental considerations, such as the early identification of his hearing loss followed by amplification and Auditory-Verbal training. The combination of these factors may have significantly benefited this participant's speech and phonological development. Yoshinago-Itano et al. (1998) demonstrated that children identified early and fitted with hearing aids under 6 months of age perform better than those fitted later. Access to early intervention before 6 months of age along with appropriately fitted amplification produces greater benefit in children with hearing loss.

Differences in phonological skills also were revealed for the remaining participants (2, 3 and 4), who, when compared to other participants, had shorter HAs ranging from 22 months to 28 months. Information gleaned from parent questionnaires revealed participants 2 and 4 experienced late identification of hearing loss, which subsequently resulted in later amplification, receipt of CIs and initiation of rehabilitative services. The combination of these factors may have resulted in less sophisticated phonological skills.

Conversely, participant 3, who had the second longest HA (28 months), had her hearing loss identified at birth, received amplification and started early intervention services at 6 months. However, unlike participants 1 and 5, her articulation and phonological abilities still were delayed. Her lower articulation and phonological abilities are attributed to the fact that she resides with parents who are profoundly deaf and who use Total Communication in the home. Therefore, she was not exposed to speech as frequently as the other participants. This finding was consistent with the findings of Chin and Finnegan (2000), who noted that children who use Total Communication do not develop speech as efficiently as children who use oral communication. Furthermore, Chin and Finnegan found that children who used oral communication responded three times more accurately to speech production tasks than children using Total Communication. Therefore, it is plausible that users

of Total Communication may suppress phonological processes at a slower rate than children who use oral communication, as evidenced by participant 3 in this study.

Clinically, the findings of this research are consistent with previous findings and imply that many factors must be considered when examining the phonological systems of young children with CIs. The proximity of CA to HA appears to indicate a more ideal outcome for pediatric CI users. A narrow gap between CA and HA indicates the presence of auditory input during the most crucial period of speech development. Tye-Murray, Spencer and Woodworth (1995) highlighted the significance of improved speech perception and comprehension in children with CIs, as it positively influenced speech production skills and contributes to the development of a more adult-like phonological system.

The study also concurs with previous researchers' findings that there is considerable individual variability in CI recipients' spoken language development. Contributing variability factors include age at receipt of CI, onset of deafness, duration of device use, motivation and type of linguistic environment (Kirk, 2000). The consideration of HA and CA is essential when assessing the phonological abilities of children with CIs. The results of this research suggest that a child with a short duration of hearing (i.e., less than 3 years) may utilize sound simplification patterns used by their peers. Children with CIs typically produce less speech than their typically developing peers. Therefore, they may use phonological processes as a means of simplifying their speech production. It is important to note that for many children with CIs, the development of more adult-like phonological skills may occur over time.

Although the present study provides valuable information regarding the phonological systems of young children with CIs, it is important to acknowledge certain limitations. The sample size was small ($n = 5$) and may not be fully representative of the phonological skills of young children who received CIs prior to age 3. Additionally, all children in this study used the same CI device; therefore, results may be different for children of comparable ages who use different technology. Furthermore, children with CIs were not included in the GFTA-2 (2000) and KLPA-2 (2002) standardization populations, indicating that scores from these assessment tools should be interpreted with caution. Consonant deletions, especially in the final position, are common in the hearing loss (HL) population, as these sounds typically have reduced intensity and are at a high frequency (e.g., /s/). The GFTA-2 and KLPA-2 do not assess vowel production and distortion or neutralization of vowels, all common errors found within the HL population (Bauman-Waengler, 2004).

The identification of phonological processes can be a preliminary step for school-based clinicians in the phonological analysis of children with CIs.

Additional components might include a language sample. A second cautionary consideration is that although segmental errors can contribute to intelligibility (Klein & Flint, 2006; Morris, Wilcox, & Schooling, 1995), such errors in suprasegmental factors such as prosody, pitch, loudness, rate and voice have been shown also to affect intelligibility (Shriberg & Kwiatkowski, 1992) and need to be included in the assessment of the CI population.

As the articulation and phonological development of CI recipients continue to improve, it may become increasingly more fitting to use standardized tests designed for children with typical hearing to assess articulation and phonological competence, both for individual CI users and the general CI population (Chin & Kaiser, 2002). Also, as additional CI recipients are mainstreamed into typical classroom settings, school-based clinicians may want to utilize ready-made and easily accessible diagnostic tools to assess the articulation and phonological skills of CI users.

Future research in this area should include a larger sample size and studies to determine the differences in phonological process use of children who received their CIs at various ages (e.g., before age 2 and after age 4). Other factors to consider could include speech-processing strategies used with CIs, nonverbal intelligence, auditory memory, vocabulary skills and specific measures of speech production and speech perception abilities. Furthermore, studies evaluating the effects of mode of communication on phonological development and idiosyncratic process use in children with CIs also are warranted.

References

- Bankson, J., & Bernthal, N. (2004). *Articulation and phonological disorders* (5th ed.). Boston: Allyn & Bacon.
- Bauman-Waengler, J. (2004). *Articulatory and phonological impairments: A clinical focus*. Boston: Pearson Education, Inc.
- Blamey, P.J., Barry, J.G., Bow, C., Sarant, J., Paatsch, L., & Wales, R. (2001a). The development of speech production following cochlear implantation. *Clinical Linguistics & Phonetics*, 15(5), 363–382.
- Blamey, P.J., Barry, J.G., & Jacq, P. (2001b). Phonetic inventory development in young cochlear implant users 6 years postoperation. *Journal of Speech, Language, and Hearing Research*, 44, 73–79.
- Chin, S.B. (2002). Aspects of stop consonant production by pediatric users of cochlear implants. *Language, Speech, and Hearing Services in Schools*, 33, 38–51.
- Chin, S.B. (2003). Children's consonant inventories after extended cochlear implant use. *Journal of Speech, Language, and Hearing Research*, 46, 849–862.
- Chin, S.B., & Finnegan, K.R. (2000). Consonant cluster production by pediatric users of cochlear implants. *The Volta Review*, 102, 157–174.

- Chin, S.B., Finnegan, K.R., & Chung, B.A. (2001). Relationships among types of speech intelligibility in pediatric users of cochlear implants. *Journal of Communication Disorders*, 34, 187–205.
- Chin, S.B., & Kaiser, C.L. (2000). Measurement of articulation in pediatric users of cochlear implants. *The Volta Review*, 102, 145–156.
- Chin, S.B., & Pisoni, D.B. (2000). A phonological system at 2 years after cochlear implantation. *Clinical Linguistics & Phonetics*, 14(1), 53–73.
- Chin, S.B., Tsai, P.L., & Gao, S. (2003). Connected speech intelligibility of children with cochlear implants and children with normal hearing. *American Journal of Speech-Language Pathology*, 12, 440–451.
- Colletti, V., Carner, M., Miorelli, V., Guida, M., Colletti L., & Fiorino, F.G. (2005). Cochlear implantation at under 12 months: Report on 10 patients. *Laryngoscope*, 115(3), 445–449.
- Dawson, P.W., Blamey, P.J., Dettman, S.J., Rowland, L.C., Barker, E.J., Tobey, E.A., et al. (1995). A clinical report on speech production of cochlear implant users. *Ear and Hearing*, 16, 551–561.
- Dyson, A.T. (1988). Phonemic inventories of 2- and 3-year-old children. *Journal of Speech and Hearing Disorders*, 53, 89–93.
- Edwards, M.I. (1983). Selection criteria for developing therapy goals. *Journal of Childhood Communication Disorders*, 7, 36–45.
- Fisher, H., & Logemann, J. (1971). *Test of Articulation Competence*. New York: Houghton and Mifflin.
- Goldman, R., & Fristoe, M. (1986). *Goldman-Fristoe Test of Articulation*. Circle Pines, MN: American Guidance Service.
- Goldman, R., & Fristoe, M. (2000). *Goldman-Fristoe Test of Articulation-2*. Circle Pines, MN: American Guidance Service.
- Grunwell, P. (1987). *Clinical phonology* (2nd ed.). Baltimore: Williams & Wilkins.
- Hodson, B., & Paden, E. (1991). *Targeting intelligible speech: A phonological approach to remediation* (2nd ed.). Austin, TX: Pro-Ed.
- Horn, D.L., Davis, R.A.O., Pisoni, D.B., & Miyamoto, R.T. (2005). Development of visual attention skills in prelingually deaf children who use cochlear implants. *Ear & Hearing*, 26, 389–408.
- Ingram, D. (2002). The measurement of whole-word productions. *Journal of Child Language*, 29, 713–733.
- International Phonetic Association. (1999). *Handbook of the International Phonetic Association: A guide to the use of the International Phonetic Alphabet*. Cambridge, UK: Cambridge University Press.
- Khan, L.M., & Lewis, N.P. (1984). A practical guide to phonological assessment and the development of treatment goals. *Communicative Disorders*, 9, 51–66.
- Khan, L.M., & Lewis, N.P. (2002). *Khan-Lewis Phonological Analysis-2*. Circle Pines, MN: American Guidance Service.

- Kirk, K., & Hill-Brown, C. (1985). Speech and language results in children with a CI. *Ear & Hearing*, 6, S36–S47.
- Kirk, K.I., Miyamoto, R.T., Ying, E.A., Perdew, A.E., & Zuganelis, H. (2002). Cochlear implantation in young children: Effects of age at implantation and communication mode. *The Volta Review*, 102, 127–144.
- Klein, E.S., & Flint, C.B. (2006). Measurement of intelligibility in disordered speech. *Language, Speech, and Hearing Services in Schools*, 37(3), 191–199.
- Ling, D. (1998). *Foundations of spoken language for the hearing-impaired child*. Washington, DC: Alexander Graham Bell Association for the Deaf.
- Lowe, R.J. (1996). *Assessment link between phonology and articulation: ALPHA* (rev. ed.). Mifflinville, PA: Speech and Language Resources.
- Miyamoto, R.T., Kirk, K.I., Robbins, A.M., Todd, S., & Riley, A. (1996). Speech perception and speech production skills of children with multichannel cochlear implants. *Acta Otolaryngologica*, 116, 240–243.
- Morris, S.R., Wilcox, K.A., & Schooling, T.L. (1995). The preschool speech intelligibility measure. *American Journal of Speech-Language Pathology*, 4, 22–28.
- Osberger, M.J., Maso, M., & Sam, L.K. (1993). Speech intelligibility of children with cochlear implants, tactile aids, or hearing aids. *Journal of Speech and Hearing Research*, 36, 186–203.
- Peng, S., Spencer, L.J., & Tomblin, J.B. (2004). Speech intelligibility of pediatric cochlear implant recipients with 7 years of device experience. *Journal of Speech, Language, and Hearing Research*, 47, 1227–1236.
- Serry, T.A., & Blamey, P.J. (1999). A 4-year investigation into phonetic inventory development in young cochlear implant users. *Journal of Speech, Language, and Hearing Research*, 42, 141–154.
- Shriberg, L.D., & Kwiatkowski, J. (1992). Intelligibility assessment in developmental phonological disorders: Accuracy of caregiver gloss. *Journal of Speech, Language, and Hearing Research*, 35, 1095–1104.
- Smit, A.B. (2004). *Articulation and phonology resource guide for school-age children and adults*. New York: Thomson Delmar Learning.
- Smit, A.B. (1993b). Phonologic error distributions in the Iowa-Nebraska Articulation Norms Project: Consonant singletons. *Journal of Speech, Language and Hearing Research*, 26, 533–547.
- Smit, A.B., & Hand, L. (1997). *Smit-Hand Articulation and Phonology Evaluation*. Los Angeles: Western Psychological Services.
- Stampe, D. (1969). The acquisition of phonetic representation. Paper presented at the Fifth Regional Meeting of the Chicago Linguistic Society, Chicago, 1969.
- Stoel-Gammon, C. (1987). Phonological skills of 2-year olds. *Language, Speech, and Hearing Services in Schools*, 18, 323–29.
- Stoel-Gammon, C. & Dunn, C. (1985). *Normal and disordered phonology in children*. Baltimore: University Park Press.
- Svirsky, M.A., Robbins, A.M., Kirk, K.I., Pisoni, D.B., & Miyamoto, R.T. (2000).

- Language development in profoundly deaf children with cochlear implants. *Psychological Science*, 11, 153–158.
- Tobey, E.A., Pancamo, S., Staller, S.J., Brimacombe, J.A., & Beiter, A.L. (1991). Consonant production in children receiving a multichannel cochlear implant. *Ear & Hearing*, 12(1), 23–31.
- Tobey, E.A., Geers, A.E., Brenner, C.B., Altuna, D., & Gabbert, G. (2003). Factors associated with development of speech production skills in children implanted by age five. *Ear and Hearing*, 24, 36S–45S.
- Tomblin, J.B., Barker, B.A., Spencer, L.J., Zhang, X., & Gantz, B.J. (2005). The effect of age at cochlear implant initial stimulation on expressive language growth in infants and toddlers, *Journal of Speech, Language and Hearing Research*, 48, 853–869.
- Tye-Murray, N., Spencer, L., & Woodworth, G.G. (1995). Acquisition of speech by children who have prolonged cochlear implant experience. *Journal of Speech and Hearing Research*, 38, 327–337.
- Waltzman, S.B., & Cohen, N.L. (1998). Cochlear implantation in children younger than 2 years old. *American Journal of Otology*, 9, 158–162.
- Yoshinaga-Itano, C., Sedey, A.L., Coulter, D.K., & Mehl, A.L. (1998). Language of early- and later-identified children with hearing loss. *Pediatrics*, 102, 1161–1171.

HARMONY
HiResolution Bionic Ear System

"During an
ultrasound,
I listened to
my baby's
heartbeat for
the first time."

—Stacey McGovern,
implanted at age 29

TREASURE

every moment of your child's life.

Stacey's third pregnancy was made even more memorable after she was implanted with the HiResolution® Bionic Ear System. The new HiRes Fidelity 120™ Sound Processing feature* is designed to help you appreciate the fullness and richness of life's precious moments.

Read more about Stacey's life with HiResolution Sound at
www.BionicEar.com/RealStories.

Connect and Discover™

* In U.S., optional feature for adults only. See package insert for details.
©2007 Advanced Bionics Corporation. All Rights Reserved. Printed in U.S.A.
ALC01 2/09/07, BW

ADVANCED
BIONICS
A Boston Scientific Company