The Volta Review

Alexander Graham Bell Association
for the Deaf and Hard of Hearing

The

Research

Speech Intelligibility in Young Cochlear
Implant Recipients: Gains During Year Three 85
David J. Ertmer, Ph.D., CCC-SLP

Vowel Production in 7- to 12-Month-Old
Infants With Hearing Loss 101
*Rebecca Nelson, Au.D.,
Christine Yoshinaga-Itano, Ph.D.,
Ann Rothpletz, Ph.D., and Allison Sedey, Ph.D.*

The Role of Age and Oral Lexical
Competence in False Belief Understanding by
Children and Adolescents With Hearing Loss 123
*Antonia González, Ph.D.,
Inmaculada Quintana, Ph.D.,
Carmen Barajas, Ph.D., and
María José Linero, M.Sc.*

The Joint Committee on Infant Hearing 2007
Position Statement 141
Reproduced with permission from *Pediatrics*,
120(4), pp. 898–921. Copyright © 2007 by the
American Academy of Pediatrics (AAP).

EMMACOLLEY :: age 3
Implanted June 2003 (left), January 2005 (right)

Introducing the
Absolute Latest
in Cochlear Implant Technology:
the **NEW PULSARci¹⁰⁰**

I CAN

Tell Emma jokes in the car
Teach her new songs and watch her dance
Talk to her from the next room
Read her stories when she's on my lap
Listen as she talks with her friends
Feel pride in her age-appropriate language skills
Understand exactly what she's saying
Marvel at her abilities in a mainstreamed classroom
Whisper "I love you" to her
Believe she can do and be anything she wants

hear**LIFE**

When you choose a cochlear implant from MED-EL – either for yourself or a loved one – YOU CAN open up a world of possibilities. We not only offer today's most technologically advanced cochlear implant systems, but we believe we are on the verge of breakthroughs that will offer users a lifetime of more exciting tomorrows. For more information about MED-EL, contact us today.

TOLL FREE (888) MEDEL-CI (633-3524) :: V/TDD PHONE (919) 572-2222 :: www.medel.com

The Volta Review

Volume 107, Number 2

ISSN 0042-8639

Fall 2007

Organized in 1890 to encourage the teaching of speech, speechreading and the use of residual hearing to people who are deaf or hard of hearing, the Alexander Graham Bell Association for the Deaf and Hard of Hearing (AG Bell) welcomes to its membership all who are interested in improving educational, professional and vocational opportunities for people who are deaf or hard of hearing. Affiliated with AG Bell are the Parent Section, the Deaf and Hard of Hearing Section and the International Professional Section.

83 Editor's Preface

Patricia M. Chute, Ed.D.

Research

- 85 Speech Intelligibility in Young Cochlear Implant Recipients:
Gains During Year Three
David J. Ertmer, Ph.D., CCC-SLP
- 101 Vowel Production in 7- to 12-Month-Old Infants With Hearing Loss
*Rebecca Nelson, Au.D., Christine Yoshinaga-Itano, Ph.D., Ann Rothpletz, Ph.D.,
and Allison Sedey, Ph.D.*
- 123 The Role of Age and Oral Lexical Competence in False Belief
Understanding by Children and Adolescents With Hearing Loss
*Antonia González, Ph.D., Inmaculada Quintana, Ph.D., Carmen Barajas, Ph.D., and
María José Linero, M.Sc.*
- 141 The Joint Committee on Infant Hearing 2007 Position Statement
Reproduced with permission from *Pediatrics*, 120(4), pp. 898–921.
Copyright © 2007 by the American Academy of Pediatrics (AAP).
- 191 Information for Contributors to *The Volta Review*

Permission to Copy: The Alexander Graham Bell Association for the Deaf and Hard of Hearing, as copyright owner of this journal, allows single copies of an article to be made for personal use. This consent does not extend to posting on Web sites or other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works of any type, or for resale without the express written permission of the publisher. For more information, contact AG Bell at 3417 Volta Place, NW, Washington, DC 20007, e-mail editor@agbell.org, or call 202/337-5220 (voice) or 202/337-5221 (TTY).

Editor's Preface

Patricia M. Chute, Ed.D.

The importance of early identification of hearing loss to enabling parents to make informed decisions about habilitation options is one of the primary tenets of AG Bell. It is this fundamental belief that provided the impetus for the organization's establishment and remains the basis from which it functions today. Parents and professionals require evidence-based research to support the choices that each makes. This issue of *The Volta Review* continues to provide its readers with information to assist them in their decision-making process. Speech and vowel production along with an understanding of theory of mind of individuals with hearing loss are explored. In addition, the most recent statement from the Joint Committee on Infant Hearing is provided so that readers can assess the results of the published studies within the context of this important position paper.

In the first paper by Ertmer, the speech intelligibility of children who received cochlear implants by 3 years of age and utilized their implants for three years is assessed. The performance of this group was compared to published data on older children with cochlear implants to determine the overall effects of early receipt of cochlear implants on speech intelligibility. Although the subject pool for this study was small ($n = 6$), the performance of the younger group of children was superior to children who received cochlear implants later in life. In addition, intelligibility scores were shown to increase during the period measured in this study (three years). Overall, the results support early receipt and use of cochlear implants as one of the contributing factors for better speech production.

Research that complements that of Ertmer is authored by Nelson, Yoshinaga-Itano and Rothpeltz. Here, the authors investigate the changes in vowel production in infants with hearing loss who are 7 to 12 months of age. This group of 54 children was divided into three subsets based on degree of hearing loss. Phonetic transcriptions of a 30-minute sample were compared to children with typical development of the same age. The results indicate that children with hearing loss demonstrate similar acquisition patterns to chil-

dren with typical hearing. The authors also note some differences depending upon the degree of hearing loss.

The final investigative piece addresses the false belief understanding of children and adolescents with hearing loss relative to their oral lexical competence. González, Quintana, Barajas and Linero begin by explaining the basis of false belief and how it relates to the development of theory of mind. The 54 participants in this study ranged in age from 6 to 19 years, had parents with typical hearing and attended an oral school for the deaf in Spain. Their results indicate that false belief is resolved in the older group of children (14 years or older) who demonstrate higher levels of oral lexical competence. When this finding is viewed with respect to children with typical hearing, the age discrepancy is pronounced: Children with typical development achieved false belief understanding by 4 years of age. The implications relative to linguistic competence also are discussed.

The final portion of this issue of *The Volta Review* is dedicated to the publication of the most recent position statement of the Joint Committee on Infant Hearing. The above studies underscore the overall importance of early identification and habilitation of children with hearing loss. Knowledge of the position statement by parents and professionals provides the necessary foundation from which each group can function effectively.

The editorial board of *The Volta Review* works diligently to provide its subscribers with quality research in the area of deafness. It is only through their dedication that this journal continues to publish the excellent level of research in the field. My appreciation and acknowledgement of that dedication is ever present and always kept in mind.

Pat Chute, Ed.D.
Editor
The Volta Review

Speech Intelligibility in Young Cochlear Implant Recipients: Gains During Year Three

David J. Ertmer, Ph.D., CCC-SLP

This investigation sought to answer two questions: (1) Do children who receive cochlear implants (CIs) by 3;0 (years; months) improve speech intelligibility significantly during their third year of cochlear implant use? (2) How do the intelligibility scores of young CI recipients compare to those reported in the literature for children who receive CIs at older ages? These questions were addressed through a prospective, longitudinal research study in which children imitated short sentences at 24, 30 and 36 months post-activation. The sentences were played for adult listeners who were instructed to orthographically transcribe the words they recognized. Percent intelligibility scores then were calculated for each speech sample. Mean intelligibility scores increased significantly from 28% to 62% during the third year of CI use. The participants also reached comparable or higher levels of intelligibility within fewer months of CI experience than reported for children who received a CI at older ages. These findings support the notion that receiving a CI by 3 years of age is advantageous for developing connected speech that can be understood by unfamiliar listeners.

Introduction

Two recent events have revolutionized communication intervention for children with hearing loss in the United States. First, the widespread implementation of newborn hearing screening procedures has made it possible to identify children with hearing loss and begin aural habilitation within the first year of life. Second, decreases in age criterion for cochlear implantation have enabled children to begin to hear within an age range when many foundational spoken language skills typically are acquired. As a result, hearing loss now is identified routinely before a child's first birthday, and cochlear implants (CIs), if indicated, are provided commonly within the first three years of life.

The primary benefits of cochlear implantation are increased auditory sensitivity and improved speech perception ability. Greatly increased access to

David J. Ertmer is an associate professor in the department of speech, language, and hearing sciences at Purdue University.

adult speech models and auditory feedback also has led to secondary improvements in spoken language development in children (see American Speech-Language-Hearing Association, 2004, for a review). In particular, children who receive CIs before their third birthdays have been shown to make accelerated progress in prelinguistic vocal development (e.g., Ertmer, Young, & Nathani, 2007; Moore & Bass-Ringdahl, 2002), phonological development (e.g., Connor, Craig, Raudenbush, Heavner, & Zwolan, 2006) and spoken language development (e.g., Svirsky, Teoh, & Neuberger, 2004). To date, however, very little is known about the time course for developing intelligible speech in this population.

Speech intelligibility has been defined as “the degree to which the speaker’s intended message is recovered by the listener” (Kent, Weismer, Kent, & Rosenbeck, 1989; p. 483). As Subtelney (1977) noted 30 years ago, “Intelligibility is considered the most practical single index to apply in assessing competence in oral communication” (p. 183). This very functional communication ability commonly is measured through rating scales, transcription analysis and word identification procedures (see Osberger, 1992, for review). The current investigation used word identification procedures to assess the intelligibility of sentences produced by children with prelingual deafness who received a CI before their third birthdays. The studies reviewed below used similar methods with children who received a CI at various ages. The purpose of this review is to establish benchmarks for comparison with the outcomes of the present study.

A mixed-model research design (i.e., cross-sectional and longitudinal) was used by Miyamoto, Svirsky, Kirk, Robbins, Todd and Riley (1997) to track improvements in speech intelligibility in children who received a CI at an average age of 5;0. Speech samples were collected prior to implantation and at six-month intervals through 90 months (7;6) of CI experience. Between 18 and 34 children gave samples at each interval. The children’s spoken sentences were played for three judges who wrote down what they understood, and a percentage intelligibility score was calculated for each sample. The results showed that mean scores increased gradually to approximately 25% during the first three years of CI use, and that scores approximated 40% for children with 3.5 to 7.5 years of CI experience. Although improvements during the first two years of CI use were relatively small, subsequent scores exceeded the widely reported level of speech intelligibility associated with children who are deaf and hard of hearing and who use hearing aids (20%) (Osberger, 1992).

Another study (Miyamoto, Kirk, Svirsky, & Sehgal, 1999) was among the first to explore the effects of age at receipt of CI on speech intelligibility. Three groups were examined: those receiving CIs before 3;0 ($n = 14$), those receiving CIs between 3;0 and 3;11 ($n = 11$) and those receiving CIs between 4;0 and 5;3 ($n = 8$). All the children were tested once between 4;0 and 5;0 years of age, thus differing in the amount of CI experience at the time of testing. The

children also were grouped as oral (speech only) or simultaneous (speech and sign) communicators. The children imitated sentences from the Beginner's Intelligibility Test (BIT) (Osberger, Robbins, Todd, & Riley, 1994). Recordings of the sentences then were played for three judges and a percentage intelligibility score was obtained for each sample. Children receiving CIs before 3;0 were found to have a higher mean score (20%) than children in either of the groups of children who received a CI at an older age (approximately 9%), although the difference between the scores did not reach statistical significance. Oral communicators achieved higher intelligibility levels than those who used sign and speech together (20% versus 4.2%). The main findings showed that initial gains in speech intelligibility were limited during the first two years of CI experience, and that oral communication promoted greater speech intelligibility gains than simultaneous communication. However, because of the disparity in the amount of CI experience for each group (e.g., 0 and 2.3 years for the oldest and youngest groups, respectively), the long-term effects of age at receipt of CI could not be assessed.

A study by Peng, Spencer and Tomblin (2004) examined speech intelligibility in 24 simultaneous communicators who received a CI between 2.5 and 11 years of age ($M = 5;1$). The participants had used their CIs for seven years with either SPEAK or MPEAK speech coding strategies. The children produced 14 sentences from the Short-Long Sentence Test (Tye-Murray, 1998). These sentences were played two times each for a panel of adult listeners. Data analysis revealed that the mean for the first presentation was 67.86% (range of scores: 5.5%–100%); the mean for the second was slightly higher (71.54%) with a similar range of scores. Higher scores were noted for users of the SPEAK versus MPEAK strategy, and for children who received a CI at relatively younger ages. In relation to the latter finding, it was estimated that a decrease of one year of age at receipt of CI was equivalent to a 5.5% increase in intelligibility.

Advances in technology and decreases in age at receipt of CI have made it reasonable to compare the speech of children with CIs to that of their peers with typical hearing. Chin, Tsai and Gao (2003) used typical development (TD) as a benchmark as they compared the intelligibility of 51 children with CIs to that of 47 children with TD. The children in the CI group had an onset of deafness between birth and 2;5 ($M = 0;2$), an age at receipt of CI between 1;5 and 5;10 ($M = 3;2$) and 0;6 to 5;7 years of CI experience ($M = 2;4$ years). Approximately 55% were oral communicators; the rest were simultaneous communicators. The children with CIs were between 2;8 and 10;8 at testing. The children with TD were between 2;6 and 6;9 years old. Both groups produced short sentences as speech samples.

For the CI group, a mean score of 34.5% intelligibility was obtained and a very wide range of scores was observed (0%–98%). Regression analyses revealed that intelligibility scores increased with age and continued CI use. In contrast, data from the TD group showed much higher levels of intelligibility,

with steady increases from approximately 54% intelligibility for 2-year-olds to 95% intelligibility for 4-year-olds. Small increases were observed at 5 and 6 years of age, when scores approximated the 100% level. In summary, after an average of 28 months of CI experience, the CI group surpassed the 20% level of intelligibility associated with older children who are deaf who use hearing aids (Osberger, 1992) but, as a group, did not reach the high levels achieved by 4-year-olds with TD.

In a related study, Tobey, Geers, Brenner, Altuna and Gabbert (2003) examined the intelligibility of 181 subjects who received a CI before 6;0 ($M = 3;5$), had an average of 5;6 of CI experience (range: 3;9-7;6) and were between 7;11 and 9;11 at the time of testing. Approximately half of the children were oral communicators and half were simultaneous communicators. The children were asked to say sentences of varying length (3, 5 or 7 syllables) in which key words had been selected as targets for measuring intelligibility. These sentences were recorded and played for adult judges, and intelligibility scores were calculated. Data analysis revealed that the children had achieved a mean intelligibility score of 63.5% at the time of testing. Thus, after 5;6 of CI experience, the children's average score far exceeded the 20% level associated with children who are deaf who use hearing aids but did not reach the level of 4-year-olds with TD (Chin et al., 2003; Copland & Gleason, 1998). Age at receipt of CI did not influence intelligibility scores.

Although the above-cited studies included children who varied greatly in chronological age, age at receipt of CI, communication modality and amount of CI experience, they share three common findings: (1) substantial increases in speech intelligibility were observed following receipt of CI; (2) scores varied greatly across children; and (3) considerable room for improvement remained in the speech of children with up to 7 years of CI experience.

Of particular interest to the current study, the findings of Peng et al. (2004), as well as the results of a rating scale study by Tye-Murray and Spencer (1995) and a transcriptional analysis study by Flipsen and Colvard (2006), suggest that receipt of a CI at younger ages may lead to more rapid improvements than receipt of a CI later in childhood. However, the largest study to date (Tobey et al., 2003) did not find an advantage related to age at receipt of CI. In light of this discrepancy and the increasing numbers of infants and toddlers who receive implants, further research is needed to determine whether receipt of a CI by 3;0 is more beneficial for developing intelligible speech than receipt of a CI later in life.

This investigation employed a prospective longitudinal research design to measure speech intelligibility across the third year of CI use in children who received a CI before 3;0. The third year of CI experience was examined because it was anticipated that young children would require approximately two years to make gains in speech perception, prelinguistic vocal development, phonological development and early linguistic development before increases in connected speech intelligibility would be evident. Thus, it was

hypothesized that the participants would make significant increases in speech intelligibility during the third year of CI use. Another aim of the study was to determine whether the intelligibility scores of the participants exceeded those reported in earlier studies of children who received a CI at older ages who had comparable or greater amounts of CI experience.

Methods

Participants

Two boys and four girls participated in this study. All had received unilateral, multichannel cochlear implants between 10 and 36 months of age ($M = 21.8$ months), and all had participated in previous studies of prelinguistic vocal development (Ertmer & Mellon, 2001; Ertmer, Young, & Nathani, 2007). Table 1 contains information on hearing status, cochlear implant technology and data collection for each child. The children are listed in order of age at receipt of CI; their names have been changed to protect confidentiality.

All the children were from American English-speaking families and, aside from hearing loss, were reported to be without physical or learning disabilities. All received family-centered intervention until their third birthdays. After that age, Danny, Diane, Kathy, Hannah and Michael were enrolled in an oral preschool program, and Ellen attended a university preschool program for children with language disorders. Although manual signs were used intermittently in the latter setting, Ellen's parents and teachers relied mainly on spoken English to communicate with her after she received her CI. Michael attended a regular preschool and received speech-language services twice weekly.

Data Collection and Stimulus Materials

The Beginner's Intelligibility Test (BIT) (Osberger et al., 1994) was administered to assess speech intelligibility after 24, 30 and 36 months of CI use. The BIT consists of four lists of 10 simple sentences incorporating vocabulary likely to be familiar to young children. For each child, a different BIT list was used during each recording session. The BIT was administered by having the tester say each sentence while demonstrating its meaning with objects. For example, for "Mommy sits," the tester placed a small female figure in a toy chair while saying the sentence. After the tester said each sentence and performed the corresponding action, the child was asked to repeat the sentence.

Each speech sample was video- and audio-recorded using a Sony Handicam 8 mm camcorder coupled with Realistic or Optimus Boundary microphones placed within 2 feet of the child. Recordings of the elicited sentences then were isolated from the tester's models, digitized at a sampling rate of 20 KHZ and stored as computer files. Each child's digitized sentences were

Table 1. Hearing History, Audiometric and Cochlear Implant Information for Each Participant

<i>Child</i>	<i>Age identified/receipt of CI (months)</i>	<i>Unaided pure-tone averages (dB HL)</i>	<i>Implant type (processing strategy)</i>	<i>Mean implant-aided pure-tone averages (dB HL) during years 1-3 of CI use</i>	<i>CA intervals (months) after receipt of CI³ Intervals: 24 30 36</i>
Danny	32/36	NR ¹	Clarion Multi-strategy (CIS)	27 (range: 22-32)	59 65 71
Diane	23/28	NR ¹	Clarion Multi-strategy (CIS)	30 (range 25-34)	51 57 63
Kathy	10/20	93	Clarion Multi-strategy (CIS)	29 (range 28-29)	43 49 55
Hannah	1/19	NR ¹	Clarion Multi-strategy (CIS)	25 (range 23-27)	42 48 54
Ellen	15/18	103	Nucleus 24 (ACE)	37 (range 32-39)	41 47 53
Michael	2 days/10	NA ²	Clarion Multi-strategy (CIS)	24 (range 19-29)	33 39 45

¹NR: No responses at maximum audiometer output levels.

²NA: Not available. Child could not be conditioned for behavioral testing. Auditory Brainstem Response testing predicted severe to profound bilateral hearing loss.

³Based on initial activation one month after implant surgery

Note. ACE = advanced combination encoder; CA = chronological age; CI = cochlear implant; CIS = continuous interleaved sampling

copied onto audiocassette tapes or burned to compact discs (CDs) so they could be presented to listener-judges. Audiocassettes were used for samples recorded earlier in the investigation. CDs were used for later samples because of the ease with which they could be produced. The quality of the two media was judged to be comparable and satisfactory.

Data Analysis

Intelligibility Judgments

The intelligibility of BIT sentences was judged by randomly recruited university students who had typical hearing, no formal training in phonological disorders and little or no exposure to the speech of children who are deaf and hard of hearing. The judges wore headphones as they listened to each sentence twice and wrote down the words they understood. They were encouraged to guess whenever they were unsure. Three judges listened to each set of sentences produced by each child. The judges were exposed to each BIT list only once.

The judges' written responses were scored by counting the number of BIT words that were correctly identified. The total number of correctly identified words was then divided by the total number of words that the child attempted to produce for a given list. For example, if a child said "Rabbit eating" instead of the target "The rabbit is eating," only two words were used in calculating the percentage of words correctly identified. Children were given credit for producing a word if the judge identified the root of the word correctly. For example, if a child said "Mommy jump" instead of "Mommy jumps," the word "jump" was counted as correct if identified by a judge. The mean percentage of words correctly identified by the three judges was used as the intelligibility score for each sample.

Statistical Analyses

Comparisons of mean scores from the three intervals were conducted with repeated measures analysis of variance (RMANOVA) using the amount of implant experience (24, 30 and 36 months) as the independent variable and children's intelligibility scores as the dependent variable. Percent correct scores were converted to Rationalized Arcsine Units (Studebaker, 1985) before the RMANOVA and post hoc tests were run.

Results

Figure 1 shows the mean intelligibility scores and standard deviations for the six participants at 24, 30 and 36 months postactivation. The means at these

Figure 1. Mean intelligibility scores and standard deviations at 24, 30 and 36 months post-CI activation

intervals increased from 28.33 (SD 24.94) to 49.83 (SD 28.42) to 61.66 (SD = 27.02), respectively. The main effect of implant experience was found to be significant, $F(2) = 21.0794$, $p \leq 0.0003$. Mean arcsine values from the three testing intervals then were compared using Tukey-Kramer adjusted t -tests. The results showed that scores from the 30- and 36-month intervals were significantly greater than those of the 24-month interval, $t(10) = -4.00$, $p \leq 0.0065$ and $t(10) = -6.36$, $p \leq 0.0002$, respectively. Marginal evidence of improvement was noted between the 30- and 36-month intervals, $t(10) = -2.37$, $p \leq 0.0918$.

Figure 2 shows individual scores during the third year of CI use. Considerable variability can be seen. Individual scores ranged from a low of 8% (Ellen, 24 months) to a high of 95% (Danny, 36 months). Danny (74%) and Kathy (39%) received the highest scores at the start of the study; the remaining scores were $\leq 21\%$ at this interval.

The amount of improvement between 24 and 36 months also varied across children. Five of the six children made relatively large gains. Danny's initial score increased 21 percentage points from 74% to 95%; Kathy's 24-month score (39%) more than doubled to 81%; Diane's and Michael's first scores (21% and 18%, respectively) tripled in value to reach the 64% level; Hannah increased her initial score (10%) by almost 40 percentage points to 49% at 36 months. Much smaller increases were noted for Ellen, who began the study with the lowest level of intelligibility (8%) and had raised her score to 17% one year later.

Individualized patterns of increase were observed during the third year of CI use. Diane, Hannah and Kathy made the greatest gains between 24 and 30 months; Ellen and Michael made their greatest progress between 30 and 36 months; and Danny made equivalent progress during both periods. Although different patterns were observed, trends toward improvement at each succeeding interval were noted for each child.

Figure 2. Individual intelligibility scores during the third year of CI use

Discussion

Group Outcomes

On average, the children in the study made substantial progress during the third year of CI use (Figure 1). Increases were noted at each succeeding interval as mean scores rose from 28% at 24 months to almost 50% at 30 months and to 62% after 36 months of CI use. With the exception of Danny, each child made greater progress during the third year of CI use than during the first and second years combined. These findings confirmed the hypothesis that the children would make significant increases in speech intelligibility during their third year of CI use. When compared with the scores of the children with TD in the Chin et al. (2003) study, scores at the 36-month interval fell between those of a 2-year-old with TD (54%) and a 3-year-old with TD (72%). The similarity of the scores for children with implants and children with TD suggests that speech was developing at a near-typical rate. Because of the small number of participants, however, the increases seen in the current study must be considered preliminary.

The following paragraphs compare group scores from the current study with those of the word identification studies summarized in the introduction. Recall that the children in the current study were oral communicators, whereas the other cited studies included either simultaneous communicators only (Peng et al., 2004) or a mix of oral and simultaneous communicators (Chin, et al., 2003); Miyamoto et al., 1997, 1999; Tobey et al., 2003). Previous investigations have shown higher intelligibility scores for oral than for simultaneous communicators (Miyamoto et al., 1999; Osberger et al., 1994). Thus, the intelligibility scores obtained in earlier studies reflect the influence of communication modality as well as age at receipt of CI.

Comparisons With Children Who Receive CIs Around 5 Years of Age

The children in the current study appeared to make more rapid progress than reported for older implant recipients. Miyamoto et al. (1997) examined the speech of children who received a CI around 5;0. Their participants had reached the 40% intelligibility level after 41 to 89 months of CI experience. By comparison, the younger implant recipients in the current study exceeded this level by 23 percentage points after only 36 months of CI use.

Children in the Peng et al. (2004) study also received a CI close to 5;0. They achieved a mean intelligibility score of 71.54% after 84 months of CI experience. This score was approximately 10 percentage points greater than the 62% achieved at 36 months in the current study. On the basis of the increasing trends seen in Figure 1, it seems reasonable to predict that the younger implant recipients will exceed the intelligibility level of the children in the Peng et al. (2004) study in fewer months of CI experience. In sum, oral communicators who received a CI at an average age of 21.8 months appeared to make more rapid progress than simultaneous communicators who received a CI around 5 years of age.

Comparisons With Children Implanted Before 5 Years

Children in the Tobey et al. (2003) study received a CI by 41 months on average and had achieved an intelligibility score of 63.5% after 66 months of CI experience. Children in the current study had reached the 62% level after 36 months of CI use. Thus, the younger recipients required 30 fewer months of implant experience to reach a level of intelligibility comparable to that of those who received a CI approximately 19 months later in life. On the basis of trends toward further increases in intelligibility, it seems reasonable to expect that the current participants will reach higher intelligibility levels with fewer months of CI experience than the children studied by Tobey et al. (2003).

The most intelligible talkers in the Miyamoto et al. (1999) study were oral communicators who received a CI before 3;0. These children achieved an intelligibility score of 20% after an average of 27 months of CI experience. The children in the current study received a CI within the same age range and had achieved a slightly higher score (28%) with three fewer months of CI use. Thus, both studies support the notion that gains in speech intelligibility are gradual during the first two years after receipt of a CI. Further comparisons with the older implant recipients in the Miyamoto et al. (1999) study cannot be made because of the children's limited amount of CI experience at the time of testing.

The methods and the ages of the children in the current study are fairly well aligned with the Chin et al. study (2003). In that investigation, BIT sentences also were used to assess intelligibility, and the participants received

their implants at a relatively young age ($M = 38$ months). The children achieved a mean intelligibility score of 34.5% after an average of 28 months of CI use. If this score were to be plotted in Figure 1, it would fall approximately midway between data from the 24- and 30-month intervals. Thus, the scores of the current study appear to be commensurate with those of the 51 children studied by Chin et al. (2003). The outcomes of both studies also suggest that receipt of a CI close to 3;0 can lead to greater gains during the first 30 months of CI experience than are seen in children who receive CIs at older ages (e.g., Miyamoto et al., 1997). It is interesting to note that the children in these studies had comparable scores despite differences in their communication modality. The similarity of the scores suggests that receipt of a CI around 3;0 might promote equivalent intelligibility gains for oral and simultaneous communicators alike during the first three years of CI experience. Further research is needed to explore this possibility.

In summary, group scores from the current study revealed substantial improvements in connected speech intelligibility during the third year of CI use. The children's improvements were found to be more rapid than those reported for children who received CIs between 3;6 and 5;1 (Miyamoto et al., 1997; Peng et al., 2004; Tobey et al., 2003) and commensurate with the scores of children who received a CI at similar (Miyamoto et al., 1999) and slightly older ages (3;2) (Chin et al., 2003). After three years of CI experience, the participants' mean intelligibility level was comparable to the individual scores of some 3- and 4-year-olds with TD studied by Chin et al. (2003).

Individual Performance

Scores at the 24-month interval indicate different rates of improvement during the first two years of CI use (Figure 2). The intelligibility levels of four children (Diane, Ellen, Hannah and Michael) approximated 20% or lower after two years of CI experience, indicating that they had not yet exceeded the level commonly found for older children who use hearing aids (Osberger, 1992). A similar finding also was seen in the children with CIs studied by Miyamoto et al. (1997, 1999). Relatively low scores at the 24-month interval support an earlier contention by Robbins, Kirk, Osberger and Ertmer (1995) that gains in intelligibility begin to appear after two years of CI experience.

Danny's and Kathy's scores at the 24-month interval (71.8% and 39%, respectively) show, however, that children can make relatively large gains during the first two years of CI use. Although Kathy's clinical history was quite similar to that of the other children, Danny's background offers some insight into factors that might promote rapid speech gains. An earlier study showed that Danny had begun to produce canonical syllables prior to receiving a CI and that he completed the process of prelinguistic vocal development relatively quickly (within six months of activation; Ertmer et al., 2007). His rapid improvements in intelligibility provide further evidence that he was adept at integrating auditory information for speech development.

At least two factors might have contributed to Danny's superior progress. To begin with, his hearing loss was identified relatively late (32 months), and, although it cannot be confirmed, it is possible that he had some hearing experience prior to receiving his CI. Second, his relatively greater physical, cognitive and motor maturity at the time of implant surgery might have enabled him to integrate auditory models and feedback for speech development more readily than younger children who had not yet developed similar abilities in these areas.

In contrast, several factors might have contributed to Ellen's limited progress. First, her average CI-aided thresholds were 8 to 12 dB HL poorer than those of the other children. Reduced access to speech models and auditory feedback could have affected speech development negatively. Second, unlike the other children, Ellen's interventionists and parents used signs intermittently to supplement spoken language after she received her CI. This method might have been adverse for speech development if Ellen focused on signed communication to a greater extent than speech, or if expectations for speech learning were less than those for oral communication (Osberger et al., 1994). Third, Ellen used a different CI than did the other children. However, the device had been shown to be efficacious for speech development in other intelligibility studies (e.g., Tobey et al., 2003) and functioned appropriately throughout the study.

Factors such as nonverbal intelligence, implant characteristics and communication modality have been shown to account for relatively large portions of variability in the oral communication skills of children who receive CIs by 5;0 (Geers, 2002). These and other personal, technological, interventional and family variables might affect speech development in younger implant recipients as well. Further studies are needed to explore these possibilities.

Clinical Implications

Integration into mainstream elementary classrooms is one of the key motivations for CI use in very young children. Successful integration requires the development of many social, communicative, pre-academic and academic skills prior to a regular school placement (Wilkins & Ertmer, 2002). Monsen (1981) concluded that listeners are "confronted by overwhelming difficulty in understanding" when speech is less than 59% intelligible (p. 850). Thus, with the exception of Ellen, the children in the current study had reached, or made considerable progress toward, speech that is "understandable" to others. Children who develop intelligible speech during the preschool years increase their chances of successful integration in the primary grades. The findings of this study suggest that receipt of a CI by 3;0 can help to promote integration into mainstream primary-level classrooms by giving children an earlier start in auditory-based speech development.

Summary

This preliminary investigation demonstrated that oral communicators who received a CI before 3;0 made substantial progress toward intelligible speech during their third year of CI experience. These gains were more rapid than those reported for children who received a CI after 3;6 who were both oral and simultaneous communicators. Future investigations with larger numbers of children, direct comparisons of ages at receipt of CI and controls for influential factors are planned to further explore speech intelligibility in young CI recipients. These investigations will provide much-needed insight into the benefits of newborn hearing screening and early intervention for functional speech development in young CI recipients.

Acknowledgments

This study was completed through grants from the National Institutes on Deafness and Other Communication Disorders (NIDCD 1R03DC04226-03; 1R01DC007863). Special thanks to the children and parents who made this study possible. The contributions of Le'Ann Scott, Jennifer Quesenberry, Chrissy Miller, Jadie Dayton, Carrie Garrard, Kendall McEwan, Jetta Martin and Jennifer Engstrom are gratefully recognized. Thanks also to Bruce Craig, Yong Wang and Benjamin Tyner for their advice and assistance in completing statistical analyses.

References

- American Speech-Language-Hearing Association. (2004). *Technical report on cochlear implants*. Rockville, MD: Author.
- Chin, S.B., Tsai, P.L. & Gao, S. (2003). Connected speech intelligibility of children with cochlear implants and children with normal hearing. *American Journal of Speech-Language Pathology*, 12, 440–451.
- Connor, C.M., Craig, H.K., Raudenbush, S.W., Heavner, K., & Zwolan, T. (2006). The age at which young deaf children receive cochlear implants and their vocabulary and speech growth: Is there added value for early implantation? *Ear and Hearing*, 27, 628–644.
- Copland, J., & Gleason, J.R. (1988). Unclear speech: Recognition and significance of unintelligible speech in preschool children. *Pediatrics*, 82, 447–452.
- Ertmer, D.J., & Mellon, J.A. (2001). Beginning to talk at 20 months: Early vocal development in a young cochlear implant recipient. *Journal of Speech, Language, and Hearing Research*, 44, 192–206.
- Ertmer, D.J., Young, N.M., & Nathani, S. (2007). Profiles of vocal development in young cochlear implant recipients. *Journal of Speech, Language, and Hearing Research*, 50, 393–407.
- Flipsen, P., & Colvard, L.G. (2006). Intelligibility of conversational speech

- produced by children with cochlear implants. *Journal of Communication Disorders*, 39, 93–108.
- Geers, A.E. (2002). Factors affecting the development of speech, language, and literacy in children with early cochlear implantation. *Language, Speech, and Hearing Services in the Schools*, 33, 172–183.
- Kent, R.D., Weismer, G., Kent, J.F., & Rosenbeck, J.C. (1989). Toward phonetic intelligibility testing in dysarthria. *Journal of Speech and Hearing Disorders*, 54, 482–499.
- Miyamoto, R.T., Svirsky, M., Kirk, K., Robbins, A.M., Todd, S., & Riley, A. (1997). Speech intelligibility of children with multichannel cochlear implants. *Annals of Otology, Rhinology, and Laryngology*, 106, 35–36.
- Miyamoto, R.T., Kirk, K., Svirsky, M., & Sehgal, S.T. (1999). Communication skills in pediatric cochlear implant recipients. *Acta Otolaryngologica*, 119, 219–224.
- Monsen, R.B. (1981). A usable test for the speech intelligibility of deaf talkers. *American Annals of the Deaf*, 126, 181–184.
- Moore, J.A. & Bass-Ringdahl, S. (2002). Infant vocal development in candidacy for and efficacy of cochlear implantation. *Annals of Otology, Rhinology, and Laryngology*, 111, 52–55.
- Osberger, M.J. (1992). Speech intelligibility in the hearing impaired: Research and clinical implications. In R.D. Kent (Ed.), *Intelligibility in speech disorders* (pp. 233–265). Philadelphia: John Benjamins Publishing.
- Osberger, M.J., Robbins, A., Todd, S., & Riley, A. (1994). Speech intelligibility of children with cochlear implants. *Volta Review*, 96, 169–180.
- Peng, S.C., Spencer, L.J., & Tomblin, J.B. (2004). Speech intelligibility of pediatric cochlear implant recipients with 7 years of device experience. *Journal of Speech, Language, and Hearing Research*, 47, 1227–1236.
- Robbins, A.M., Kirk, K.I., Osberger, M.J., & Ertmer, D. (1995). Speech intelligibility of implanted children. *The Annals of Otology, Rhinology and Laryngology*, 106(Suppl.), 399–401.
- Studebaker, G.A. (1985). A “rationalized” arcsine transform. *Journal of Speech and Hearing Research*, 28, 455–462.
- Subtelney, J. (1977). Assessment of speech with implications for training. In F. Bess (Ed.), *Childhood deafness* (pp. 183–194). New York: Grune & Stratton.
- Svirsky, M., Teoh, S-W., & Neuberger, H. (2004). Development of language and speech perception in congenitally, profoundly deaf children as a function of age at cochlear implantation. *Audiology and Neuro-otology*, 9, 224–233.
- Tobey, E.A., Geers, A.E., Brenner, C., Altuna, D., & Gabbert, G. (2003). Factors associated with development of speech production skills in children implanted by age five. *Ear and Hearing*, 24(Suppl.), 36S–45S.
- Tye-Murray, N. (1998). Speech, language, and literacy development. In N.

- Tye-Murray & W. Clark (Eds.), *Foundations of aural rehabilitation: Children, adults, and their family members* (pp. 415–446). San Diego: Singular.
- Tye-Murray, N., & Spencer, L (1995). Acquisition of speech by children who have prolonged cochlear implant experience. *Journal of Speech, Language, and Hearing Research*, 38, 327–337.
- Wilkins, M. & Ertmer, D.J. (2002). Introducing young children who are deaf or hard of hearing to spoken language: Child's Voice, an oral school. *Language, Speech, and Hearing Services in the Schools*, 33, 196–204.

Vowel Production in 7- to 12-Month-Old Infants With Hearing Loss

Rebecca Nelson, Au.D., Christine Yoshinaga-Itano, Ph.D.,
Ann Rothpletz, Ph.D., and Allison Sedey, Ph.D.

The purpose of this study was to examine vowel production in 7- to 12-month-old infants with hearing loss. Fifty-four infants were divided into three groups according to degree of hearing loss (mild-to-moderate, moderately severe-to-severe, profound), and their vocalizations were phonetically transcribed from 30-minute videotaped samples. These transcriptions were analyzed to determine the percentage of infants in each group who produced the various English vowel phones. Results showed that vowel acquisition patterns in infants of this age with all degrees of hearing loss were similar to those of children of the same age with typical hearing. Established production of the /i/ and /e/ vowels was seen only in the mild-to-moderate hearing loss group, which may demonstrate the emerging role of auditory feedback in vowel production. Findings are discussed in terms of clinical implications and directions for future research.

Introduction

The prelinguistic stage of phonological development occurs during the first year of life. Unfortunately, there is not an abundance of documented research examining vowel development during these first 12 months. Even fewer research studies have investigated vowel development in infants with hearing loss. Although research has shown that vowel development is affected by

Rebecca Nelson, Au.D., is a doctoral student in the department of speech, language and hearing sciences at the University of Colorado–Boulder, where she studies cultural and linguistic factors affecting audiologic service provision. Christine Yoshinaga-Itano, Ph.D., is a professor in the department of speech, language and hearing sciences at the University of Colorado–Boulder and specializes in intervention services for infants, children and adults with hearing loss. Ann Rothpletz, Ph.D. is a research assistant professor in the department of psychological and brain sciences at the University of Louisville. Allison Sedey, Ph.D. is a research associate at the University of Colorado–Boulder specializing in speech and language development in children who are deaf and hard of hearing.

hearing loss, the majority of these studies explored phonological productions occurring after the first year of life (Dagenais & Critz-Crosby, 1992; Ertmer, 2001; Ertmer et al., 2002; Yoshinaga-Itano & Sedey, 2001). It is known that perception and correct production of most vowels are developed by 3 years of age in infants who are developing typically (Larkins, 1983) and that vowel production in children who are deaf and hard of hearing usually is delayed in comparison to children with typical hearing or lesser degrees of hearing loss (Yoshinaga-Itano & Apuzzo, 1998; Yoshinaga-Itano & Sedey, 2001). However, vowel acquisition in children with hearing loss during the first year of life has remained largely uninvestigated. This paper considers the effects of hearing loss on the development of vowel phones in 7- to 12-month-old infants. Specifically, we were interested in determining if vowel acquisition was similar across groups of infants with various degrees of hearing loss and similar to that of infants with typical hearing.

General Characteristics of English Vowels

The English language contains 16 vowel sounds /i, ɪ, e, ɛ, æ, a, u, ʊ, o, ɔ, ɑ, ʌ, ə, ɜ, ɝ, ɞ/ and five diphthongs /aɪ, ɔɪ, aʊ, eɪ, oʊ/. This number, however, varies according to the dialect. Many sources cite only 15 English vowel sounds. Phonetically speaking, vowels are speech sounds shaped within the oral cavity that do not obstruct the flow of air through the cavity (Howard & Heselwood, 2002). Their production comes from articulatory movements that are associated with certain acoustic characteristics that allow for their differentiation (Donegan, 2002). All vowels are phonated (voiced), and they commonly are differentiated according to tongue location in the vertical and horizontal dimensions of the mouth.

Vowel sounds can be categorized as high (tongue body is elevated from the neutral position), mid (tongue body height is neutral) or low (tongue body is lowered). Neutral position here refers to the tongue being situated in a middle position in the oral cavity, not high or low. Additionally, vowels are front (tongue extended forward in the oral cavity), central (neutral tongue location) and back (tongue retracted in the oral cavity) (Yavas, 1998). The vowel quadrilateral illustrates the relative tongue positioning during the production of each vowel (Figure 1). Along the vertical axis of the oral cavity, /i, ɪ, u, ʊ/ may be considered high and high-mid vowels, /e, ɛ, o, ɔ, ə/ considered mid and low-mid vowels and /æ, a, ʌ/ considered low vowels, although /ʌ/ sometimes is also considered a mid or low-mid vowel (Shriberg & Kent, 2003; Yavas, 1998). Along the horizontal plane, the front vowels are /i, ɪ, e, ɛ, æ/, central vowels are /ə, ʌ/ and back vowels are /u, ʊ, o, ɔ, ɑ/ (Shriberg & Kent, 2003; Yavas, 1998).

Another way to classify vowels is according to their first and second formant frequency components (F1 and F2, respectively). Generally speaking, decreasing the size of the oral cavity and/or enlarging the pharyngeal cavity

Modified from Shriberg, L. & Kent, R. (2003)

Figure 1. English vowel quadrilateral.

lowers the F1 frequency (Donegan, 2002). Similarly, moving the tongue forward and upward in the mouth raises the F2 frequency because it decreases the size of the oral cavity (Donegan, 2002). These relationships allow vowels' acoustic characteristics to be estimated on the basis of their location in the vowel quadrilateral (i.e., the tongue's location during their production).

Typical Prelinguistic Vocalic Development

During their first year of life, infants who are developing typically go through various stages of prelinguistic vocalization, which likely reflect their developing auditory, neural and vocal motor systems. Between 7 and 12 months of age, infants usually begin to produce well-formed syllables and a greater diversity of sounds (Oller & Eilers, 1988; Smith, Goffman, & Stark, 1995). Studies of the vocalic productions of infants who are developing typically have found that between approximately 7 and 13 months of age, the most frequently produced vowel phones were front, central and mid-to-low vowel phones (Buhr, 1980; Kent & Bauer, 1985; Smith & Oller, 1981). Conversely, back vowel-like and high vowel-like sounds were produced much less frequently (Buhr, 1980; Kent & Bauer, 1985; Smith & Oller, 1981). These findings indicate a trend of early vocalic production that corresponds to place of location, with mid-to-low and front-to-central vowels occurring more frequently than high or back vowels.

Theoretical Considerations in Prelinguistic Vocalic Development

Numerous physiological and environmental factors, such as cognitive and motor development, neurological maturation, native language and other linguistic factors, vision, auditory abilities and social reinforcement all have

been hypothesized to affect phonological development in some way. Various theoretical models relating to these factors, each with its own strengths and weaknesses, have been developed in attempt to explain the phenomenon of phonological acquisition. Vihman (1996) provided a comprehensive discussion of these theoretical models of phonological development. Of greatest interest to the current study are models pertaining to the roles of auditory perception and oral motor development in vowel acquisition between 7 and 12 months of age. As tongue location in the oral cavity corresponds to vocalic formant frequencies, patterns found in vowel acquisition could be related either to oral motor development or auditory perception of the different vowel formant frequencies. Therefore, the discussion of theoretical models here focuses on examining auditory and oral motor development.

One theory of phonological acquisition that considers infants' articulatory-auditory feedback loop development was proposed by Olmsted (1966), although he placed much more emphasis on the perceptual component than the articulatory aspect. Olmsted postulated that pronunciation of phones is learned in relation to their ease of perception. He argued that children learn to use auditory cues from their own productions as feedback regarding whether or not their articulations (based on their kinesthetic feedback) match those they have previously heard in adult speech. This theory hypothesized that some phones contain more easily discriminable components (e.g., voicing, nasality) than others, and these phones will be produced earlier in development (Olmsted, 1966).

The ability to imitate speech sounds requires an understanding of the connection between the articulatory movements produced and the sounds that are perceived. Neurological growth in the auditory cortex occurs between 4.5 and 12 months of age, coinciding with the emergence of language-specific phoneme perception (Moore, 2002). It is possible that maturation of the neural pathways enables developing infants to perceive, classify and store the sounds heard into mental representations of the vowels (and consonants) of their ambient language. These representations then may be used as prototypes for vowel productions as the infants begin to babble and explore the sounds of speech (Donegan, 2002).

In addition to evidence regarding the role of auditory input in phonological acquisition, the oral-motor abilities of developing infants must be considered when discussing prelinguistic phonological development. It is known that over the first four to five years of life, infants' vocal tracts progress toward resembling those of adults (Buhr, 1980; Lieberman, 1975; Mackenzie Beck, 1997). It has been suggested that the right-angle bend found in adult vocal tracts does not appear until around 9 or 10 months of age (Buhr, 1980). This bend separates the vocal tract into two distinct cavities (pharyngeal and oral), allowing for the greater first and second formant resonance contrasts that are characteristic of the high back and front vowels /u/ and /i/, respectively (Lieberman, 1975).

In summary to this point, the articulatory and acoustic characteristics of

vowels appear to be related to vowel acquisition in infancy. Low-to-mid, front and central vowels are the most commonly produced vowels early in life (Buhr, 1980; Kent & Bauer, 1985; Smith & Oller, 1981). Front vowels and low vowels can be formed simply by adjusting the position of the jaw, suggesting that jaw control develops early and facilitates the production of these vowels (Buhr, 1980). As infants mature, improvements in vowel accuracy likely are attributable to the development of better tongue control, which allows for better manipulation of the oral cavity (Buhr, 1980).

Impact of Hearing Loss on Prelinguistic Vowel Development

Examining children with hearing loss provides an interesting perspective on the process of phonological acquisition, given their more limited auditory experiences. Various studies have found reduced vocalic inventories within the child's ambient language and/or reduced formant frequency regions in young children with hearing loss as compared to their peers with typical hearing (Ertmer, 2001; Ertmer et al., 2002; Kent, Osberger, Netsell, & Hustedde, 1987; Yoshinaga-Itano & Apuzzo, 1998; Yoshinaga-Itano & Sedey, 2001). In addition, comparisons across groups of children with various hearing ranges have indicated that young children with lesser degrees of hearing loss exhibited larger phonemic repertoires than those with greater degrees of hearing loss (Yoshinaga-Itano & Apuzzo, 1998; Yoshinaga-Itano & Sedey, 2001). Another study showed that with appropriate amplification and intervention, most children with hearing loss were able to produce almost all of the vowels of the English language by 5 years of age (Yoshinaga-Itano & Sedey, 2001). Given that infants who are developing typically have complete vowel repertoires by approximately 3 years of age (Larkins, 1983), the Yoshinaga-Itano and Sedey (2001) findings show a significant delay in the vowel development of children with hearing loss.

Although the aforementioned studies indicated reduced vocalic diversity in the productions of children with hearing loss, information regarding specific phonemic difficulties for children with hearing loss would provide insight into the effects of auditory perception on vowel acquisition. It has been found that neutral vowels (i.e., /ə/ and /ʌ/) were the most commonly produced vowels by children (age 6 to 42 months) with hearing loss, regardless of their specific degree of hearing loss (Yoshinaga-Itano, Stredler-Brown, & Jancosek, 1992). The only vowel that differed significantly in production by degree of hearing loss was /e/, which was produced by a significantly larger percentage of children who were hard of hearing (i.e., better ear pure tone average [BPTA] of less than 70 dB hearing level [HL]) than children who were deaf (i.e., BPTA of greater than 70 dB HL) (Yoshinaga-Itano et al., 1992).

Studies investigating the development of the vowel formant regions and tongue positions in children with hearing loss have found reduced vowel formant regions and less distinct tongue positioning compared to those of

children with typical hearing (Dagenais & Critz-Crosby, 1992; Kent et al., 1987). Interestingly, Kent et al. (1987) found that an infant with bilateral severe-to-profound hearing loss continually produced a more restricted range of vocalic formant frequencies between 8 and 15 months of age. The reduced formant regions he exhibited may be related to less distinct tongue positioning noted in older children with hearing loss (Dagenais & Critz-Crosby, 1992). Furthermore, these reduced formant regions likely correspond to the reduced vowel inventories found in the studies previously mentioned. On the other hand, a young girl who received a cochlear implant at 19 months of age increased her vocalic diversity and was able to produce vowels from all vowel categories (low-back, central, mid-front, etc.) within five months of receiving her implant (Ertmer, 2001), demonstrating vowel productions much more similar to those of young children with typical hearing. These results indicate that early auditory stimulation may minimize the reduction in F1–F2 regions characteristic of vowel productions by children with profound hearing loss.

Purpose

Phonological development during the first year of life is the foundation for speech and oral language acquisition. Because 6 to 12 months of age appears to be the time when language-specific perceptions and vocalizations emerge (Oller & Eilers, 1988; Polka & Werker, 1994), it is important to know how limitations in auditory input affect vocal development at this age. It is well established that hearing loss reduces vowel diversity in infancy (Ertmer 2001; Yoshinaga-Itano & Apuzzo, 1998; Yoshinaga-Itano & Sedey, 2001); however, the only phoneme-specific information regarding vowel inventories in this population has come from case studies or studies with small sample sizes. The purpose of this study was to investigate the production of individual vowel phones by a large number of infants with hearing loss, as well as how differing degrees of hearing loss (mild-to-moderate, moderately severe-to-severe, profound) affect vocalic production.

This study addressed the following questions:

1. Within each degree of hearing loss group (mild-to-moderate, moderately severe-to-severe, profound), which vowels are established in production by 12 months of age?
2. How does the vowel production of infants with hearing loss compare to that of infants with typical hearing between 7 and 12 months of age?
3. What is the effect of degree of hearing loss on vowel production in 7- to 12-month-old infants?
4. How do the results relate to the theoretical considerations of audition and oral motor development?
5. What are the implications of the results for clinicians and researchers working with children with hearing loss?

Method

Participants

Fifty-four infants with hearing loss, ages 7 to 12 months, participated in this study. Infants were divided into three groups according to degree of hearing loss: mild-to-moderate, moderately severe-to-severe and profound. Infants in the mild-to-moderate group had a BPTA between 26 and 55 dB HL. A BPTA between 56 and 90 dB HL was classified as a moderately severe to severe hearing loss, and a BPTA greater than 90 dB HL was considered a profound hearing loss. The sample sizes for the groups were 21, 21 and 12, respectively.

The infants in this study were participants in a larger longitudinal project examining speech and language development in young children with hearing loss from birth to age 7 years. Thus, audiologic records were available for review when the children were older than they were when the data for the current study were obtained. This allowed for a more accurate determination of thresholds than if only their infant hearing results had been used. Only infants with a confirmed degree of hearing loss, as determined by two certified audiologists, were included in this study. Audiologists reviewed the longitudinal audiologic data for the participants to determine their degree of hearing loss. Frequency-specific behavioral audiologic data were available for 51 (94.4%) of the 54 infants. Degree of hearing loss classifications were based on audiologic records at the oldest age available (range = 8 to 84 months of age). For the majority of these infants, Auditory Brainstem Response (ABR) data or Visual Reinforcement Audiometry (VRA) data at a young age were available to confirm degree of hearing loss classifications and serve as an indicator of stable hearing thresholds. Audiologic records for five of the infants (7.41%) provided evidence of progressive hearing loss. For these infants, degree of hearing loss was classified according to the audiologic records available at the age closest to 7 to 12 months. Three of the infants (5.56%) had only ABR data available; one of these infants had tone burst ABR data available in addition to click ABR data. All of the infants were from Colorado, New Mexico or Wyoming. The demographic characteristics of the subjects are described below and summarized in Table 1 according to degree of hearing loss group.

Gender

The sample contained approximately half females and half males, with this same distribution found within each degree of hearing loss group.

Additional Disabilities

According to parent reports, 72.2% of the 54 infants had no known additional disabilities, while 27.8% exhibited one or more disabilities in addition to hearing loss, including cognitive, motor, neurological, social-emotional

Table 1. Demographic Characteristics of Subjects According to Degree of Hearing Loss Group (percentage of participants)

<i>Characteristic</i>	<i>Degree of Hearing Loss Group</i>			
	<i>Mild to Moderate (n = 21)</i>	<i>Moderately Severe to Severe (n = 21)</i>	<i>Profound (n = 12)</i>	<i>All Infants (n = 54)</i>
<i>Gender</i>				
Boy	47.6%	47.6%	50.0%	48.1%
Girl	52.4%	52.4%	50.0%	51.9%
<i>Additional disabilities</i>				
Present	19.0%	47.6%	8.3%	27.8%
Absent	81.0%	52.4%	91.7%	72.2%
<i>Mother's education level</i>				
Less than high school	14.3%	4.8%	8.3%	9.3%
High school	23.8%	42.8%	41.7%	35.2%
Post-high school	61.9%	52.4%	50.0%	55.5%
<i>Ethnicity</i>				
Minority	23.8%	23.8%	25.0%	24.1%
Not a minority	76.2%	76.2%	75.0%	75.9%
<i>Communication mode</i>				
Oral only	33.3%	28.6%	25.0%	29.6%
Oral + sign	42.9%	47.6%	58.3%	48.2%
No data	23.8%	23.8%	16.7%	22.2%
<i>Hearing status of mother</i>				
Typical	100.0%	95.2%	100.0%	98.1%
Hearing loss	0.0%	4.8%	0.0%	1.9%
<i>Hearing status of father</i>				
Typical	85.7%	100.0%	100.0%	94.4%
Hearing loss	14.3%	0.0%	0.0%	5.6%
<i>Etiology</i>				
Unknown	42.8%	61.9%	75.0%	57.4%
Heredity	28.5%	14.3%	8.3%	18.5%
Cytomegalovirus	14.3%	0.0%	0.0%	5.6%
Goldenhar syndrome	4.8%	4.8%	0.0%	3.7%
Other	9.6%	19.0%	16.7%	14.8%
<i>Spoken language</i>				
English only	85.7%	90.5%	91.7%	88.9%
English & Spanish	14.3%	9.5%	8.3%	11.1%
<i>Regular intervention</i>				
CHIP ^a	61.9%	68.4%	50.0%	61.5%
CHIP+ ^b	38.1%	21.1%	41.7%	32.7%
Other	0.0%	10.5%	8.3%	5.8%

^aColorado Home Intervention Program

^bColorado Home Intervention Program and other therapy services

and/or visual disabilities. The presence of additional disabilities for the moderately severe-to-severe hearing loss group was 47.6%, which was higher than that of the mild-to-moderate and profound groups, whose percentages were 19% and 8.3%, respectively.

Mother's Level of Education

The highest educational level completed by the mothers of the infants was used as an index for socioeconomic status. In each degree of hearing loss group, at least half of the mothers had completed some post-high school education. Overall, 9.3% of the mothers did not finish high school, 35.2% had a high school diploma and 55.5% had completed at least one year of post-secondary education. The moderately severe-to-severe group had the smallest percentage of mothers who did not finish high school (4.8%), while the mild-to-moderate group had the largest percentage (14.3%). A Kruskal-Wallis H Test revealed no statistically significant difference between degree of hearing loss groups in the mean mother's education level ($\chi^2 = 1.39$, $df = 2$, $p = 0.5$). Previous research has shown no significant correlation between mother's education level and speech production by children with hearing loss (Yoshinaga-Itano & Sedey, 2001).

Ethnicity

In each degree of hearing loss group, Caucasian infants accounted for approximately 75% of the sample. Most of the remaining infants were Latino. This distribution is representative of that found in Colorado (U.S. Census Bureau, n.d.), the home state for the majority of the infants in the study.

Communication Mode used by Primary Caregiver

The communication mode typically used with the infants by their primary caregivers was recorded from parent report. Data were not available for 12 (22.2%) of the infants. Of the 42 infants with available communication mode information, approximately 40% of the caregivers communicated only orally with their infants, while roughly 60% communicated through a combination of sign and oral methods. None of the caregivers communicated using sign language only. Slightly more caregivers in the mild-to-moderate group (33.3%) used only oral methods than those in the moderately severe-to-severe group (28.6%), who used oral methods slightly more than the caregivers in the profound group (25%).

Hearing Status of Parents

One mother and three fathers of the infants who participated in the study were hard of hearing. All other parents had typical hearing, and all infants had at least one parent with typical hearing.

Etiology

Genetic syndromes, cytomegalovirus and heredity were the most common known causes of hearing loss in this sample. However, the etiology was unknown for 57.4% of the infants.

Spoken Languages Used at Home

English was spoken in the homes of all of the infants in this study. For almost 90% of the infants, English was the only language used in the home. Within the mild to moderate group, 85.7% of the families used only English in the home, while 14.3% communicated through both English and Spanish. Within the profound and moderately severe-to-severe groups, just under 10% of the families used both Spanish and English to communicate.

Intervention Program

Information regarding intervention services was available for 52 of the 54 infants. Ninety-four percent of the infants received services through the Colorado Home Intervention Program (CHIP). CHIP is a family-centered intervention program that provides services that focus on hearing, speech, language and social-emotional development in children with hearing loss birth to age 3 years. Typically, a CHIP facilitator is paired with a family and provides intervention services in their home once a week for approximately 1 to 1.5 hours. Therapy is parent-centered, meaning that it focuses on educating the parents so that they can implement intervention techniques in their everyday routines.

In addition to receiving services through CHIP, 32.7% of families received therapy from other sources. These sources included intervention services at a children's hospital, private therapy from speech-language pathologists or other organizations, and community-centered early intervention programs. These services were generally more child-centered than the CHIP program. Approximately 6% of the infants did not participate in the CHIP program but did receive other intervention services. Within the mild-to-moderate group, all families received CHIP services. Between the moderately severe-to-severe and profound groups, a total of three families received services from other providers without any CHIP services.

Age at Identification of Hearing Loss

Table 2 contains information on the average age at which the infants in this study had their hearing loss confirmed, began wearing amplification, began receiving intervention services and were videotaped interacting with their parents. This information was available for 53 of the 54 infants through parent report. On average, the infants' hearing losses were identified around 9 weeks

Table 2. Demographic Characteristics of Subjects According to Degree of Hearing Loss Group (mean and standard deviations in months of age)

Characteristic	Degree of Hearing Loss Category							
	Mild to Moderate n = 21		Moderately Severe to Severe n = 21		Profound n = 12		All Infants n = 54	
	Mean	S.D.	Mean	S.D.	Mean	S.D.	Mean	S.D.
Age at identification	2.68	2.64	1.90	1.33	2.06	2.42	2.24	2.17
Age aided	4.65	2.61	3.72	1.99	4.40	2.29	4.26	2.32
Age at intervention	4.76	2.81	4.39	2.59	3.46	2.54	4.33	2.66
Age at videotaping	9.71	1.27	9.00	1.10	9.58	1.00	9.41	1.17

of age. The infants in the mild-to-moderate group had a slightly older average age of identification (2.68 months of age) than the infants with greater degrees of hearing loss (approximately 2 months of age).

Age Aided

Parents of 50 of the 54 infants provided information regarding the age when amplification was first received. The only amplification devices used by the infants in this study were hearing aids. On average, the infants were fitted with hearing aids at 4.26 months of age. The infants in the moderately severe-to-severe group received their amplification devices slightly earlier than those in the other two groups, on average at 3.72 months of age. Nonetheless, as seen in Table 2, there was significant variability within each hearing loss group in terms of the age that these infants were first aided.

Age at Intervention

The infants' age when intervention services began also was obtained through parent report. There was much variability within each hearing loss group in terms of the age at which the infants began receiving intervention services. Across hearing loss categories, the infants began receiving services at an average age of 4.33 months. More specifically, the infants with profound hearing loss began receiving services at an average age of 3.46 months, whereas the infants with hearing loss ranging from mild to severe began receiving services at an average age of 4.5 months.

Age at Videotaping

All infants in the study were videotaped to monitor their developmental progress. This process will be discussed further in the following section. Infants in all degree of hearing loss groups were videotaped, on average,

between 9 and 10 months of age. Analysis of variance of age (in months) at which the infants were videotaped revealed no significant differences across the three hearing loss groups ($F = 2.214$, $p = .120$).

Procedures

All but three of the infants received early intervention services through CHIP. As part of CHIP, a comprehensive assessment of the child's development, including speech production, was conducted every six months. The infants were videotaped for approximately 30 minutes while participating in naturalistic play with a parent or primary caregiver. A sibling also was occasionally present in the recording. After the taping, the parents determined whether the video sample was representative of the infant's typical behavior. If not, the recording was repeated on a different day to obtain an accurate representation of the infant's abilities. The videotaped interactions then were analyzed to determine the infant's developmental progress.

A linguistics doctoral student at the University of Colorado at Boulder phonetically transcribed the infant vocalizations from the videotape samples. This graduate student had extensive experience transcribing speech samples of children with hearing loss. Only vowels produced in isolation or in babbled, nonword utterances were included in this study. Vocalizations such as cries, screams, raspberries and vegetative sounds were not transcribed, and consonant phone analyses were not included in this study. Only vowel productions determined to be identifiable English phones and produced on an egressive air stream were included. Non-English vowel productions were not included in the analyses because only 5 of the 54 infants (9.3%) produced any non-English vowel-type sounds. All of these infants produced only one to two non-English vowel sounds in their speech samples, which on average consisted of 99.5 vowel tokens (range = 3 to 341 vowel phones per session). The vast majority of vowel-like sounds produced by these infants were identifiable English phones.

Trained graduate students from the Linguistics Department at the University of Colorado at Boulder retranscribed just over 20% of the transcripts used in this study in order to assess coding reliability. Mean intercoder reliability was 77.8% (range = 62.5% to 93.75%) and was assessed by calculating the percentage agreement between the vowel types identified in the original transcriptions and those identified in the retranscriptions.

Data were analyzed descriptively to determine the percentage of infants in each hearing loss group that produced the various English vowel phones in nonword utterances. These percentages then were categorized to show which vowel phones were most commonly produced by the infants in each degree of hearing loss group. The phones were categorized as established, emerging or not yet emerging. Established vowel sounds were defined as those produced at least two times by 50% or more of the infants in the degree of hearing loss group. Vowel phones produced two or more times by 25% to 49%

of the infants in a degree of hearing loss group were classified as emerging. If fewer than 25% of the infants in a degree of hearing loss group produced the vowel sound two or more times, that phone was considered to be not yet emerging. The criterion of at least two productions of the vowel phone was used to decrease the likelihood that it appeared during the sample by chance. This criterion has been used in previous studies (Stoel-Gammon & Cooper, 1984).

Results

Established and Emerging Production

Table 3 shows the percentage of infants within each degree of hearing loss group that produced each individual vowel phone two or more times. Asterisks denote phones that were considered emerging (one asterisk) or established (two asterisks) for each degree of hearing loss group. Established production of the vowel phones /ɪ, ɛ, æ, a, ə, ʌ/ was demonstrated by all three degree of hearing loss groups. The phones /i, e/ were also produced by more than 50% of the infants in the mild-to-moderate hearing loss group, whereas these vowels were at most just beginning to emerge in the other two groups. No other phones reached established production in the moderately severe-to-severe or profound groups.

Table 3. Percentage of Infants by Degree of Hearing Loss Category Who Demonstrated Two or More Productions for Each Vowel Phone

Vowel phone		Degree of hearing loss category		
		Mild to Moderate n = 21	Moderately Severe to Severe n = 21	Profound n = 12
/i/	see	52.4%**	19.0%	16.7%
/ɪ/	sit	90.5%**	71.4%**	91.7%**
/e, eɪ/	say	57.1%**	28.6%*	33.3%*
/ɛ/	step	90.5%**	85.7%**	100%**
/æ/	sad	81.0%**	71.4%**	83.3%**
/a/	ball	71.4%**	76.2%**	83.3%**
/ə/	sum	66.7%**	71.4%**	83.3%**
/ʌ/	cut	61.9%**	71.4%**	100%**
/u/	suit	19.0%	19.0%	8.3%
/ʊ/	cook	42.9%*	33.3%*	25%*
/o, ou/	sew	0.0%	9.5%	0.0%
/ɔ/	caught	0.0%	4.8%	0.0%
/ɜ/	stern	0.0%	0.0%	0.0%
/ɔɪ/	soy	0.0%	0.0%	0.0%
/aʊ/	cow	0.0%	4.8%	8.3%
/aɪ/	cry	28.6%*	23.8%	25%*

*Emerging production

**Established production

Two additional vowel phones, /ʊ/ and /aɪ/, were emerging within the mild-to-moderate hearing loss group. Approximately 43% of the infants in the mild-to-moderate group produced the vowel phone /ʊ/. This phone also was emerging in the moderately severe-to-severe and profound hearing loss groups; however, the percentage of infants producing this phone was lower than for the mild-to-moderate group. The diphthong /aɪ/ also was emerging or near emerging in all three groups. In addition, /e/ was emerging in both the moderately severe-to-severe and profound hearing loss groups, while the /i/ phone was not emerging in either group. The remaining vowel phones were not yet emerging in any of the groups. The /u/ phone was just approaching emerging status in the mild-to-moderate and moderately severe-to-severe groups, whereas the phones /o, ɔ, ɜ, ɔɪ, aʊ/ were never or rarely produced by any of the infants.

Location of Production

Figure 1 illustrates the vowel quadrilateral, which depicts the relative location of the tongue in the oral cavity during the production of English vowel phones. The vowel phones (/ɪ, ε, æ, a, ə, ʌ/), which reached established production in all of the degree of hearing loss groups, are located in the front or center portions of the oral cavity and do not require the tongue to be raised to the highest portions of the cavity or retracted posteriorly. The two phones established only in the mild-to-moderate group, /i, e/, are located in the high to mid-frontal portion of the vowel quadrilateral. The phones that were scarcely produced by any of the infants in this study were back vowels (/u, o, ɔ/), rhotic vowels (/ɜ/) and diphthongs (/aɪ, aʊ/).

Production Trends

Differences in the general trends in percentage of infants producing the vowel phones were noted across degree of hearing loss groups. Within the profound hearing loss group, either the vast majority of the infants were producing the phone, or very few of the infants were producing the phone. More specifically, all of the phones either were produced by more than 80% of the infants or by 33.3% or fewer of the infants in the profound hearing loss group (see Table 3). The mild-to-moderate and moderately severe-to-severe groups were more similar in that a fuller array of percentages was present in their results.

Discussion

Similarities and Differences in Vowel Production Between Degree of Hearing Loss Categories and Infants With Typical Hearing

Many similarities were found in the vowel samples of the infants in this study. Across all three degree of hearing loss groups, the trends in established

production included vocalization of front and central-low to mid-range vowels, with an absence of production of back vowels and diphthongs. This suggests that order of vowel acquisition in the first year of life does not differ significantly according to degree of hearing loss. These findings are consistent with those of previous research on vowel development in infants with typical hearing (Buhr, 1980; Kent & Bauer, 1985; Smith & Oller, 1981), as well as with hearing loss (Smith, 1982; Yoshinaga-Itano et al., 1992).

Two main differences by degree of hearing loss were noted in this study. First, differences in the production of the high- and mid-front vowels /i/ and /e/ were found. In contrast to the infants with greater degrees of hearing loss, those in the mild-to-moderate hearing loss category demonstrated established production of these vowels. These differences may be attributed, at least in part, to differences in age when the videotapes were gathered. Infants with mild-to-moderate hearing loss on average were approximately three weeks older than those with moderate-to-profound losses. This slight difference in age could have been enough to allow the infants with mild-to-moderate hearing loss to establish production of these two phones. However, analysis of variance showed no statistical differences among the three groups in terms of age in months. Furthermore, Yoshinaga-Itano et al. (1992) found similar results in that /e/ was produced significantly more often in their study by the children between birth and 3 years of age who were hard of hearing than those who were deaf. This vowel was the only one in that study for which significant differences in production were noted. Yoshinaga-Itano et al. (1992) credited the children with production if the phone was present one time or more.

The other main difference noted among the three groups is related to the production trends found in each group. Although there appeared to be an all-or-none production pattern in the profound hearing loss group, the mild-to-moderate and moderately severe-to-severe groups both demonstrated a fuller array of percentages, indicating a more gradual accumulation of the vowel sounds. This finding may indicate that as a group, infants with profound hearing loss use a different processing strategy for speech production than those with lesser degrees of hearing loss. Yoshinaga-Itano et al. (1992) found wider variability in the percentage of infants with severe-to-profound hearing loss producing the vowel phones than the current study. A major difference between the two studies is that Yoshinaga-Itano et al. (1992) included children with severe hearing loss and those with profound hearing loss in the same group. Additionally, the age of the infants at the time of videotaping ranged from 0 to 36 months of age in their study, and the criterion of one rather than two productions of the individual vowel phones was used. Therefore, it could be argued that the all-or-none production pattern was not evidenced in the earlier investigation because of the differing production criteria or the influence of a wider range of ages. Another possible

explanation could be that the inclusion of children with severe hearing loss in the same group as the children with profound hearing loss minimized the presence of this pattern in their results because this different processing strategy is present only in children with profound hearing loss. If the pattern noted in the current study is a valid trend in vowel acquisition in infants with profound hearing loss, it could have great implications for future research. Research examining infants' phonological approximations of the vowels, as well as audibility and discriminability in these infants, may provide a better understanding of this production trend. However, further investigation with a larger sample is necessary before any conclusions should be reached because of the small sample size ($n = 12$) of infants with profound hearing loss in this study. In addition, this trend may disappear as hearing aid technology continues to advance.

Although the infants with profound hearing loss demonstrated a different production trend than those with milder degrees of hearing loss, the pattern of established versus emerging vowels in the moderately severe-to-severe group was more similar to that of the profound group than that of the mild-to-moderate group. One plausible explanation for this pattern is that better hearing thresholds allowed the infants in the mild-to-moderate group to perceive and subsequently acquire the vowels more quickly than the infants with moderately severe to profound hearing loss. Demographic differences between the samples also should be considered. For example, there was a higher incidence of additional disabilities in the moderately severe-to-severe category than in either of the other categories. Other disabilities can significantly affect a child's development. The high percentage of additional disabilities in the moderately severe-to-severe hearing loss group may well have confounded the results for this group. Future studies should take care to minimize differences in this characteristic. Only 21.1% of the infants with moderately severe-to-severe hearing loss received intervention services in addition to CHIP, whereas approximately 40% of the infants with mild-to-moderate and profound hearing losses received additional services. Both factors may have affected the overall results for the moderately severe-to-severe group, causing them to resemble those of the profound group rather than those of the mild-to-moderate group. If these demographic characteristics were more similar to those of the other groups, the results might have more closely resembled those of the mild-to-moderate group rather than the profound group. One other demographic characteristic to consider is that almost 50% of subjects in this study used sign in addition to oral methods of communication, which could have the potential to affect their speech production. However, the similar trends seen between these infants and infants with typical hearing suggest that use of manual communication in addition to oral communication does not have a negative impact on infants' speech production at this age.

Although the exact role of moto-kinesthetic, auditory and neurological development in the acquisition of premeaningful speech vocalizations is not yet fully understood, the general production trends of the infants in this study appear to support the theory that infants still are developing vowel sounds moto-kinesthetically between 7 and 12 months of age. As has been found in younger infants (Buhr, 1980; Smith & Oller, 1981), the infants in this study produced front and central vowels much more frequently than back vowels, despite their reduced auditory input. Still, it should be remembered that all infants investigated in this study were identified and began receiving intervention services and amplification at an early age. If audition does play a role in prelinguistic speech development, these services likely would minimize the effects of reduced audition on speech development and may be at least partially responsible for similarities in vowel production found across hearing loss categories, as well as similarities to the vowel production of infants with typical hearing. It is possible that children whose hearing losses were identified at older ages might exhibit delays in their production of these vowels.

The fact that only the mild-to-moderate degree of hearing loss group reached established production of /i/ and /e/ calls into question the assumption of a solely moto-kinesthetic theory of speech development. If early speech development is mainly moto-kinesthetically-based, as some research has suggested, this difference may indicate that the influence of auditory perception on vowel production begins to emerge between 7 and 12 months of age. These two vowels have the highest F2 frequencies (Hillenbrand, Getty, Clark, & Wheeler, 1995) and among the lowest relative intensity levels (Levitt, 1978) of the vowels examined in this study. Therefore, it is possible that better hearing abilities allowed the infants with mild-to-moderate hearing loss to perceive and produce these two vowel phones, whereas most of the infants with greater degrees of hearing loss were not able to do so. According to Olmsted's (1966) theory, the lack of auditory feedback during the production of these two phones (for infants with moderately severe to profound hearing losses) may cause these vowels to be less discriminable and therefore learned later than other vowel phones with more discernable auditory and kinesthetic feedback. It seems plausible that infants' abilities to perceive these vowel sounds would reinforce their production. Because production of these mid- to high-front vowel sounds requires more oral motor coordination than production of the neutral vowels, infants with greater degrees of hearing loss who are not able to distinguish mid- to high-front vowels may not receive the auditory reinforcement necessary to stimulate their production. Therefore, the differing production abilities of /i/ and /e/ found in these three groups of infants may be the result of auditory perception interacting with moto-kinesthetic development. The presence or absence of these two vowels in infant

productions may be an early indicator of the effects of auditory perception on vowel production; however, the underlying cause of differences in production of these vowels is not understood well enough to use this as an indicator at this time.

While differing perceptual abilities may account for these differences among the three groups, it should be kept in mind that /i/ and /e/ are not among the vowels that typically are considered well-established between 7 and 12 months of age, even among children with typical hearing (Buhr, 1980; Kent & Bauer, 1985). Although both of these studies contained only a small number of participants, they provide evidence that these two phones are not well developed at this age. A case study investigating one infant's vocalic productions showed that between 29 and 48 weeks of age the child did not produce /i/ or /e/ two or more times in the majority of the recordings (Buhr, 1980). Based on these findings, it is quite impressive that the infants with mild-to-moderate hearing loss in this study were able to reach established production of these two phones. It may be that the early identification and provision of amplification, coupled with specialized intervention services, facilitated the development of these phones slightly earlier than would be expected.

Implications for Intervention and Future Research

The results of this study indicate that prior to 12 months of age, acquisition of vowel phones generally develops in the same order regardless of hearing ability. This information may serve as a benchmark for interventionists working with infants with hearing loss and be useful in the development of intervention goals. In the presence of early identification and intervention, a reduced vocalic inventory in this age range could be a preliminary indication of other disabilities, such as oral motor delays. Yoshinaga-Itano et al. (1992) suggested including the goal of increasing the diversity of vowel productions as part of intervention programs for children with hearing loss. Although insignificant differences in the number of vowels produced among degree of hearing loss groups indicate that inventorying the number of vowel productions may not be the best indicator of speech development, it is a method that is understood easily and may be motivating for parents to observe (Yoshinaga-Itano et al., 1992). This observation is important because parent motivation is essential for consistent hearing aid usage and successful intervention. Vowel diversity information also can be useful when developing intervention plans, such as targeting syllables and sounds that are within the child's vocalic repertoire if focusing on consonant development.

Although there are numerous studies related to consonant development, research in the area of vocalic acquisition in infants with hearing loss, as well as in the general population, is still minimal. Results of this investigation demonstrated an interesting all-or-none trend in the production patterns of

the profound hearing loss group. Although this finding may reflect utilization of a unique processing strategy by these infants, it may also be attributed to the small sample ($n = 12$) of infants with profound hearing loss. Further research into vowel development with a larger sample of infants with profound hearing loss would help to clarify whether the results were due to true differences in speech production processing.

In addition, investigations into the vowel development as well as the consonant development of infants with hearing loss at older ages will help to expand understanding of how speech develops in these children. The results of this study indicate that between 7 and 12 months of age, infants with hearing loss exhibit vowel repertoires similar to those of children with typical hearing. Nonetheless, it has been shown that vowel formant regions may decrease continually as these infants mature (Kent et al., 1987), and vowel productions of older children with profound hearing loss are more centralized than those of children with typical hearing (Dagenais & Critz-Crosby, 1992). It was not until 15 months of age that the infant studied by Kent et al. (1987) no longer produced the high F2 frequencies necessary for /i/ and /e/. Although Kent et al. (1987) did not discuss the specific vowel-like phones produced by this infant, the formant frequency information provided suggests that the infant may have been able to produce /i/ and /e/ at 8 months of age but lost that ability as he matured. Therefore, longitudinal investigations of vowel acquisition in infants with hearing loss would provide valuable information regarding the progression of vocalic development in these infants.

Finally, future studies could include formant structure analysis in addition to phonetic transcription techniques. Acoustic analysis of the infants' vocalizations was not possible in this study because of the naturalistic nature of the videotapes, which included overlapping vocalizations by the infant and his or her caregiver. Additionally, the videotapes were recorded in the infants' natural home environments. These environments vary in ambient noise levels and could inhibit the ability to perform acoustic analyses. Inclusion of formant structure analysis in future studies would provide a more complete picture of the vowel productions of these infants.

Acknowledgments

This research was supported in part by National Institutes of Health Contract NO1-DC-4-2141, Centers for Disease Control and Prevention Grant/Cooperative UR3/CCU824219, Office of Education # H325D030031A & H324C030074, Maternal and Child Health, the Colorado Department of Education, the Colorado Department of Public Health and Environment and the University of Colorado, Boulder. The authors also wish to give special thanks to Bill Gavin, CHIP and STEP*HI regional coordinators, CHIP and STEP*HI

parent facilitators and participating families for their contributions to this project.

References

- Buhr, R.D. (1980). The emergence of vowels in an infant. *Journal of Speech and Hearing Research*, 23, 56–72.
- Dagenais, P.A., & Critz-Crosby, P. (1992). Comparing tongue positioning by normal-hearing and hearing-impaired children during vowel production. *Journal of Speech and Hearing Research*, 35, 35–44.
- Donegan, P. (2002). Normal vowel development. In M.J. Ball & F.E. Gibbon (Eds.), *Vowel disorders* (pp. 1–35). Boston: Butterworth-Heinemann.
- Ertmer, D.J. (2001). Emergence of a vowel system in a young cochlear implant recipient. *Journal of Speech, Language, and Hearing Research*, 44, 803–813.
- Ertmer, D.J., Young, N., Grohne, K., Mellon, J.A., Johnson, C., Corbett, K., et al. (2002). Vocal development in young children with cochlear implants: Profiles and implications for intervention. *Language, Speech, and Hearing Services in Schools*, 33, 184–195.
- Hillenbrand, J., Getty, L.A., Clark, M.J., & Wheeler, K. (1995). Acoustic characteristics of American English vowels. *Journal of the Acoustical Society of America*, 97, 3099–3111.
- Howard, S.J., & Heselwood, B. (2002). The contribution of phonetics to the study of vowel development and disorders. In M.J. Ball & F.E. Gibbon (Eds.), *Vowel disorders* (pp. 37–82). Boston: Butterworth-Heinemann.
- Kent, R.D., & Bauer, H.R. (1985). Vocalizations of one-year-olds. *Journal of Child Language*, 12, 491–526.
- Kent, R.D., Osberger, M.J., Netsell, R., & Hustedde, C.G. (1987). Phonetic development in identical twins differing in auditory function. *Journal of Speech and Hearing Disorders*, 52, 64–75.
- Larkins, P. (1983). Development at 3 years. In J. Irwin & S. Wong (Eds.), *Phonological development in children 18 to 72 months* (pp. 87–105). Carbondale, IL: Southern Illinois University Press.
- Levitt, H. (1978). The acoustics of speech production. In M. Ross & T.G. Giolas (Eds.), *Auditory management of hearing-impaired children* (pp. 45–115). Baltimore: University Park Press.
- Lieberman, P. (1975). *On the origins of language*. New York: MacMillan.
- Mackenzie Beck, J. (1997). Organic variation of the vocal apparatus. In W. Hardcastle & J. Laver (Eds.), *The handbook of phonetic sciences* (pp. 256–297). Oxford: Blackwell.
- Moore, J.K. (2002). Maturation of human auditory cortex: Implications for speech perception. *The Annals of Otology, Rhinology, & Laryngology*, 111, 7–10.
- Oller, D.K., & Eilers, R.E. (1988). The role of audition in infant babbling. *Child Development*, 59, 441–449.

- Olmsted, D.L. (1966). A theory of the child's learning of phonology. *Language*, 42, 531–535.
- Polka, L., & Werker, J. (1994). Developmental changes in perception of non-native vowel contrasts. *Journal of Experimental Psychology: Human Perception and Performance*, 20, 421–436.
- Shriberg, L., & Kent, R. (2003). *Clinical phonetics* (3rd ed.). Boston: Allyn & Bacon.
- Smith, A., Goffman, L., & Stark, R.E. (1995). Speech motor development. *Seminars in Speech and Language*, 16, 87–99.
- Smith, B.L. (1982). Some observations concerning premeaningful vocalizations of hearing-impaired infants. *Journal of Speech and Hearing Disorders*, 47, 439–441.
- Smith, B.L., & Oller, D.K. (1981). A comparative study of premeaningful vocalizations produced by normal and Down's syndrome infants. *Journal of Speech and Hearing Disorders*, 46, 46–51.
- Stoel-Gammon, C., & Cooper, J.A. (1984). Patterns of early lexical and phonological development. *Journal of Child Language*, 11, 247–271.
- United States Census Bureau. (n.d.). United States census 2000. Retrieved November 6, 2004, from <http://quickfacts.census.gov/qfd/states/08000.html>
- Vihman, M.M. (1996). *Phonological development*. Cambridge, MA: Blackwell Publishers.
- Yavas, M. (1998). *Phonology development and disorders*. San Diego, CA: Singular Publishing Group.
- Yoshinaga-Itano, C., & Apuzzo, M.L. (1998). The development of deaf and hard of hearing children identified early through the high-risk registry. *American Annals of the Deaf*, 143, 416–424.
- Yoshinaga-Itano, C., & Sedey, A. (2001). Early speech development in children who are deaf or hard of hearing: Interrelationships with language and hearing. In C. Yoshinaga-Itano & A. Sedey (Eds.), *Language, speech, and social-emotional development of children who are deaf or hard of hearing: The early years* [Monograph]. *The Volta Review*, 100, 181–211.
- Yoshinaga-Itano, C., Stredler-Brown, A., & Jancosek, E. (1992). From phone to phoneme: What can we understand from babble. *The Volta Review*, 94, 283–314.

The Role of Age and Oral Lexical Competence in False Belief Understanding by Children and Adolescents With Hearing Loss

Antonia González, Ph.D., Inmaculada Quintana, Ph.D.,
Carmen Barajas, Ph.D., and María José Linero, M.Sc.

In the past decade, most studies have reported that children who are deaf and hard of hearing who have parents with typical hearing experience a serious delay in the understanding of false belief. False belief understanding consists of the ability to infer that someone else believes that something is true when one knows it to be wrong. This ability has been considered a reference point in studying the development of theory of mind (ToM). The main aim of this work is to evaluate the relationship between age, oral language and understanding of false belief in subjects who are deaf and hard of hearing. We investigated the relationship between age and ToM when the maximum age of participants included in the study was increased to 19 years. We also studied the relationship between the oral linguistic level of the participants and their performance in ToM. The sample consisted of 54 participants who are deaf and hard of hearing (ages 6 to 19 years) and who come from families with typical hearing. The results show that only from age 14 years onward is there a high percentage of success in the resolution of false belief tasks. Besides age, the level of oral lexical competence

Antonia González, Ph.D., is an assistant professor of developmental psychology in the departament of developmental and educational psychology, University of Málaga (Spain), who specializes in language development and deafness. Inmaculada Quintana, Ph.D., is an assistant professor of development disabilities in the departament of developmental and educational psychology, University of Málaga (Spain) who specializes in development disabilities. Carmen Barajas, Ph.D., is an assistant professor of developmental psychology in the departament of developmental and educational psychology, University of Málaga (Spain) who specializes in cognitive development. María José Linero, M.Sc., is an assistant professor of language disabilities in the departament of developmental and educational psychology, University of Málaga (Spain) who specializes in language development and deafness.

is a variable that can help to explain the differences in ToM performance among individuals with hearing loss.

Introduction

The ability to understand that other people have mental states such as beliefs, intentions, memories and desires that may differ from one's own is known as theory of mind (ToM) (Premack & Woodruff, 1978). How and when children acquire the ability to think about and take account of others' mental states, as well as their own, is a topic of much interest to developmental psychologists. It is easy to see why: The ability influences almost every aspect of human social behavior. To be a good conversationalist, to participate appropriately in a full range of social relationships and to reason intelligently about behavior, an individual must know something about others, how they are similar to and different from oneself, what they are likely to know and how they are likely to behave in various circumstances. ToM enables people to make sense of and predict others' behavior; as such, its development is an important key to social, communicative and affective life.

There is no doubt that the main indicator of the development of ToM is the understanding of false belief: people's ability to infer that someone else believes that something is true when they themselves know it to be wrong. In the false belief paradigm, a story is told in which the protagonist is deceived by the situation and holds a false belief regarding an object's whereabouts. The evaluated children have to respond to questions about what the main character would do in the face of his or her false beliefs. Wimmer and Perner (1983) were the first to design this task within this paradigm. This task later was adapted by Baron-Cohen, Leslie and Frith (1985) and became the classic false belief task. Specifically, the task is as follows: A female puppet (Sally) hides a marble in a basket and leaves the scene. While she is gone, the other female puppet (Anne) shifts the marble to a covered box. Sally returns, and the child taking the test is asked, "Where will Sally look for her marble?" (correct answer: basket) followed by two control questions: "Where is the marble really?" and "Where did Sally put the marble in the beginning?" to make sure both that the child has taken note of the marble's new location and that he or she is able to recall the original hiding place.

This ability is used to attribute false beliefs to others in a great variety of everyday situations. For example, after arriving home without warning, Juan knows that Maria thinks he is still at work; or Arturo, after eating his sister Lucia's sweets, knows that when she looks at the box of sweets she will think that it is full and be angry when she finds it empty.

There is an almost unanimous consensus that children with typical development reach this ability around the age of 4 ½ (Gopnik & Astington, 1988; Perner, Leekam & Wimmer, 1987; Wellman, Cross & Watson, 2001; Wimmer & Perner, 1983). However, in certain conditions of task presentation, correct

performances are observed in younger children when resolving classic false belief tasks (Yazdi, German, Defeyter, & Siegal, 2005). In addition to this observation, other abilities associated with understanding false belief, such as attributing to a person the disposition to carry out an action and predicting his or her behavior, also can be observed in even younger children, including those 13 ½ months old (Song, Baillargeon, & Fisher, 2005).

From a theoretical sociocultural viewpoint, social experience and the development of language together are indispensable factors for the development of ToM (Garfield, Peterson, & Perry, 2001). Within this perspective, three types of work can be differentiated. The first type focuses on the daily interactions of children in the home by describing activities and conversations (between adults and children and between peers) that stimulate the development of ToM (Brown, Donelan-McCall, & Dunn, 1996; Dunn, 1994, 1999). The second type shows the relationships between the language used by adults when telling stories and playing with their children and the development of ToM in the children (Astington & Baird, 2005; de Rosnay, Pons, Harris, & Morrell, 2004; Ruffman, Perner, & Parkin, 1999; Ruffman, Slade, & Crowe, 2002). The third type focuses on the relationship between child linguistic development and the development of ToM (Astington & Jenkins, 1999; De Villiers & Pyers, 2002; Happé, 1995; Ziatas, Durkin, & Pratt, 1998).

The first research into the third type was done by Happé (1995), who found that the verbal age of children, as measured by a receptive vocabulary test, is predictive of success in false belief tasks in both children with autism and children with typical development, and that this effect is independent of chronological age. Recent research has provided more specific data concerning the relationships between language and ToM. Ziatas et al. (1998) found a relationship between the understanding of false belief and the use of mental terms (thinking, knowing and guessing) in children with autism, children with Asperger syndrome, children with specific language impairments and children with typical development. Astington and Jenkins (1999) carried out a longitudinal study and verified the effect of early linguistic development (lexicon and syntax) on the development of ToM. De Villiers and Pyers (2002) established a relationship between children's proficiency in the syntax of complementation and their performance in false belief tasks. They argued that the child needs the full syntax of mental verbs plus sentential complements in order to represent in his or her own mind the belief states of other people, not simply to report about them in speech.

Slade and Ruffman (2005) examined the effect of semantic and syntactic development separately in the development of ToM in preschool children. The authors concluded that ToM has more to do with general linguistic ability than syntax or semantics, and that this ability contains both semantic and syntactic components. However, they acknowledged that receptive vocabulary tests, such as the British Picture Vocabulary Scale (BPVS) or Peabody

Picture Vocabulary Test (PPVT), usually correlate with ToM and found that semantic measures correlate highly with those of syntax and age.

The study of the development of ToM in children who are deaf and hard of hearing has been carried out in tandem with the works cited. In their pioneering study, Peterson and Siegal (1995) reported that a very low percentage of children between the ages of 8 and 13 who are deaf and hard of hearing succeeded in the classic false belief task from Baron-Cohen et al. (1985) described earlier. This low performance was attributed to a deficit in ToM, resulting from the lack of access to communication between adults with typical hearing and children who are deaf and hard of hearing in the home (conversational hypothesis). The concept of deficit was endorsed by the lack of significant differences between the age of the group of children who succeeded in the false belief task and the age of the group who did not succeed.

Later studies did not find any relationship between the age of children who are deaf and hard of hearing and their performance on ToM tasks (Courtin, 2000; Courtin & Melot, 1998; Woolfe, Want, & Siegal, 2002). All of these studies share the fact that their samples were made up of children younger than 8 years old with hearing loss, which led to the idea that it would be necessary to evaluate older children with hearing loss for age to be a predicting variable. The research of Russell et al. (1998), using a sample of participants up to 16 years old who are deaf and hard of hearing, helps to clarify this point, as they found that improvements in false belief understanding begin from the age of 13 onward.

Subsequently, Peterson and Siegal (1999) found correlations between age and success in false belief tasks with a wider sample of children who are deaf and hard of hearing than in their first study. These data led them to switch from their concept of deficit to delay. Another key contribution by Peterson and Siegal (1999) was the introduction of the variable "linguistic development." The authors found a correlation between the linguistic ability of children who are deaf and hard of hearing who have parents with typical hearing and their success in false belief tasks. Nevertheless, it is necessary to point out that the role of language is relative because, when its effect is parcelled out, the differences in ToM performance observed between children with hearing loss whose parents have hearing loss and use sign language (native signers) and children with hearing loss whose parents have typical hearing (non-native signers) continue to be significant.

The conclusions reached by Woolfe et al. (2002) are very similar to those of Peterson and Siegal (1999). When two groups of children who are deaf and hard of hearing and who have similar "signing" age are compared, native signers still present better performance. In this study, as in others (Courtin, 2000; Jackson, 2001; Peterson, Wellman, & Liu, 2005; Woolfe et al., 2002), children with hearing loss who were native signers did not have difficulties in solving false beliefs tasks. These results led the authors to support the

conversational hypothesis that Peterson and Siegal (1995) had already formulated.

De Villiers and de Villiers (2000) based their work on deafness and ToM on a linguistic rather than a conversational hypothesis. Starting from the hypothesis that children with hearing loss who use spoken language have limited language, their vocabulary and syntax at the age of 4 are still insufficient to refer to others' beliefs. These authors evaluated the understanding of false belief using, besides classic tasks, a task that they called "less verbal." This task is basically a version of the unexpected content concept, but rather than verbally reporting what someone thinks is in the box, the child must predict whether a character is surprised or not when he sees the unusual content, choosing the right facial expression. Their results showed that the utterance of complex sentences, including verbs referring to cognitive states, predicted the performance of children who are deaf and hard of hearing in their sample in verbal and "less verbal" false belief tasks and that this relation was maintained even when the effect of age was parcelled out. However, the authors did not find any correlation between children's verbal age, measured with the Peabody Picture Vocabulary Test-Revised (PPVT-R) and their performance in false belief tasks.

Jackson's study (2001) provided data that supported the hypothesis of de Villiers and de Villiers (2000). To evaluate the linguistic level of native and non-native signing children who are deaf and hard of hearing, she used the British Sign Language receptive skills test of Herman, Holmes and Woll (1999). The receptive linguistic level of children with hearing loss who use spoken language was assessed with the BPVS. These results showed a positive correlation between language and ToM, which was maintained for children with hearing loss who either were non-native signers or who used spoken language, even when the effect of age was parcelled out. Jackson indicated, in line with de Villiers and de Villiers (2000), that properties specific to language are required for ToM development over and above experiences related to age and neurological maturation.

The study by Pyers (2005) also supports this thesis. In a study of Nicaraguan adolescents and adults who are deaf and hard of hearing, the variable associated with false belief understanding was the lexical and syntactic development of their sign language, more than age and experience combined. Pyers (2005) defended the view that language is key to the uniquely human capacity to represent the thoughts and belief of others; without adequate language, humans fail to acquire a mature ToM.

In summary, the main findings of our review are as follows:

1. Most children with hearing loss who are non-native signers present very low performance in false belief understanding.
2. When the participants are under 13 years old, few studies found correlations between age and performance in false belief tasks.

3. There is a relationship between the dominance of sign language and the understanding of false belief in children who are deaf and hard of hearing who use sign language and also between oral linguistic development and the understanding of false belief in children with hearing loss who use spoken language.

The first of these conclusions is widely supported by empirical studies, but little work has been done on the second and third.

In addition, most available studies include children up to the age of 10; a few deal with older children up to 13 years old and, on rare occasions, up to 16 years old. However, as already pointed out, only with a wide age range can any significant relationship between age and ToM be found. Furthermore, we found just one study that included a sample of Spanish participants who were deaf and hard of hearing (Figueras-Costa & Harris, 2001); however, the ToM assessment was conducted using a nonverbal version of standard false belief tasks.

The main aim of our study was to evaluate the relationship between age, language and understanding of false belief in participants with hearing loss who learned sign language at a later age. We investigated the relationship between age and ToM when the age of participants includes individuals up to 19 years old and the relationship between the linguistic level of the participants and their performance in ToM.

Methods

Participants

The original sample consisted of 61 participants. Seven participants were excluded because of low IQ or difficulties in understanding ToM tasks (see Assessment), so the final sample consisted of 54 participants. The participants' ages ranged from 6 to 19 years; 27 were between 6 and 13 years old (mean age 9.9) and the remaining 27 were between 14 and 19 years (mean age 16.3). All of the children had parents with typical hearing, most of whom used the oral modality with their children. The participants attended a school for children who are deaf and hard of hearing with an oral approach in Málaga (Spain). These children acquired sign language at a later stage through contact with peer children at school. Around 81.5% began to attend a special school when they were older than 4 years of age. Of these, 27.3% began when they were older than 7 years of age, having previously been educated in ordinary oralist state schools. All of the subjects presented prespeech deafness with losses of 70 dB or more in their better ear (38.9% severe losses and 61.1% profound losses), and none had a cochlear implant. Of the participants,

50% were male and 50% were female; 50% had a low sociocultural level, 43.1% an average level and only 6.9% a high one.

Assessment

All participants were evaluated with the Leiter International Performance Scale–Revised (Leiter-R) for nonverbal intelligence (Roid & Miller, 1997) to verify that they did not have associated intellectual deficiencies. Four participants were excluded from the original sample because they scored lower than 70 (mental deficiency according to Leiter-R). Furthermore, all the participants were evaluated in the following variables:

1. Oral Lexical Competence: This was evaluated through the Test de Vocabulario en Imágenes Peabody (TVIP), the Hispanic-American adaptation by Dunn, Padilla, Lugo and Dunn (1986) of PPTV-R by Dunn and Dunn (1981). The PPTV-R already has been used in some studies dealing with the oral language–ToM relationship in people who are deaf and hard of hearing, for example, in de Villiers and de Villiers (2000). This test was chosen for several reasons: Its application fit the age range of our sample; it was easy and fast to administer and to correct; and it provided the verbal age of subjects. The test assesses the level of oral receptive vocabulary of subjects by having participants identify words through lip-reading and then point to the drawing from a set of four that corresponds to the target word. Words were articulated correctly and clearly to ensure recognition via lip-reading.
2. Theory of Mind: Two classic false belief tasks were used. They were narrated in sign language (because all the study participants preferred this modality), and each sequence was acted out. To ascertain whether, in addition to attributing a false belief to others, participants understood what motivates it, we introduced a justification question to the classic experimental question.
 - 2.1. *Change of Location* (“marble task” of Baron-Cohen et al., 1985). The task was adapted by including a character of each sex to facilitate the participants’ identification with the characters in the story (as in Peterson & Siegal, 1995). In this way, the participants could resort to the signs for *boy* or *girl* if they forgot a character’s name. The Spanish names Paco and María also were substituted. In brief, the procedure consisted of two trials, each involving two puppets. In the first trial, a puppet, María, hides a marble in a basket and leaves the scene. While she is gone, the other puppet, Paco, shifts the marble to a covered box. María returns, and the child taking the test is asked, “Where will María look for her marble?” (correct answer: basket) and “Why?” followed by two control questions: “Where is

the marble really?" and "Where did María put the marble in the beginning?"

- 2.2. *Unexpected Contents Task* ("Smarties task" of Hogrefe, Wimmer, & Perner, 1986). Following the modification introduced by Gopnik and Astington (1988), we used different container-material pairs to prevent children from communicating information about recipients and contents to each other. Each participant was presented with just one container-material pair. All containers used had clear pictures of the expected contents and were familiar to the children. In brief, the task involved a misleadingly familiar sweets container (Smarties) that actually held pencils. The participants were shown a container and asked what they thought was inside. All participants replied "Smarties", "chocolates" or "sweets." They were then shown that the container really contained pencils. Then, the participants were asked what a naive classmate would say on first seeing the closed container and why, followed by two control questions: "What was really in the container?" and "What did you first think was in the container?"

In each story the participants replied to two experimental questions: The first one (attribution question) helped evaluate whether the participants attributed a false belief to the character; the second (justification question) helped verify whether participants understood why the character had such a false belief. As in Perner et al. (1987), we introduced a series of control questions to ensure that the participants understood and recalled the essential aspects of the story. Only the participants who replied correctly to the control questions were entered into the sample. Three participants (age 6) were excluded from the original sample because they failed the control questions.

Taking into account the two tasks and the two experimental questions per task, each participant could attain one of the following overall scores:

- 0: He/she did not answer any of the questions correctly.
- 1: He/she replied correctly to the attribution question in one task but did not justify it correctly.
- 2: He/she replied correctly to the two questions in one task.
- 3: He/she replied correctly to the two questions in the two tasks.

There were no cases in which the two tasks were successfully done and only one of them was correctly justified. The scores are shown in Table 1.

The tests were administered as follows: Leiter-R in the first session and TVIP and ToM tasks in the second. All the tasks were administered by the researchers, with the assistance of an interpreter in sign language for ToM tasks.

Table 1. Distribution of ToM Scores Across All Subjects

	ToM			
	0	1	2	3
N	10	6	6	32
%	18.5	11.1	11.1	59.3

Results

Before analyzing the relationships between age, language and ToM performance in the children and adolescents with hearing loss in this study, we tested whether other variables specific to the sample (sex, level of hearing loss and sociocultural level) had an effect on their performance in false belief tasks. To this end, we carried out a one-way analysis of variance (univariate), taking as the dependent variable the ToM score, and as the independent variable each of the cited variables. Upon comparing the groups of participants defined by these variables, we found no significant differences in ToM scores: sex, $F(1,52) = 0.030$, $p = \text{ns}$; level of hearing loss, $F(1,52) = 0.023$, $p = \text{ns}$; and sociocultural level, $F(2,51) = 0.917$, $p = \text{ns}$. Therefore, we considered it unnecessary to control for the effect of sex, hearing loss level, or sociocultural level of the participants in subsequent analyses.

To analyze the effects of the variables “age” and “oral linguistic level” on ToM, we carried out a 3×2 (age groups $\times$ oral level groups) univariate analysis of variance. For the variable “age”, participants were divided into three groups: young (age range 6–10, mean age 8.6), middle (age range 11–13, mean age 12) and old (age range 14–19, mean age 16.3). This age choice was based on the fact that most studies included participants whose age was up to 10 years old, that only one study included those age 16 (finding improvement from the age of 13 onward) and that no studies included participants over age 16. We obtained oral linguistic level by converting the direct scores into equivalent ages provided by TVIP. In light of the low scores obtained by participants in oral linguistic level, we assigned them to two groups: 4 years old or younger and older than 4 years old.

Of the two factors included in the model (age and oral linguistic level), the groups defined by the variable age, $F(2,48) = 5.83$, Mean Squared Error (MSE) = 4.03, $p = 0.005$, and by the variable oral linguistic level, $F(1,48) = 12.16$, MSE = 8.41, $p = 0.001$, had significantly different scores in ToM. There was no significant interaction between age and oral linguistic level, $F(2,48) = 2.43$, MSE = 1.68, $p = 0.09$.

Multiple comparisons with the Games-Howell post-hoc tests ($p < 0.05$) were carried out to help us learn in more detail whether there were significant differences in ToM between the three age groups. Data regarding the age

Table 2. ToM Performance According to Age

Age	Theory of Mind (ToM)						S.D.
	N	0	1	2	3	Mean	
6–10 years	14	8 (57.2)	3 (21.4)	0 (0)	3 (21.4)	0.86	1.231
11–13 years	13	1 (7.7)	3 (23)	4 (30.8)	5 (38.5)	2	1.000
14–19 years	27	1 (3.7)	0 (0)	2 (7.4)	24 (88.9)	2.81	0.62

Note. Percentage in parentheses.

factor revealed significant differences in ToM scores among all the age groups, and the differences were always in favor of the older group. The descriptive data above provide more detail on these differences.

As can be seen in Table 2, the participants usually did not succeed in either of the tasks up to 10 years of age. Between the ages of 11 and 13, most participants (69.3%) succeeded in and correctly justified one or two of the tasks. From the age of 14, the vast majority of participants (88.9%) attained the highest scores. The mean scores in ToM increased in relation to age. Furthermore, the most heterogeneous group regarding the mean is the one with a younger chronological age (6–10 years).

To provide further details regarding the effect of oral linguistic level on ToM, we grouped the descriptive data of the participants' performance in ToM presented as a function of their oral linguistic level.

As can be seen in Table 3, more than half the participants with an oral linguistic level equivalent to 4 years old or younger (57.5%) did not reach the highest score, whereas the majority of children with oral linguistic levels above age 4 (85.7%) reached the highest score. The mean scores in ToM increased in relation to the linguistic age of the participants, and the most heterogeneous group regarding the mean was the one with a low oral age.

Also provided is descriptive data on the participants' performance in ToM grouped as a function of their age and oral linguistic level.

As seen in Figure 1, the participants with a higher oral linguistic level performed better in ToM than did those with a lower level, independently of whether they belong to the young, middle or older group. This difference in

Table 3. ToM Performance According to Oral Level

Oral Level	Theory of Mind (ToM)						S.D.
	N	0	1	2	3	Mean	
≤4 years	33	10 (30.3)	5 (15.1)	4 (12.1)	14 (42.5)	1.67	1.315
>4 years	21	0 (0)	1 (4.8)	2 (9.5)	18 (85.7)	2.81	0.112

Note. Percentage in parentheses.

Figure 1. *Note.* ToM = Theory of Mind

ToM performance is very pronounced in the young group and decreases in the middle and older groups; the figure shows only a small difference between scores in ToM between those with higher and lower oral levels among the older group.

Discussion

The first aim of our study was to test whether the relationship between age and ToM identified by most authors changes when the age of the subjects studied is expanded to include those up to the age of 19. The first observation of interest is that the percentage of participants attaining maximum performance was high (59%) compared with other studies; for example, in studies by Peterson and Siegal (1998, 1999), with a sample of 5- to 12-year-old participants, the percentage solving false belief tasks did not reach 50%; in the research by Russell et al. (1998), with participants ranging from 4 to 16 years old, only 28% were successful in the task. This difference seems due to the fact that the age range of our sample was expanded, thereby increasing the percentage of participants who succeeded in the tasks.

Another finding of interest is that 22% of the participants showed a certain ability to understand false belief even if they did not attain maximum performance (those obtaining intermediate scores of 1 or 2). Moreover, our data indicate that succeeding in the two tasks before the age of 11 was infrequent, whereas intermediate scores abounded among 11- and 13-year-old children. This information is of note because the data can lead to the idea that the

understanding of false belief by participants who are deaf and hard of hearing can be progressive, instead of dichotomous (pass/fail) as was considered conventionally. It seems that, first, the attribution is made, then the participant reflects on it and justifies it accordingly, and finally, the ability is generalized to different situations.

This partial understanding of false belief observed in some participants led to further consideration of research on this issue.

First, it is possible that, before the conceptual change taking place that enables full development of ToM, defended by Wellman et al. (2001), children who are deaf and hard of hearing may pass through a prior transitional period in the same way as children with typical hearing. The children who are deaf and hard of hearing in the sample who scored a 1 (those carrying out false belief attributions in a task but not managing to explain their attributions) may be in this transitional period.

Second, there seems to be a sequence of steps in ToM evolution in preschoolers who are developing typically that is also observed in subjects who are deaf and hard of hearing, albeit at older ages than among children with typical hearing (Peterson et al., 2005). This sequence evolves from understanding diverse desires to understanding hidden emotions. Other achievements occur between these two steps, such as understanding diverse belief, knowledge access and false belief. Our data suggest that if there are general stages in ToM, there could also be a specific sequence in the understanding of false belief. Within the setting of this sequence, before attributing false beliefs in a general and justified way, some children who are deaf and hard of hearing manage to attribute false belief correctly in some of the tasks without apparent prior analysis or make attributions and offer justifications in one task but not in others.

Regarding the age at which better performance in ToM is found, by the age of 14, the participants in the sample who were deaf and hard of hearing reached a level of performance similar to that of children 4 to 6 years old with typical hearing; 88% of the participants between 14 and 19 years old reached the maximum scores. In fact, all participants older than 16 reached the maximum scores. In this regard, our data confirm the findings of Russell et al. (1998) who reported participants improving their performance in ToM from the age of 13 onwards. It is unclear what competencies might be present in older children who are deaf and hard of hearing that facilitate their development of a mature ToM. Up to the age of 14, communicative experiences with their families, peers and teachers, aided by their own linguistic development, probably have contributed to achieving this development in ToM. This observation leads to the idea that it may be possible to accelerate the development of ToM.

The other aim of this study was to test whether the level of oral lexical competence had an effect on ToM performance among the participants. The results show that oral linguistic age is a significant variable in explaining the

variance in participants' ToM performance. In this regard, the data are consistent with those of Happé (1995), who reported that oral linguistic age (as measured with the PPVT) correlated with performance in ToM in a sample of autistic children and children who are developing typically. The data also agree with those of Jackson (2001), who found positive correlations in children with hearing loss who use spoken language between ToM and linguistic age measured through receptive vocabulary (BPVS). However, the data do not match those of de Villiers and de Villiers (2000), who did not report a significant relationship between the oral linguistic age of children who are deaf and hard of hearing (using PPVT-R) and the understanding of false belief. A reason for this discrepancy might be found in the age of the participants from their sample (4 to 9 years old), who were younger than those in the current study and represent a more limited age range. It is likely that, at these ages, there is not enough variability in the oral linguistic level of participants to explain the variance in the performance of false belief tasks.

It has to be pointed out that because we are dealing with children and adolescents who are deaf and hard of hearing but who live in families with typical hearing, it seems that the experiences suited to stimulating the development of ToM are provided by the oral modality most widely used in the communicative exchanges. In this study, the differences in ToM performance related to the oral linguistic level of the participants were significant when participants with a linguistic age of 4 or younger were compared to those older than 4. As indicated in the introduction, children with typical hearing achieve false belief understanding at 4 ½ years old; thus, it is not surprising that better ToM performance was found in the participants in this study who were deaf and hard of hearing when their oral lexical competence was higher than 4 years. We think that reaching a linguistic age higher than 4 years old in TVIP involves wider linguistic abilities than simple lexical understanding. Other authors share this assumption; Ruffman, Slade, Rowlandson, Rumsey and Garnham (2003), when analyzing the relationship between the measures in receptive vocabulary and ToM, pointed out the possibility that these measures of lexical development also reflect syntactic development. Happé (1995) considered that the verbal age obtained with the PPVT can represent a measure of general ability. It seems that these abilities facilitate the understanding of false belief, since they make possible communication and social interaction and provide the necessary representation tools.

Therefore, chronological age—involving a personal history of social experiences—and level of oral lexical competence are variables that explain the differences in ToM performance in participants who are deaf and hard of hearing. These data confirm the idea that the delay in ToM traditionally attributed to the population who are deaf and hard of hearing arises from the following fact: The communicative experiences and the language development necessary for the development of ToM converge at a later age in children who are deaf and hard of hearing than in children with typical hearing.

This conclusion agrees with that of de Rosnay et al. (2004), who stated that children with better linguistic development can include and incorporate information about mental states from communicative experiences better than can those who are at an earlier stage of language development.

In light of the practical importance of a functional ToM as a social tool facilitating communication and social interaction with others, children who are deaf and hard of hearing are likely to face communication problems in everyday social situations, in addition to those stemming from difficulties with speech and language (Russell et al., 1998). This fact indicates the importance of a suitable intervention. In contrast to younger children, for whom early linguistic intervention could ensure a typical development of ToM, it seems that intervention in older children—in particular, the study participants who, even at age 15, still had not reached false belief understanding—should be based on situations that simulate the ordinary interaction contexts in which the development of ToM occurs (e.g., solving conflicts or understanding other peoples' intentions, desires and feelings).

Finally, other variables, such as the quality of the interaction in the family, social experience and sign linguistic development, should be taken into account to complete the analysis of the factors accounting for inter-individual differences in the development of ToM in people who are deaf and hard of hearing. It would be of interest to study the development of ToM in people who are deaf and hard of hearing beyond false belief understanding in contexts such as attributing complex communicative intentions like irony, lying or hiding emotions.

Acknowledgment

This study was carried out with financial help from the Ministry of Science and Technology of Spain (reference BSO 2000-1202). We express our gratitude to the teachers and students from the school for children who are deaf and hard of hearing in Málaga for their collaboration.

References

- Astington, J.W., & Baird, J.A. (2005). Why language matters for theory of mind. Oxford: University Press.
- Astington, J.W., & Jenkins, J.M. (1999). A longitudinal study of the relation between language and theory-of-mind development. *Developmental Psychology*, 35(5), 1311–1320.
- Baron-Cohen, S., Leslie, A.M., & Frith, U. (1985). Does the autistic child have theory of mind? *Cognition*, 21, 37–46.
- Brown, J.R., Donelan-McCall, N., & Dunn, J. (1996). Why talk about mental states? The significance of children's conversations with friends, siblings, and mothers. *Child Development*, 67, 836–849.

- Courtin, C. (2000). The impact of sign language on the cognitive development of deaf children: The case of theories of mind. *Journal of Deaf Studies and Deaf Education*, 5(3), 266–276.
- Courtin, C., & Melot, A. (1998). Development of theories of mind in deaf children. In M. Marschark and M.D. Clark (Eds.), *Psychological Perspectives on Deafness*. Volume 2. London: LEA.
- De Rosnay, M., Pons, F., Harris, P., & Morrell, J. (2004). A lag between understanding false belief and emotion attribution in young children: Relationships with linguistic ability and mothers' mental-state language. *The British Journal of Developmental Psychology*, 22(2), 197–219.
- De Villiers, J., & de Villiers, P. (2000). Linguistic determinism and the understanding of false beliefs. In P. Mitchell & K. Riggs (Eds.), *Children's reasoning and the mind*. New York: Psychology Press.
- De Villiers, J., & Pyers, J.E. (2002). Complements to cognition: A longitudinal study of the relationship between complex syntax and false-belief understanding. *Cognitive Development*, 17, 1037–1060.
- Dunn, J. (1994). Changing minds and changing relationships. In C. Lewis & P. Mitchell (Eds.), *Children's early understanding of mind*. London: LEA.
- Dunn, J. (1999). Making sense of the social world: Mindreading, emotion and relationships. In D.P. Zelazo, J.N. Astington, & D.R. Olson (Eds.), *Developing theories of intention*. London: LEA.
- Dunn, L.L. M., & Dunn, L.M. (1981). Peabody Picture Vocabulary Test-Revised. Circle Pines, MN.: American Guidance Service.
- Dunn, L.L. M., Padilla, E.R., Lugo, D.E., & Dunn, L.M. (1986). Test de Vocabulario en Imágenes Peabody-R. Circle Pines, MN.: American Guidance Service.
- Figueras-Costa, B., & Harris, P. (2001). Theory of mind development in deaf children: A non-verbal test of false-belief understanding. *Journal of Deaf Studies and Deaf Education*, 6(2), 92–102.
- Garfield, J.L., Peterson, C., & Perry, T. (2001). Social cognition, language acquisition and the development of the theory of mind. *Mind and Language* 16, 494–541.
- Gopnik, A., & Astington, J.W. (1988). Children's understanding of representational change and its relation to the understanding of false belief and the appearance-reality distinction. *Child Development*, 59, 26–37.
- Happé, F. (1995). The role of age and verbal ability in the theory of mind task performance of subjects with autism. *Child Development*, 66, 843–855.
- Herman, R., Holmes, S., & Woll, B. (1999). Assessing BSL development: Receptive skills test. London: Forest Bookshop.
- Hogrefe, J., Wimmer, H., & Perner, J. (1986). Ignorance versus false belief: A developmental lag in attribution of epistemic states. *Child Development*, 57, 567–582.
- Jackson, A.L. (2001). Language facility and theory of mind development in deaf children. *Journal of Deaf Studies and Deaf Education*, 6(3), 161–176.

- Perner, J., Leekam, S., & Wimmer, H. (1987). Three-year-olds' difficulty with false belief: The case for a conceptual deficit. *British Journal of Developmental Psychology*, 5, 125–137.
- Peterson, C., & Siegal, M. (1995). Deafness, conversation and theory of mind. *Journal of Child Psychology and Psychiatry*, 36, 459–474.
- Peterson, C., & Siegal, M. (1998). Changing focus on the representational mind: Deaf, autistic and normal children's concepts of false photos, false drawings and false beliefs. *British Journal of Developmental Psychology*, 16, 301–320.
- Peterson, C., & Siegal, M. (1999). Representing inner worlds: Theory of mind in autistic, deaf, and normal hearing children. *Psychological Science*, 10(2), 126–129.
- Peterson, C., Wellman, H.M., & Liu, D. (2005). Steps in theory-of-mind development for children with deafness or autism. *Child Development*, 76(2), 502–517.
- Premack, D., & Woodruff, G. (1978). Does the chimpanzee have a theory of mind? *Behavioral and Brain Sciences*, 1, 515–526.
- Pyers, J.E. (2005). The relationship between language and false-belief understanding: Evidence from learners of an emerging sign language in Nicaragua. *Dissertation Abstracts International*, 66(2B), 1200.
- Roid, G.H. & Miller, L.J. (1997). *Leiter International Performance Scale-Revised*. Wood Dale, IL: Stoelting Co.
- Ruffman, T., Perner, J., & Parkin, L. (1999). How parenting style affects false belief understanding. *Social Development*, 8, 395–411.
- Ruffman, T., Slade, L., & Crowe, E. (2002). The relation between children's and mothers' mental state language and theory-of-mind understanding. *Child Development*, 73(3), 734–751.
- Ruffman, T., Slade, L., Rowlandson, K., Rumsey, C., & Garnham, A. (2003). How language relates to belief, desire and emotion understanding. *Cognitive Development*, 18, 139–158.
- Russell, P., Hosie, J., Gray, C., Scott, C., Hunter, N., Banks, J., & Macaulay, M. (1998). The development of theory of mind in deaf children. *Journal of Child Psychology and Psychiatry*, 39, 903–910.
- Slade, L., & Ruffman, T. (2005). How language does (and does not) relate to theory of mind: A longitudinal study of syntax, semantics, working memory and false belief. *The British Journal of Developmental Psychology*, 23, 117–141.
- Song, H., Baillargeon, R., & Fisher, C. (2005). Can infants attribute to an agent a disposition to perform a particular action? *Cognition*, 98(2), B45–B55.
- Wellman, H.M., Cross, D., & Watson, J. (2001). Meta-analysis of theory-of-mind development: The truth about false belief. *Child Development*, 72(3), 655–684.
- Wimmer, H., & Perner, J. (1983). Beliefs about beliefs: Representation and the

- constraining function of wrong beliefs in young children's understanding of deception. *Cognition*, 13, 103–128.
- Woolfe, T., Want, S.C., & Siegal, M. (2002). Signposts to development: Theory of mind in deaf children. *Child Development*, 73(3), 768–778.
- Yazdi, A.A., German, T.P., Defeyter, M.A., & Siegal, M. (2005). Competence and performance in belief-desires reasoning across two cultures: The truth, the whole truth and nothing but the truth about false belief? *Cognition*, 100(2), 343–368.
- Ziatas, K., Durkin, K., & Pratt, C. (1998). Belief term development in children with autism, Asperger syndrome, specific language impairment, and normal development: Links to theory of mind development. *Journal of Child Psychology and Psychiatry*, 39, 755–763.

Joint Committee on Infant Hearing

Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs

Reproduced with permission from *Pediatrics*, 120(4), pp. 898–921.
Copyright © 2007 by the American Academy of Pediatrics (AAP).

TABLE OF CONTENTS

ABBREVIATIONS

THE POSITION STATEMENT

JCIH 2007 POSITION STATEMENT UPDATES

1. Definition of Targeted Hearing Loss
2. Hearing Screening and Rescreening Protocols
3. Diagnostic Audiology Evaluation
4. Medical Evaluation
5. Early Intervention
6. Surveillance and Screening in the Medical Home
7. Communication
8. Information Infrastructure

I. BACKGROUND

II. PRINCIPLES

III. GUIDELINES FOR EARLY HEARING DETECTION AND INTERVENTION PROGRAMS

A. ROLES AND RESPONSIBILITIES

B. HEARING SCREENING

1. Screening Protocols in the Well-Baby Nursery
2. Screening Protocols in the NICU
3. Conveying Test Results
4. Outpatient Rescreening or Infants Who Do Not Pass the Birth Admission Screening

C. CONFIRMATION OF HEARING LOSS IN INFANTS REFERRED FROM UNHS

1. Audiologic Evaluation
 - i. Evaluation: Birth to 6 Months of Age
 - ii. Evaluation: 6 to 36 Months of Age
 - iii. Other Audiologic Test Procedures

2. Medical Evaluation
 - i. Pediatrician/Primary Care Physician
 - ii. Otolaryngologist
 - iii. Other Medical Specialists

D. EARLY INTERVENTION

1. Principles of Early Intervention
 - i. Designated Point of Entry
 - ii. Regular Developmental Assessment
 - iii. Opportunities for Interaction With Other Parents of Children With Hearing Loss
 - iv. Opportunities for Interaction With Individuals Who Are Deaf or Hard of Hearing
 - v. Provision of Communication Options
 - vi. Skills of the Early Intervention Professional
 - vii. Quality of Intervention Services
 - viii. Intervention for Special Populations of Infants and Young Children

2. Audiologic Habilitation

3. Medical and Surgical Intervention

4. Communication Assessment and Intervention

E. CONTINUED SURVEILLANCE, SCREENING, AND REFERRAL OF INFANTS AND TODDLERS

F. PROTECTING THE RIGHTS OF INFANTS AND FAMILIES

G. INFORMATION INFRASTRUCTURE

H. BENCHMARKS AND QUALITY INDICATORS

1. Quality Indicators for Screening
2. Quality Indicators for Confirmation of Hearing Loss
3. Quality Indicators for Early Intervention

IV. CURRENT CHALLENGES, OPPORTUNITIES, AND FUTURE DIRECTIONS

1. Challenges
2. Opportunities for System Development and Research

V. CONCLUSIONS

ACKNOWLEDGEMENTS

APPENDIX 1. Risk Indicators Associated with Permanent Hearing Loss

APPENDIX 2: Algorithm for Hearing Screening

REFERENCES

ABBREVIATIONS: JCIH, Joint Committee on Infant Hearing; EHDI, early hearing detection and intervention; NICU, neonatal intensive care unit; ABR, auditory brainstem response; CMV, cytomegalovirus; ECMO, extracorporeal membrane oxygenation; AAP, American Academy of Pediatrics; MCHB, Maternal and Child Health Bureau; HRSA, Health Resources and Services Administration; NIH, National Institutes of Health; NIDCD, National Institute

on Deafness and Other Communication Disorders; CDC, Centers for Disease Control and Prevention; UNHS, universal newborn hearing screening; OAE, otoacoustic emissions testing; IFSP, individualized family service plan; OME, otitis media with effusion; DSHPSHWA, Directors of Speech and Hearing Programs in State Health and Welfare Agencies; GPRA, Government Performance and Results Act; OMB, Office of Management and Budgets.

THE POSITION STATEMENT

The Joint Committee on Infant Hearing (JCIH) endorses early detection of and intervention for infants with hearing loss. The goal of early hearing detection and intervention (EHDI) is to maximize linguistic competence and literacy development for children who are deaf or hard of hearing. Without appropriate opportunities to learn language, these children will fall behind their hearing peers in communication, cognition, reading, and social-emotional development. Such delays may result in lower educational and employment levels in adulthood.¹ To maximize the outcome for infants who are deaf or hard of hearing, the hearing of all infants should be screened no later than 1 month of age. Those not passing screening should have a comprehensive audiologic evaluation no later than 3 months of age. Infants with confirmed hearing loss should receive appropriate intervention no later than 6 months of age from health care and education professionals with expertise in hearing loss and deafness in infants and young children. Regardless of previous hearing-screening outcomes, all infants with or without risk factors should receive ongoing surveillance of communicative development beginning at 2 months of age during well-child visits in the medical home.² EHDI systems should guarantee seamless transitions for infants and their families through this process.

JCIH 2007 POSITION STATEMENT UPDATES

The following are highlights of updates made since the JCIH 2000 statement³:

1. Definition of Targeted Hearing Loss

- The definition has been expanded from congenital permanent bilateral, unilateral sensory, or permanent conductive hearing loss to include neural hearing loss (eg, “auditory neuropathy/dyssynchrony”) in infants admitted to the neonatal intensive care unit (NICU).

2. Hearing Screening and Rescreening Protocols

- Separate protocols are recommended for NICU and well-baby nurseries. NICU babies admitted for greater than 5 days are to have auditory brainstem response (ABR) included as part of their screening so that neural hearing loss will not be missed.

- For infants who do not pass automated ABR in the NICU, referral should be made directly to an audiologist for rescreening and, when indicated, comprehensive evaluation including ABR.
- For rescreening, a complete screening on both ears is recommended, even if only one ear failed the initial screening.
- For readmissions in the first month of life for all infants (NICU or well baby) when there are conditions associated with potential hearing loss (eg, hyperbilirubinemia requiring exchange transfusion or culture-positive sepsis), a repeat hearing screening is recommended before discharge.

3. Diagnostic Audiology Evaluation

- Audiologists with skills and expertise in evaluating newborn and young infants with hearing loss should provide audiology diagnostic and auditory habilitation services (selection and fitting of amplification device).
- At least one ABR test is recommended as part of a complete audiology diagnostic evaluation for children younger than 3 years for confirmation of permanent hearing loss.
- The timing and number of hearing re-evaluations for children with risk factors should be customized and individualized depending on the relative likelihood of a subsequent delayed-onset hearing loss. Infants who pass the neonatal screening but have a risk factor should have at least 1 diagnostic audiology assessment by 24 to 30 months of age. Early and more frequent assessment may be indicated for children with cytomegalovirus (CMV) infection, syndromes associated with progressive hearing loss, neurodegenerative disorders, trauma, or culture-positive postnatal infections associated with sensorineural hearing loss; for children who have received ECMO or chemotherapy; and when there is caregiver concern or a family history of hearing loss.
- For families who elect amplification, infants in whom permanent hearing loss is diagnosed should be fitted with an amplification device within 1 month of diagnosis.

4. Medical Evaluation

- For infants with confirmed hearing loss, a genetics consultation should be offered to their families.
- Every infant with confirmed hearing loss should be evaluated by an otolaryngologist with knowledge of pediatric hearing loss and have at least 1 examination to assess visual acuity by an ophthalmologist experienced in evaluating infants.
- The risk factors for congenital and acquired hearing loss have been combined in a single list, rather than grouped by time of onset.

5. Early Intervention

- All families of infants with any degree of bilateral or unilateral permanent hearing loss should be considered eligible for early intervention services.
- There should be recognized central referral points of entry that ensure specialty services for infants with confirmed hearing loss.
- Early intervention services for infants with confirmed hearing loss should be provided by professionals with expertise in hearing loss, including educators of the deaf, speech-language pathologists, and audiologists.
- In response to a previous emphasis on “natural environments,” the committee recommends that both home-based and center-based intervention options should be offered.

6. Surveillance and Screening in the Medical Home

- For all infants, regular surveillance of developmental milestones, auditory skills, parental concerns, and middle ear status should be performed in the medical home, consistent with the American Academy of Pediatrics (AAP) pediatric periodicity schedule. All infants should have an objective standardized screening of global development with a validated assessment tool at 9, 18, and 24 to 30 months of age or at any time if the health care professional or family has concern.
- Infants who do not pass the speech-language portion of a medical home global screening or for whom there is a concern regarding hearing or language should be referred for speech-language evaluation and audiology assessment.

7. Communication

- The birth hospital, in collaboration with the state EHDI coordinator, should ensure that the hearing screening results are conveyed to the parents and the medical home.
- Parents should be provided with appropriate follow-up and resource information, and hospitals should ensure that each infant is linked to a medical home.
- Information at all stages of the EHDI process is to be communicated to the family in a culturally sensitive and understandable format.
- Individual hearing screening information and audiology diagnostic and habilitation information should be promptly transmitted to the medical home and the state EHDI coordinator.
- Families should be made aware of all communication options and available hearing technologies (presented in an unbiased manner). Informed family choice and desired outcome guide the decision-making process.

8. Information Infrastructure

- States should implement data-management and tracking systems as part of an integrated child health information system to monitor the quality of EHDI services and provide recommendations for improving systems of care.
- An effective link between health and education professionals is needed to ensure successful transition and to determine outcomes of children with hearing loss for planning and establishing public health policy.

I. BACKGROUND

It has long been recognized that unidentified hearing loss at birth can adversely affect speech and language development as well as academic achievement and social-emotional development. Historically, moderate-to-severe hearing loss in young children was not detected until well beyond the newborn period, and it was not unusual for diagnosis of milder hearing loss and unilateral hearing loss to be delayed until school age.

In the late 1980s, Dr. C. Everett Koop, then US Surgeon General, on learning of new technology, encouraged that detection of hearing loss be included in the Healthy People 2000: goals for the nation.⁴ In 1988, the Maternal and Child Health Bureau (MCHB), a division of the US Health Resources and Services Administration (HRSA), funded pilot projects in Rhode Island, Utah, and Hawaii to test the feasibility of a universal statewide screening program to screen newborn infants for hearing loss before hospital discharge. The National Institutes of Health (NIH), through the National Institute on Deafness and Other Communication Disorders (NIDCD), issued in 1993 the “Consensus Statement on Early Identification of Hearing Impairment in Infants and Young Children.”⁵ The statement concluded that all infants admitted to the NICU should be screened for hearing loss before hospital discharge and that universal screening should be implemented for all infants within the first 3 months of life.⁴ In its 1994 position statement, the JCIH endorsed the goal of universal detection of infants with hearing loss and encouraged continuing research and development to improve methods for identification of and intervention for hearing loss.^{6,7} The AAP released a statement recommending newborn hearing screening and intervention in 1999.⁸ In 2000, citing advances in screening technology, the JCIH endorsed the universal screening of all infants through an integrated, interdisciplinary system of EHDI.³ The Healthy People 2010 goals included an objective to “increase the proportion of newborns who are screened for hearing loss by one month, have audiologic evaluation by 3 months, and are enrolled in appropriate intervention services by 6 months.”⁹

The ensuing years have seen remarkable expansion in newborn hearing screening. At the time of the NIH consensus statement, only 11 hospitals in

the United States were screening more than 90% of their newborn infants. In 2000, through the support of Representative Jim Walsh (R-NY), Congress authorized HRSA to develop newborn hearing screening and follow-up services, the Centers for Disease Control and Prevention (CDC) to develop data and tracking systems, and NIDCD to support research in early hearing detection and intervention. By 2005, every state had implemented a newborn hearing-screening program, and approximately 95% of newborn infants in the United States were screened for hearing loss before hospital discharge. Congress recommended cooperation and collaboration among several federal agencies and advocacy organizations to facilitate and to support the development of state EHDI systems.

EHDI programs throughout the nation have demonstrated not only the feasibility of universal newborn hearing screening (UNHS) but also the benefits of early identification and intervention. There is a growing body of literature indicating that when identification and intervention occur no later than 6 months of age for newborn infants who are deaf or hard of hearing, the infants perform as much as 20 to 40 percentile points higher on school-related measures (vocabulary, articulation, intelligibility, social adjustment, and behavior).^{10–13}

Still, many important challenges remain. Despite the fact that approximately 95% of newborn infants have their hearing screened in the United States, almost half of newborn infants who do not pass the initial screening fail to have appropriate follow-up to either confirm the presence of a hearing loss and/or initiate appropriate early intervention services (<http://www.infanthearing.org/>; <http://www.cdc.gov/ncbddd/ehdi/>; <http://www.nidcd.nih.gov/health/>).

State EHDI coordinators report system-wide problems, including failure to communicate information to families in a culturally sensitive and understandable format at all stages of the EHDI process, lack of integrated state data-management and tracking systems, and a shortage of facilities and personnel with the experience and expertise needed to provide follow-up for infants referred from newborn screening programs.¹⁴ Available data indicate that a significant number of children who need further assessment do not receive appropriate follow-up evaluations. However, the outlook is improving as EHDI programs focus on the importance of strengthening follow-up and intervention.

II. PRINCIPLES

All children with hearing loss should have access to resources necessary to reach their maximum potential. The following principles provide the foundation for effective EHDI systems and have been updated and expanded since the JCIH 2000 position statement.

1. All infants should have access to hearing screening using a physiologic measure no later than 1 month of age.
2. All infants who do not pass the initial hearing screening and the subsequent rescreening should have appropriate audiologic and medical evaluations to confirm the presence of hearing loss no later than 3 months of age.
3. All infants with confirmed permanent hearing loss should receive early intervention services as soon as possible after diagnosis but no later than 6 months of age. A simplified, single point of entry into an intervention system appropriate for children with hearing loss is optimal.
4. The EHDI system should be family centered with infant and family rights and privacy guaranteed through informed choice, shared decision making, and parental consent in accordance with state and federal guidelines. Families should have access to information about all intervention and treatment options and counseling regarding hearing loss.
5. The child and family should have immediate access to high-quality technology, including hearing aids, cochlear implants, and other assistive devices when appropriate.
6. All infants and children should be monitored for hearing loss in the medical home.¹⁵ Continued assessment of communication development should be provided by appropriate professionals to all children with or without risk indicators for hearing loss.
7. Appropriate interdisciplinary intervention programs for infants with hearing loss and their families should be provided by professionals knowledgeable about childhood hearing loss. Intervention programs should recognize and build on strengths, informed choices, traditions, and cultural beliefs of the families.
8. Information systems should be designed and implemented to interface with electronic health records and should be used to measure outcomes and report the effectiveness of EHDI services at the patient, practice, community, state, and federal levels.

III. GUIDELINES FOR EARLY HEARING DETECTION AND INTERVENTION PROGRAMS

The 2007 guidelines were developed to update the 2000 JCIH position statement principles and to support the goals of universal access to hearing screening, evaluation, and intervention for newborn and young infants embodied in Healthy People 2010.⁹ The guidelines provide current information on the development and implementation of successful EHDI systems.

Hearing screening should identify infants with specifically defined hearing loss on the basis of investigations of long-term, developmental consequences of hearing loss in infants, currently available physiologic screening techniques, and availability of effective intervention in concert with established

principles of health screening.^{15–18} Studies have demonstrated that current screening technologies are effective in identifying hearing loss of moderate and greater degree.¹⁹ In addition, studies of children with permanent hearing loss indicate that moderate or greater degrees of hearing loss can have significant effects on language, speech, academic, and social-emotional development.²⁰ High-risk target populations also include NICU infants, because research data indicates that this population is at highest risk of having neural hearing loss.^{21–23}

The JCIH, however, is committed to the goal of identifying all degrees and types of hearing loss in childhood and recognizes the developmental consequences of even mild degrees of permanent hearing loss. Recent evidence suggests, however, that current hearing screening technologies fail to identify some infants with mild forms of hearing loss.^{24,25} Additionally, depending on the screening technology selected, infants with hearing loss related to neural conduction disorders or “auditory neuropathy/auditory dyssynchrony” may not be detected through a UNHS program. Although the JCIH recognizes that these disorders may result in delayed communication,^{26–28} currently recommended screening algorithms (ie, use of otoacoustic emissions testing [OAE] alone) preclude universal screening for these disorders. Because these disorders typically occur in children who require NICU care,²¹ the JCIH recommends screening this group with the technology capable of detecting auditory neuropathy/dyssynchrony: automated ABR.

All infants, regardless of newborn hearing screening outcome, should receive ongoing monitoring for development of age-appropriate auditory behaviors and communication skills. Any infant who demonstrates delayed auditory and/or communication skills development, even if they passed newborn hearing screening, should receive audiologic evaluation to rule out hearing loss.

A. Roles and Responsibilities

The success of EHDI programs depends on families working in partnership with professionals as a well-coordinated team. The roles and responsibilities of each team member should be well defined and clearly understood. Essential team members are the birth hospital, families, pediatricians or primary health care professionals (ie, the medical home), audiologists, otolaryngologists, speech-language pathologists, educators of children who are deaf or hard of hearing, and other early intervention professionals involved in delivering EHDI services.^{29,30} Additional services including genetics, ophthalmology, developmental pediatrics, service coordination, supportive family education, and counseling should be available.³¹

The birth hospital is a key member of the team. The birth hospital, in collaboration with the state EHDI coordinator, should ensure that parents and primary health care professionals receive and understand the hearing

screening results, that parents are provided with appropriate follow-up and resource information, and that each infant is linked to a medical home.² The hospital ensures that hearing screening information is promptly transmitted to the medical home and appropriate data are submitted to the state EHDI coordinator.

The most important role for the family of an infant who is deaf or hard of hearing is to love, nurture, and communicate with the baby. From this foundation, families usually develop an urgent desire to understand and meet the special needs of their infant. Families gain knowledge, insight, and experience by accessing resources and through participation in scheduled early intervention appointments including audiological, medical, habilitative, and educational sessions. This experience can be enhanced when families choose to become involved with parental support groups, individuals who are deaf or hard of hearing, and/or their children's deaf or hard-of-hearing peers. Informed family choices and desired outcomes guide all decisions for these children. A vital function of the family's role is ensuring direct access to communication in the home and the daily provision of language learning opportunities. Over time, the child benefits from the family's modeling of partnerships with professionals and advocating for their rights in all settings. The transfer of responsibilities from families to their child develops gradually and increases as their child matures, growing in independence and self-advocacy.

Pediatricians, family physicians, and other allied health care professionals, working in partnership with parents and other professionals such as audiologists, therapists, and educators, constitute the infant's medical home.² A *medical home* is defined as an approach to providing health care services where care is accessible, family centered, continuous, comprehensive, coordinated, compassionate, and culturally competent. The primary health care professional acts in partnership with parents in a medical home to identify and access appropriate audiology, intervention, and consultative services needed to develop a global plan of appropriate and necessary health and habilitative care for infants identified with hearing loss and infants with risk factors for hearing loss. All children undergo surveillance for auditory skills and language milestones. The infant's pediatrician, family physician, or other primary health care professional is in a position to advocate for the child and family.^{2,16}

An audiologist is a person who, by virtue of academic degree, clinical training, and license to practice, is qualified to provide services related to the prevention of hearing loss and the audiologic diagnosis, identification, assessment, and nonmedical and nonsurgical treatment of persons with impairment of auditory and vestibular function, and to the prevention of impairments associated with them. Audiologists serve in a number of roles. They provide newborn hearing screening program development, management, quality assessment, service coordination and referral for audiologic

diagnosis, and audiologic treatment and management. For the follow-up component, audiologists provide comprehensive audiologic diagnostic assessment to confirm the existence of the hearing loss, ensure that parents understand the significance of the hearing loss, evaluate the infant for candidacy for amplification and other sensory devices and assistive technology, and ensure prompt referral to early intervention programs. For the treatment and management component, audiologists provide timely fitting and monitoring of amplification.³² Other audiologists may provide diagnostic and auditory treatment and management services in the educational setting and provide a bridge between the child/family and the audiologist in the clinic setting as well as other service providers. Audiologists also provide services as teachers, consultants, researchers, and administrators.

Otolaryngologists are physicians whose specialty includes determining the etiology of hearing loss; identifying related risk indicators for hearing loss, including syndromes involving the head and neck; and evaluating and treating ear diseases. An otolaryngologist with knowledge of childhood hearing loss can determine whether medical and/or surgical intervention may be appropriate. When medical and/or surgical intervention is provided, the otolaryngologist is involved in the long-term monitoring and follow-up with the infant's medical home. The otolaryngologist provides information and participates in the assessment of candidacy for amplification, assistive devices, and surgical intervention, including reconstruction, bone-anchored hearing aids, and cochlear implantation.

Early intervention professionals are trained in a variety of academic disciplines, such as speech-language pathology, audiology, education of children who are deaf or hard of hearing, service coordination, or early childhood special education. All individuals who provide services to infants with hearing loss should have specialized training and expertise in the development of audition, speech, and language. Speech-language pathologists provide both evaluation and intervention services for language, speech, and cognitive-communication development. Educators of children who are deaf or hard of hearing integrate the development of communicative competence within a variety of social, linguistic, and cognitive/academic contexts. Audiologists may provide diagnostic and habilitative services within the individualized family service plan (IFSP) or school-based individualized education plan. To provide the highest quality of intervention, more than one provider may be required.

The care coordinator is an integral member of the EHDI team who facilitates the family's transition from screening to evaluation to early intervention.³³ This individual must be a professional (eg, social worker, teacher, nurse) who is knowledgeable about hearing loss. The care coordinator incorporates the family's preferences for outcomes into an IFSP as required by federal legislation. The care coordinator supports the family members in their choice of the infant's communicative development. Through the IFSP review,

the infant's progress in language, motor, cognitive, and social-emotional development is monitored. The care coordinator assists the family in advocating for the infant's unique developmental needs.

The deaf and hard-of-hearing community includes members with direct experience with signed language, spoken language, hearing aid and cochlear implant use, and other communication strategies and technologies. Optimally, adults who are deaf or hard-of-hearing should play an integral part in the EHDI program. Both adults and children in the deaf and hard-of-hearing community can enrich the family's experience by serving as mentors and role models. Such mentors have experience in negotiating their way in a hearing world, raising infants or children who are deaf or hard of hearing, and providing families with a full range of information about communication options, assistive technology, and resources available in the community.

A successful EHDI program requires collaboration between a variety of public and private institutions and agencies assuming responsibility for specific components (eg, screening, evaluation, intervention). Roles and responsibilities may differ from state to state. Each state has defined a lead coordinating agency with oversight responsibility. The lead coordinating agency in each state should be responsible for identifying the public and private funding sources available to develop, implement, and coordinate EHDI systems.

B. Hearing Screening

Multidisciplinary teams of professionals, including audiologists, physicians, and nursing personnel, are needed to establish the UNHS component of EHDI programs. All team members work together to ensure that screening programs are of high quality and are successful. An audiologist should be involved in each component of the hearing screening program, particularly at the level of statewide implementation and, whenever possible, at the individual hospital level. Hospitals and agencies should also designate a physician to oversee the medical aspects of the EHDI program.

Each team of professionals responsible for the hospital-based UNHS program should review the hospital infrastructure in relationship to the screening program. Hospital-based programs should consider screening technology (ie, OAE or automated ABR testing); validity of the specific screening device; screening protocols, including the timing of screening relative to nursery discharge; availability of qualified screening personnel; suitability of the acoustical and electrical environments; follow-up referral criteria; referral pathways for follow-up; information management; and quality control and improvement. Reporting and communication protocols must be well defined and include the content of reports to physicians and parents, documentation of results in medical records, and methods for reporting to state registries and national data sets.

Physiologic measures must be used to screen newborns and infants for hearing loss. Such measures include OAE and automated ABR testing. Both OAE and automated ABR technologies provide noninvasive recordings of physiologic activity underlying normal auditory function, both are easily performed in neonates and infants, and both have been successfully used for UNHS.^{19,34–37} There are, however, important differences between the 2 measures. OAE measurements are obtained from the ear canal using a sensitive microphone within a probe assembly that records cochlear responses to acoustic stimuli. Thus, OAEs reflect the status of the peripheral auditory system extending to the cochlear outer hair cells. In contrast, ABR measurements are obtained from surface electrodes that record neural activity generated in the cochlea, auditory nerve, and brainstem in response to acoustic stimuli delivered via an earphone. Automated ABR measurements reflect the status of the peripheral auditory system, the eighth nerve, and the brainstem auditory pathway.

Both OAE and ABR screening technologies can be used to detect sensory (cochlear) hearing loss¹⁹; however, both technologies may be affected by outer or middle ear dysfunction. Consequently, transient conditions of the outer and middle ear may result in a “fail” screening test result in the presence of normal cochlear and/or neural function.³⁸ Moreover, because OAEs are generated within the cochlea, OAE technology cannot be used to detect neural (eighth nerve or auditory brainstem pathway) dysfunction. Thus, infants with neural conduction disorders or auditory neuropathy/dyssynchrony without concomitant sensory dysfunction will not be detected by OAE testing.

Some infants who pass newborn hearing screening will later demonstrate permanent hearing loss.²⁵ Although this loss may reflect delayed-onset hearing loss, both ABR and OAE screening technologies will miss some hearing loss (eg, mild or isolated frequency region losses).

Interpretive criteria for pass/fail outcomes should reflect clear scientific rationale and should be evidenced based.^{39,40} Screening technologies that incorporate automated response detection are necessary to eliminate the need for individual test interpretation, to reduce the effects of screener bias or operator error on test outcome, and to ensure test consistency across infants, test conditions, and screening personnel.^{41–45} When statistical probability is used to make pass/fail decisions, as is the case for OAE and automated ABR screening devices, the likelihood of obtaining a pass outcome by chance alone is increased when screening is performed repeatedly.^{46–48} This principle must be incorporated into the policies of rescreening.

There are no national standards for the calibration of OAE or ABR instrumentation. Compounding this, there is a lack of uniform performance standards. Manufacturers of hearing-screening devices do not always provide sufficient supporting evidence to validate the specific pass/fail criteria and/or automated algorithms utilized in their instruments.⁴⁹ In the absence of

national standards, audiologists must obtain normative data for the instruments and protocols they use.

The JCIH recognizes that there are important issues differentiating screening performed in the well-baby nursery from that performed in the NICU. Although the goals in each nursery are the same, numerous methodologic and technological issues must be considered in program design and pass/fail criteria.

1. Screening Protocols in the Well-Baby Nursery

Many inpatient well-baby screening protocols provide one hearing screening and, when necessary, a repeat screening no later than the time of discharge from the hospital, using the same technology both times. Use of either technology in the well-baby nursery will detect peripheral (conductive and sensory) hearing loss of 40 dB or greater.¹⁹ When automated ABR is used as the single screening technology, neural auditory disorders can also be detected.⁵⁰ Some programs use a combination of screening technologies (OAE testing for the initial screening, followed by automated ABR for rescreening (ie, 2-step protocol⁵), to decrease the fail rate at discharge and the subsequent need for outpatient follow-up.^{34,35,37,51–53} Using this approach, infants who do not pass an OAE screening but subsequently pass an automated ABR are considered a screening “pass.” Infants in the well-baby nursery who fail automated ABR should not be rescreened by OAE and “passed,” because such infants are presumed to be at risk of having a subsequent diagnosis of auditory neuropathy/dyssynchrony.

2. Screening Protocols in the NICU

A NICU is defined as a facility in which a neonatologist provides primary care for the infant. Newborn units are divided into categories as follows:

- Level I: basic care, well-baby nurseries
- Level II: specialty care by a neonatologist for infants at moderate risk of serious complications
- Level III: a unit that provides both specialty and subspecialty care including the provision of life support (mechanical ventilation)

A total of 120 level-II NICUs and 760 level-III NICUs have been identified in the United States by survey, and infants who have spent time in the NICU represent 10% to 15% of the newborn population.⁵⁴

The JCIH 2007 position statement includes neonates at risk of having neural hearing loss (auditory neuropathy/auditory dyssynchrony) in the target population to be identified in the NICU,^{55–57} because there is evidence that

neural hearing loss results in adverse communication outcomes.^{22,50} Consequently, the JCIH recommends ABR technology as the only appropriate screening technique for use in the NICU. For infants who do not pass automated ABR testing in the NICU, referral should be made directly to an audiologist for rescreening and, when indicated, comprehensive evaluation, including diagnostic ABR, rather than for general outpatient rescreening.

3. Conveying Test Results

Screening results should be conveyed immediately to families so they understand the outcome and the importance of follow-up when indicated. To facilitate this process for families, primary health care professionals should work with EHDI team members to ensure that:

- Communications with parents are confidential and presented in a caring and sensitive manner, preferably face-to-face.
- Educational materials are developed and disseminated to families that provide accurate information at an appropriate reading level and in a language they are able to comprehend.
- Parents are informed in a culturally sensitive and understandable manner that their infant did not pass screening and informed about the importance of prompt follow-up. Before discharge, an appointment should be made for follow-up testing.

To facilitate this process for primary care physicians, EHDI systems should ensure that medical professionals:

- Receive the results of the screening test (pass, did not pass, or missed) as documented in the hospital medical record.
- Receive communication directly from the hospital screening program regarding each infant in their care who did not pass or is missed and recommendations for follow-up.

4. Outpatient Rescreening for Infants Who Do Not Pass the Birth Admission Screening

Many well-baby screening protocols will choose to incorporate an outpatient rescreening within 1 month of hospital discharge to minimize the number of infants referred for follow-up audiologic and medical evaluation. The outpatient rescreening should include the testing of both ears, even if only 1 ear failed the inpatient screening.

Outpatient screening no later than 1 month of age should also be available to infants who were discharged before receiving the birth admission screening or who were born outside a hospital or birthing center. State EHDI coordinators should be aware of some of the following situations under which infants may be lost to the UNHS system:

- Home births and other out-of-hospital births: States should develop a mechanism to systematically offer newborn hearing screening for all out-of-hospital births.
- Across state border births: States should develop written collaborative agreements among neighboring states for sharing hearing screening results and follow-up information.
- Hospital missed screenings: When infants are discharged before the hearing screening is performed, a mechanism should be in place for the hospital to contact the family and arrange for an outpatient hearing screening.
- Transfers to in-state or out-of-state hospitals: Discharge and transfer forms should contain the information of whether a hearing screening was performed and the results of any screening. The recipient hospital should complete a hearing screening if not previously performed or if there is a change in medical status or a prolonged hospitalization.
- Readmits: For readmissions in the first month of life when there are conditions associated with potential hearing loss (eg, hyperbilirubinemia requiring exchange transfusion or culture-positive sepsis), a screening ABR should be performed before discharge.

Additional mechanisms for states to share hearing screening results and other medical information include: 1) incorporating the hearing screening results in a statewide child health information system; and 2) providing combined metabolic screening and hearing screening results to the primary care physician.

C. Confirmation of Hearing Loss in Infants Referred From UNHS

Infants who meet the defined criteria for referral should receive follow-up audiologic and medical evaluations with fitting of amplification devices, as appropriate, no later than 3 months of age. Once hearing loss is confirmed, coordination of services should be expedited by the infant's medical home and Part C coordinating agencies for early intervention services, as authorized by the Individuals with Disabilities Education Act, following the EHDI algorithm developed by the AAP (Appendix 2).

1. Audiologic Evaluation

Comprehensive audiologic evaluation of newborn and young infants who fail newborn hearing screening should be performed by audiologists experienced in pediatric hearing assessment. The initial audiologic test battery to confirm a hearing loss in infants must include physiologic measures and, when developmentally appropriate, behavioral methods. Confirmation of an infant's hearing status requires a test battery of audiologic test procedures to

assess the integrity of the auditory system in each ear, to estimate hearing sensitivity across the speech frequency range, to determine the type of hearing loss, to establish a baseline for further monitoring, and to provide information needed to initiate amplification device fitting. Comprehensive assessment should be performed on both ears even if only one ear failed the screening test.

i. Evaluation: Birth to 6 Months of Age

For infants from birth to a developmental age of approximately 6 months, the test battery should include a child and family history, an evaluation of risk factors for congenital hearing loss, and a parental report of the infant's responses to sound. The audiologic assessment should include the following:

- Child and family history
- A frequency-specific assessment of the ABR using air-conducted tone bursts and bone-conducted tone bursts when indicated. When permanent hearing loss is detected, frequency-specific ABR is needed to determine the degree and configuration of hearing loss in each ear for fitting of amplification devices.
- Click-evoked ABR using both condensation and rarefaction single-polarity stimulus, if there are risk indicators for neural hearing loss (auditory neuropathy/auditory dyssynchrony) such as hyperbilirubinemia or anoxia, to determine whether a cochlear microphonic is present.²⁸ Furthermore, because some infants with neural hearing loss have no risk indicators, any infant who demonstrates "no response" on ABR elicited by tone burst stimuli must be evaluated by a click-evoked ABR, as previously described.
- Distortion product or transient evoked otoacoustic emissions.
- Tympanometry using a 1000-Hz probe tone.
- Clinician observation of the infant's auditory behavior as a cross-check, in conjunction with electrophysiologic measures. Behavioral observation alone is not adequate for determining whether hearing loss is present in this age group, nor is it adequate for the fitting of amplification devices.

ii. Evaluation: 6 to 36 Months of Age

For subsequent testing of infants and toddlers at developmental ages of 6 to 36 months, the confirmatory audiologic test battery includes the following:

- Child and family history
- Parental report of auditory and visual behaviors and communication milestones.
- Behavioral audiometry (either visual reinforcement or conditioned-play

audiometry, depending on the child's developmental level), including pure-tone audiometry across the frequency range for each ear and speech detection and recognition measures.

- OAE testing.
- Acoustic immittance measures (tympanometry and acoustic reflex thresholds).
- ABR, if responses to behavioral audiometry are not reliable or if ABR has not been performed in the past.

iii. Other Audiologic Test Procedures

At this time, there is insufficient evidence for use of the auditory steady state response (ASSR) as the sole measure of auditory status in newborn and infant populations.⁵⁸ Auditory steady-state response is a new evoked potential test that can accurately measure auditory sensitivity beyond the limits of other test methods. It can determine frequency specific thresholds from 250 Hz to 8 kHz. Clinical research is investigating its potential use in the standard pediatric diagnostic test battery. Similarly, there are insufficient data for routine use of acoustic middle ear muscle reflexes in the initial diagnostic assessment of infants younger than 4 months.⁵⁹ Both tests could be used to supplement the battery or could be included at older ages. Emerging technologies, such as broad-band reflectance, may be used to supplement conventional measures of middle ear status (tympanometry and acoustic reflexes) as the technology becomes more widely available.⁵⁹

2. Medical Evaluation

Every infant with confirmed hearing loss and/or middle ear dysfunction should be referred for otologic and other medical evaluation. The purpose of these evaluations is to determine the etiology of hearing loss, to identify related physical conditions, and to provide recommendations for medical/surgical treatment as well as referral for other services. Essential components of the medical evaluation include clinical history, family history of childhood-onset permanent hearing loss, identification of syndromes associated with early- or late-onset permanent hearing loss, a physical examination, and indicated radiologic and laboratory studies (including genetic testing). Portions of the medical evaluation, such as urine culture for CMV, a leading cause of hearing loss, might even begin in the birth hospital, particularly for infants spending time in the NICU.⁶⁰⁻⁶²

i. Pediatrician/Primary Care Physician

The infant's pediatrician or other primary health care professional is responsible for monitoring the general health, development, and well-being of

the infant. In addition, the primary care physician must assume responsibility to ensure that the audiologic assessment is conducted on infants who do not pass screening and must initiate referrals for medical specialty evaluations necessary to determine the etiology of the hearing loss. Middle-ear status should be monitored, because the presence of middle-ear effusion can further compromise hearing. The primary care physician must partner with other specialists, including the otolaryngologist, to facilitate coordinated care for the infant and family. Because 30% to 40% of children with confirmed hearing loss will demonstrate developmental delays or other disabilities, the primary care physician should closely monitor developmental milestones and initiate referrals related to suspected disabilities.⁶³ The medical home algorithm for management of infants with either suspected or proven permanent hearing loss is provided in Appendix 2.¹⁵

The pediatrician or primary care physician should review every infant's medical and family history for the presence of risk indicators that require monitoring for delayed-onset or progressive hearing loss and should ensure that an audiologic evaluation is completed for children at risk of hearing loss at least once by 24 to 30 months of age, regardless of their newborn screening results.²⁵ Infants with specific risk factors, such as those who received ECMO therapy and those with CMV infection, are at increased risk of delayed-onset or progressive hearing loss^{64–67} and should be monitored closely. In addition, the primary care physician is responsible for ongoing surveillance of parent concerns about language and hearing, auditory skills, and developmental milestones of all infants and children regardless of risk status, as outlined in the pediatric periodicity schedule published by the AAP.¹⁶

Children with cochlear implants may be at increased risk of acquiring bacterial meningitis compared with children in the general US population.⁶⁸ The CDC recommends that all children with, and all potential recipients of, cochlear implants follow specific recommendations for pneumococcal immunization that apply to cochlear implant users and that they receive age-appropriate *Haemophilus influenzae* type b vaccines. Recommendations for the timing and type of pneumococcal vaccine vary with age and immunization history and should be discussed with a health care professional.⁶⁹

ii. Otolaryngologist

Otolaryngologists are physicians and surgeons who diagnose, treat, and manage a wide range of diseases of the head and neck, specializing in treating hearing and vestibular disorders. They perform a full medical diagnostic evaluation of the head and neck, ears, and related structures, including a comprehensive history and physical examination, leading to a medical diagnosis and appropriate medical and surgical management. Often, a hearing or balance disorder is an indicator of, or related to, a medically treatable con-

dition or an underlying systemic disease. Otolaryngologists work closely with other dedicated professionals, including physicians, audiologists, speech-language pathologists, educators, and others, in caring for patients with hearing, balance, voice, speech, developmental, and related disorders.

The otolaryngologist's evaluation includes a comprehensive history to identify the presence of risk factors for early-onset childhood permanent hearing loss, such as family history of hearing loss, having been admitted to the NICU for >5 days, and having received ECMO (see Appendix 1).^{70,71}

A complete head and neck examination for craniofacial anomalies should document defects of the auricles, patency of the external ear canals, and status of the eardrum and middle ear structures. Atypical findings on eye examination, including irises of 2 different colors or abnormal positioning of the eyes, may signal a syndrome that includes hearing loss. Congenital permanent conductive hearing loss may be associated with craniofacial anomalies seen in disorders such as Crouzon disease, Klippel-Feil syndrome, and Goldenhar syndrome.⁷² The assessment of infants with these congenital anomalies should be coordinated with a clinical geneticist.

In large population studies, at least 50% of congenital hearing loss has been designated as hereditary, and nearly 600 syndromes and 125 genes associated with hearing loss have already been identified.^{72,73} The evaluation, therefore, should include a review of family history of specific genetic disorders or syndromes, including genetic testing for gene mutations such as GJB2 (connexin-26), and syndromes commonly associated with early-onset childhood sensorineural hearing loss^{72,74-76} (Appendix 1). As the widespread use of newly developed conjugate vaccines decreases the prevalence of infectious etiologies such as measles, mumps, rubella, *H influenzae* type b, and childhood meningitis, the percentage of each successive cohort of early-onset hearing loss attributable to genetic etiologies can be expected to increase, prompting recommendations for early genetic evaluations. Approximately 30% to 40% of children with hearing loss have associated disabilities, which can be of importance in patient management. The decision to obtain genetic testing is dependent on informed family choice in conjunction with standard confidentiality guidelines.⁷⁷

In the absence of a genetic or established medical cause, a computed tomography scan of the temporal bones may be performed to identify cochlear abnormalities, such as Mondini deformity with an enlarged vestibular aqueduct, which have been associated with progressive hearing loss. Temporal bone imaging studies may also be used to assess potential candidacy for surgical intervention, including reconstruction, bone-anchored hearing aid, and cochlear implantation. Recent data have shown that some children with electrophysiologic evidence suggesting auditory neuropathy/dyssynchrony may have an absent or abnormal cochlear nerve that may be detected on magnetic resonance imaging.⁷⁸

Historically, an extensive battery of laboratory and radiographic studies was routinely recommended for newborn infants and children with newly diagnosed sensorineural hearing loss. However, emerging technologies for the diagnosis of genetic and infectious disorders have simplified the search for a definitive diagnosis, obviating the need for costly diagnostic evaluations in some instances.^{70,71,79}

If, after an initial evaluation, the etiology remains uncertain, an expanded multidisciplinary evaluation protocol including electrocardiography, urinalysis, testing for CMV, and further radiographic studies is indicated. The etiology of neonatal hearing loss, however, may remain uncertain in as many as 30% to 40% of children. Once hearing loss is confirmed, medical clearance for hearing aids and initiation of early intervention should not be delayed while this diagnostic evaluation is in process. Careful longitudinal monitoring to detect and promptly treat coexisting middle ear effusions is an essential component of ongoing otologic management of these children.

iii. Other Medical Specialists

The medical geneticist is responsible for the interpretation of family history data, the clinical evaluation and diagnosis of inherited disorders, the performance and assessment of genetic tests, and the provision of genetic counseling. Geneticists or genetic counselors are qualified to interpret the significance and limitations of new tests and to convey the current status of knowledge during genetic counseling. All families of children with confirmed hearing loss should be offered and may benefit from a genetics evaluation and counseling. This evaluation can provide families with information on etiology of hearing loss, prognosis for progression, associated disorders (eg renal, vision, cardiac), and likelihood of recurrence in future offspring. This information may influence parents' decision making regarding intervention options for their child.

Every infant with a confirmed hearing loss should have an evaluation by an ophthalmologist to document visual acuity and rule out concomitant or late-onset vision disorders, such as Usher syndrome.^{1,80} Indicated referrals to other medical subspecialists, including developmental pediatricians, neurologists, cardiologists, and nephrologists, should be facilitated and coordinated by the primary health care professional.

D. Early Intervention

Before newborn hearing screening was instituted universally, children with severe to profound hearing loss, on average, completed the 12th grade with a third- to fourth-grade reading level and language levels of a 9- to 10-year-old hearing child.⁸¹ In contrast, infants and children with mild to profound

hearing loss who are identified in the first 6 months of life and provided with immediate and appropriate intervention have significantly better outcomes than later-identified infants and children in vocabulary development,^{82,83} receptive and expressive language,^{12,84} syntax,⁸⁵ speech production,^{13,86–88} and social-emotional development.⁸⁹ Children enrolled in early intervention within the first year of life have also been shown to have language development within the normal range of development at 5 years of age.^{31,90}

Therefore, according to federal guidelines, once any degree of hearing loss is diagnosed in a child, a referral should be initiated to an early intervention program within 2 days of confirmation of hearing loss (CFA 303.321d). The initiation of early intervention services should begin as soon as possible after diagnosis of hearing loss, but no later than 6 months of age. Even when the hearing status is not determined to be the primary disability, the family and child should have access to intervention with a provider knowledgeable about hearing loss.⁹¹

UNHS programs have been instituted throughout the United States for the purpose of preventing the significant and negative effects of hearing loss on the cognitive, language, speech, auditory, social-emotional, and academic development of infants and children. To achieve this goal, hearing loss must be identified as quickly as possible after birth, and appropriate early intervention must be available to all families and infants with permanent hearing loss. Some programs have demonstrated that most children with hearing loss and no additional disabilities can achieve and maintain language development within the typical range of children with normal hearing.^{12,13,85,90} Because these studies are descriptive and not causal studies, the efficacy of specific components of intervention cannot be separated from the total provision of comprehensive services. Thus, the family-centered philosophy, the intensity of services, the experience and training of the provider, the method of communication, the curricula, the counseling procedures, the parent support and advocacy, and the deaf and hard-of-hearing support and advocacy are all variables with unknown effects on the overall outcomes of any individual child. The key component of providing quality services is the expertise of the provider specific to hearing loss. These services may be provided in the home or in a center, or a combination of the 2 locations.

The term “intervention services” is used to describe any type of habilitative, rehabilitative, or educational program provided to individuals with hearing loss. In some cases with mild hearing losses, amplification technology may be the only service provided. Some parents choose only developmental assessment or occasional consultation, such as parents with infants who have unilateral hearing losses. Children with high-frequency losses and normal hearing in the low frequencies may only be seen by a speech-language pathologist, and those with significant bilateral sensorineural hearing losses might be seen by an educator of the deaf and receive additional services.

1. Principles of Early Intervention

To ensure informed decision making, parents of infants with newly diagnosed hearing loss should be offered opportunities to interact with other families who have infants or children with hearing loss as well as adults and children who are deaf or hard of hearing. In addition, parents should also be offered access to professional, educational, and consumer organizations and provided with general information on child development, language development, and hearing loss. A number of principles and guidelines have been developed that offer a framework for quality early intervention service delivery systems for children who are deaf or hard of hearing and their families.⁹² Foundational characteristics of developing and implementing early intervention programs include a family-centered approach, culturally responsive practices, collaborative professional-family relationships and strong family involvement, developmentally appropriate practice, interdisciplinary assessment, and community-based provision of services.

i. Designated Point of Entry

States should develop a single point of entry into intervention specific for hearing impairment to ensure that, regardless of geographic location, all families who have infants or children with hearing loss receive information about a full range of options regarding amplification and technology, communication and intervention, and accessing appropriate counseling services. This state system, if separate from the state's Part C system, should integrate and partner with the state's Part C program. Parental consent must be obtained, according to state and federal requirements, to share the IFSP information with providers and transmit data to the state EHDI coordinator.

ii. Regular Developmental Assessment

To ensure accountability, individual, community, and state health and educational programs should assume the responsibility for coordinated, ongoing measurement and improvement of EHDI process outcomes. Early-intervention programs must assess the language, cognitive skills, auditory skills, speech, vocabulary, and social-emotional development of all children with hearing loss at 6-month intervals during the first 3 years of life, using assessment tools standardized on children with normal hearing and norm-referenced assessment tools that are appropriate to measure progress in verbal and visual language.

The primary purpose of regular developmental monitoring is to provide valuable information to parents about the rate of their child's development as well as programmatic feedback concerning curriculum decisions. Families also become knowledgeable about expectations and milestones of typical

development of hearing children. Studies have shown that valid and reliable documentation of developmental progress is possible through parent questionnaires, analysis of videotaped conversational interactions, and clinically administered assessments.^{10-13,51,85,87-90,93-96} Documentation of developmental progress should be provided on a regular basis to parents and, with parental release of information, to the medical home and audiologist. Although criterion-referenced checklists may provide valuable information for establishing intervention strategies and goals, these assessment tools alone are not sufficient for parents and intervention professionals to determine whether a child's developmental progress is comparable with his or her hearing peers.

iii. Opportunities for Interaction With Other Parents of Children With Hearing Loss

Intervention professionals should seek to involve parents at every level of the EHDI process and develop true and meaningful partnerships with parents. To reflect the value of the contributions that selected parents make to development and program components, these parents should be paid as contributing staff members. Parent representatives should be included in all advisory board activities. In many states, parents have been integral and often have taken leadership roles in the development of policy, resource material, communication mechanisms, mentoring and advocacy opportunities, dissemination of information, and interaction with the deaf community and other individuals who are deaf or hard of hearing. Parents, often in partnership with individuals who are deaf and hard of hearing, have also participated in the training of professionals. They should be participants in the regular assessment of program services to ensure ongoing improvement and quality assurance.

iv. Opportunities for Interaction With Individuals Who Are Deaf or Hard of Hearing

Intervention programs should include opportunities for involvement of individuals who are deaf or hard of hearing in all aspects of EHDI programs. Because intervention programs serve children with mild to profound, unilateral or bilateral, permanent conductive, and sensory or neural hearing disorders, role models who are deaf or hard of hearing can be significant assets to an intervention program. These individuals can serve on state EHDI advisory boards and can be trained as mentors for families and children with hearing loss who choose to seek their support. Almost all families choose at some time during their early childhood programs to seek out both adults and child peers with hearing loss. Programs should ensure that these opportunities are available and can be delivered to families through a variety of communications means, such as Web sites, e-mail, newsletters, videos, retreats, picnics and other social events, and educational forums for parents.

v. Provision of Communication Options

Research studies thus far of early-identified infants with hearing loss have not found significant differences in the developmental outcomes by method of communication when measured at 3 years of age.^{10-13,85,87,88,90,93,96} Therefore, a range of options should be offered to families in a nonbiased manner. In addition, there are reports of children with successful outcomes for each of the different methods of communication. The choice is a dynamic process on a continuum, differs according to the individual needs of each family and can be adjusted as necessary on the basis of a child's rate of progress in developing communication skills. Programs need to provide families with access to skilled and experienced early intervention professionals to facilitate communication and language development in the communication option chosen by the family.

vi. Skills of the Early Intervention Professional

All studies with successful outcomes reported for early-identified children who are deaf or hard of hearing have intervention provided by specialists trained in parent-infant intervention services.^{12,90,97} Early intervention programs should develop mechanisms to ensure that early intervention professionals have special skills necessary for providing families with the highest quality of service specific to children with hearing loss. Professionals with a background in deaf education, audiology, and speech-language pathology will typically have the skills needed for providing intervention services. Professionals should be highly qualified in their respective fields and should be skilled communicators who are knowledgeable and sensitive to the importance of enhancing families' strengths and supporting their priorities. When early intervention professionals have knowledge of the principles of adult learning, it increases their success with parents and other professionals.

vii. Quality of Intervention Services

Children with confirmed hearing loss and their families have the right to prompt access to quality intervention services. For newborn infants with confirmed hearing loss, enrollment into intervention services should begin as soon after hearing loss confirmation as possible, and no later than 6 months of age. Successful early intervention programs 1) are family centered; 2) provide families with unbiased information on all options regarding approaches to communication; 3) monitor development at 6-month intervals using norm-referenced instruments; 4) include individuals who are deaf or hard of hearing; 5) provide services in a natural environment in the home or in the center; 6) offer high-quality service regardless of where the family lives; 7) obtain informed consent; 8) are sensitive to cultural and language differences and

provide accommodations as needed; and 9) conduct annual surveys of parent satisfaction.

viii. Intervention for Special Populations of Infants and Young Children

Developmental monitoring should also occur at regular 6-month intervals for special populations of children with hearing loss, including minimal and mild bilateral hearing loss,⁹⁸ unilateral hearing loss,^{99,100} and neural hearing loss,²² because these children are at risk of having speech and language delay. Research findings indicate that approximately one third of children with permanent unilateral loss experience significant language and academic delays.^{99,100}

2. Audiologic Habilitation

Most infants and children with bilateral hearing loss and many with unilateral hearing loss benefit from some form of personal amplification.³² If the family chooses personal amplification for their infant, hearing aid selection and fitting should occur within 1 month of initial confirmation of hearing loss even when additional audiologic assessment is ongoing. Audiologic habilitation services should be provided by an audiologist experienced with these procedures. Delay between confirmation of the hearing loss and fitting of an amplification device should be minimized.^{51,101}

Hearing aid fitting proceeds optimally when the results of physiologic audiologic assessment including diagnostic ABR, OAE and tympanometry and medical examination are in accord. For infants below a developmental age of six months, hearing aid selection will be based on physiologic measures alone. Behavioral threshold assessment using visual reinforcement audiometry should be obtained as soon as possible to cross-check and augment physiologic findings (<http://www.audiology.org/>).

The goal of amplification device fitting is to provide the infant with maximum access to all of the acoustic features of speech within an intensity range that is safe and comfortable. That is, amplified speech should be comfortably above the infant's sensory threshold, but below the level of discomfort across the speech frequency range for both ears. To accomplish this in infants, amplification device selection, fitting, and verification should be based on a prescriptive procedure that incorporates individual real-ear measures that account for each infant's ear canal acoustics and hearing loss.³² Validation of the benefits of amplification, particularly for speech perception, should be examined in the clinical setting as well as in the child's typical listening environments. Complementary or alternative technology, such as FM systems or cochlear implants, may be recommended as the primary and/or secondary listening device, depending on the degree of the infant's hearing loss, the goals of auditory habilitation, the infant's acoustic environments, and the

family's informed choices.³ Monitoring of amplification, as well as the long-term validation of the appropriateness of the individual habilitation program, requires ongoing audiologic assessment along with electroacoustic, real-ear, and functional checks of the hearing instruments. As the hearing loss becomes more specifically defined through audiologic assessments and as the child's ear canal acoustics change with growth, refinement of the individual prescriptive hearing aid gain and output targets is necessary. Monitoring also includes periodic validation of communication, social-emotional, and cognitive development and, later, academic performance to ensure that progress is commensurate with the child's abilities. It is possible that infants and young children with measurable residual "hearing" (auditory responses) and well-fit amplification devices may fail to develop auditory skills necessary for successful oral communication. Ongoing validation of the amplification device is accomplished through interdisciplinary evaluation and collaboration with the early intervention team and family.

Cochlear implantation should be given careful consideration for any child who appears to receive limited benefit from a trial with appropriately fitted hearing aids. According to US Food and Drug Administration guidelines, infants with profound bilateral hearing loss are candidates for cochlear implantation at 12 months of age and children with bilateral severe hearing loss are eligible at 24 months of age. The presence of developmental conditions (eg, developmental delay, autism) in addition to hearing loss should not, as a rule, preclude the consideration of cochlear implantation for an infant or child who is deaf. Benefits from hearing aids and cochlear implants in children with neural hearing loss have also been documented. The benefit of acoustic amplification for children with neural hearing loss is variable.^{28,102} Thus, a trial fitting is indicated for infants with neural hearing loss until the usefulness of the fitting can be determined. Neural hearing loss is a heterogeneous condition; the decision to continue or discontinue use of hearing aids should be made on the basis of the benefit derived from amplification. Use of cochlear implants in neural hearing loss is growing, and positive outcomes have been reported for many children.²⁸

Infants and young children with unilateral hearing loss should also be assessed for appropriateness of hearing aid fitting. Depending on the degree of residual hearing in unilateral loss, a hearing aid may or may not be indicated. Use of "contralateral routing of signals" amplification for unilateral hearing loss in children is not recommended.¹⁰³ Research is currently underway to determine how to best manage unilateral hearing loss in infants and young children.

The effect of otitis media with effusion (OME) is greater for infants with sensorineural hearing loss than for those with normal cochlear function.⁷³ Sensory or permanent conductive hearing loss is compounded by additional transient conductive hearing loss associated with OME. OME further reduces access to auditory cues necessary for the development of spoken English.

OME also negatively affects the prescriptive targets of the hearing aid fitting, decreasing auditory awareness and requiring adjustment of the amplification characteristics. Prompt referral to either the primary care physician or an otolaryngologist for treatment of persistent OME is indicated in infants with sensorineural hearing loss.¹⁰⁴ Definitive resolution of OME should never delay the fitting of an amplification device.^{73,105}

3. Medical and Surgical Intervention

Medical intervention is the process by which a physician provides medical diagnosis and direction for medical and/or surgical treatment options for hearing loss and/or related medical disorder(s) associated with hearing loss. Treatment varies from the removal of cerumen and the treatment of OME to long-term plans for reconstructive surgery and assessment of candidacy for cochlear implants. If necessary, surgical treatment of malformation of the outer and middle ears, including bone-anchored hearing aids, should be considered in the intervention plan for infants with permanent conductive or mixed hearing loss when a child reaches an appropriate age.

4. Communication Assessment and Intervention

Language is acquired with greater ease during certain sensitive periods of infant and toddler development.^{106–108} The process of language acquisition includes learning the precursors of language, such as the rules pertaining to selective attention and turn taking.^{20,109,110} Cognitive, social, and emotional development are influenced by the acquisition of language. Development in these areas is synergistic. A complete language evaluation should be performed at regular intervals for infants and toddlers with hearing loss. The evaluation should include an assessment of oral, manual, and/or visual mechanisms as well as cognitive abilities.

A primary focus of language intervention is to support families in fostering the communication abilities of their infants and toddlers who are deaf or hard of hearing.²⁰ Spoken and/or sign language development should be commensurate with the child's age and cognitive abilities and should include acquisition of phonologic (for spoken language), visual/spatial/motor (for signed language), morphologic, semantic, syntactic, and pragmatic skills, depending on the family's preferred mode of communication.

Early intervention professionals should follow family-centered principles to assist in developing communicative competence of infants and toddlers who are deaf or hard of hearing.^{111–113} Families should be provided with information specific to language development and access to peer and language models as well as family-involved activities that facilitate language development of children with normal hearing and children who are hard of

hearing or deaf.^{114,115} Depending on family choices, families should be offered access to children and adults with hearing loss who are appropriate and competent language models. Information on spoken language and signed language, such as American Sign Language¹¹⁶ and cued speech, should be provided.

E. Continued Surveillance, Screening, and Referral of Infants and Toddlers

Appendix 1 presents 11 risk indicators associated with either congenital or delayed-onset hearing loss. A single list of risk indicators is presented in the current JCIH statement, because there is significant overlap among those indicators associated with congenital/neonatal hearing loss and those associated with delayed-onset/acquired or progressive hearing loss. Heightened surveillance of all infants with risk indicators is, therefore, recommended. There is a significant change in the definition of risk indicator 3, which has been modified from NICU stay >48 hours to NICU stay >5 days. Consistent with JCIH 2000,³ the 2007 position statement recommends use of risk indicators for hearing loss for 3 purposes. Historically, the first use of risk indicators is for the identification of infants who should receive audiologic evaluation but who live in geographic locations (eg, developing nations, remote areas) where universal hearing screening is not yet available.^{3,19,21,24,25,64,117–123} This use has become less common as a result of the expansion of UNHS. The second purpose of risk indicator identification is to help identify infants who pass the neonatal screening but are at risk of developing delayed-onset hearing loss and, therefore, should receive ongoing medical, speech and language, and audiologic surveillance. Third, the risk indicators are used to detect infants who may have passed neonatal screening but have mild forms of permanent hearing loss.²⁵

Because some important indicators, such as family history of hearing loss, may not be determined during the course of UNHS,^{14,72} the presence of all risk indicators for acquired hearing loss should be determined in the medical home during early well-baby visits. Risk indicators that are marked with an asterisk in Appendix 1 are of greater concern for delayed-onset hearing loss. Early and more frequent assessment may be indicated for children with cytomegalovirus (CMV) infection,^{117,124,125} syndromes associated with progressive hearing loss,⁷² neurodegenerative disorders,⁷² trauma,^{126–128} or culture-positive postnatal infections associated with sensorineural hearing loss^{129,130}; for children who have received ECMO⁶⁴ or chemotherapy¹³¹; and when there is caregiver concern or a family history of hearing loss.¹⁶

For all infants with and without risk indicators for hearing loss, developmental milestones, hearing skills, and parent concerns about hearing, speech, and language skills should be monitored during routine medical care consistent with the AAP periodicity schedule.

The JCIH has determined that the previously recommended approach to

follow-up of infants with risk indicators for hearing loss only addressed children with identifiable risk indicators and failed to consider the possibility of delayed-onset hearing loss in children without identifiable risk indicators. In addition, concerns were raised about feasibility and cost associated with the JCIH 2000 recommendation for audiologic monitoring of all infants with risk indicators at 6-month intervals. Because approximately 400 000 infants are cared for annually in NICUs in the United States, and the JCIH 2000 recommendation included audiology assessments at 6-month intervals from 6 months to 36 months of age for all infants admitted to a NICU for >48 hours, an unreasonable burden was placed on both providers of audiology services and families. In addition, there was no provision for identification of delayed-onset hearing loss in infants without an identifiable risk indicator. Data from 2005 for 12 388 infants discharged from NICUs in the National Perinatal Information Network indicates that 52% of infants were discharged within the first 5 days of life and these infants were significantly less likely to have an identified risk indicator for hearing loss other than NICU stay. Therefore, the JCIH 2007 recommends an alternative, more inclusive strategy of surveillance of all children within the medical home based on the pediatric periodicity schedule. This protocol will permit the detection of children with either missed neonatal or delayed-onset hearing loss, irrespective of the presence or absence of a high-risk indicator.

The JCIH recognizes that an optimal surveillance and screening program within the medical home would include the following:

- At each visit, consistent with the AAP periodicity schedule, infants should be monitored for auditory skills, middle ear status, and developmental milestones (surveillance). Concerns elicited during surveillance should be followed by administration of a validated global screening tool.¹³² A validated global screening tool is administered at 9, 18, and 24 to 30 months to all infants or, if there is physician or parental concern about hearing or language, sooner.¹³²
- If an infant does not pass the speech-language portion of the global screening in the medical home or if there is physician or caregiver concern about hearing or spoken language development, the child should be referred immediately for further evaluation by an audiologist and a speech-language pathologist for a speech and language evaluation with validated tools.¹³²
- Once hearing loss is diagnosed in an infant, siblings who are at increased risk of having hearing loss should be referred for audiologic evaluation.^{14,75,133,134}
- All infants with a risk indicator for hearing loss (Appendix 1), regardless of surveillance findings, should be referred for an audiologic assessment at least once by 24 to 30 months of age. Children with risk indicators

highly associated with delayed-onset hearing loss, such as having received ECMO or having CMV infection, should have more frequent audiologic assessments.

- All infants for whom the family has significant concerns regarding hearing or communication should be promptly referred for an audiologic and speech-language assessment.
- A careful assessment of middle ear status (using pneumatic otoscopy and/or tympanometry) should be completed at all well-child visits, and children with persistent middle ear effusion lasting 3 months or longer should be referred for otologic evaluation.¹³⁵

F. Protecting the Rights of Infants and Families

Each agency or institution involved in the EHDI process shares responsibility for protecting infant and family rights in all aspects of UNHS, including access to information including potential benefits and risks in the family's native language, input into decision making, and confidentiality.⁷⁷ Families should receive information about childhood hearing loss in easily understood language. Families have the right to accept or decline hearing screening or any follow-up care for their newborn infant within the statutory regulations, just as they have for any other screening or evaluation procedures or intervention.

EHDI data merit the same level of confidentiality and security afforded all other health care and education information in practice and law. The infant's family has the right to confidentiality of the screening and follow-up assessments and the acceptance or rejection of suggested intervention(s). In compliance with federal and state laws, mechanisms should be established that ensure parental release and approval of all communications regarding the infant's test results, including those to the infant's medical home and early intervention coordinating agency and programs. The Health Insurance Portability and Accountability Act (Pub L No. 104-191 [1996]) regulations permit the sharing of health information among health care professionals.

G. Information Infrastructure

In the 2000 position statement,³ the JCIH recommended development of uniform state registries and national information databases incorporating standardized methodology, reporting, and system evaluation. EHDI information systems are to provide for the ongoing and systematic collection, analysis, and interpretation of data in the process of measuring and reporting associated program services (eg, screening, evaluation, diagnosis, and/or intervention). These systems are used to guide activities, planning, implementation, and evaluation of programs and to formulate research hypotheses.

EHDI information systems are generally authorized by legislators and

implemented by public health officials. These systems vary from a simple system collecting data from a single source to electronic systems that receive data from many sources in multiple formats. The number and variety of systems will likely increase with advances in electronic data interchange and integration of data, which will also heighten the importance of patient privacy, data confidentiality, and system security. Appropriate agencies and/or officials should be consulted for any projects regarding public health surveillance.⁶⁹

Federal and state agencies are collaborating in the standardization of data definitions to ensure the value of data sets and to prevent misleading or unreliable information. Information management is used to improve services to infants and their families; to assess the quantity and timeliness of screening, evaluation, and enrollment into intervention; and to facilitate collection of demographic data on neonatal and infant hearing loss.

The JCIH endorses the concept of a limited national database to permit documentation of the demographics of neonatal hearing loss, including prevalence and etiology across the United States. The information obtained from the information management system should assist both the primary health care professional and the state health agency in measuring quality indicators associated with program services (eg, screening, diagnosis, and intervention). The information system should provide the measurement tools to determine the degree to which each process is stable and sustainable and conforms to program benchmarks. Timely and accurate monitoring of relevant quality measures is essential.

Since 1999, the CDC and the Directors of Speech and Hearing Programs in State Health and Welfare Agencies (DSHPSHWA) have collected annual aggregate EHDI program data needed to address the national EHDI goals. In 1999, a total of 22 states provided data for the DSHPSHWA survey. Participation had increased to 48 states, 1 territory, and the District of Columbia in 2003. However, many programs have been unable to respond to all the questions on the survey because of lack of a statewide comprehensive data-management and reporting system.

The Government Performance and Results Act (GPRA) of 1993 (Public Law 103-62) requires that federal programs establish measurable goals approved by the US Office of Management and Budget (OMB) that can be reported as part of the budgetary process, thus linking future funding decisions with performance. The HRSA has modified its reporting requirements for all grant programs. The GPRA measures that must be reported to the OMB by the MCHB annually for the EHDI program are:

- The number of infants screened for hearing loss before discharge from the hospital
- The number of infants with a confirmed with hearing loss no later than 3 months of age

- The number of infants enrolled in a program of early intervention no later than 6 months of age
- The number of infants with confirmed or suspected hearing loss referred to an ongoing source of comprehensive health care (ie, medical home)
- The number of children with nonsyndromic hearing loss who have developmentally appropriate language and communication skills at school entry

One GPRA measure that must be reported to the OMB by the CDC annually for the EHDI program is:

- The percentage of newborn infants with a positive screening result for hearing loss who are subsequently lost to follow-up.

EHDI programs have made tremendous gains in their ability to collect, analyze, and interpret data in the process of measuring and reporting associated program services. However, only a limited number of EHDI programs are currently able to accurately report the number of infants screened, evaluated, and enrolled in intervention, the age of time-related objectives (eg, screening by 1 month of age), and the severity or laterality of hearing loss. This is complicated by the lack of data standards and by privacy issues within the regulations of the Family Educational Rights and Privacy Act of 1974 (Pub L No. 93-380).

Given the current lack of standardized and readily accessible sources of data, the CDC EHDI program, in collaboration with the DSHPSHWA, developed a revised survey to obtain annual EHDI data from states and territories in a consistent manner to assess progress toward meeting the national EHDI goals and the Healthy People 2010 Objectives. In October 2006, the OMB, which is responsible for reviewing all government surveys, approved the new Early Hearing Detection and Intervention Hearing Screening and Follow-Up Survey. To facilitate this effort, the CDC EHDI Data Committee is establishing the minimum data elements and definitions needed for information systems to be used to assess progress toward the national EHDI goals.

The JCIH encourages the CDC and HRSA to continue their efforts to identify barriers and explore possible solutions with EHDI programs to ensure that children in each state who seek hearing-related services in states other than where they reside receive all recommended screening and follow-up services. EHDI systems should also be designed to promote the sharing of data regarding early hearing loss through integration and/or linkage with other child health information systems. The CDC currently provides funds to integrate the EHDI system with other state/territorial screening, tracking, and surveillance programs that identify children with special health care needs. Grantees of the MCHB are encouraged to link hearing screening data

with such child health data sets as electronic birth certificate, vital statistics, birth defects registries, metabolic or newborn dried “blood spot” screening, immunization registries, and others.

To promote the best use of public health resources, EHDI information systems should be periodically evaluated, and such evaluations should include recommendations for improving quality, efficiency, and usefulness. The appropriate evaluation of public health surveillance systems becomes paramount as these systems adapt to revise case definitions, address new health-related events, adopt new information technology, ensure data confidentiality, and assess system security.⁶⁹

Currently, federal sources of systems support include Title V block grants to states for maternal and child health care services, Title XIX (Medicaid) federal and state funds for eligible children, and competitive US Department of Education personnel preparation and research grants. The NIDCD provides grants for research related to early identification and intervention for children who are deaf or hard of hearing.¹³⁹

Universities should assume responsibility for special-track, interdisciplinary, professional education programs for early intervention for infants and children with hearing loss. Universities should also provide training in family systems, the grieving process, cultural diversity, auditory skill development, and deaf culture. There is a critical need for in-service and preservice training of professionals related to EHDI programs, which is particularly acute for audiologists and early interventionists with expertise in hearing loss. This training will require increased and sustained funding for personnel preparation.

H. Benchmarks and Quality Indicators

The JCIH supports the concept of regular measurements of performance and recommends routine monitoring of these measures for interprogram comparison and continuous quality improvement. Performance benchmarks represent a consensus of expert opinion in the field of newborn hearing screening and intervention. The benchmarks are the minimal requirements that should be attained by high-quality EHDI programs. Frequent measures of quality permit prompt recognition and correction of any unstable component of the EHDI process.¹⁴⁰

1. Quality Indicators for Screening

- Percentage of all newborn infants who complete screening by 1 month of age. Recommended benchmark is >95% (age correction for preterm infants is acceptable).
- Percentage of all newborn infants who fail initial screening and fail any subsequent rescreening before comprehensive audiologic evaluation. Recommended benchmark is <4%.

2. Quality Indicators for Confirmation of Hearing Loss

- Of infants who fail initial screening and any subsequent rescreening, the percentage who complete a comprehensive audiologic evaluation by 3 months of age. Recommended benchmark is 90%.
- For families who elect amplification, the percentage of infants with confirmed bilateral hearing loss receiving amplification devices within 1 month of confirmation of hearing loss. Recommended benchmark is 95%.

3. Quality Indicators for Early Intervention

- For infants with confirmed hearing loss who qualify for Part C, the percentage for whom parents have signed an IFSP no later than 6 months of age. Recommended benchmark is 90%.
- For children with acquired or late-identified hearing loss, the percentage for whom parents have signed an IFSP within 45 days of the diagnosis. Recommended benchmark is 95%.
- The percentage of infants with confirmed hearing loss who receive the first developmental assessment using standardized assessment protocols (not criterion reference checklists) for language, speech, and nonverbal cognitive development no later than 12 months of age. Recommended benchmark is 90%.

IV. CURRENT CHALLENGES, OPPORTUNITIES, AND FUTURE DIRECTIONS

Despite the tremendous progress made since 2000, the following are challenges to the success of the EHDI system:

1. Challenges

All of the following listed challenges are considered important for the future development of successful EHDI systems:

- Too many children are lost between the failed screening and the rescreening and between the failed rescreening and the diagnostic evaluation.
- There is a shortage of professionals with skills and expertise in both pediatrics and hearing loss, including audiologists, deaf educators, speech-language pathologists, early intervention professionals, and physicians.
- There is often a lack of timely referral for diagnosis of, and intervention for, suspected hearing loss in children.
- Consistent and stable state and federal funding is needed for program sustainability.

- When compared with services provided for adults, pediatric services in all specialties are poorly reimbursed.
- Access to uniform Part C services is inadequate among states and within states.
- There is a lack of integrated state data-management and tracking systems.
- Demographics and cultural diversity are rapidly changing.
- Funding for hearing aids, loaner programs, cochlear implants, and FM systems is needed.
- There is a lack of specialized services for children with multiple disabilities and hearing loss.
- Children may not qualify for services (state Part C guidelines) before demonstrating language delays (prevention model vs deficit model).
- Children may not qualify for assistive technology (prevention model vs deficit model).
- There is a lack of in-service education for key professionals.
- There are regulatory barriers to sharing information among providers and among states.
- No national standards exist for the calibration of OAE or ABR instrumentation, and there is a lack of uniform performance standards.

2. Opportunities for System Development and Research

- Establish programs to ensure the development of communication for infants and children with all degrees and types of hearing loss, allowing them access to all educational, social, and vocational opportunities throughout their life span.
- Develop improved, rapid, reliable screening technology designed to differentiate specific types of hearing loss.
- Develop and validate screening technologies for identifying minimal hearing loss.
- Develop state data-management systems with the capacity for the accurate determination of the prevalence for delayed-onset or progressive hearing loss.
- Develop state data-tracking systems to follow infants with suspected and confirmed hearing loss through individual state EHDI programs.
- Track the certification credentials of the service providers for children with confirmed hearing loss receiving Part C early intervention services and early childhood special education.
- Track genetic, environmental, and pharmacologic factors that contribute to hearing loss, thus allowing for tailored prevention and intervention strategies.
- Continue to refine electrophysiologic diagnostic techniques, algorithms, and equipment to enable frequency-specific threshold assessment for use with very young infants.

- Continue to refine techniques to improve the selection and fitting of appropriate amplification devices in infants and young children.
- Conduct translational research pertaining to young children with hearing loss, in particular, genetic, diagnostic, and outcomes studies.
- Initiate prospective population-based studies to determine the prevalence and natural history of auditory neural conduction disorders.
- Conduct efficacy studies to determine appropriate early intervention strategies for infants and children with all degrees and types of hearing loss.
- Conduct additional studies on the efficacy of intervention for infants and children who receive cochlear implants younger than 2 years.
- Conduct additional studies on the efficacy of hearing aid use in infants and children younger than 2 years.
- Conduct additional studies of the auditory development of children who have appropriate amplification devices in early life.
- Expand programs within health, social service, and education agencies associated with early intervention and Head Start programs to accommodate the needs of the increasing numbers of early-identified children.
- Adapt education systems to capitalize on the abilities of children with hearing loss who have benefited from early identification and intervention.
- Develop genetic and medical procedures that will determine more rapidly the etiology of hearing loss.
- Ensure transition from Part C (early intervention) to Part B (education) in ways that encourage family participation and ensure minimal disruption of child and family services.
- Study the effects of parents' participation in all aspects of early intervention.
- Test the utility of a limited national dataset and develop nationally accepted indicators of EHDI system performance.
- Encourage the identification and development of centers of expertise where specialized care is provided in collaboration with local service providers.
- Obtain the perspectives of individuals who are deaf or hard of hearing in developing policies regarding medical and genetic testing and counseling for families who carry genes associated with hearing loss.¹⁴¹

V. CONCLUSIONS

Since the JCIH 2000 statement, tremendous and rapid progress has been made in the development of EHDI systems as a major public health initiative. The percentage of infants screened annually in the United States has increased from 38% to 95%. The collaboration at all levels of professional organizations, federal and state government, hospitals, medical homes, and

families has contributed to this remarkable success. New research initiatives to develop more sophisticated screening and diagnostic technology, improved digital hearing aid and FM technologies, speech-processing strategies in cochlear implants, and early intervention strategies continue. Major technological breakthroughs have been made in facilitating the definitive diagnosis of both genetic and nongenetic etiologies of hearing loss. In addition, outcomes studies to assess the long-term outcomes of special populations, including infants and children with mild and unilateral hearing loss, neural hearing loss, and severe or profound hearing loss managed with cochlear implants, are providing information on the individual and societal impact and the factors that contribute to an optimized outcome. It is apparent, however, that there are still serious challenges to be overcome and system barriers to be conquered to achieve optimal EHDI systems in all states in the next 5 years. Follow-up rates remain poor in many states, and funding for amplification in children is inadequate. Funding to support outcome studies is necessary to guide intervention and to determine factors other than hearing loss that affect child development. The ultimate goal, to optimize communication, social, academic, and vocational outcomes for each child with permanent hearing loss, must remain paramount.

ACKNOWLEDGEMENTS

The Year 2007 Position Statement was developed by the Joint Committee on Infant Hearing (JCIH). Joint committee member organizations and their respective representatives who prepared this statement include (in alphabetical order): the Alexander Graham Bell Association for the Deaf and Hard of Hearing (Jackie Busa, BA and Judy Harrison, MA); the American Academy of Audiology (Jodi Chappell; Christine Yoshinaga-Itano, PhD; and Alison Grimes, AuD); the American Academy of Otolaryngology-Head and Neck Surgery (Patrick E. Brookhouser, MD, and Stephen Epstein, MD); the American Academy of Pediatrics (Albert Mehl, MD, and Betty Vohr, MD [Chairperson, March 2005–present]); the American Speech-Language-Hearing Association (Judith Gravel, PhD [Chair March 2003–March 2005]; Jack Roush, PhD; and Judith Widen, PhD); the Council of Education of the Deaf, whose member organizations include the Alexander Graham Bell Association for the Deaf and Hard of Hearing, American Society for Deaf Children, Conference of Educational Administrators of Schools and Programs for the Deaf, Convention of American Instructors of the Deaf, National Association of the Deaf, and Association of College Educators of the Deaf and Hard of Hearing (Beth S. Benedict, PhD; and Bobbie Scoggins, EdD); and the Directors of Speech and Hearing Programs in State Health and Welfare Agencies (Michelle King, MS; Linda Pippins, MCD; and David H. Savage, MSC). Ex officio contributors to the JCIH include Jill Ackermann, MS; Amy Gibson, MS, RN; and Thomas Tonniges, MD (American Academy of Pediatrics); and Pamela Mason, MEd

(American Speech-Language-Hearing Association). We also acknowledge the contribution of John Eichwald, MA, and Irene Forsman, MS, RN.

Joint committee member organizations that adopt this statement include (in alphabetical order): the Alexander Graham Bell Association for the Deaf and Hard of Hearing, the American Academy of Audiology, the American Academy of Otolaryngology-Head and Neck Surgery, the American Academy of Pediatrics, the American Speech-Language-Hearing Association, the Council on Education of the Deaf (see individual organizations listed above), and the Directors of Speech and Hearing Programs in State Health and Welfare Agencies.

Appendix I. Risk Indicators Associated with Permanent Congenital, Delayed-Onset or Progressive Hearing Loss in Childhood

1. Caregiver concern* regarding hearing, speech, language, or developmental delay.⁶²
2. Family history* of permanent childhood hearing loss.^{24,136}
3. Neonatal intensive care of >5 days, or any of the following regardless of length of stay: ECMO,* assisted ventilation, exposure to ototoxic medications (gentamycin and tobramycin) or loop diuretics (furosemide/lasix), and hyperbilirubinemia requiring exchange transfusion.^{64,130}
4. In-utero infections, such as CMV,* herpes, rubella, syphilis, and toxoplasmosis.^{64-67,124,125}
5. Craniofacial anomalies, including those involving the pinna, ear canal, ear tags, ear pits, and temporal bone anomalies.²⁴
6. Physical findings, such as white forelock, associated with a syndrome known to include a sensorineural or permanent conductive hearing loss.²⁴
7. Syndromes associated with hearing loss or progressive or late-onset hearing loss,* such as neurofibromatosis, osteopetrosis, and Usher syndrome.¹³⁰ Other frequently identified syndromes include Waardenburg, Alport, Pendred, and Jervell and Lange-Nielson.⁷²
8. Neurodegenerative disorders,* such as Hunter syndrome, or sensory motor neuropathies, such as Friedreich ataxia and Charcot-Marie-Tooth syndrome.¹³⁰
9. Culture-positive postnatal infections associated with sensorineural hearing loss,* including confirmed bacterial and viral (especially herpes viruses and varicella) meningitis.^{129,130,137,138}
10. Head trauma, especially basal skull/temporal bone fracture* requiring hospitalization.¹²⁶⁻¹²⁸
11. Chemotherapy¹³¹

* Risk indicators that are marked with an asterisk are of greater concern for delayed-onset hearing loss.

Appendix 2: Algorithm for Hearing Screening. Available at: <http://www.medicalhomeinfo.org/screening/>
Screen%20Materials/Algorithm.pdf

Universal Newborn Hearing Screening, Diagnosis, and Intervention Guidelines for Pediatric Medical Home Providers

January 2003

REFERENCES

1. Holden-Pitt L, Diaz J. Thirty years of the annual survey of deaf and hard of hearing children and youth: a glance over the decades. *Am Ann Deaf*. 1998;143:72–76
2. American Academy of Pediatrics, Medical Home Initiatives for Children With Special Needs Project Advisory Committee. The medical home. *Pediatrics*. 2002;110:184–186
3. Joint Committee on Infant Hearing. Year 2000 position statement: principles and guidelines for early hearing detection and intervention programs. *Pediatrics*. 2000;106:798–817
4. US Department of Health and Human Services, Office of Disease Prevention and Health Promotion. *Healthy People 2000: National Health Promotion and Disease Prevention Objectives*. Washington, DC: US Government Printing Office; 1991. Available at: <http://odphp.osophs.dhhs.gov/pubs/hp2000/hppub97.htm>. Accessed January 24, 2007
5. National Institutes of Health. *Early Identification of Hearing Impairment in Infants and Young Children*. NIH Consensus Development Conference Statement. Bethesda, MD: National Institutes of Health; 1993:1–24. Available at: <http://consensus.nih.gov/1993/1993HearingInfantsChildren092html.htm>. Accessed January 24, 2007
6. Joint Committee on Infant Hearing. 1994 position statement. *AAO-HNS Bulletin*. 1994;12:13
7. Joint Committee on Infant Hearing. 1994 position statement. *ASHA*. 1994;36:38–41
8. American Academy of Pediatrics, Task Force on Newborn and Infant Hearing. Newborn and infant hearing loss: detection and intervention. *Pediatrics*. 1999;103:527–530
9. US Department of Health and Human Services, Office of Disease Prevention and Health Promotion. *Healthy People 2010. Vol II: Objectives for Improving Health*. 2nd ed. Rockville, MD: Office of Disease Prevention and Health Promotion, US Department of Health and Human Services; 2000
10. Yoshinaga-Itano C. Efficacy of early identification and early intervention. *Semin Hear*. 1995;16:115–123
11. Yoshinaga-Itano C. Levels of evidence: universal newborn hearing screening (UNHS) and early hearing detection and intervention systems (EHDI). *J Commun Disord*. 2004;37:451–465
12. Yoshinaga-Itano C, Sedey AL, Coulter DK, Mehl AL. Language of early- and later-identified children with hearing loss. *Pediatrics*. 1998;102:1161–1171
13. Yoshinaga-Itano C, Coulter D, Thomson V. The Colorado Newborn

- Hearing Screening Project: effects on speech and language development for children with hearing loss. *J Perinatol*. 2000;20:S132–S137
14. White K. The current status of EHDI programs in the United States. *Ment Retard Dev Disabil Res Rev*. 2003;9:79–88
 15. American Academy of Pediatrics, Task Force on Improving the Effectiveness of Newborn Hearing Screening, Diagnosis, and Intervention. Universal newborn hearing screening, diagnosis, and intervention: guidelines for pediatric medical home providers. Elk Grove Village, IL: American Academy of Pediatrics; 2003. Available at: [http://www.medicalhomeinfo.org/screening/Screen% 20Materials/Algorithm.pdf](http://www.medicalhomeinfo.org/screening/Screen%20Materials/Algorithm.pdf). Accessed January 23, 2007
 16. American Academy of Pediatrics, Committee on Practice and Ambulatory Medicine. Recommendations for preventive pediatric health care. *Pediatrics*. 2000;105:645–646
 17. Fletcher RH, Fletcher SW, Wagner EW. *Clinical Epidemiology: The Essentials*. 2nd ed. Baltimore, MD: Williams & Wilkins; 1988
 18. Sackett DL, Hayes RB, Tugwell P. *Clinical Epidemiology: A Basic Science for Clinical Medicine*. 2nd ed. Boston, MA: Little Brown & Co; 1991
 19. Norton SJ, Gorga MP, Widen JE, Folsom RC, Sininger Y, Cone-Wesson B, et al. Identification of neonatal hearing impairment: evaluation of transient evoked otoacoustic emission, distortion product otoacoustic emission, and auditory brain stem response test performance. *Ear Hear*. 2000;21:508–528
 20. Carney AE, Moeller MP. Treatment efficacy: hearing loss in children. *J Speech Lang Hear Res*. 1998;41:S61–S84
 21. D’Agostino JA, Austin L. Auditory neuropathy: a potentially under-recognized neonatal intensive care unit sequela. *Adv Neonatal Care*. 2004;4:344–353
 22. Sininger YS, Hood LJ, Starr A, Berlin CI, Picton TW. Hearing loss due to auditory neuropathy. *Audiol Today*. 1995;7:10–13
 23. Starr A, Sininger YS, Pratt H. The varieties of auditory neuropathy. *J Basic Clin Physiol Pharmacol*. 2000;11:215–230
 24. Cone-Wesson B, Vohr BR, Sininger YS, et al. Identification of neonatal hearing impairment: infants with hearing loss. *Ear Hear*. 2000;21:488–507
 25. Johnson JL, White KR, Widen JE, et al. A multicenter evaluation of how many infants with permanent hearing loss pass a two-stage otoacoustic emissions/automated auditory brainstem response newborn hearing screening protocol. *Pediatrics*. 2005;116:663–672
 26. Berlin CI, Hood L, Morlet T, Rose K, Brashears S. Auditory neuropathy / dys-synchrony: diagnosis and management. *Ment Retard Dev Disabil Res Rev*. 2003;9:225–231
 27. Doyle KJ, Sininger Y, Starr A. Auditory neuropathy in childhood. *Laryngoscope*. 1998;108:1374–1377

28. Rance G. Auditory neuropathy/dys-synchrony and its perceptual consequences. *Trends Amplif.* 2005;9:1–43
29. American Speech-Language-Hearing Association. The use of FM amplification instruments for infants and preschool children with hearing impairment. *ASHA.* 1991;33(Suppl 5):1–2. Available at: http://www.asha.org/NR/rdonlyres/226A8C6D-5275-44CC-BFB5-7E0AEA133849/0/18847_1.pdf. Accessed January 24, 2007
30. American Speech-Language Hearing Association, Joint Committee of ASHA and Council on Education of the Deaf. Service provision under the Individuals with Disabilities Education Act (IDEA–Part H) to children who are deaf and hard of hearing ages birth to 36 months. *ASHA.* 1994;36:117–121
31. Calderon R, Barges J, Sidman S. Characteristics of hearing families and their young deaf and hard of hearing children. Early intervention follow-up. *Am Ann Deaf.* 1998;143:347–362
32. Pediatric Working Group. Amplification for infants and children with hearing loss. *Am J Audiol.* 1996;5:53–68
33. American Academy of Pediatrics, Committee on Children With Disabilities. Care coordination in the medical home: integrating health and related systems of care for children with special health care needs. *Pediatrics.* 2005;116:1238–1244
34. Finitzo T, Albright K, O’Neal J. The newborn with hearing loss: detection in the nursery. *Pediatrics.* 1998;102:1452–1460
35. Mason JA, Herrmann KR. Universal infant hearing screening by automated auditory brainstem response measurement. *Pediatrics.* 1998;101:221–228
36. Prieve B, Dalzell L, Berg A, et al. The New York State universal newborn hearing screening demonstration project: outpatient outcome measures. *Ear Hear.* 2000;21:104–117
37. Vohr BR, Carty LM, Moore PE, Letourneau K. The Rhode Island Hearing Assessment Program: experience with statewide hearing screening (1993–1996). *J Pediatr.* 1998;133:353–357
38. Doyle KJ, Burggraaff B, Fujikawa S, Kim J, MacArthur CJ. Neonatal hearing screening with otoscopy, auditory brain stem response, and otoacoustic emissions. *Otolaryngol Head Neck Surg.* 1997;116:597–603
39. Hyde ML, Davidson MJ, Alberti PW. Auditory test strategy. In: Jacobson JT, Northern JL, eds. *Diagnostic Audiology.* Austin, TX: Pro-ed; 1991: 295–322
40. Hyde MD, Sininger YS, Don M. Objective detection and analysis of auditory brainstem response: an historical perspective. *Semin Hear.* 1998; 19:97–113
41. Eilers RE, Miskiel E, Ozdamar O, Urbano R, Widen JE. Optimization of automated hearing test algorithms: simulations using an infant response model. *Ear Hear.* 1991;12:191–198

42. Herrmann BS, Thornton AR, Joseph JM. Automated infant hearing screening using the ABR. Development and validation. *Am J Audiol.* 1995;4:6–14
43. McFarland WH, Simmons FB, Jones FR. An automated hearing screening technique for newborns. *J Speech Hear Disord.* 1980;45:495–503
44. Ozdamar O, Delgado RE, Eilers RE, Urbano RC. Automated electrophysiologic hearing testing using a threshold-seeking algorithm. *J Am Acad Audiol.* 1994;5:77–88
45. Pool KD, Finitzo T. Evaluation of a computer-automated program for clinical assessment of the auditory brain stem response. *Ear Hear.* 1989; 10:304–310
46. Benjamini Y, Yekutieli D. Quantitative trait Loci analysis using the false discovery rate. *Genetics.* 2005;171:783–790
47. Hochberg Y, Benjamini Y. More powerful procedures for multiple significance testing. *Stat Med.* 1990;9:811–818
48. Zhang JH, Chung TD, Oldenburg KR. A simple statistical parameter for use in evaluation and validation of high throughput screening assays. *J Biomol Screen.* 1999;4:67–73
49. Gravel JS, Karma P, Casselbrant ML, Marchisio P, Andalibi A, Passali D, et al. Recent advances in otitis media. 7. Diagnosis and screening. *Ann Otol Rhinol Laryngol Suppl.* 2005;194:104–113
50. Sininger YS, Abdala C, Cone-Wesson B. Auditory threshold sensitivity of the human neonate as measured by the auditory brainstem response. *Hear Res.* 1997;104:27–38
51. Arehart KH, Yoshinaga-Itano C, Thomson V, Gabbard SA, Brown AS. State of the states: the status of universal newborn screening, assessment, and intervention systems in 16 states. *Am J Audiol.* 1998;7:101–114
52. Gravel J, Berg A, Bradley M, et al. New York State universal newborn hearing screening demonstration project: effects of screening protocol on inpatient outcome measures. *Ear Hear.* 2000;21:131–140
53. Mehl AL, Thomson V. Newborn hearing screening: the great omission. *Pediatrics.* 1998;101(1):e4. Available at: www.pediatrics.org/cgi/content/full/101/1/e4. Accessed January 23, 2007
54. Stark AR and American Academy of Pediatrics, Committee on Fetus and Newborn. Levels of neonatal care. *Pediatrics.* 2004;114:1341–1347
55. Berg AL, Spitzer JB, Towers HM, Bartosiewicz C, Diamond BE. Newborn hearing screening in the NICU: profile of failed auditory brainstem response/passed otoacoustic emission. *Pediatrics.* 2005;116:933–938
56. Shapiro SM. Bilirubin toxicity in the developing nervous system. *Pediatr Neurol.* 2003;29:410–421
57. Starr A, Picton TW, Sininger Y, Hood LJ, Berlin CI. Auditory neuropathy. *Brain.* 1996;119:741–753
58. Stapells DR, Gravel JS, Martin BA. Thresholds for auditory brain stem responses to tones in notched noise from infants and young children

- with normal hearing or sensorineural hearing loss. *Ear Hear.* 1995;16:361–371
59. Keefe DH, Gorga MP, Neely ST, Zhao F, Vohr BR. Ear-canal acoustic admittance and reflectance measurements in human neonates. II. Predictions of middle-ear dysfunction and sensorineural hearing loss. *J Acoust Soc Am.* 2003;113:407–422
 60. Boppana SB, Fowler KB, Pass RF, et al. Congenital cytomegalovirus infection: association between virus burden in infancy and hearing loss. *J Pediatr.* 2005;146:817–823
 61. Nagy A, Endreffy E, Streitman K, Pinter S, Pusztai R. Incidence and outcome of congenital cytomegalovirus infection in selected groups of preterm and full-term neonates under intensive care. *In Vivo.* 2004;18:819–823
 62. Roizen NJ. Etiology of hearing loss in children. Nongenetic causes. *Pediatr Clin North Am.* 1999;46:49–64, x
 63. Karchmer MA, Allen TE. The functional assessment of deaf and hard of hearing students. *Am Ann Deaf.* 1999;144:68–77
 64. Fligor BJ, Neault MW, Mullen CH, Feldman HA, Jones DT. Factors associated with sensorineural hearing loss among survivors of extracorporeal membrane oxygenation therapy. *Pediatrics.* 2005;115:1519–1528
 65. Fowler K, Stagno S, Pass R, Britt W, Boll T, Alford C. The outcome of congenital cytomegalovirus infection in relation to maternal antibody status. *N Engl J Med.* 1992;326:663–667
 66. Madden C, Wiley S, Schleiss M, et al. Audiometric, clinical and educational outcomes in a pediatric symptomatic congenital cytomegalovirus (CMV) population with sensorineural hearing loss. *Int J Pediatr Otorhinolaryngol.* 2005;69:1191–1198
 67. Rivera LB, Boppana SB, Fowler KB, Britt WJ, Stagno S, Pass RF. Predictors of hearing loss in children with symptomatic congenital cytomegalovirus infection. *Pediatrics.* 2002;110:762–767
 68. Reefhuis J, Honein MA, Whitney CG, et al. Risk of bacterial meningitis in children with cochlear implants. *N Engl J Med.* 2003;349:435–445
 69. Centers for Disease Control and Prevention. Pneumococcal vaccination for cochlear implant candidates and recipients: updated recommendations of the Advisory Committee on Immunization Practices. *MMWR Morb Mortal Wkly Rep.* 2003;52:739–740
 70. Morzaria S, Westerberg BD, Kozak FK. Evidence-based algorithm for the evaluation of a child with bilateral sensorineural hearing loss. *J Otolaryngol.* 2005;34:297–303
 71. Preciado DA, Lawson L, Madden C, et al. Improved diagnostic effectiveness with a sequential diagnostic paradigm in idiopathic pediatric sensorineural hearing loss. *Otol Neurotol.* 2005;26:610–615
 72. Nance WE. The genetics of deafness. *Mental Retard Dev Disabil Res Rev.* 2003;9:109–119

73. Brookhouser P, Worthington D, Kelly W. Fluctuating and/or progressive sensorineural hearing loss in children. *Laryngoscope*. 1994;104: 958–964
74. Denoyelle F, Marlin S, Weil D, et al. Clinical features of the prevalent form of childhood deafness, DFNB1, due to a connexin-26 gene defect: implications for genetic counselling. *Lancet*. 1999;353(9161):1298–1303
75. Nance WE, Kearsey MJ. Relevance of connexin deafness (DFNB1) to human evolution. *Am J Hum Genet*. 2004;74:1081–1087
76. Santos RL, Aulchenko YS, Huygen PL, et al. Hearing impairment in Dutch patients with connexin 26 (GJB2) and connexin 30 (GJB6) mutations. *Int J Pediatr Otorhinolaryngol*. 2005;69:165–174
77. National Institute on Deafness and Other Communicating Disorders. *Communicating Informed Consent to Individuals Who Are Deaf or Hard-of-Hearing*. Bethesda, MD: National Institute on Deafness and Other Communicating Disorders, National Institutes of Health; 1999. NIH Publication No. 00-4689
78. Buchman CA, Roush PA, Teagle HF, Brown CJ, Zdanski CJ, Grose JH. Auditory neuropathy characteristics in children with cochlear nerve deficiency. *Ear Hear*. 2006;27:399–408
79. Preciado DA, Lim LH, Cohen AP, et al. A diagnostic paradigm for childhood idiopathic sensorineural hearing loss. *Otolaryngol Head Neck Surg*. 2004;131:804–809
80. Johnson DH. *Deafness and Vision Disorders: Anatomy and Physiology, Assessment Procedures, Ocular Anomalies, and Educational Implications*. Springfield, IL: Charles C. Thomas Publisher; 1999
81. Traxler CB. The Stanford Achievement Test, 9th Edition: national norming and performance standards for deaf and hard-of-hearing students. *J Deaf Stud Deaf Educ*. 2000;5:337–348
82. Mayne A, Yoshinaga-Itano C, Sedey AL, Carey A. Expressive vocabulary development of infants and toddlers who are deaf or hard of hearing. *The Volta Review*. 1998;100:1–28
83. Mayne AM, Yoshinaga-Itano C, Sedey AL. Receptive vocabulary development of infants and toddlers who are deaf or hard of hearing. *The Volta Review*. 1998;100:29–52
84. Pipp-Siegel S, Sedey AL, VanLeeuwen AM, Yoshinaga-Itano C. Mastery motivation and expressive language in young children with hearing loss. *J Deaf Stud Deaf Educ*. 2003;8:133–145
85. Yoshinaga-Itano C, Coulter D, Thomson V. Developmental outcomes of children with hearing loss born in Colorado hospitals with and without universal newborn hearing screening programs. *Semin Neonatol*. 2001;6: 521–529
86. Apuzzo ML, Yoshinaga-Itano C. Early identification of infants with significant hearing loss and the Minnesota Child Development Inventory. *Semin Hear*. 1995;16:124–137

87. Yoshinaga-Itano C, Apuzzo ML. The development of deaf and hard of hearing children identified early through the high-risk registry. *Am Ann Deaf*. 1998;143:416–424
88. Yoshinaga-Itano C, Apuzzo ML. Identification of hearing loss after age 18 months is not early enough. *Am Ann Deaf*. 1998;143:380–387
89. Yoshinaga-Itano C. The social-emotional ramifications of universal newborn hearing screening: early identification and intervention of children who are deaf or hard of hearing. In: *Proceedings of the Second International Pediatric Conference: A Sound Foundation Through Early Amplification; November 8–10, 2001; Chicago, IL*. Stafa, Switzerland: Phonak Inc; 2001. Available at: www.phonak.com/professional/informationpool/proceedings2001.htm. Accessed January 23, 2007
90. Moeller MP. Early intervention and language development in children who are deaf and hard of hearing. *Pediatrics*. 2000;106(3):e43. Available at: www.pediatrics.org/cgi/content/full/106/3/e43. Accessed January 25, 2007
91. Kennedy C, McCann D, Campbell MJ, Kimm L, Thornton R. Universal newborn screening for permanent childhood hearing impairment: an 8-year follow-up of a controlled trial. *Lancet*. 2005;366:660–662
92. Bodner-Johnson B, Sass-Lehrer M. *The Young Deaf or Hard of Hearing Child*. Baltimore, MD: Paul H Brookes; 2003
93. Yoshinaga-Itano C, Sedey A. Early speech development in children who are deaf or hard-of-hearing: interrelationships with language and hearing. *The Volta Review*. 1998;100:181–211
94. Yoshinaga-Itano C. Early intervention after universal neonatal hearing screening: impact on outcomes. *Ment Retard Dev Disabil Res Rev*. 2003;9:252–266
95. Yoshinaga-Itano C. From screening to early identification and intervention: discovering predictors to successful outcomes for children with significant hearing loss. *J Deaf Stud Deaf Educ*. 2003;8:11–30
96. Yoshinaga-Itano C, Abdala de Uzategui C. Early identification and social emotional factors of children with hearing loss and children screened for hearing loss. In: Kurtzer-White E, Luterman D, eds. *Early Childhood Deafness*. Baltimore, MD: York Press; 2001:13–28
97. Calderon R. Parental involvement in deaf children's education programs as a predictor of child's language, early reading, and social-emotional development. *J Deaf Stud Deaf Educ*. 2000;5:140–155
98. Bess FH, Dodd-Murphy J, Parker RA. Children with minimal sensorineural hearing loss: prevalence, educational performance, and functional status. *Ear Hear*. 1998;19:339–354
99. Bess FH, Tharpe AM. An introduction to unilateral sensorineural hearing loss in children. *Ear Hear*. 1986;7:3–13
100. Bess FH, Tharpe AM. Unilateral hearing impairment in children. *Pediatrics*. 1984;74:206–216

101. American Speech-Language-Hearing Association. *Guidelines for the Audiologic Assessment of Children From Birth to 5 Years of Age*. Rockville, MD: American Speech-Language-Hearing Association; 2004. Available at: <http://www.asha.org/NR/rdonlyres/0BB7C840-27D2-4DC6-861B-1709ADD78BAF/0/v2GLAudAssessChild.pdf>. Accessed January 24, 2007
102. Rance G, Cone-Wesson B, Wunderlich J, Dowell R. Speech perception and cortical event related potentials in children with auditory neuropathy. *Ear Hear*. 2002;23:239–253
103. American Academy of Audiology. *Pediatric Amplification Protocol*. Reston, VA: American Academy of Audiology; 2003. Available at: <http://www.audiology.org/NR/rdonlyres/53D26792-E321-41AF-850F-CC253310F9DB/0/pedamp.pdf>. Accessed January 24, 2007
104. Rosenfeld RM, Culpepper L, Doyle KJ, et al. Clinical practice guideline: otitis media with effusion. *Otolaryngol Head Neck Surg*. 2004;130(5 Suppl):S95–S118
105. Diefendorf AO, Gravel JS. Behavioral observation and visual reinforcement audiometry. In: Gerber SE, ed. *Handbook of Pediatric Audiology*. Washington, DC: Gallaudet University Press; 1996:55–83
106. Clark T. SKI*HI: applications for home-based intervention. In: Roush J, Matkin ND, eds. *Infants and Toddlers with Hearing Loss: Family-Centered Assessment and Intervention*. Baltimore, MD: York Press Inc; 1994:237–251
107. Mahshie SN. *Educating Deaf Children Bilingually*. Washington, DC: Gallaudet University Press; 1995
108. Sharma A, Tobey E, Dorman M, et al. Central auditory maturation and babbling development in infants with cochlear implants. *Arch Otolaryngol Head Neck Surg*. 2004;130:511–516
109. Kuhl PK, Andruski JE, Chistovich IA, et al. Cross-language analysis of phonetic units in language addressed to infants. *Science*. 1997;277:684–686
110. Kuhl PK, Williams KA, Lacerda F, Stevens KN, Lindblom B. Linguistic experience alters phonetic perception in infants by 6 months of age. *Science*. 1992;255:606–608
111. Baker-Hawkins S, Easterbrooks S. *Deaf and Hard of Hearing Students: Educational Service Delivery Guidelines*. Alexandria, VA: National Association of State Directors of Special Education; 1994
112. Bamford JM. Early intervention . . . what then? In: Bess FH, ed. *Children with Hearing Impairment: Contemporary Trends*. Nashville, TN: The Vanderbilt Bill Wilkerson Center Press; 1998:353–358
113. Fischer RM. The Mama Lere Home: Vanderbilt University. In: Roush J, Matkin ND, eds. *Infants and Toddlers with Hearing Loss: Family-Centered Assessment and Intervention*. Baltimore, MD: York Press Inc; 1994:195–213
114. Marschark M. *Raising and Educating a Deaf Child*. New York, NY: Oxford University Press; 1997

115. Thompson M. ECHI: the University of Washington, Seattle. In: Roush J, Natkin ND, eds. *Infants and Toddlers with Hearing Loss: Family-Centered Assessment and Intervention*. Baltimore, MD: York Press Inc; 1994:253–275
116. Pollack D, Goldberg D, Caleffe-Schenck N. *Educational Audiology for the Limited-Hearing Infant and Preschooler: An Auditory Verbal Program*. 3rd ed. Springfield, IL: Charles C. Thomas Publisher; 1997
117. Barbi M, Binda S, Caroppo S, et al. Multicity Italian study of congenital cytomegalovirus infection. *Pediatr Infect Dis J*. 2006;25:156–159
118. Barrenas ML, Jonsson B, Tuvemo T, Hellstrom PA, Lundgren M. High risk of sensorineural hearing loss in men born small for gestational age with and without obesity or height catch-up growth: a prospective longitudinal register study on birth size in 245,000 Swedish conscripts. *J Clin Endocrinol Metab*. 2005;90:4452–4456
119. Davis A, Hind S. The newborn hearing screening programme in England. *Int J Pediatr Otorhinolaryngol*. 2003;67(Suppl 1):S193–S196
120. Jacobson J, Jacobson C. Evaluation of hearing loss in infants and young children. *Pediatr Ann*. 2004;33:811–821
121. Mestan KK, Marks JD, Hecox K, Huo D, Schreiber MD. Neurodevelopmental outcomes of premature infants treated with inhaled nitric oxide. *N Engl J Med*. 2005;353:23–32
122. Robertson CM, Tyebkhan JM, Peliowski A, Etches PC, Cheung PY. Ototoxic drugs and sensorineural hearing loss following severe neonatal respiratory failure. *Acta Paediatr*. 2006;95:214–223
123. Vohr BR, Widen JE, Cone-Wesson B, et al. Identification of neonatal hearing impairment: characteristics of infants in the neonatal intensive care unit and well-baby nursery. *Ear Hear*. 2000;21:373–382
124. Nance WE, Lim BG, Dodson KM. Importance of congenital cytomegalovirus infections as a cause for pre-lingual hearing loss. *J Clin Virol*. 2006;35:221–225
125. Pass RF, Fowler KB, Boppana SB, Britt WJ, Stagno S. Congenital cytomegalovirus infection following first trimester maternal infection: symptoms at birth and outcome. *J Clin Virol*. 2006;35:216–220
126. Lew HL, Lee EH, Miyoshi Y, Chang DG, Date ES, Jerger JF. Brainstem auditory-evoked potentials as an objective tool for evaluating hearing dysfunction in traumatic brain injury. *Am J Phys Med Rehabil*. 2004;83:210–215
127. Vartialnen E, Karjalainen S, Karja J. Auditory disorders following head injury in children. *Acta Oto-Laryngologica*. 1985;99:529–536
128. Zimmerman WD, Ganzel TM, Windmill IM, Nazar GB, Phillips M. Peripheral hearing loss following head trauma in children. *Laryngoscope*. 1993;103:87–91
129. Arditi M, Mason EO Jr, Bradley JS, et al. Three-year multicenter surveillance of pneumococcal meningitis in children: clinical characteristics,

- and outcome related to penicillin susceptibility and dexamethasone use. *Pediatrics*. 1998;102:1087–1097
130. Roizen NJ. Nongenetic causes of hearing loss. *Ment Retard Dev Disabil Res Rev*. 2003;9:120–127
 131. Bertolini P, Lassalle M, Mercier G, et al. Platinum compound-related ototoxicity in children: long-term follow-up reveals continuous worsening of hearing loss. *J Pediatr Hematol Oncol*. 2004;26:649–655
 132. American Academy of Pediatrics, Council on Children With Disabilities, Section on Developmental Behavioral Pediatrics, Bright Futures Steering Committee, Medical Home Initiatives for Children With Special Needs Project Advisory Committee. Identifying infants and young children with developmental disorders in the medical home: an algorithm for developmental surveillance and screening. *Pediatrics*. 2006;118:405–420
 133. Fortnum H, Davis A. Epidemiology of permanent childhood hearing impairment in Trent Region, 1985–1993. *Br J Audiol*. 1997;31:409–446
 134. Orzan E, Polli R, Martella M, Vinanzi C, Leonardi M, Murgia A. Molecular genetics applied to clinical practice: the Cx26 hearing impairment. *Br J Audiol*. 1999;33:291–295
 135. American Academy of Pediatrics, Subcommittee on Otitis Media With Effusion, American Academy of Family Physicians, American Academy of Otolaryngology-Head and Neck Surgery. Otitis media with effusion. *Pediatrics*. 2004;113:1412–1429
 136. Morton CC, Nance WE. Newborn hearing screening—a silent revolution. *N Engl J Med*. 2006;354:2151–2164
 137. Bess FH. Children with unilateral hearing loss. *J Acad Rehab Audiol*. 1982;15:131–144
 138. Biernath KR, Reefhuis J, Whitney CG, Mann EA, Costa P, Eichwald J, et al. Bacterial meningitis among children with cochlear implants beyond 24 months after implantation. *Pediatrics*. 2006;117:284–289
 139. Roush J, Bess FH, Gravel J, Harrison M, Lenihan S, Marvelli A. Preparation of personnel to serve children with hearing loss and their families: current status and future needs. In: *2004 Summit on Deafness Proceedings: Spoken Language in the 21st Century. Predicting Future Trends in Deafness*; February 26–29, 2004; Washington, DC
 140. Agency for Health Care Policy and Research. *Using Clinical Practice Guidelines to Evaluate Quality of Care. Vol II: Methods*. Rockville, MD: US Department of Health and Human Services, Public Health Service; 1995. AHCPR Publication No. 95-0046
 141. Brick K. Genetics of deafness, deaf people and the past, present and future. Presented at the Workshop on the Genetics of Congenital Hearing Impairment; Centers for Disease Control and Prevention and Gallaudet University; June 7, 1999; Atlanta GA

"During an ultrasound, I listened to my baby's heartbeat for the first time."

—Stacey McGovern,
implanted at age 29

TREASURE

every moment of your child's life.

Stacey's third pregnancy was made even more memorable after she was implanted with the HiResolution® Bionic Ear System. The new HiRes Fidelity 120™ Sound Processing feature* is designed to help you appreciate the fullness and richness of life's precious moments.

Read more about Stacey's life with HiResolution Sound at
www.BionicEar.com/RealStories.

Connect and Discover™

*In U.S., optional feature for adults only. See pdaSpec insert for details.
©2007 Advanced Bionics Corporation. All Rights Reserved. Printed in U.S.A.
AUC07 2/09/2007 8/07

ADVANCED
BIONICS
A Boston Scientific Company