The Volta Review

Alexander Graham Bell Association
for the Deaf and Hard of Hearing

The

Is Auditory-Verbal Therapy Effective for Children with
Hearing Loss?

361

*By Dimity Dornan, Ba.Sp.Th., F.S.P.A.A., LSLS Cert. AVT;
Louise Hickson, B.Sp.Thy. (Hons), M.Aud., Ph.D.;
Bruce Murdoch, B.Sc. (Hons), Ph.D., D.Sc.;
Todd Houston, Ph.D., CCC-SLP, LSLS Cert. AVT;
and Gabriella Constantinescu, B.S.Path (Hons), Ph.D.*

Venturing Beyond the Sentence Level:
Narrative Skills in Children with Hearing Loss

389

*By Christina Reuterskiöld, Ph.D.;
Tina Ibertsson, Ph.D.; and Birgitta Sahlén, Ph.D.*

Effects of Auditory-Verbal Therapy for School-Aged
Children with Hearing Loss: An Exploratory Study

407

*By Elizabeth Fairgray, M.Sc., LSLS Cert. AVT;
Suzanne C. Purdy, Ph.D.; and Jennifer L. Smart, Ph.D.*

Use of Differential Reinforcement to Increase Hearing
Aid Compliance: A Preliminary Investigation

435

*By Sandie M. Bass-Ringdahl, Ph.D., CCC-A;
Joel E. Ringdahl, Ph.D.; and Eric W. Boelter, Ph.D.*

2010 AG Bell Research Symposium Proceedings

465

*"Re-Modeling the Deafened Cochlea for Auditory
Sensation: Advances and Obstacles"*

The Volta Review

Volume 110, Number 3

Fall 2010

ISSN 0042-8639

The Alexander Graham Bell Association for the Deaf and Hard of Hearing helps families, health care providers and education professionals understand childhood hearing loss and the importance of early diagnosis and intervention. Through advocacy, education, research and financial aid, AG Bell helps to ensure that every child and adult with hearing loss has the opportunity to listen, talk and thrive in mainstream society. With chapters located in the United States and a network of international affiliates, AG Bell supports its mission: Advocating Independence through Listening and Talking!

359 Editors' Preface

Joseph Smaldino, Ph.D., and Kathryn L. Schmitz, Ph.D.

Research

361 Is Auditory-Verbal Therapy Effective for Children with Hearing Loss?

By Dimity Dornan, Ba.Sp.Th., F.S.P.A.A., LSLS Cert. AVT;

Louise Hickson, B.Sp.Thy. (Hons), M.Aud., Ph.D.;

Bruce Murdoch, B.Sc. (Hons), Ph.D., D.Sc.;

Todd Houston, Ph.D., CCC-SLP, LSLS Cert. AVT; and

Gabriella Constantinescu, B.S.Path (Hons), Ph.D.

389 Venturing Beyond the Sentence Level: Narrative Skills in Children with Hearing Loss

By Christina Reuterskiöld, Ph.D., Tina Ibertsson, Ph.D., and

Birgitta Sahlén, Ph.D.

407 Effects of Auditory-Verbal Therapy for School-Aged Children with Hearing Loss: An Exploratory Study

By Elizabeth Fairgray, M.Sc., LSLS Cert. AVT; Suzanne C. Purdy, Ph.D.; and

Jennifer L. Smart, Ph.D.

435 Use of Differential Reinforcement to Increase Hearing Aid Compliance: A Preliminary Investigation

By Sandie M. Bass-Ringdahl, Ph.D., CCC-A; Joel E. Ringdahl, Ph.D.; and

Eric W. Boelter, Ph.D.

Literature Review

447 Concerns Regarding Direct-to-Consumer Hearing Aid Purchasing

By Suzanne H. Kimball, Au.D.

Book Reviews

459 Auditory-Verbal Practice: Toward a Family Centered Approach

By Kathryn Ritter, Ph.D., LSLS Cert. AVT

461 Developmental Language Disorders: Learning, Language, and the Brain

By Alan G. Kamhi, Ph.D.

2010 AG Bell Research Symposium Proceedings

- 465 Re-Modeling the Deafened Cochlea for Auditory Sensation:
Advances and Obstacles

Regular Features

- 487 Directory of Professional Programs
- 489 Information for Contributors to *The Volta Review*

Permission to Copy: The Alexander Graham Bell Association for the Deaf and Hard of Hearing, as copyright owner of this journal, allows single copies of an article to be made for personal use. This consent does not extend to posting on Web sites or other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works of any type, or for resale without the express written permission of the publisher. For more information, contact AG Bell at 3417 Volta Place, NW, Washington, DC 20007, email editor@agbell.org, or call (202) 337-5220 (voice) or (202) 337-5221 (TTY).

Joseph Smaldino, Ph.D.,
Editor

Kathryn L. Schmitz, Ph.D.,
Senior Associate Editor

Editors' Preface

For over 110 years *The Volta Review* has presented rigorous, scientific research exploring the listening and spoken language development of individuals who are deaf or hard of hearing. This edition of *The Volta Review* presents research on speech and language development as well as the effectiveness of a listening and spoken language approach.

"Venturing Beyond the Sentence Level" explores language growth of school-aged children with hearing loss within the context of oral narrative skill development. Results indicated that children with hearing loss were able to develop narratives similar to peers with typical hearing, although analysis did find poorer development of higher level language skills. A correlation was also found between the age of identification and narration abilities. "Use of Differential Reinforcement to Increase Hearing Aid Compliance" discusses a strategy used to encourage hearing aid use among small children, and how the successful use of hearing aids improved speech and language skills. And "Concerns Regarding Direct-to-Consumer Hearing Aids" summarizes recent research exploring the validity and accuracy of hearing aids purchased through independent distributors instead of a licensed audiologist. The results of each study will surprise you. Finally, a book review of "Developmental Language Disorders: Learning, Language, and the Brain" opens up a discussion about the focus of language development: should a practitioner's focus be on the child, or the end results?

The rest of this edition offers compelling evidence regarding the effectiveness of auditory-verbal practice. "Is Auditory-Verbal Therapy Effective for Children with Hearing Loss?" completes a 50-month longitudinal study following the language, literacy, and emotional development of children with hearing loss. The end results indicate that children with hearing loss who succeed with auditory-verbal therapy are well-adjusted and have language skills on par with their peers who have typical hearing. "Effects of Auditory-Verbal Therapy for School-Aged Children" offers the exploratory findings of a

study looking at the effectiveness of listening and spoken language intervention, indicating significant improvements after beginning therapy. Finally, a review of the new text, "Auditory-Verbal Practice: Toward a Family-Centered Approach," discusses the merits of engaging the family unit in a child's development of listening and spoken language.

As an added bonus, we have included the proceedings of the 2010 AG Bell Research Symposium, "Re-Modeling the Deafened Cochlea for Auditory Sensation: Advances and Obstacles." The presenters offer an overview on the advancements and limitations of stem cell and cochlear sensory cell regeneration, providing insights into the future of this line of research.

As we wrap up 2010, exploring the language development of individuals who are deaf and hard of hearing has never been more important. And while a large number of studies exist that specifically focused on auditory-verbal practice, the results do not provide conclusive evidence of success on a broader level. We encourage you and your colleagues to consider contributing to this body of research with a large-scale case study that explores the positive, and negative, results of auditory-verbal practice to provide rigorous, scientific evidence of the strength of this communication approach.

We hope you enjoy this issue, and please don't hesitate to contribute to *The Volta Review*.

Sincerely,

A handwritten signature in black ink that reads "Joseph Smaldino" with a stylized flourish at the end.

Joseph Smaldino, Ph.D.
Professor and Chair, Department of Communication Sciences
Illinois State University
jsmaldi@ilstu.edu

A handwritten signature in black ink that reads "Kathryn L. Schmitz" with a stylized flourish at the end.

Kathryn L. Schmitz, Ph.D.
Assistant Professor and Interim Chair, Liberal Studies Department
National Technical Institute for the Deaf/Rochester Institute of Technology
kls4344@rit.edu

Is Auditory-Verbal Therapy Effective for Children with Hearing Loss?

Dimity Dornan, Ba.Sp.Th., F.S.P.A.A., LSLS Cert. AVT;
Louise Hickson, B.Sp.Thy. (Hons), M.Aud., Ph.D.;
Bruce Murdoch, B.Sc. (Hons), Ph.D., D.Sc.;
Todd Houston, Ph.D., CCC-SLP, LSLS Cert. AVT; and
Gabriella Constantinescu, B.S.Path (Hons), Ph.D.

A longitudinal study reported positive speech and language outcomes for 29 children with hearing loss in an auditory-verbal therapy program (AVT group) (aged 2 to 6 years at start; mean PTA 79.39 dB HL) compared with a matched control group with typical hearing (TH group) at 9, 21, and 38 months after the start of the study. The current study investigates outcomes over 50 months for 19 of the original pairs of children matched for language age, receptive vocabulary, gender, and socioeconomic status. An assessment battery was used to measure speech and language over 50 months, and reading, mathematics, and self-esteem over the final 12 months of the study. Results showed no significant differences between the groups for speech, language, and self-esteem ($p > 0.05$). Reading and mathematics scores were comparable between the groups, although too few for statistical analysis. Auditory-verbal therapy has proved to be effective for this population of children with hearing loss.

Dimity Dornan, Ba.Sp.Th., F.S.P.A.A., LSLS Cert. AVT, is a postgraduate student in the School of Health and Rehabilitation Sciences, University of Queensland in Australia, and the Managing Director and Founder of the Hear and Say Centre in Brisbane, Australia. Louise Hickson, B.Sp.Thy. (Hons), M.Aud., Ph.D., is a Professor of Audiology in the School of Health and Rehabilitation Sciences, University of Queensland in Australia. Bruce Murdoch, B.Sc. (Hons), Ph.D., D.Sc., is a Professor of Speech Pathology and Director of the Motor Speech Research Centre in the School of Health and Rehabilitation Sciences, University of Queensland in Australia. Todd Houston, Ph.D., CCC-SLP, LSLS Cert. AVT, is the Director of Graduate Studies Program in Auditory Learning & Spoken Language in the Department of Communicative Disorders and Deaf Education, Utah State University in the United States. Gabriella Constantinescu, B.S.Path. (Hons), Ph.D., is the lead researcher at the Hear and Say Centre in Brisbane, Australia. Correspondence concerning this article should be directed to Ms. Dornan at dimity@hearandsaycentre.com.au.

Introduction

This longitudinal study was designed to investigate the effectiveness of auditory-verbal therapy (AVT) for a group of children with hearing loss (AVT group). Since the introduction of universal newborn hearing screening, digital hearing aids, and cochlear implants, there has been increased debate about educational options for children with hearing loss. Appropriate and timely information is needed in order to guide parent and professional decision-making. However, rigorous evidence for the outcomes of any of the educational approaches in use today, including AVT, is minimal (Gravel & O'Gara, 2003; Sussman, et al., 2004). Existing research studies on AVT outcomes have been criticized as being few in number and lacking in rigor (Eriks-Brophy, 2004; Rhoades, 2006). These studies also had limited generalizability because of their retrospective nature, inconsistency in the use of standardized assessments, the possibility of self-selected populations, and lack of control groups.

A review of research findings on outcomes of AVT was conducted by Dornan, Hickson, Murdoch, and Houston (2008). In this review, several studies were found to demonstrate a typical rate of progress for the language development of children in AVT programs (Hogan, Stokes, White, Tyszkiewicz, & Woolgar, 2008; Rhoades, 2001; Rhoades & Chisolm, 2000). However, these findings needed substantiation with a controlled study. Such comparisons were undertaken in the earlier stages of this research where outcomes of the AVT group were compared with those for a matched group of children with typical hearing (TH group) (Dornan, Hickson, Murdoch, & Houston, 2007; 2009). At baseline, there were 29 children in the AVT group, between 2 and 6 years old (mean pure tone average of 76.17dB HL). The children were matched with the TH group for initial language age, receptive vocabulary, gender, and socioeconomic status (as measured by head of the household's education level). Both groups were assessed over time using a battery of assessments. Speech and language outcomes for the AVT group were compared with those for the TH group from a baseline measure (referred to as the pretest) to 9, 21, 38, and 50 months after the baseline tests (referred to as the posttests). Results of these earlier studies have been positive, with the AVT group showing significant progress for speech and language at the same rate as the TH group (Dornan, et al., 2007; 2009). An exception was receptive vocabulary, for which the AVT group achieved the same progress as the TH group at the 9 months posttest ($p > 0.05$) (Dornan, et al., 2007), but the TH group scored significantly higher than the children in the AVT group at the 21 and 38 months posttests ($p \leq 0.05$) (Dornan, et al., 2009). Despite this difference, the AVT children's mean age equivalent was within the typical range for their chronological age.

Further study on the outcomes of the AVT group is important because few controlled longitudinal studies of speech and language outcomes are available for children with hearing loss. In addition, an extension of the study time

allowed us to include measures of academic outcomes for the children. It is widely acknowledged that significant hearing loss in children also impacts academic achievement, which usually lags behind the norm for children with typical hearing (Powers, 2003). Academic success for a child with hearing loss in the mainstream has been linked with a number of factors, including education using listening and spoken language, a shorter period of hearing loss prior to amplification or cochlear implantation, and level of intelligence (Damen, van den Oever-Goltstein, Langereis, Chute, & Mylanus, 2006; Geers, et al., 2002). The fundamental academic skill of reading is often severely affected by significant hearing loss, with many children never achieving functional literacy skills (Moeller, Tomblin, Yoshinaga-Itano, Connor, & Jerger, 2007; Vermeulen, van Bon, Schreuder, Knoors, & Snik, 2007). Traxler (2000) found that children with severe-to-profound hearing loss typically completed 12th grade with language levels of 9- to 10-year-old children who had typical hearing, and 50% of those students read at a 4th-grade reading level or less. Reports such as these are in contrast to two studies on the reading abilities of children in an AVT program (Robertson & Flexer, 1993; Wray, Flexer, & Vaccaro, 1997), which found that children were able to read at or above age-appropriate levels. However, as the previous research did not include control groups and used different assessments, interpretation of findings and close comparison of results are difficult.

Poor consistency has also been reported among studies on the reading development of children with hearing loss using a variety of education approaches (Marschark, Rhoten, & Fabich, 2007). Spencer and Oleson (2008) studied the reading abilities of 72 children with hearing loss who had used unspecified education approaches (after 48 months of cochlear implant use) and concluded that early access to sound helped build better phonological processing skills, one of the likely contributors to reading success (National Institute of Child Health & Human Development, 2000). Spencer and Oleson (2008) also found that 59% of variance in reading skills for these children could be explained by early speech perception and speech production performance. However, other researchers found that although early cochlear implantation had a long-term positive impact on listening and spoken language development, it did not result in age-appropriate reading levels in high school for the majority of students (Geers, Tobey, Moog, & Brenner, 2008).

Another important academic skill is mathematics and, as with reading, research on the mathematical achievements of children with hearing loss is generally inadequate because studies are rare and seldom use the same measures. Furthermore, no data is available on mathematics outcomes for children in AVT programs. However, mathematics skill levels below that of their peers with typical hearing are consistently reported for children with hearing loss (Kritzer, 2009). Mathematics ability for children with hearing loss has shown to be related to a child's skills in reading, language, and morphological knowledge regarding word segmentation and meaning (Kelly & Gaustad, 2007). Nunes and Moreno (2002) reported two aspects of the functioning ability of

children with hearing loss that place them at risk for underachievement in mathematics, over and above reduced access to hearing: (1) reduced opportunities for incidental learning and (2) difficulty in making inferences involving time sequences. Traxler (2000) found that the mathematics performance of school-aged students with hearing loss indicated only partial mastery of mathematical knowledge and skills. High school graduates were found to have computational skills comparable to 6th grade students with typical hearing, and mathematics problem solving skills comparable to 5th grade students with typical hearing (Traxler, 2000). Low academic attainments in mathematics, as well as reading, may have significant economic impact on the child's future because of the relationship that exists between education level and income (Nunes & Moreno, 2002).

In addition to reading, mathematics, and overall academic achievement, the way children with hearing loss perceive themselves and their abilities is an important outcome. No research on the self-esteem of children educated in AVT programs is available. Researchers have found that for children with significant hearing loss who do not develop language skills commensurate with their peers, self-esteem and emotional development are often severely affected (Bat-Chava, Martin, & Kosciw, 2005; Hintermair, 2006; Nicholas & Geers, 2003). Self-esteem measures usually take the form of either a child or a parent-reported questionnaire or survey, either oral or written (e.g., Percy-Smith, et al., 2006; Schorr, Roth, & Fox, 2009). During the 38 months posttest we added a self-esteem questionnaire in which parents responded to questions regarding their child's sense of self, sense of belonging, sense of personal power, and overall self-esteem. Results showed that self-esteem levels were not significantly different between groups. It was important to investigate whether these positive self-esteem results would continue as the group advanced through school. Hence, the self-esteem questionnaire was repeated at the 50 months posttest.

This entire study used a battery of assessments to investigate the effectiveness of AVT over a 50-month time period for a group of children with hearing loss. We studied whether the promising outcomes for listening and spoken language for the AVT group shown in earlier stages of this longitudinal study (Dornan, et al., 2007; 2009) were maintained over 50 months by 19 of the same children who remained in the study for the full 50 months. Reading, mathematics, and self-esteem were also investigated over the last 12 months of the study, by which time most of the 2 groups had reached school age. Outcomes for the AVT group were compared with those for the 19 matched children in the TH group over 50 months.

Method

This study employed a matched group, repeated measures design. At the start of the study, the TH group was individually matched to the AVT group

for total language, receptive vocabulary, gender, and socioeconomic level (as measured by the education level of the head of the household).

Participants

Auditory-Verbal Therapy Group (AVT Group)

Selection criteria for the participants were: Pure-Tone Average (PTA) at 500 Hz, 1000 Hz, 2000 Hz, and 4000 Hz of ≥ 40 dB hearing threshold levels in the better ear; prelingually deafened (at ≤ 18 months old); attended the educational program weekly for intensive one-on-one, parent-based AVT for a minimum of 6 months; wore hearing devices consistently (hearing aids and/or cochlear implants) and aided hearing was within the speech range or had received a cochlear implant; no other significant cognitive or physical disabilities reported by parents or educators; 2 to 6 years of age at the first test session; and both parents spoke only English to the child.

The children attended one of the five regional centers of an AVT program in Queensland, Australia, which offers a range of services including audiology, early intervention, and a cochlear implant program. This program adheres to the Principles of Auditory-Verbal Therapy (adapted from Pollack, 1970; endorsed by the AG Bell Academy for Listening and Spoken Language, 2007). Even though a particular AVT program may adhere to all of these principals, programs may vary in the operational details. A description of the AVT program in this study can be found at <http://www.hearandsaycenter.com.au/mission-delivery.html>.

Of the 10 children who left the study between the 38-month and 50-month posttests, 2 had left the program because of diagnosis of additional disabilities, 6 had moved away or were unavailable for testing, and the departure of 2 TH group children from the study necessitated omitting their matched AVT group pair. The remaining AVT group participants had bilateral sensorineural hearing loss ranging from moderate to profound (mean PTA 79.39 dB HL; range = 45 dB to >110 dB). All children were fitted with hearing aids, commencing intervention within 3 months of diagnosis. Of these, 13 children received unilateral Cochlear Nucleus CI 24 implants and used an Advanced Combined Encoder (ACE) processing strategy. The median age for receiving a cochlear implant was 23.04 months (mean = 27.54 months, SD = 15.24). During the study, 6 of these children received a bilateral cochlear implant. All but 1 of the unilateral cochlear implant users in the study also wore a hearing aid in the contra-lateral ear. Both hearing devices were balanced by their audiologist according to the recommendation of Ching, Psarros, and Incerti (2003). All children wore their hearing devices consistently throughout the study. A battery of speech perception tests was administered by an audiologist to ensure that the children's listening skills were developing optimally. The mean age of the AVT group at the start of the study was 3.80 years

(SD = 1.15) and the mean age at the 50 month follow-up was 8.02 years (SD = 1.28) (Table 1).

Typical Hearing Group (TH group)

The TH group was recruited by families and staff of the AVT program, and their characteristics are also available in Table 1. Selection criteria for the participants were: hearing threshold levels within the range of 0 to 20 dB at 500 Hz, 1000 Hz, 2000 Hz, and 4000 Hz for both ears; typical articulation as measured by the Goldman-Fristoe Test of Articulation (Goldman & Fristoe, 2001) and using Australian norms (Kilminster & Laird, 1978); no significant cognitive or physical disabilities (as evidenced by case history or parent report); and both parents spoke only English to the child.

Sixty four children were initially tested to ensure controlled matching with the AVT group. The TH group children who remained in the study were matched at the initial assessment with the AVT group for total language age (± 3 months) on the Preschool Language Scale (PLS-4) (Zimmerman, Steiner, & Pond, 2002) or the Clinical Evaluation of Language Fundamentals

Table 1. Characteristics of the AVT group and the TH group at 50 months posttest

	<i>AVT Group</i>	<i>TH Group</i>
N	19	19
Mean Age in months (SD)	96.26 (15.32)	87.84 (16.68)
Gender		
Male	14	14
Female	5	5
Age at identification in months	22.29 (11.82)	n/a
Mean PTA hearing loss in better ear in dB (SD)	79.39 (23.79)	n/a
Onset of Loss		
Congenital	17	n/a
Prelingual	2	n/a
Age at CI , if applicable, in months (SD)	27 (5.8)	n/a
Time spent in AVT Program in months (SD)	70 (16.34)	n/a
Hearing Device:		
Bilateral HA's	5	n/a
Unilateral hearing aid	1	n/a
HA and CI in contra-lateral ears	6	n/a
Unilateral CI only	1	n/a
Bilateral CI's	6	n/a
Parents educated beyond high school	18	18
Occupation category of head of household		
Professional	14%	65%
Manager	43%	15%
Trade/technical	29%	5%

CI = Cochlear Implant; HA = Hearing Aid

(CELF-3) (Semel, Wiig, & Secord, 1995), as well as for receptive vocabulary on the Peabody Picture Vocabulary Test (PPVT-3) (Dunn & Dunn, 1997). Matching criteria also included gender and socioeconomic level, as assessed by highest education level of the head of the household. The rationale for matching for language age rather than chronological age has been discussed in an earlier paper (Dornan, et al., 2009). Had chronological age been used for matching (instead of language age), the children with typical hearing generally would have had a higher language level than the children with hearing loss, introducing the possibility that the children in the TH group might progress faster. Deciding how to define socioeconomic level for matching purposes was difficult because there are many different perspectives and a number of different possible measures (Kumar, et al., 2008). Some factors that might have been measured include family income, education level of the parents, and parental occupation (Marschark & Spencer, 2003). It was thought that questions about family income may deter parents from long-term commitment to the longitudinal study before it had commenced. Consequently, the highest level of education of the head of the household was used. As an added check, the occupations of both groups were placed in categories according to those developed by Jones (2003), as occupation category has been found to impact the vocabulary learning of a child with hearing loss (Hart & Risley, 1995) (see Table 1). It was concluded that both AVT group and TH group parents had a moderate to high socioeconomic status. The mean age of the TH group at the start of the study was 3.11 years ($SD = 1.22$) and the mean age at the 50 month follow-up was 7.32 years ($SD = 1.39$) (Table 1).

Materials

To assess total language, receptive vocabulary, speech, reading, mathematics, and self-esteem at pre- and posttest for participants in both the AVT and TH groups, a battery of assessments was used (Table 2). As some assessments are recorded differently, (i.e. standard scores, percentile ranks, and raw scores), this is also represented in Table 2. Additional information on the assessments for reading, mathematics, and self esteem is included in the Appendix.

Procedure

Appropriate ethical clearance and parent consent was gained for this study (Dornan, et al., 2007, 2009). Assessments of children in the AVT group took place at the child's program center. For the TH group, testing was performed either at the center, at the child's education setting in a quiet room, or at the child's home. Speech, language, reading, and mathematics testing was performed by experienced, qualified speech-language pathologists. Because of geographic constraints and for convenience, available qualified staff performed the testing and, frequently, different speech-language pathologists

Table 2. Battery of assessments

<i>Test</i>	<i>Description of Test</i>	<i>Scoring</i>
Language		
Preschool Language Scale-Fourth Edition (PLS-4) (Zimmerman, et al., 2002).	Measures young child's receptive and expressive language from birth to 6 years, 11 months. Australian norms were not available. Used at pretest for all children but one pair. CELF-3 was used for this pair. Not used at 50 months posttest.	The scoring ceiling used was five consecutive items incorrect. Receptive language and oral expression were expressed as standard scores because the CELF-3 does not have age equivalents for comparison. Total language score is expressed as an age equivalent.
Clinical Evaluation of Language Fundamentals (CELF-3) (Semel, et al., 1995).	Measures child's receptive and expressive language from 6 years to 21 months. CELF-3 used for all children at posttest. Six subtests were administered only to children who achieved higher than the top score for the PLS-4. Subtests were Sentence Structure, Word Structure, Concepts and Directions, Formulated Sentences, Word Classes, and Sentence Recalling. Australian norms were not available.	If a child scored the highest possible score on the PLS-4, the CELF-3 was administered. Receptive language and oral expression are expressed as standard scores (age equivalents are not available) and total language is expressed as an age equivalent.
Receptive Vocabulary		
Peabody Picture Vocabulary Test (PPVT-3) (Dunn & Dunn, 1997).	Measures child's receptive vocabulary. Because this test was developed in the United States, Australian alternatives for some items were used by the testers: (a) cupboard for closet, (b) rubbish for garbage, (c) biscuit for cookie, and (d) jug for pitcher. Australian norms were not available.	Child's score is expressed as an age equivalent.
Speech		
Goldman-Fristoe Test of Articulation-2 (GFTA-2) (Goldman & Fristoe, 2001).	Assesses articulation of consonants and was administered to participants in both AVT and TH groups. Australian norms were not available.	Child's score is expressed as an age equivalent.

<p>Reading Reading Progress Tests (RPT) (Vincent, Crumpler, & de la Mare, 1997).</p>	<p>Stage I is used in the first 3 years of school and assesses pre-reading and early reading skills in first year of school and reading comprehension in the second and third years of school. Stage 2 is used for school years 3–6 and assesses outcomes for reading by assessing a range of literal and inferential skills and reading vocabulary. Australian norms were available.</p>	<p>One mark is awarded for each correct answer. No marks are awarded for multiple choice questions where more than one choice has been selected. Score is expressed as a percentile rank.</p>
<p>Mathematics I Can Do Maths (Doig & de Lemnos, 2000). Progressive Achievement Tests in Mathematics (PATMaths) (Australian Council of Educational Research, 2005).</p>	<p>This test assesses numeracy development in first 3 years of school. Australian norms are available. This test assesses mathematic achievement levels in school years 3 to 11. Australian norms are available.</p>	<p>One mark is awarded for each correct answer. Score is expressed as a percentile rank. One mark is awarded for each correct answer. Score is expressed as a percentile rank.</p>
<p>Self-esteem Insight (Morris, 2003).</p>	<p>This questionnaire assesses development of self-esteem from 3–19 years of age (preschool and primary). Parents were asked to complete this questionnaire, and the 36 questions were divided into 3 different areas, which included their child's sense of self, sense of belonging, and sense of personal power. Parents were asked to report whether the skill was evident "Most of the Time" (3 points), "Quite Often" (2 points), "Occasionally" (1 point), or "Almost Never" (0 points).</p>	<p>The sum of the scores for the 3 areas studied (sense of self, sense of belonging, and sense of personal power) were totalled (maximum possible score = 108) and then rated as "High," "Good," "Vulnerable," or "Very Low" according to score-based criteria: "High" = 87–108; "Confident and at ease with self, other people, and the world most of the time." "Good" = 64–86; "Feels good about self, but takes knocks now and again." "Vulnerable" = 40–63; "Tends not to feel very confident." "Very Low" = 0–39; "Depressed or very challenging behaviour to cover this up."</p>

assessed the children at pre- and posttest. Tester reliability was not examined in the study; however, all tests were administered according to the standardized instructions in the test manuals. Language and speech tests were administered over two or three sessions according to the needs of each child with a rest break between assessments, and were discontinued if a child showed fatigue or distress. The children's responses to the GFTA-2 were judged to be correct or incorrect at the time of testing. The order of presentation of the standardized tests to the TH group was different than the AVT group to account first for screening, and then to establish a match with a child in the AVT group before the child was unnecessarily tested.

For the AVT group, assessments were performed in the best aided condition. For all children with cochlear implants, the optimally functioning MAP (assessed by the child's audiologist and auditory-verbal therapist) was used for assessments. Both "T" levels (threshold, or minimum amount of current causing sound to be detected) and "C" levels (maximum amount of current causing discomfort) for the child's MAP were measured behaviorally and confirmed objectively. Optimal implant performance was verified by stability of the MAP, consistent identification by the child of the seven sound test (i.e. the Australian adaptation of Ling's Six Sound Test; Romanik, 1990), other speech perception tests, and the cochlear implant-assisted audiogram (a record of the child's cochlear implant aided thresholds for responses at 250 Hz, 500 Hz, 1000 Hz, 2000 Hz, and 4000 Hz). The Ling sounds are a range of speech sounds encompassing frequencies widely used clinically to verify effectiveness of hearing aid fitting in children (Agung, Purdy, & Kitamura, 2005). The Ling Six Sound Test was developed for the North American population (Ling, 2002), and /o/ was added (seven sound test) to account for differences in production and spectral content of Australian vowels (Agung, et al., 2005). For the children with hearing aids, best aided condition was determined by the audiologist and auditory-verbal therapist, performance of the seven sound test, speech perception tests, and the child's aided audiogram.

For speech and language assessments, the mean time between the pretest and 50 months posttest was 51.16 months for the AVT group ($SD = 1.12$) and 51.37 months for the TH group ($SD = 0.94$), which was not significantly different ($t = -0.335$, $p = 0.742$). Similarly, mean times between pretest (38 months) and posttest (50 months) for reading, mathematics, and self-esteem assessments ($M = 12.73$ months, $SD = 2.03$ for the AVT group; $M = 13.26$ months, $SD = 1.91$ for the TH group) were also not significantly different for the two groups ($t = -1.398$, $p = 0.171$).

Results

Preliminary analysis was carried out to ensure the validity of matching participant groups at the pretest, that is, the matching of total language on the PLS-4 or CELF-3 and receptive vocabulary on the PPVT-3. A Mann-Whitney

test showed that the total mean language ages of the AVT ($M = 3.58$ years; $SD = 1.46$) and TH groups ($M = 3.5$ years; $SD = 1.52$) were comparable ($z = -0.307$; $p = 0.759$). Similarly, there were no significant differences ($z = -0.197$; $p = 0.844$) for mean receptive vocabulary ages at pretest on the PPVT-3 between the AVT ($M = 3.06$ years; $SD = 1.56$) and TH groups ($M = 2.97$ years; $SD = 1.46$). Overall, both groups were found to be matched for total language age and receptive vocabulary at the pretest.

Speech and Language

Table 3 displays the pre- and posttest mean age equivalents, standard deviations, z and p values for total language, receptive vocabulary, and speech for the 19 children in the AVT and TH groups at pretest and at the 50 months posttest.

For total language age assessed using the PLS-4 or CELF-3, both groups made significant progress over 50 months and the change in scores over this period of time was not significantly different between the groups. Further comparisons of receptive and expressive language results were made using standard scores on the CELF-3 as age equivalence is not calculated on this assessment. For receptive language, no significant changes in standard scores were found from pretest to posttest for each group because standard scores

Table 3. Pre- and posttest mean age equivalents, standard deviations (in parentheses), and z and p values for total language, receptive vocabulary, and speech for 19 children in the AVT and TH groups at pretest and at 50 months posttest

<i>Test</i>	<i>Group</i>	<i>Pretest Mean Age Equivalent (months) (SD)</i>	<i>Posttest Mean Age Equivalent (months) (SD)</i>	<i>z</i>	<i>p</i>
PLS-4/CELF-3	AVT	42.95 (17.59)	94.26 (34.60)	-3.824	<0.001*
	TH	42.00 (18.27)	98.05 (12.37)	-3.825	<0.001*
<i>Group Comparison</i>				1.550	0.121
PPVT-3	AVT	36.74 (18.56)	93.95 (43.88)	-3.824	<0.001*
	TH	35.57 (17.55)	100 (19.01)	-3.825	<0.001*
<i>Group Comparison</i>				-2.921	0.235
GFTA-2	AVT	36.74 (17.38)	80.42 (15.76)	-3.824	<0.001*
	TH	42.11 (17.04)	86.00 (11.76)	-3.765	<0.001*
<i>Group Comparison</i>				-0.336	0.737

Note 1: * = significant difference (acceptable level of significance is ≤ 0.05); progress with time for each group analysed using the Wilcoxon Signed Rank Test; between group comparisons of progress analysed using the Mann-Whitney Test.

Note 2: AVT group chronological age at pretest was 45.6 months ($SD = 13.8$) and at posttest was 96.26 months ($SD = 15.32$).

TH group chronological age at pretest was 37.32 months ($SD = 14.64$) and at posttest was 87.84 months ($SD = 16.68$).

are age corrected ($z = -1.808, p = 0.071$ for the AVT group; $z = -1.7, p = 0.089$ for the TH group). Also, the amount of change displayed by both groups was not significantly different ($z = 0.599, p = 0.549$). Similarly, for expressive language, there was no significant change in standard scores between pre- and posttest for each group ($z = -1.002, p = 0.316$ for the AVT group; $z = -1.373, p = 0.170$ for the TH group) and no significant difference for amount of progress between the two groups ($z = -1.131, p = 0.895$).

Receptive Vocabulary

For receptive vocabulary, as assessed using the PPVT-3, both groups showed significant changes in age equivalents over the 50 months with no significant difference in the amount of change between the groups (Table 3). The mean age equivalent for the AVT group was within the typical range for their chronological age.

Speech

On the GFTA-2, significant increases in age equivalents were evident for both groups over 50 months with no significant differences in the changes between the two groups (Table 3). At the 50 months posttest, 42% of the AVT group and 63% of the TH group scored at the ceiling of 7 years, 8 months on this test.

Reading and Mathematics

For these assessments, a smaller sample of 7 pairs of children in each group was available for comparison. This is because a number of children in both groups had not yet entered school or begun formal reading and mathematics by the 50 months posttest (1 child in the AVT group; 3 children in the TH group) or had not reached this stage by the 38 months posttest (4 children in the AVT group; 13 children in the TH group children). Table 4 shows the pre- and posttest percentile ranks for reading and mathematics assessments for the 14 eligible children in the AVT group and TH group at 38 and 50 months posttest.

Table 4. Pre- and posttest percentile ranks for reading and mathematics for the 14 children in the AVT group and TH group at 38 and 50 months posttest

<i>Name of Test</i>	<i>Group</i>	<i>N</i>	<i>Pretest Percentile</i>		<i>N</i>	<i>Posttest Percentile</i>	
			<i>Rank (SD)</i>	<i>Range</i>		<i>Rank (SD)</i>	<i>Range</i>
Reading	AVT	7	83.57 (17.74)	51–98	7	88.14 (10.90)	46–99
	TH	7	88.14 (7.90)	75–98	7	90.14 (9.81)	79–99
Mathematics	AVT	7	60.43 (35.02)	23–98	7	77.57 (28.54)	32–99
	TH	7	81.28 (24.88)	67–96	7	80.86 (19.35)	77–92

The numbers of children in each group were considered too small for statistical comparison. For reading, the AVT group scores were in the 83rd percentile at the 38 months posttest and in the 88th percentile at the 50 months posttest. Similarly, for mathematics, the AVT group scores were in the 60th percentile at the 38 months posttest and in the 77th percentile at the 50 months posttest. The percentile ranks at the 50 months posttest for both groups were comparable (see Table 4).

Self-Esteem

Eighteen parents of the AVT group and 16 parents of the TH group responded to the self-esteem questionnaire at the 50 months posttest, and 10 matched pairs were identified with scores both at the 38 months posttest and the 50 months posttest. Table 5 shows the results for sense of self, sense of belonging, and sense of personal power subscales for both groups at 50 months, the highest possible score being 36 in each category.

Mann-Whitney tests showed no significant differences between the groups for sense of self, sense of belonging, and sense of personal power components of the questionnaire. Furthermore, at the 50 months posttest, the total self-esteem scores between the two groups were not significantly different. The majority of children in both participant groups (80% in the AVT group and 70% in the TH group) were rated as having “high” self-esteem while the remainder had “good” self-esteem. No children from either group were rated in the “vulnerable” or “very low” categories.

Discussion

This study reported speech perception outcomes for the AVT group from the 38 months to the 50 months posttests and also compared outcomes for the AVT group for receptive, expressive, and total language, receptive vocabulary, and speech over 50 months with outcomes of the matched TH group. In addition, the study also compared reading, mathematics, and self-esteem outcomes between the AVT and TH groups over the last 12 months of the study. The AVT group’s promising earlier outcomes of typical rate of progress for total language and speech skills to those of hearing controls (Dornan, et al., 2007, 2009) has been maintained over the 50 months. Furthermore, receptive vocabulary progress, reported to be slower for the AVT group than for the TH group at earlier posttests (Dornan, et al., 2009), was found to have accelerated significantly at the 50 months posttest to develop at the same rate as the TH group. The AVT group maintained standard score levels for receptive and expressive language, which were similar to results for the TH group. Self-esteem levels were not significantly different between the groups, with predominantly high self-esteem reported for both groups. We will further discuss and compare these results with our previous research.

Table 5. Pre- and posttest raw scores for self-esteem for the AVT group and the TH group for primary Insight at 38 and 50 months posttests

		Sense of Self (SD)		Sense of Belonging (SD)		Sense of Personal Power (SD)		TOTAL Self-esteem (SD)	
		Mean 38 months (SD)	Mean 50 months (SD)	Mean 38 months (SD)	Mean 50 months (SD)	Mean 38 months (SD)	Mean 50 months (SD)	Mean 38 months (SD)	Mean 50 months (SD)
AVT Group		31.07 (3.93)	32.1 (3.11)	31.64 (2.98)	32.4 (3.09)	30.29 (3.79)	31.60 (3.72)	92.86 (9.46)	96.1 (9.22)
TH Group		31.93 (4.8)	32.94 (2.81)	29.92 (5.27)	33.8 (2.53)	27.36 (7.17)	31.5 (2.88)	89.93 (12.82)	98.4 (7.44)
Group	<i>z</i>	-0.324	-0.609	-0.949	-0.996	-0.946	-0.229	-0.621	-0.417
Comparison	<i>p</i>	0.746	0.542	0.343	0.319	0.344	0.819	0.534	0.677

Acceptable level of significance = ≤ 0.05 . Progress with time for each group analysed using the Wilcoxon Signed Rank Test. Between group comparisons of progress analysed using the Mann-Whitney Test.

Total language growth for the AVT group was at a rate of 12.31 months per year, comparing favorably to a rate of 13.45 months for the TH group. The majority of the AVT group (79%) and the entire TH group scored within the typical range or above for language at the 50 months posttest. The AVT group achieved mean total language scores, which were 2.1 months less than their mean chronological age or within one standard deviation of the mean for their age. The only other studies that have indicated such positive language growth results have included children fitted with hearing aids at less than 6 months of age (Yoshinaga-Itano, Sedey, Coulter, & Mehl, 1998) or children receiving cochlear implants before 18 months of age (Ching, et al., 2009; Dettman, Pinder, Briggs, Dowell, & Leigh, 2007; Svirsky, Teoh, & Neuburger, 2004). In the present study, 1 child had been fitted with hearing aids before 6 months and 2 children had been fitted with cochlear implants at less than 18 months. Nevertheless, the group as a whole achieved age appropriate language.

These positive results for language of the AVT group are similar to those obtained previously for children in AVT programs (e.g. Rhoades, 2001; Rhoades & Chisolm, 2000) in which the majority of children were reported to show no significant chronological age and language age gaps when entering mainstream school. Results obtained here are superior to a number of other studies of children with hearing loss educated using a range of different interventions (e.g. Blamey, Barry, et al., 2001; Geers, Nicholas, & Sedey, 2003; Sarant, Holt, Dowell, Rickards, & Blamey, 2008).

The AVT group progressed in receptive vocabulary development at a rate of 13.73 months per year over the 50 months of the study, compared to the TH group at 15.46 months, with no significant difference in progress between the two groups. For the AVT group, 68% had scores within the typical range or above for receptive vocabulary, compared to 100% of the TH group. At the 50 months posttest, the gap between chronological age and age equivalence for the AVT group for receptive vocabulary was 2.4 months. This suggests that the AVT group were functioning as expected for their age for receptive vocabulary. The receptive vocabulary results for the AVT group are superior to those found in the literature, which have reported levels of receptive vocabulary for children with hearing loss lower than children with typical hearing (e.g. Blamey, Sarant, et al., 2001; Eisenberg, Kirk, Martinez, Ying, & Miyamoto, 2004; Fagan & Pisoni, 2010; Hayes, Geers, Treiman, & Moog, 2009; Schorr, Roth, & Fox, 2008; Uziel, et al., 2007).

Similar to earlier stages of the study, the AVT group achieved intelligible speech with the same scores as the TH group (Dornan, et al., 2007, 2009). The rate of change in scores per year for correct articulation of consonants in words was 10.48 months for the AVT group and 10.53 months for the TH group. The lack of a significant difference between the changes in speech scores for the AVT and TH groups is surprising because children with hearing loss typically have difficulty with articulation of speech sounds (Marschark, Lang, & Albertini, 2002; Schorr, et al., 2008; Uziel, et al., 2007). An increase in accuracy of

consonant production for children with implants (like most of the AVT group in this study) has been reported, as well as an increasing ability with longer implant experience and use of oral communication (Tobey, Geers, Brenner, Altuna, & Gabbert, 2003). A high correlation between speech perception and speech production has also been reported for children with cochlear implants (Phillips, et al., 2009). It is likely that the combination of cochlear implant use and AVT may have positively influenced the level of speech skills achieved by the AVT group in this study.

In this paper we report preliminary results for reading and mathematics over a 12-month period for a small sample of children ($n = 7$). Over the last 12 months of this study, the AVT group results for reading improved from the 84th percentile to the 88th percentile (as compared to from the 88th percentile to the 90th percentile for the TH group). The AVT group results for mathematics improved from the 60th percentile to the 77th percentile (as compared with remaining around the 81st percentile for the TH group). Since percentile ranks are already normalized scores, the improvement in percentile ranks for the AVT group indicates that the children were progressing at a faster rate than is typical. The positive results for reading and mathematics, although for a very small group, show the potential for this group of children to be successful in the mainstream. As the AVT group was relatively young (8.02 years) at the 50 months posttest, it will be important to follow up this study, particularly for reading, over a longer term as it has been found that reading scores for a group of 85 adolescents with cochlear implants studied from ages 8–9 years did not keep pace with their language development at ages 15–18 years (Geers, et al., 2008). In addition, further large-scale studies are needed to investigate reading progress for children in AVT programs. Positive reading achievement for children with hearing loss educated using AVT has been reported in a number of studies (Durieux-Smith, et al., 1998; Goldberg & Flexer, 1993, 2001; Robertson & Flexer, 1993; Wray, et al., 1997), and has been related to speech perception and speech production performance (Spencer & Oleson, 2008). Together, these findings are in contrast to unfavorable reports on reading ability for children with hearing loss in some studies (e.g. Boothroyd & Boothroyd-Turner, 2002; Moeller, et al., 2007; Traxler, 2000; Vermeulen, et al., 2007). The good levels of speech perception and speech production achieved by the AVT group in this research (Dornan, et al., 2009) may have had an influence on their reading achievement. The addition of an assessment of phonological processing in future research may add to the information on reading skills for children in AVT programs.

In relation to mathematics, the same inherent problems of sample size made interpretation of the results difficult. At the 50 months posttest, however, the mean percentile rank for the AVT group for mathematics was high (78th percentile), as was the percentile rank for the TH group (81st percentile), which suggests that the results are relatively comparable for both groups. Since the mathematics assessment was both read by the AVT group and presented to them verbally, these outcomes represent positive skill levels for listening and

reading as well as mathematics for this group. The current AVT group performed better than the group studied by Traxler (2000), who found that the mathematics ability of high school students with hearing loss was at a “basic level” or below. The findings for the AVT group may be explained by their good reading and language skills, as Kelly and Gaustad (2007) found that levels of reading and language skills influenced the ability of a child with hearing loss to achieve in mathematics. In addition, the good listening ability of the AVT group may well have influenced their mathematics ability, as their listening ability allowed them opportunities for incidental learning of early mathematics concepts, unlike the study reported by Nunes and Moreno (2002). More studies on mathematics outcomes for children with hearing loss are needed to add to the body of knowledge in this area.

The self-esteem results for the AVT group are better than those obtained by a number of other researchers who reported adversely affected self-esteem (Nicholas & Geers, 2003), mental health (Laurenzi & Monteiro, 1997), and socio-emotional development (Prizant & Meyer, 1993) for children with significant hearing loss. In the present study, there was no significant difference between the AVT group and the TH group for self-esteem. These results are in agreement with those in a Danish parent survey of children with hearing loss (Percy-Smith, et al., 2006), which reported a satisfactory or very satisfactory level of well-being for children with cochlear implants. The AVT group results are also in agreement with those of Schorr et al. (2009) who found that 37 children (ages 5–14 years) who received a cochlear implant and used listening and spoken language reported improved quality of life; positive self-esteem was also related to receiving a cochlear implant at a younger age. The high results for self-esteem for the AVT group could be a factor of their good use of their hearing device(s), their good listening skills, and speech and language development, but their mean age of cochlear implantation was not particularly early (27 months). It is significant that these results for self-esteem were based on a parent rating, showing that at the 50 months posttest, the parents perceived that there was little detrimental impact on the child’s self-esteem as a result of the hearing loss.

Although this study’s findings are promising, the outcomes cannot be generalized for a number of reasons. First, both the AVT and TH groups were mainly from a moderate to high socioeconomic level. This may have caused a self selection of both groups of children, making some interpretation difficult. Similarly, a number of studies on outcomes of AVT have reported the predominance of well-educated parents (Dornan, et al., 2007, 2009; Easterbrooks, O’Rourke, & Todd, 2000; Rhoades & Chisolm, 2000). Socioeconomic status has been found to be a significant predictor of better speech perception performance for children with hearing loss (Hodges, Dolan Ash, Balkany, Schloffman, & Butts, 1999), and has also been associated with better language for children with TH (Hart & Risley, 1995; Hoff-Ginsberg, 1991). Higher socioeconomic levels have also been found to be associated with higher reading and writing scores and a

lower risk of academic delays (Geers, 2003; Martineau, Lamarche, Marcoux, & Bernard, 2001). Low socioeconomic status has been reported as being associated with reduced academic opportunity and underachievement (Connor & Zwolan, 2004). Therefore it is suggested that if only children from high socioeconomic groups attended an education program, better outcomes for speech perception, language, reading, and writing would possibly result. As AVT is becoming more available to diverse family groups, the limitations of the generalized outcomes of this study must be acknowledged. Another limitation of this research is the fact that even though one child with mild cerebral palsy was included, two children had left the program in the first 9 months of the study because of the diagnosis of additional disabilities. It is acknowledged that the outcomes for the AVT group may not be applicable to today's growing cohort of children newly diagnosed with hearing loss as a result of newborn hearing screening who also have other disabilities (Larroque, et al., 2008). In addition, similar comments are applicable to the generalizability of outcomes data for children who are not native English speakers, which is another increasing demographic group in the population of children with hearing loss.

Whether AVT is effective for children and families across a wide socioeconomic range remains an important empirical question for future research. A further study limitation included the relatively small numbers of participants, particularly for reading and mathematics comparisons. Despite these limitations, the research goes some way towards providing a benchmark for minimum rate of progress for children with hearing loss acquiring listening and spoken language.

Summary

The results described here provide evidence that AVT is an effective intervention option for the AVT group. Speech perception improved significantly with moderate to high levels at 50 months after the start of the study. Although the group was identified at a mean age of 22.29 months, much later than the current "international gold standard" of 6 months of age (Joint Committee on Infant Hearing, 2007; Yoshinaga-Itano, et al., 1998), their language and speech attainments have been the same as a matched control group of children with TH over a 50 month time period. Reading, mathematics, and self-esteem outcomes were also comparable for both groups over the last 12 months of the study period. This study has provided a research model, utilizing a control group matched for language age, which could also be replicated across different languages, cultures, and countries and with different education approaches.

Acknowledgements

Financial support for this study has been provided by the Hear and Say Centre; School of Health and Rehabilitation Sciences, University of Queensland,

Brisbane, Australia; Queensland Council of Allied Health Professionals; and the Commonwealth of Australia through the Cooperative Research Centre for Cochlear Implant and Hearing Aid Innovation (CRC HEAR, Australia). The authors also wish to acknowledge the support of the staff and parents of the Hear and Say Centre, Ellen McKeering, Dr. Melody Harrison, and Peter Dornan. Statistical analysis was performed by Dr. Ross Darnell of the School of Health and Rehabilitation Sciences, University of Queensland, and Dr. Gabriella Constantinescu of the Hear and Say Centre. Special thanks are due to Jane Thompson and Renee O’Ryan who have helped in manuscript preparation.

References

- AG Bell Academy for Listening and Spoken Language. (2007). *Principles of LSLS auditory-verbal therapy*. Retrieved November 29, 2010, from <http://www.agbellacademy.org>.
- Agung, K.B., Purdy, S.C., & Kitamura, C. (2005). The Ling sound test revisited. *The Australian and New Zealand Journal of Audiology*, 27(1), 33–41.
- Australian Council of Education Research. (2005). *Progressive achievement tests in mathematics (PAT Maths)* (3rd ed.), Melbourne, Australia: ACER Press.
- Bat-Chava, Y., Martin, D., & Kosciw, J. (2005). Longitudinal improvements in communication and socialization of deaf children with cochlear implants and hearing aids: Evidence from parental reports. *The Journal of Child Psychology and Psychiatry*, 46(12), 1287–1296.
- Blamey, P.J., Barry, J., Bow, C., Sarant, J., Paatsch, L., & Wales, R. (2001). The development of speech production following cochlear implantations. *Clinical Linguistics and Phonetics*, 15(5), 363–382.
- Blamey, P.J., Sarant, J.Z., Paatsch, L.E., Barry, J.G., Bow, C.P., Wales, R.J., et al. (2001). Relationships among speech perception, production, language, hearing loss, and age in children with impaired hearing. *Journal of Speech, Language, and Hearing Research*, 44(2), 264–285.
- Boothroyd, A., & Boothroyd-Turner, D. (2002). Postimplantation audition and educational attainment in children with prelingually acquired profound deafness. *Annals of Otology, Rhinology & Laryngology*, 111, 79–84.
- Ching, T., Dillon, H., Day, J., Crowe, K., Close, L., Chisolm, K., et al. (2009). Early language outcomes of children with cochlear implants: Interim findings of the NAL study on longitudinal outcomes of children with hearing impairment. *Cochlear Implants International*, 10(S1), 28–32.
- Ching, T., Psarros, C., & Incerti, P. (2003). Management of children using cochlear implants and hearing aids. *The Volta Review*, 103(1), 39–57.
- Connor, C.M., & Zwolan, T.A. (2004). Examining multiple sources of influence on the reading comprehensive skills of children who use cochlear implants. *Journal of Speech, Language and Hearing Research*, 47(3), 509–526.

- Damen, G.W., van den Oever-Goltstein, M.H., Langereis, M.C., Chute, P.M., & Mylanus, E.A. (2006). Classroom performance of children with cochlear implants in mainstream education. *Annals of Otology, Rhinology, and Laryngology*, 115(7), 542–552.
- Dettman, S.J., Pinder, D., Briggs, R.J.S., Dowell, R.C., & Leigh, J.R. (2007). Communication development in children who receive the cochlear implant younger than 12 months: Risks versus benefits. *Ear & Hearing*, 28(2 Suppl), 11S–18S.
- Doig, B., & de Lemnos, M. (2000). *I can do maths*. Melbourne, Australia: ACER Press.
- Dornan, D., Hickson, L., Murdoch, B., & Houston, T. (2007). Outcomes of an auditory-verbal program for children with hearing loss: A comparative study with a matched group of children with typical hearing. *The Volta Review*, 107(1), 37–54.
- Dornan, D., Hickson, L., Murdoch, B., & Houston, T. (2008). Speech and language outcomes for children with hearing loss educated in auditory-verbal therapy programs: A review of the evidence. *Communicative Disorders Review*, 2(3–4), 157–172.
- Dornan, D., Hickson, L., Murdoch, B., & Houston, T. (2009). Longitudinal study of speech and language for children with hearing loss in auditory-verbal therapy programs. *The Volta Review*, 109(2–3), 1–25.
- Dunn, L., & Dunn, L. (1997). *Peabody picture vocabulary test* (3rd ed.). Circle Pines, MN: American Guidance Service.
- Durieux-Smith, A., Olds, J., Fitzpatrick, E., Eriks-Brophy, A., Duquette, C., Cappelli, M., et al. (1998). Outcome of AVT: Results of a follow-up study. *American Speech and Hearing Association Leader*, 3, 99.
- Easterbrooks, S.R., O'Rourke, C.M., & Todd, N.W. (2000). Child and family factors associated with deaf children's success in auditory-verbal therapy. *The American Journal of Otology*, 21, 341–344.
- Eisenberg, L.S., Kirk, K.I., Martinez, A.S., Ying, E., & Miyamoto, R.T. (2004). Communication abilities of children with aided residual hearing: Comparison with cochlear implant users. *Archives of Otolaryngology – Head & Neck Surgery*, 130, 563–569.
- Eriks-Brophy, A. (2004). Outcomes of auditory-verbal therapy: A review of the evidence and a call for action. *The Volta Review*, 104(1), 21–35.
- Fagan, M.K., & Pisoni, D.B. (2010). Hearing experience and receptive vocabulary development in deaf children with cochlear implants. *Journal of Deaf Studies and Deaf Education*, 15(2), 149–161.
- Geers, A.E. (2003). Predictors of reading skill development in children with early cochlear implantation. *Ear and Hearing*, 24(1), 59S–68S.
- Geers, A., Brenner, C., Nicholas, J., Uchanski, R., Tye-Murray, N., & Tobey, E. (2002). Rehabilitation factors contributing to implant benefit in children. *Annals of Otology, Rhinology and Laryngology*, 111(5), Suppl. 189, Part 2, 127–130.

- Geers, A.E., Nicholas, J.G., & Sedey, A.L. (2003). Language skills of children with early cochlear implantation. *Ear and Hearing*, 24(Suppl. 1), S46–S58.
- Geers, A., Tobey, E., Moog, J., & Brenner, C. (2008). Long-term outcomes of cochlear implantation in the preschool years: From elementary grades to high school. *International Journal of Audiology*, 47(Suppl. 2), S21–S30.
- Goldberg, D.M., & Flexer, C. (1993). Outcome survey of auditory-verbal graduates: Study of clinical efficacy. *Journal of the American Academy of Audiology*, 4, 189–200.
- Goldberg, D.M., & Flexer, C. (2001). Auditory-verbal graduates: An updated outcome survey of clinical efficacy. *Journal of the American Academy of Audiology*, 12(8), 406–414.
- Goldman, R., & Fristoe, M. (2001). *Goldman-Fristoe test of articulation*. Shoreview, MN: American Guidance Service.
- Gravel, J., & O’Gara, J. (2003). Communication options for children with hearing loss. *Mental Retardation and Developmental Disabilities Research Reviews*, 9, 243–251.
- Hart, B., & Risley, T.R. (1995). *Meaningful differences in the everyday experience of young American children*. Baltimore, MD: Paul H. Brooks.
- Hayes, H., Geers, A.E., Treiman, R., & Moog, J.S. (2009). Receptive vocabulary development in deaf children with cochlear implants: Achievement in an intensive auditory-oral educational setting. *Ear and Hearing*, 30, 128–135.
- Hintermair, M. (2006). Parental resources, parental stress, and socioemotional development of deaf and hard of hearing children. *Journal of Deaf Studies & Deaf Education*, 11(4), 493–513.
- Hodges, A.V., Dolan Ash, M., Balkany, T.J., Schloffman, J.J., & Butts S.L. (1999). Speech perception results in children with cochlear implants: Contributing factors. *Otolaryngology, Head and Neck Surgery*, 121(1), 31–34.
- Hoff-Ginsberg, E. (1991). Mother-child conversation in different social classes and communicative settings. *Child Development*, 62, 782–796. Retrieved September 26, 2010, from <http://onlinelibrary.wiley.com/doi/10.1111/j.1467-8624.1991.tb01569.x>.
- Hogan, S., Stokes, J., White, C., Tyszkiewicz, E., & Woolgar, A. (2008). An evaluation of auditory verbal therapy using rate of early language development as an outcome measure. *Deafness and Education International*, 10(3), 143–167.
- Joint Committee on Infant Hearing. (2007). Year 2007 position statement: Principles and guidelines for early hearing detection and intervention. *Pediatrics*, 120(4), 898–921.
- Jones, R. (2003). *Parental occupation coding*. Report submitted to Performance Measurement and Reporting Taskforce, Ministerial Council on Education, Employment Training and Youth Affairs, Appendix 4 (pp. 33–34). Retrieved June 8, 2009, from http://www.curriculum.edu.au/verve/_resources/parentaloccupation_file.pdf.

- Kelly, R.R., & Gaustad, M.G. (2007). Deaf college students' mathematical skills relative to morphological knowledge, reading level and language proficiency. *Journal of Deaf Studies and Deaf Education*, 12(1), 25–37.
- Kilminster, M.G.E., & Laird, E.M. (1978). Articulation development in children aged three to nine years. *Australian Journal of Human Communication Disorders*, 6(1), 23–30.
- Kritzer, K. (2009). Barely started and already left behind: A descriptive analysis of the mathematics ability demonstrated by young deaf children. *Journal of Deaf Studies and Deaf Education*, 14(4), 409–421.
- Kumar, S., & Centre for Allied Health Evidence Review Team. (2008). *A systematic review of the literature on early intervention for children with a permanent hearing loss*. Technical report produced for Queensland Health. Retrieved June 7, 2009, from <http://www.health.qld.gov.au/healthyhearing/pages/publications.asp>.
- Larroque, B., Ancel, P., Marret, S., Marchand, L., André, M., Arnaud, C., et al. (2008). Neurodevelopmental disabilities and special care of 5-year-old children born before 33 weeks of gestation (the EPIPAGE study): A longitudinal cohort study. *The Lancet*, 371(9615), 813–820.
- Laurenzi, C., & Monteiro, B. (1997). Mental health and deafness – the forgotten specialism? *ENT News*, 6, 22–24.
- Ling, D. (2002). *Speech and the hearing impaired child: Theory and practice* (2nd ed.). Washington, DC: Alexander Graham Bell Association for the Deaf.
- Marschark, M., Lang, H.G., & Albertini, J.A., (2002). *Educating deaf students: From research to practice*. New York: Oxford University Press.
- Marschark, M., Rhoten, C., & Fabich, M. (2007). Effects of cochlear implants on children's reading and academic achievements. *Journals of Deaf Studies and Deaf Education*, 12, 269–282.
- Marschark, M., & Spencer, P.E. (2003). *Oxford handbook of deaf studies, language, and education*. New York: Oxford University Press.
- Martineau, G., Lamarche, P.A., Marcoux, S., & Bernard P-M. (2001). The effect of early intervention on academic achievement of hearing-impaired children. *Early Education and Development*, 12(2), 275–289.
- Moeller, M.P., Tomblin, J.B., Yoshinaga-Itano, C., Connor, C.M., & Jerger, S. (2007). Current state of knowledge: Language and literacy of children with hearing impairment. *Ear and Hearing*, 28(6), 729–739.
- Morris, E. (2003). *Insight*. London: Nfer Nelson.
- National Institute of Child Health & Human Development. (2000). *Report of the National Reading Panel. Teaching children to read: An evidence-based assessment of the scientific research literature on reading and its implications for reading instruction*. Retrieved June 16, 2007, from <http://www.nichd.nih.gov/publications/nrp/smallbook.htm>.
- Nicholas, J.G., & Geers, A.E. (2003). Hearing status, language modality, and young children's communicative and linguistic behavior. *Journal of Deaf Studies and Deaf Education*, 8(4), 422–437.

- Nunes, T., & Moreno, C. (2002). An intervention program for promoting deaf pupil's achievement in mathematics. *Journal of Deaf Studies and Deaf Education*, 7(2), 120–133.
- Percy-Smith, L., Jensen, J.H., Josvassen, J.L., Jønsson, M.H., Andersen, J., Samar, C.F., et al. (2006). Parent's perceptions of their children's speech, language and social outcome after cochlear implantation. *Ugeskrift for Læger*, 168(33), 2659–2664.
- Phillips, L., Hassanzadeh, S., Kosaner, J., Martin, J., Deibl, M., & Anderson, I. (2009). Comparing auditory perception and speech production outcomes: Non-language specific assessment of auditory perception and speech production in children with cochlear implants. *Cochlear Implants International*, 10(2), 92–102.
- Pollack, D. (1970). *Educational audiology for the limited-hearing infant and pre-schooler*. Springfield, IL: Charles C. Thomas Publisher, Ltd.
- Powers, S. (2003). Influences of student and family factors on academic outcomes of mainstream secondary school deaf students. *Journal of Deaf Studies and Deaf Education*, 8, 57–78.
- Prizant, B.M., & Meyer, E.C. (1993). Socioemotional aspects of language and social-communication disorders in young children and their families. *American Journal of Speech-Language Pathology*, 2, 56–71.
- Rhoades, E.A. (2001). Language progress with an auditory-verbal approach for young children with hearing loss. *International Pediatrics*, 16(1), 1–7.
- Rhoades, E.A. (2006). Research outcomes of auditory-verbal intervention: Is the approach justified? *Deafness and Education International*, 8(3), 125–143.
- Rhoades, E.A., & Chisholm, T.H. (2000). Global language progress with an auditory-verbal approach for children who are deaf and hard of hearing. *The Volta Review*, 102(1), 5–24.
- Robertson, L., & Flexer, C. (1993). Reading development: A parent survey of children with hearing impairment who developed speech and language through the auditory-verbal method. *The Volta Review*, 95(3), 253–261.
- Romanik, S. (1990). *Auditory skills curriculum*. Moorebank, Australia: New South Wales Department of Education and Training.
- Sarant, J., Holt, C.M., Dowell, R.C., Rickards, F.W., & Blamey, P. (2008). Spoken language development in oral preschool children with permanent childhood deafness. *Journal of Deaf Studies and Deaf Education*, 14(2), 205–217.
- Schorr, E.A., Roth, F.P., & Fox, N.A. (2008). A comparison of the speech and language skills of children with cochlear implants and children with typical hearing. *Communicative Disorders Quarterly*, 29(4), 195–210.
- Schorr, E.A., Roth, F.P., & Fox, N.A. (2009). Quality of life for children with cochlear implants perceived benefits and problems and the perception of single words and emotional sounds. *Journal of Speech, Language and Hearing Research*, 52, 141–152.
- Semel, E., Wiig, E.H., & Secord, W.A. (1995). *Clinical evaluation of language fundamentals* (3rd ed.). San Antonio, TX: The Psychological Corporation.

- Spencer, L.J., & Oleson, J.J. (2008). Early listening and speaking skills predict later reading proficiency in pediatric cochlear implant users. *Ear and Hearing*, 29(2), 270–280.
- Sussman, K.D., Duncan, J., Estabrooks, W., Hulme, J., Moog, J.S., & McConkey Robbins, A. (2004). The option of spoken communication. *2004 Summit on Deafness Proceedings - Spoken Language Options in the 21st Century: Predicting Future Trends in Deafness*, pp. 14–17. Washington, DC: Alexander Graham Bell Association for Deaf and Hard of Hearing.
- Svirsky, M.A., Teoh, S.W., & Neuburger, H. (2004). Development of language and speech perception in congenitally, profoundly deaf children as a function of age at cochlear implantation. *Audiology and Neuro-Otology*, 9, 224–233.
- Tobey, E., Geers, A., Brenner, C., Altuna, D., & Gabbert, G. (2003). Factors associated with development of speech production skills in children implanted by age five. *Ear and Hearing*, 24(1 Suppl), 36S–45S.
- Traxler, C.B. (2000). The Stanford Achievement Test, 9th Edition: National norming and achievement standards for deaf and hard-of-hearing students. *Journal of Deaf Studies and Deaf Education*, 5, 337–348.
- Uziel, A.S., Sillon, M., Vieu, A., Artieres, F., Piron, J.P., Daures, J.P., et al. (2007). Ten-year follow-up of a consecutive series of children with multichannel cochlear implants. *Otology and Neurotology*, 28(5), 615–628.
- Vermeulen, A., van Bon, W., Schreuder, R., Knoors, H., & Snik, A. (2007). Reading comprehension of deaf children with cochlear implants. *The Journal of Deaf Studies and Deaf Education*, 12(3), 283–302.
- Vincent, D., Crumpler, M., & de la Mare, M. (1997). *Reading progress tests*. Berkshire, UK: Nfer Nelson.
- Wray, D., Flexer, C., & Vaccaro, V. (1997). Classroom performance of children who are deaf or hard of hearing and who learned spoken communication through the auditory-verbal approach: An evaluation of treatment efficacy. *The Volta Review*, 99(2), 107–119.
- Yoshinaga-Itano, C., Sedey, A.L., Coulter, D.K., & Mehl, A.L. (1998). Language of early and later identified children with hearing loss. *Pediatrics*, 102, 1161–1171.
- Zimmerman, I.L., Steiner, V.G., & Pond, R.E. (2002). *Preschool Language Scale - 4*. San Antonio, TX: The Psychological Corporation.

Appendix: Further Description of Reading, Mathematics and Self-Esteem Tests

Reading Tests

Reading Progress Tests (RPT) (Vincent, Crumpler, & de la Mare, 1997)

The Reading Progress Tests are a series of 7 British tests for ages 5 to 11. They comprise a Literacy Baseline Test of prereading and early literacy skills and 6 tests of reading comprehension (Reading Progress Test 1 to Reading Progress Test 6).

These tests provide individual or group measures of reading progress in two stages, Stage 1 (5–7 years) and Stage 2 (7–11 years) through the first 6 years of schooling. Tests are administered in a manner that ensures that children comprehend the nature of the task. Feedback is given when the method of response is incorrect, with time to correct their response. Feedback on whether the response is correct or incorrect is not given. Up to three repeats of a target word in instructions, plus practice items, are allowable. Children are reminded to refer back to the text to prevent relying on memory for their responses. There are no time limits but they usually take up to 45–50 minutes to administer. Australian norms were available and percentile ranks have been used to describe a child's level of ability.

The Literacy Baseline Test has three purposes: to provide a baseline from which to measure subsequent progress, as a screening procedure designed to identify a child likely to face difficulties in development of early reading skills, and as an appraisal of early literacy development. This test assesses existing reading and spelling ability, identification of initial sounds in spoken words and the identification of rhymes in spoken words (phonological awareness), familiarity with literacy concepts (such as knowing which words on the cover of a book are likely to be the name of the book, or which is the first word in a line of print), knowledge of letter names, and letter sounds. The child is asked to underline, or otherwise mark or point to, the correct response. This test was administered to Grade 1 children in the present study.

RPT1 and RPT2 are tests of reading comprehension that have two main purposes: to allow a standardized assessment of the child's reading comprehension and to monitor a child's progress in reading comprehension from one assessment point to the next in comparison with the progress made by other children in the same age-group. Both tests include three main types of comprehension question: (1) identifying the meaning of individual words, (2) selection of the correct answer from a number of choices after reading a short story, nonfiction passage or poem, and (3) choosing or supplying missing words in a short story or non-fiction passage. The majority of responses required consisted of marking one of a multiple choice selection. These tests were administered to Grade 2 and Grade 3 children in the present study.

RPT3, RPT4, RPT5, and RPT6 are similar to RPT1 and RPT2 in construction and administration, but are of a higher reading comprehension level. They were administered to Grade 4, 5, 6, and 7 children, respectively, in the present study.

Mathematics Tests

I Can Do Maths (Doig & de Lemnos, 2000)

I Can Do Maths is an Australian test of numeracy development in the early years of schooling. Children are requested to write, draw, count, and measure in response to the questions, which cover three main areas of numeracy (number, measurement, and space) and are ordered by increasing level of difficulty. The complete set of questions is covered in two books: Level A (30 questions), which was administered to children in Grade 1 in the present study, and Level B (33 questions), which was administered to a child in Grade 2. All questions are read to the children to avoid performance being affected by reading ability. The test is untimed but usually takes 30–40 minutes. A short break can be given. Australian norms were available and scores were expressed as a percentile rank.

Progressive Achievement Tests in Mathematics (PATMaths)
(Australian Council of Educational Research, 2005)

PATMaths is an Australian test of mathematics consisting of 8 tests (Test A and Tests 1 to 7), each in a separate book and each containing separate assessments of number, space, measurement, chance, and data with later tests containing questions on patterns and algebra. Test A required 20 minutes of testing time plus time for administration, and Tests 1 to 7 required 40 minutes of testing time. Test A was administered to children in Grade 3 while Test 1 was given to Grade 4 children, Test 2 to Grade 5, Test 3 to Grade 6, and Test 4 to Grade 7. Australian norms were available and scores were expressed as a percentile rank.

Self-Esteem Tests

Insight Preschool and Insight Primary (Morris, 2002)

Insight Preschool and Insight Primary are self-esteem indicators which can be used to explore the three key elements of a child's self-esteem: their sense of self, belonging, and personal power. Insight Pre-School covers ages 3–5 years and Insight Primary covers ages 5–11 years. Self-esteem is seen as a highly personal experience unique to each individual, which can mean how a person believes in themselves, how they feel when they are with other people, or how

they feel when they tackle something new or difficult. Insight Preschool consists of 24 questions and Insight Primary consists of 36 questions that the parent reads and responds to in writing. This can also be answered by a teacher but in this study, the parent was asked to respond. The form of the test chosen was according to whether the child attended preschool or primary school. Examples of questions included "Is your child usually contented?" and "Does your child try something first before asking for help?" The scoring was according to categories of whether the behaviour was observed "Most of the time" (3 points), "Quite often" (2 points), "Occasionally" (1 point), or "Almost never" (0 points). Table 2 shows the categories for interpretation of these scores.

Venturing Beyond the Sentence Level: Narrative Skills in Children with Hearing Loss

Christina Reuterskiöld, Ph.D., Tina Ibertsson, Ph.D., and
Birgitta Sahlén, Ph.D.

This study explores the differences in oral narrative skills between school-age children with mild-to-moderate sensorineural hearing loss (HL) and children who have typical hearing and language development. Narrative samples were collected following a picture-elicited storytelling task. Language samples were transcribed and coded for a number of measures, including narrative content, syntax, and grammar as well as amount of relevant information shared with the listener. Results indicated that the most vulnerable aspect of narration in children with HL is sharing information that is relevant for the task and context. Children with a sensorineural HL diagnosed during their preschool years are at risk for poorer development of higher level language skills, such as narrative production, compared with same-age peers.

Introduction

This study explores the differences in oral narrative skills between school-age children with mild-to-moderate sensorineural hearing loss (HL) (better ear hearing level [BEHL] 30–70 dB) and children who have typical hearing and language development (TD). It is well known that a HL can have a negative effect on a child's development of language and reading (Yoshinaga-Itano, 1986). Limited auditory input, even at the level of a mild-to-moderate HL, may potentially affect narrative skills as well. Students with HL are likely to be

Christina Reuterskiöld, Ph.D., is an Associate Professor in the Department of Communicative Sciences and Disorders at New York University. Tina Ibertsson, Ph.D., is a Senior Lecturer in the Department of Logopedics, Phoniatrics and Audiology at Lund University in Sweden. Birgitta Sahlén, Ph.D., is a Professor in the Department of Logopedics, Phoniatrics and Audiology at Lund University in Sweden. Correspondence concerning this manuscript should be directed to Dr. Reuterskiöld at ecw4@nyu.edu.

part of a school-based, speech-language pathologist's (SLP) caseload, although access to an SLP may differ among countries and regions. Furthermore, children with milder forms of HL may show subtle language and communication problems, which may elude the SLP (Brackett, 1997).

The present study was part of a more comprehensive project focusing on children with HL and children with language disabilities. Data on other skills, such as reading, working memory, and novel word learning, have been published elsewhere (Hansson, Forsberg, Löfqvist, Mäki-Torkko, & Sahlén, 2004; Sahlén, Hansson, Ibertsson, & Reuterskiöld-Wagner, 2005).

Narrative Skills in Children

Narrative skills are crucial higher language abilities, which are important for academic success and, in particular, for reading comprehension (Bishop & Edmundson, 1987; Feagans & Applebaum, 1986; for an extensive discussion, also see *Topics in Language Disorders*, 28, 2008). Narration is considered an ecologically valid way to assess language skills in school-aged children (Botting, 2002). From preschool age, students are expected to share personal and fictional narratives with other children and adults. Cognitive processing demands are high in a narrative task since speakers need to function at different levels of processing simultaneously. According to Hedberg and Westby (1993), producing a narrative is more difficult than participating in a conversation for three reasons. First, the storyteller has to formulate sentences that relate to a central theme or topic and that follow one another temporally or logically (centering and chaining). These mental operations must be performed simultaneously, thus forcing the storyteller to function at two different levels. Second, during storytelling, the narrator does not receive the same kind of support from listeners as from a speaking partner in a conversation. Third, during narration, cues from the environment are less readily available than during a conversation, which is usually more context-bound than a narrative. A narrative task is thus an example of a higher level language skill, which requires considerable amounts of cognitive resources.

Researchers have suggested that problems experienced by children with language disabilities may be a result of a limited processing capacity (Johnston, 1994, 2006; Kahneman, 1973; Leonard, 1998). Instead of focusing on a lack of knowledge in one or several specific areas, this view focuses on the simultaneous and synergistic interaction of different processes with each other, with context, and with the nature of the material to be processed (Lahey & Bloom, 1994). Johnston (1994) identified three aspects of processing capacity: rate, efficiency, and power. Kail and Salthouse (1994) likewise used three metaphors for aspects of information processing, but called them time, space, and energy. Time refers to the rate at which information can be processed, and space to the availability of space in memory. Energy refers to whether or not the child possesses sufficient mental resources to complete tasks of varying complexity

(Leonard, 1998). As Kail and Salthouse (1994) pointed out, aspects of energy, space, and time are interrelated and may sometimes be interchangeable, an issue that is related to the complexity of the task. Previous research has shown that children with language disabilities demonstrate difficulties with several levels of narrative production, such as content organization, grammatical accuracy, and the ability to share relevant content with the listener when compared with children with TD (Fey, Catts, Proctor-Williams, Tomblin, & Zhang, 2004; Liles, Duffy, Merritt, & Purcell, 1995; Reuterskiöld-Wagner, Sahlén, & Nettelbladt, 1999).

Language and Literacy Skills in Children with Hearing Loss

Gilbertson and Kamhi (1995) studied novel word learning in children with mild or moderate sensorineural HL. They included a battery of language tests in their study and found that only 50% of the participating children with mild or moderate sensorineural HL performed as well as children who had typical hearing. Performance on language tests was not related to the level of HL. The authors concluded that there is a tendency to view language disabilities in this population as secondary to the HL. Some children, however, may be better viewed as having language disabilities in the presence of a concurrent HL (Gilbertson & Kamhi, 1995).

In two studies comparing language and literacy skills in children with mild-to-moderate HL and children with specific language disabilities (SLD), researchers in Oxford (Briscoe, Bishop, & Norbury, 2001; Norbury, Bishop, & Briscoe, 2001) found that children with HL were not as behind with language and literacy skills as expected. The researchers discussed their results in light of current theories of SLD as a result of underlying auditory processing deficits. Their conclusion was that although some children with HL have phonological processing deficits like children with SLD, these deficits do not seem to be sufficient to cause the severe problems with language and literacy development as seen in children with SLD. Consistent with the results by the Oxford group, Sahlén et al. (2005) found that children with HL performed closer to the level of children with TD on reading measures than to that of children with SLD. There was no difference between the children with HL and the children with SLD on nonword repetition. Children with TD performed significantly better than both clinical groups on this measure. As expected, phonological processing seems to be a vulnerable area in children with HL, but as a group these children appear to compensate for their weakness in ways that children with SLD do not. The children in the present study were also participants in the Sahlén et al. (2005) study.

It is well known that there is a reciprocal relationship between the development of listening and spoken language and written language throughout childhood (e.g., Catts & Kamhi, 2005). Reading can be viewed as a receptive language skill with language presented in a written mode. In the present study,

we were interested in characterizing how children with mild-to-moderate HL developed their higher level expressive skills, in particular narrative skills.

Narrative Skills in Children with Hearing Loss

A few studies have investigated narrative skills in children with HL. A study by Yoshinaga-Itano (1986) included children representing five levels of HL, from mild to profound. Subjects were divided into five age groups, from 7 to 21 years. Although the focus of that study was the production of written, one-picture elicited narratives, it is interesting to note that children with HL, across groups, used significantly fewer cohesive devices (e.g., use of connectives and features of lexical cohesion) than children who had typical hearing. Furthermore, children with HL included fewer story-grammar units than children with typical hearing. When comparing the groups with hearing losses, a surprising result was that the moderate HL group had better stories than the mild HL group. Difficulties with higher level language skills thus seem to occur not only in children with more severe HL but also in children with mild HL.

Greenfield (2002) compared oral narrative development in 4-, 5-, and 6-year-old mainstreamed children with moderate-to-severe HL and children who had typical hearing. Results showed that the children with mild-to-moderate HL used fewer story-grammar elements in their narrations compared to the children with typical hearing. The difference between the groups was particularly apparent in a story-retelling condition compared with sequential picture-generated stories and personal narratives. Greenfield also included an analysis of the use of past tense markers and temporal terms (e.g., *and then*, *when*, *first*, *after*, and *later*). Both types of structures were used less frequently across tasks by children with HL than by children with typical hearing.

In this paper, we extend earlier work on narrative skills in children with language disabilities (Reuterskiöld-Wagner, et al., 1999; Reuterskiöld-Wagner, Nettelbladt, & Nilholm, 2000) and focus on school-aged children with mild-to-moderate HL. There is a paucity of studies on language skills in the Swedish-speaking school-aged population with HL, and to our knowledge this is the first study focusing on narrative skills. In the current study we use a set of measures based on previous research indicating areas of vulnerability in children with language disabilities. Swedish-speaking children with disabilities frequently use nonfinite verb forms, such as the infinitive, when finite forms are required. They also show difficulties with word order in sentences initiated with an element other than the subject (Hansson & Leonard, 2003; Hansson, Nettelbladt, & Leonard, 2000). In Swedish, the verb must be placed in the second position in a sentence and regular word order is subject-verb-object, as in English. For example, *Jag äter kakor.* (*I eat cookies.*) However, when a word other than the subject, such as an adverb, is placed first in the sentence, the verb has to follow. This results in an X-verb-subject (XVS) word order: *Nu äter*

jag kakor. (Now eat I cookies.) An important feature of storytelling is that utterances are commonly initiated with a connective, such as the temporal adverb *then*. In Swedish, the word order will thus be XVS, and this is used to create cohesion in stories.

Cohesion in stories is also created through the use of conjunctions or connectives. Seminal work in this area was conducted by Halliday and Hasan (1976; also see Peterson & Dodsworth, 1991), who pointed out that the function of conjunctive elements in narratives is to conjoin meaning across sentences independently of sentence grammar. The purpose of connectives is to show how the content expressed is logically connected to previous text: additively, temporary, causally, or adversely. Vion and Colas (2005) found that temporal connectives were most common, followed by additive, causal, and adversative in picture-elicited stories from 191 French-speaking children ages 7–11 years. However, according to Lahey (1988), the developmental sequence in narrative production progresses from additive narratives to temporal narratives, and finally causal narratives. The type of connective used indicates which type of relationship exists between propositions. In an additive chain, where the connective *and* is used, the order of propositions is not important. Statements can change place without a noticeable effect on the overall story. Gillam and Johnston (1992), however, cited Thompson (1984), stating that it is often difficult to distinguish between the use of *and* as a connective and as a discourse adverbial. They therefore chose to exclude *and* from their narrative analysis of oral and written narratives. In a temporal narrative, the connectives *then* or *and then* are usually used, and the order of propositions cannot shift places without the narrative changing in context. Finally, the connectives *because*, *cause*, and *so* indicate a causal relationship between content units, and constitute the most advanced level of narrative development.

Earlier studies of English-speaking children with SLD have indicated lexical variation, and particularly the use of different verbs as an area of weakness (Conti-Ramsden & Jones, 1997; Fletcher & Peters, 1984; Leonard, Miller, & Gerber, 1999; Watkins, Kelly, Harbers, & Hollis, 1995). In addition, grammatical complexity and accuracy have been found to be a persistent problem in school-age children with an early diagnosis of language disabilities (e.g., Fey et al., 2004). We are currently investigating language skills in a group of school-aged Swedish-speaking children with a preschool diagnosis of a language disability.

Reuterskiöld-Wagner et al. (1999) used a measure of relevance similar to the one described by Schneider (1996) while studying narration in children with language disabilities. Utterances were judged as relevant to the context or not relevant to the context, according to a three-step scale. Utterances were classified as “different and relevant,” “different and irrelevant,” or “different and very irrelevant.” Utterances judged as irrelevant were expressed more frequently by children with a lower level of language comprehension than their peers. In the present study, a new measure of relevance was developed.

We computed a relevance ratio by which we tapped into the child's sharing of content judged as relevant to the story line. In our previous study, we found this level of narrative performance to be particularly vulnerable in children with language disabilities, particularly in children with a low level of language comprehension.

Language comprehension takes place at different levels. According to Sperber and Wilson's *Relevance Theory* (1986), there is a gap between the meaning of sentences and the thoughts conveyed in an utterance. The gap must be filled by inference, and the object of inference during a conversation is the speaker's intentions. The listener (and speaker) must decode the meaning of the words (the coding-decoding mode) and interpret the intention behind the use of the words (the inferential mode) simultaneously. When inferential communication is not understood it might result in misinterpretations. Several studies have shown that children with language disabilities have great difficulties interpreting implicit information in stories (Bishop & Adams, 1992; Reuterskiöld-Wagner, et al., 1999). The Relevance Theory has been used as a framework in studies involving children with language disabilities. Leinonen & Kerbel (1999) suggested that children with pragmatic disabilities may have particular problems understanding meaning in open-ended communicative situations where they must select the appropriate interpretation from a number of possible ones. Children with a mild-to-moderate HL are frequently subject to degraded auditory speech signals and may therefore learn to be sensitive to other people's intentions in order to grasp the meaning of utterances (top-down processing). It is not far-fetched to suggest that this strategy may sometimes become too challenging. If language processing takes place within a system of limited cognitive capacity, where there are trade-off effects between linguistic levels, it might be expected that a task with high processing demands, such as narration, may tax children with HL beyond their capacity.

Taken together, previous research involving both English-speaking and Swedish-speaking individuals with HL and individuals with language disabilities give us reason to think that school-aged Swedish-speaking children with mild-to-moderate HL may show problems with aspects of narrative production. Higher level language skills have not been studied in depth in children with mild-to-moderate HL. Based on the literature, we expected children with HL to perform below the level of children with TD on our measures. Narrative skills have been linked to reading skills and entail a high level of processing demands, including expressive language. A HL might tax the language processing system of a child with HL to its limits. The study was guided by the following question:

Is there a difference in narrative performance between Swedish-speaking school-age children with HL and children with TD in terms of narrative content, relevant information, tense marking of verbs, percent XVS structures,

number of communication unit (C-unit) connectives, lexical variation, and grammatical accuracy?

Method

Participants

The present study included two groups of children: 18 children with HL and 17 children with TD and language skills (control group). All children were assessed in the Department of Logopedics, Phoniatrics and Audiology at Lund University in Sweden, except for 5 children with HL, who were tested in their school for geographic reasons. The study was part of a larger project, and two experienced SLPs as well as one audiologist performed the assessments. The whole procedure was audio and video recorded.

The children with HL all had a mild-to-moderate, bilateral, sensorineural HL (BEHL 30–70 dB) and had been fitted with hearing aids in at least one ear. However, 1 child (P3 in Table 1), who had a progressive, high-frequency HL, was fitted several times but used her hearing aids inconsistently since she did not consider them helpful.

Participants were recruited from ear, nose, and throat clinics in southern Sweden, had parents with typical hearing, were educated using listening and spoken language and attended mainstream schools, and were monolingual speakers of Swedish. As hearing ability was the key consideration, our subjects were selected based on results obtained from audiological testing, and their performance on language measures did not form the basis for inclusion in the study. Eighteen children between 9 years, 1 month, and 13 years, 3 months (mean age: 11 years, 1 month) participated and had a mean BEHL of 41.40 dB (average air conduction threshold at octave intervals between 500 and 4000 Hz) and a range of 30–57.5 dB HL. Sixteen children had a nonverbal IQ above 80, while 2 children had scores of 73 (as assessed with Raven's progressive matrices, RSPM; Raven, Court, & Raven, 1990). According to the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition, an IQ of 70 is the cutoff for an intellectual handicap or mental retardation. The mean nonverbal IQ was 91 (standard deviation [SD] 13.67). The age of identification of the HL varied (mean age of identification: 50.6 months; median: 52.5 months). For demographic data, see Table 1.

The control subjects had no history of any developmental or academic delay. Mean age of the control children was 10 years, 1 month (range: 8 years, 1 month – 11 years, 9 months). IQ testing of children without developmental or academic delays is not customary in Sweden, and these scores were not available. The study was approved by the University Committee on Research Involving Human Subjects at the Medical Faculty, Lund University. Parents received written and oral information about the project and signed an informed consent form stating that they and their child agreed to participate.

Table 1. Demographic information for children with HL

ID	Gender	IQ	Age at diagnosis (months)	Age at start of HA use (months)	BEHL 0.5-4 kHz* (dB)	WEHL 0.5-4kHz** (dB)	Progressive HL	Aetiology
1	F	88	36	36	58	58	No	Unknown
2	F	93	37	37	58	73	No	Unknown
3	F	118	89	89***	58	61	Yes	Hereditary
4	F	88	55	57	50	50	Yes	Unknown
5	M	87	56	57	34	35	No	Unknown
6	F	80	50	51	56	58	Yes	Unknown
7	F	100	15	15.5	45	48	No	Unknown
8	M	81	53	58	36	41	Unknown	Unknown
9	M	73	36	50	36	40	Unknown	Unknown
10	F	83	65	67	32	45	Unknown	Unknown
11	F	111	29	36	33	43	No	Hereditary
12	M	73	59	60	30	34	No	Unknown
13	M	100	53	68	31	36	No	Hereditary
14	F	117	52	53	40	43	No	Hereditary
15	F	82	52	53	31	36	No	Unknown
16	M	100	36	38	45	50	No	Hereditary
17	M	98	54	57	33	45	No	Unknown
18	M	82	84	87	37	40	No	Hereditary

* BEHL 0.5-4 kHz = Better ear hearing level, average threshold at 0.5, 1, 2, and 4 kHz

** WEHL 0.5-4 kHz = Worse ear hearing level, average threshold at 0.5, 1, 2, and 4 kHz

*** child fitted, but does not use hearing aids

HA = hearing aid

Procedure

As a practice item, the test administrator presented each child with a picture sequence, one picture at a time, and told a story. Then the examiner told the child, "Now I am going to show you some pictures, and you will tell me a story. I'll help you to get started." Pictures from *One Frog Too Many* (Mayer & Mayer, 1975) were laid out one at a time, and the child was asked to look carefully at each picture. The examiner pointed to the first picture, providing a sentence as a story starter: "This story is about a boy and his pets, who are going out on a raft," and asked the child to continue the story. No support apart from nodding and acknowledgment by a "mhm" or a "yes" was provided. Only in cases when the child was silent or did not continue the story did the examiner provide support, such as asking "and then?" or repeating the child's utterance. The full story can be found in Appendix A.

Analysis of Transcriptions

All narratives were transcribed by the first author, who has extensive experience in transcription of language samples and who followed the transcription

conventions in the SALT program (Miller & Chapman, 1983–2008), including coding of unintelligible utterances. The transcribed narratives were divided into C-units (Loban, 1976) consisting of a main clause and any attached subordinate clauses. Criteria were defined based on Hughes, McGillivray, & Schmidec's criteria (1997). C-units were coded for a number of measures.

Content was measured by computing the percentage of story-grammar units from a maximum score of 13 story-grammar units (Stein & Glenn, 1979). Carefully selected criteria for each target statement were set up for an utterance to qualify as a story-grammar unit. For target C-unit 10 (e.g., see Appendices A and B) "They can't find him," the child had to include "not find" in order for the utterance to generate a story-grammar score.

A relevance ratio was obtained by computing a story-grammar score per number of C-units. This measure was included to tap into the ability of the child to stay on topic and provide information that was relevant and crucial for the listener to understand the content of the story. Children who produced a high number of utterances (or C-units) without any target story-grammar information obtained a low relevance ratio.

Tense marking of verbs was assessed by computing unmarked verbs per total number of verbs. Unmarked verbs include nonfinite forms where the tense marker or a modal auxiliary has been omitted as well as forms of verbs of the first conjugation where past tense marking is optional in spoken language.

Percentage of XVS clauses from C-units containing a subject and a verb was included as a measure of word order.

Number of C-unit connectives (initiating C-units) was computed per C-unit. We included *och* (and), *sedan/sen*, (then/and then), *därför/så* (because/so), and *men* (but). These structures represent additive, temporal, causal, and adversative connectives (Lahey, 1988; Vion & Colas, 2005). According to Lahey, there is a developmental progression in the use of additive, temporal, and causal chains, with the first being least advanced and the last being most advanced.

Lexical variation was measured as number of different verbs per C-unit. Percentage of correct C-units was included as a holistic measure of grammatical accuracy.

Statistical Analyses

The first author of this study plus an expert on grammatical features of language disabilities in Swedish (see Hansson, et al., 2000) each independently coded 50% of the narratives from the group with TD. The first author coded all the narratives from the children with HL. The second author also coded 10% of these narratives. Inter-rater agreement on 10% of the narratives was 100% on story-grammar score and number of correct C-units. On number of XVS structures and number of C-units, reliability reached 98% and for the number of unmarked and different verbs, it reached 97%. Agreement was computed by dividing the number of codes in agreement by the number of codes in the

transcript. All coding was compared and consensus was reached through discussion. To view the narrative with target content, see Appendix B.

To be conservative, we used the unpaired, exact Wilcoxon-Mann-Whitney (WMW) test for group comparisons. The WMW test was chosen because it does not assume a normal distribution, and the exact versions were used to account for the small sample sizes. We did not use a Bonferroni test since this procedure is overly conservative (see Perneger, 1998).

Results

Table 2 shows that there were no significant differences between the group of children with HL and the group with TD, other than that the children with HL showed a lower relevance ratio than their same age-peers with TD (HL mean: 0.61, SD 0.25; TD mean: 0.79, SD 0.24; $Z = -2.531$, $p = .01$). This means that they provided less crucial content information per C-unit than their peers with TD. The variability (SD) in both groups was relatively high in terms of grammatical accuracy (percent correct C-units) and percent XVS clauses. Although not at a level of statistical significance, the group with TD produced slightly more story-grammar units, more XVS clauses, more grammatically correct C-units, and fewer unmarked verbs (e.g., / hon sitta/ for /hon sitter/ - /she sit/ for /she sits/ or /she is sitting/) than the group with HL.

Discussion

The results show that as a group, the children with HL performed at the same level as children with TD on all of our measures except one. Our review of the literature suggested that children with mild-to-moderate HL may show less developed higher order language skills than children with TD. This prediction was not confirmed, however, although a lack of significance does not necessarily mean no difference in small populations. In addition to analyzing results at the group level, we conducted a post-hoc analysis of potential outliers in the data. We did not find any such outliers, and the variation within the

Table 2. Group comparisons: HL and TD

<i>Variable</i>	<i>P-value comparing the 2 groups</i>	<i>Mean HL</i>	<i>SD HL</i>	<i>Mean TD</i>	<i>SD TD</i>
Story Gram	.171	5.89	2.05	6.7	1.8
Relevance Ratio	.01	0.61	0.25	0.79	0.14
% Corr C-units	.498	96.94	7.72	95.91	7.33
% Unmarked Verbs	.577	4.11	5.58	3.35	5.58
% XVS Clauses	.98	51.94	27.98	57.2	25.06
Connect/ C-Unit	0.258	0.75	0.24	0.82	0.19
Diff Verbs/ C-Unit	.14	0.95	0.43	1.04	0.17

group of children with HL was similar to the variation in the TD group (Table 2). This result does not corroborate the results in the Gilbertson and Kamhi (1995) study, in which a subgroup of children with HL were found to show a language profile similar to children with language disabilities. One difference between the studies, which may have affected results, is that the participating children were younger in the Gilbertson & Kamhi study than the children in the current study. Furthermore, we found that, as in Sahlén et al. (2005), this study's group of children with HL performed at the same level as children with TD on reading comprehension and reading-related tasks, such as the Rapid Automatized Naming (RAN; Denckla & Rudel, 1974), as well as on lexical and complex working memory tasks.

XVS sentences constitute about 40% of all statements in Swedish conversational speech (Jørgensen, 1976). Both groups of children in the current study showed a rather high level of within-group variation on the use of XVS structures. The group with TD showed a SD of 25.1% and the children with HL a SD of 28%. These results may be interpreted as evidence that the ability to create cohesion in narratives through word order variation is a late feature of language development acquired throughout schooling.

There was no significant difference between the groups on production of C-unit connectives (*and, then, but, because, etc.*) linking C-units together. Although not at a level of statistical significance, there was a trend for children with HL to use more additive connectives and fewer of the more advanced connectives. Both groups, however, used very few adversative (*but*) and causal (*cause, because, so*) connectives. It is possible that the short, picture-elicited story used did not encourage the use of more advanced structures.

There was one significant difference between groups in the present study. The group of children with HL conveyed less relevant content information per C-unit than their peers. In other words, they may have produced as many C-units or more, but the content was not always crucial and relevant to the story. According to Sperber and Wilson (1986), a high degree of relevance makes information worth processing for an individual. These authors pointed out that human beings are, and need to be, efficient at information processing. Information processing consumes energy and takes effort. Since efficiency is defined relative to a goal, reaching that goal at a minimal cost is desirable. It seems reasonable to think that a subtle negative impact on information processing during childhood, such as from a HL, may tax a child considerably and lead to a lower degree of efficacy in communication. If cognitive resources are consumed by an increased demand on lower level perceptual processes, higher level processes, such as judgment of intention and efficacy in communication, may suffer.

Implications of Findings

Narrative-based language intervention has been found to be useful for children with specific language disabilities and children with cochlear implants

(Justice, Swanson, & Buehler, 2008; Swanson, Fey, Mills, & Hood, 2005). Based on the group results from the current study, school-aged children with mild-to-moderate HL do not show difficulties with most aspects of narrative production, but they are not as effective in conveying relevant information to the listener as their peers with TD. Assessment of higher level language skills should thus be included in the monitoring of the academic performance of children with mild-to-moderate HL. Narrative intervention should explicitly target identification of relevant information and monitoring of listener reactions in a variety of narrative contexts. Furthermore, intervention goals should always be based on individual profiles of narrative performance, since a qualitative analysis of narrative development is necessary for appropriate goals to be formulated. Recent experimental research has shown that written narrative tasks facilitate oral narrative production more than oral narrative facilitates written narrative (Johansson, 2009). A focus on written narratives may thus be the most beneficial approach in work with school-aged children with poor narrative skills.

Methodological Considerations

Picture-elicited contexts may not be the best language elicitation tasks to really explore language skills in school-aged children. A range of narrative tasks should be used to obtain a full picture of a child's narrative abilities. A personal narrative can be used, although this is an elicitation context that demands more active participation from the clinician (Walldén & Åkerlund, 2008). To really explore school-aged children's higher level language skills, an expository task may be used. Furthermore, expository tasks have been found to elicit more grammatically complex story structures than other speaking contexts (Nippold, Mansfield, Billow, & Tomblin, 2008).

Conclusion

This study was conducted to gain insight into whether narration is a vulnerable area of language production in children with mild-to-moderate HL and, if so, which aspects of narration appear to be vulnerable in this population. Our results showed that a group of school-aged children with a HL diagnosed during their preschool years produced narratives that were similar to those produced by children with TD. One result indicated somewhat poorer development of higher level language skills, however. Children with HL were less effective in sharing crucial content information with the listener as compared with children with TD. Children with mild-to-moderate HL should be monitored and assessed at intervals by a SLP. Problems with higher level language skills can easily go unnoticed, and tasks that tax the child's language processing skills, such as narratives or expository tasks, should be included during assessment.

Acknowledgements

Our special gratitude goes to audiologists Ewa Holst and Ingrid Lennart, who performed hearing screenings. We also thank Dr. Kristina Hansson and Lena Asker-Arnason for their generous collaboration. We are very grateful to all the children and their parents for their participation, to graduate students for collecting normative data, to Jessica Forsberg for testing children with HL, and to Dr. Elina Mäki-Torkko for helping us recruit the children with HL. This study was financed by a grant (no. 2000-0171:01) from the Bank of Sweden Tercentenary Foundation. Preliminary results from the present study were presented at the 25th Annual Symposium on Research in Child Language Disorders in Madison, Wisconsin, in June 2004. Finally, the authors wish to thank Dr. Anders Löfqvist for his very helpful comments on an earlier version of the paper.

References

- Bishop, D., & Adams, C. (1992). Comprehension problems in children with specific language impairment: Literal and inferential meaning. *Journal of Speech and Hearing Research*, 35, 119–129.
- Bishop, D.V.M., & Edmundson, A. (1987). Language impaired 4-year olds: Distinguishing transient from persistent impairment. *Journal of Speech and Hearing Disorders*, 52, 156–173.
- Botting, N. (2002). Narrative as a clinical tool for the assessment of linguistic and pragmatic impairments. *Child Language Teaching and Therapy*, 18, 1–22.
- Brackett, D. (1997). Intervention for children with hearing impairment in the general education settings. *Language Speech and Hearing Services in the Schools*, 28, 355–362.
- Briscoe, J., Bishop, D.V.M., & Norbury, F.C. (2001). Phonological processing, language, and literacy: A comparison of children with mild-to-moderate sensorineural hearing loss and those with specific language impairment. *Journal of Child Psychology and Psychiatry*, 42, 329–340.
- Catts, H., & Kamhi, A. (Eds.). (2005). *Language and reading disabilities* (2nd ed.). Boston, MA: Pearson Edu, Inc.
- Conti-Ramsden, G., & Jones, M. (1997). Verb use in specific language impairment. *Journal of Speech, Language, and Hearing Research*, 40, 1298–1313.
- Denckla, M.B., & Rudel, R. (1974). Rapid “automatized” naming of pictured objects, colors, and letters, and numbers by normal children. *Cortex*, 10, 186–202.
- Feagans, L., & Applebaum, M.L. (1986). Validation of language subtypes in learning disabled children. *Journal of Educational Psychology*, 78, 358–364.
- Fey, M.E., Catts, H.W., Proctor-Williams, K., Tomblin, B., & Zhang, X. (2004). Oral and written story composition skills of children with language impairment. *Journal of Speech, Language and Hearing Research*, 47, 1301–1318.

- Fletcher, P., & Peters, A. (1984). Characterizing language impairment in children: An exploratory study. *Language Testing*, 1, 33–49.
- Gilbertson, M., & Kamhi, A.G. (1995). Novel word learning in children with hearing impairment. *Journal of Speech and Hearing Research*, 38, 630–642.
- Gillam, R.B., & Johnston, J. (1992). Spoken and written language relationships in language/learning impaired and normally achieving school-age children. *Journal of Speech and Hearing Research*, 35, 1303–1315.
- Greenfield, M. (2002). *Narrative differences of children with normal hearing in general education and mainstreamed children with hearing loss of ages four, five and six years* (Doctoral dissertation). New York: New York University. (Publication No. AAT 3031303.)
- Halliday, M.A.K., & Hasan, T. (1976). *Cohesion in English*. London: Longman.
- Hansson, K., Forsberg, J., Löfqvist, A., Mäki-Torkko, E., & Sahlén, B. (2004). Working memory and novel word learning in children with hearing impairment and children with specific language impairment. *International Journal of Language and Communication Disorders*, 39, 401–422.
- Hansson, K., & Leonard, L.B. (2003). The use and productivity of verb morphology in specific language impairment: An examination of Swedish. *Linguistics*, 41–42, 351–379.
- Hansson, K., Nettelbladt, U., & Leonard, L.B. (2000). Specific language impairment in Swedish: The status of verb morphology and word order. *Journal of Speech, Language and Hearing Research*, 43, 848–864.
- Hedberg, N.L., & Westby, C.E. (1993). *Analyzing storytelling skills: Theory to practice*. Tucson, AZ: Communication Skill Builders.
- Hughes, D., McGillivray, L., & Schmidec, M. (1997). *Guide to narrative language: Procedures for assessment*. Eau Claire, WI: Thinking Publications.
- Johansson, V. (2009). *Developmental aspects of text production in writing and speech* (Doctoral dissertation). Travaux de l'Institut de Linguistique de Lund no 48. Lund University, Lund, Sweden.
- Johnston, J. (1994). Cognitive abilities of children with language impairment. In R. Watkins & M. Rice (Eds.), *Specific language impairments in children*. Baltimore, MD: Paul H. Brookes.
- Johnston, J. (2006). *Thinking about child language. Research to practice*. Eau Claire, WI: Thinking Publications.
- Jørgensen, N. (1976). *Meningsbyggnaden i talad svenska* [Sentence structure in spoken Swedish]. Lund, Sweden: Studentlitteratur.
- Justice, E.D., Swanson, L.A., & Buehler, V. (2008). Use of narrative-based language intervention with children who have cochlear implants. *Topics in Language Disorders*, 28, 149–161.
- Kahneman, D. (1973). *Attention and effort*. Englewood Cliffs, NJ: Prentice-Hall.
- Kail, R., & Salthouse, T.A. (1994). Processing speed as a mental capacity. *Acta Psychologica*, 86, 199–225.
- Lahey, M. (1988). *Language disorders and language development*. Needham Heights, MA: Allyn & Bacon.

- Lahey, M., & Bloom, L. (1994). Variability and language learning disabilities. In G. Wallach & K. Butler (Eds.). *Language learning disabilities in school-age children and adolescents*. Boston, MA: Allyn & Bacon.
- Leinonen, E., & Kerbel, D., (1999). Relevance theory and pragmatic impairment. *International Journal of Language and Communication Disorders*, 34, 367–391.
- Leonard, L. (1998). *Children with specific language impairment*. Cambridge, MA: MIT Press.
- Leonard, B.L., Miller, C., & Gerber, E. (1999). Grammatical morphology and the lexicon in children with specific language impairment. *Journal of Speech, Language and Hearing Research*, 42, 678–689.
- Liles, B.Z., Duffy, R.J., Merritt, D.D., & Purcell, S.L. (1995). Measurement of narrative discourse ability in children with language disorders. *Journal of Speech and Hearing Research*, 38, 415–425.
- Loban, W. (1976). *Language development: Kindergarten through grade twelve*. Urbana, IL: National Council of Teachers of English.
- Mayer, M., & Mayer, M. (1975). *One frog too many*. New York: Penguin Books.
- Merritt, D.D., & Liles, B.Z. (1987). Grammar ability in children with and without language disorder: Story generation, story retelling and story comprehension. *Journal of Speech and Hearing Research*, 30, 539–552.
- Miller, J., & Chapman, R. (1983–2008). Computer programs for the systematic analysis of language transcripts SALT. Research Version 8. Language Analysis Laboratory, Waisman Center, University of Wisconsin-Madison.
- Nippold, M.A., Mansfield, T.C., Billow, J.L. & Tomblin, J.B. (2008). Expository discourse in adolescents with language impairments: Examining syntactic development. *American Journal of Speech-Language Pathology*, 17(4), 356–366.
- Norbury F.C., Bishop, D.V.M, & Briscoe, J. (2001). Production of English finite verb morphology: A comparison of SLD and mild-moderate hearing impairment. *Journal of Speech, Language, and Hearing Research*, 44, 165–178.
- Perneger, T.V. (1998). What's wrong with Bonferroni adjustments? *British Medical Journal*, 316, 1236–1238.
- Peterson, C., & Dodsworth, P. (1991). A longitudinal analysis of young children's cohesion and noun specification in narratives. *Journal of Child Language*, 18, 397–415.
- Raven, J.C., Court, J.H., & Raven, J. (1990). *Raven's standardized progressive matrices*. Oxford: Oxford Psychologists Press.
- Reuterskiöld-Wagner, C., Nettelbladt, U., & Nilholm, C. (2000). Conversation versus narration in preschool children with language impairment. *International Journal of Language and Communication Disorders*, 35, 337–352.
- Reuterskiöld-Wagner, C., Sahlén, B., & Nettelbladt, U. (1999). What's the story? Narration and comprehension in Swedish preschool children with language impairment. *Child Language Teaching and Therapy*, 15(2), 83–93.

- Sahlén, B., Hansson, K., Ibertsson, T., & Reuterskiöld-Wagner, C. (2005). Reading in primary school age—a comparative study of children with hearing impairment and children with specific language impairment. *Acta Neuropsychologia*, 1, 393–407.
- Schneider, P. (1996). Effects of pictures versus orally presented stories on story retellings by children with language impairment. *American Journal of Clinical Practice*, 5, 86–96.
- Sperber, D., & Wilson, D. (1986). *Relevance, Communication and Cognition*. Oxford: Basil Blackwell Ltd.
- Stein, N.R., & Glenn, C.G. (1979). An analysis of story comprehension in elementary school children. In R.O. Freedle (Ed.), *New directions in discourse processing* (pp. 53–120). Norwood, NJ: Ablex.
- Swanson, L.A., Fey, M.E., Mills, C.E., & Hood, L.S. (2005). Use of narrative-based language intervention with children who have specific language impairment. *American Journal of Speech–Language Pathology*, 14(2), 131–143.
- Thompson, S.A. (1984). “Subordination” in formal and informal discourse. In D. Schiffrin (Ed.), *Georgetown University Round Table on Language and Linguistics 1984* (pp. 85–94). Washington, DC: Georgetown University Press.
- Vion, M., & Colas, A. (2005). Using connectives in oral French narratives: Cognitive constraints and development of narrative skills. *First Language*, 25, 39–66.
- Walldén, R.-M., & Åkerlund, V. (2008). *Muntlig och skriftlig berättarförmåga samt läsförståelse hos barn med typisk språkutveckling i år 5 och 6* [Oral and written narration as reading comprehension in children with typical language development in grade 5 and 6]. (Master’s thesis). Lund University, Sweden.
- Watkins, R.V., Kelly, D.J., Harbers, H.M., & Hollis, W. (1995). Measuring children’s lexical diversity differentiating typical and impaired language learners. *Journal of Speech and Hearing Research*, 38, 1349–1355.
- Yoshinaga-Itano, C. (1986). Beyond the sentence level: What’s in a hearing impaired child’s story. *Topics in Language Disorders*, 6, 71–83.

Appendix A: Target Narrative

Swedish: Den stora grodan sparkar av den lilla grodan från flotten. Den lilla grodan blir rädd. Den stora grodan skrattar och tittar på den lilla grodan i vattnet, medan flotten åker vidare. Den lilla grodan blir rädd. Det ser sköldpaddan. Han skvallrar till pojken och hunden. Sköldpaddan och hunden blir arga på den stora grodan. Den stora grodan blir rädd när de är arga. "Åh nej" säger pojken. Alla säger "vi måste leta efter den lilla grodan." De letar överallt och ropar på den lilla grodan. De kan inte hitta honom. De får gå hem utan den lilla. Pojken är jätteledsen. Hunden morrar på den stora grodan. Den stora grodan skäms. Han får inte följa med hem.

English translation: The big frog kicks the little frog off the raft (or boat). The little frog is frightened. The big frog is laughing as he watches the little frog in the water, while the raft is floating away. The little frog is frightened. The turtle sees this. He tattle-tes on the big frog. The turtle and the dog get angry at the big frog. The big frog gets frightened. "Oh no," says the boy. They all say, "We have to look for the little frog." They look everywhere and call out for the little frog. They can't find him. They have to go home without the little one. The boy is very sad. The dog is growling at the big frog. The big frog feels ashamed. He is not allowed to walk home with the others.

Appendix B: Story-Grammar Scoring

Each minimal utterance yielded a score of 1, with a total possible score of 16 points.

<i>Story Element</i>	<i>Minimal Utterance</i>
<u>Initiating event:</u>	
1. The big frog kicks the little frog off the raft (or boat).	Big frog kicks little frog off. (He kicks him off).
2. The big frog is laughing as he watches the little frog in the water, while the raft is floating away. The little frog is frightened.	Big frog is laughing. Big frog happy/smiling.
<u>Reaction:</u>	
3. The turtle sees this.	Turtle watching/sees.
4. He tells the boy and the dog.	Tattle-tales on the big frog/turtle tells boy and/or dog. They are angry/upset.
5. The turtle and the dog get angry with the big frog.	
6. The big frog gets frightened	Big frog (he) is frightened.
7. "Oh no" says the boy.	Boy (he) is upset/surprised. (a negative reaction)
<u>Plan:</u>	
8. They all say "we have to look for the little frog."	Have to look/search.
<u>Action:</u>	
9. They look everywhere and call out for the little frog.	They go looking/searching. (indicate more than one character as agents)
<u>Consequence:</u>	
10. They can't find him.	Can't find him.
11. They have to go home.	Went home.
12. Without the little frog.	Without little frog.
<u>Resolution:</u>	
13. The boy is very sad.	The boy/he is sad.
14. The dog is growling at the big frog.	Dog growling/angry.
15. The big frog feels ashamed.	Big frog is ashamed/sad/feels bad.
16. He is not allowed to walk home with the others.	Big frog stays/is not coming.

Effects of Auditory-Verbal Therapy for School-Aged Children with Hearing Loss: An Exploratory Study

Elizabeth Fairgray, M.Sc., LSLS Cert. AVT; Suzanne C. Purdy, Ph.D.; and Jennifer L. Smart, Ph.D.

With modern improvements to hearing aids and cochlear implant (CI) technology, and consequently improved access to speech, there has been greater emphasis on listening-based therapies for children with hearing loss, such as auditory-verbal therapy (AVT). Speech and language, speech perception in noise, and reading were evaluated before and after 20 weeks of weekly speech and language therapy (SLT) based on AVT principles. Participants were 7 children (ages 5–17 years) with sensorineural hearing loss. Five participants had profound, bilateral sensorineural hearing loss and used 1 or 2 CIs. The remaining 2 had moderate-to-severe and severe-to-profound losses, respectively, and used hearing aids. Significant improvements were seen in speech perception and production, and in one measure of receptive language. The challenge for future research is to conduct controlled studies using a wider range of sensitive assessment tools in order to establish the real benefits of AVT-based SLT interventions to improve outcomes.

Introduction

Although the need for speech and language therapy (SLT) is widely recognized when working with children who are deaf or hard of hearing, there

Elizabeth Fairgray, M.Sc., LSLS Cert. AVT, is a Speech-Language Pathologist who runs The Listening and Language Clinic at the University of Auckland and is also a Research Fellow in the Speech Science Division of the Department of Psychology at The University of Auckland in New Zealand. Suzanne C. Purdy, Ph.D., is an Associate Professor and an Audiologist who heads the Speech Science Division of the Department of Psychology at The University of Auckland in New Zealand. Jennifer L. Smart, Ph.D., is an Assistant Professor in the Department of Audiology, Speech-Language Pathology and Deaf Studies at Towson University in Towson, Maryland. Correspondence concerning this article should be directed to Ms. Fairgray at L.Fairgray@auckland.ac.nz.

is little research evidence for improved communication outcomes after specific SLT interventions. In general, SLT approaches for children with hearing loss have focused either on listening skills or on the integration of multiple visual and auditory cues to aid communication. With improvements in hearing aid and cochlear implant (CI) technology, and consequently improved access to the speech signal, there has been greater emphasis on listening-based therapies (Connor, Craig, Raudenbush, Heavner, & Zwolan, 2006).

Auditory-verbal therapy (AVT) emphasizes listening rather than looking, early identification of hearing loss, and optimal amplification in order to enhance access to the speech signal (Caleffe-Schenck, 1992; Lim & Simser, 2005). Key elements of AVT can be summarized as (a) utilizing focused audiological management; (b) ensuring the immediate fitting of hearing technology; (c) presenting the auditory stimuli as the main sensory input; (d) providing early, intense (re)habilitation; (e) adopting a family-centered approach; (f) teaching language and speech through listening; (g) integrating listening into every aspect of daily life; and (h) promoting education in mainstream classes with peers who have typical hearing (Estabrooks, n.d.).

The Alexander Graham Bell Association for the Deaf and Hard of Hearing endorses a listening and spoken language approach, but there is paucity of research evidence for its effectiveness (Eriks-Brophy, 2004; Goldberg & Flexer, 1993; Rhoades, 1982; Rhoades & Chisholm, 2000; Wray, Flexer, & Vaccaro, 1997). Reasons for the lack of clear research findings include the variability in levels and duration of deafness, varying provision of amplification, different school settings, and different causes of deafness. Rhoades (2006) reviewed the outcomes of AVT research and concluded that there is limited evidence supporting the use of AVT.

The current study seeks to increase understanding of the impact of AVT on speech and language outcomes. It was hypothesized that children and young adults with moderate-to-profound hearing loss will benefit from SLT that is based on listening and spoken language principles. The specific aims of the project were to investigate (a) speech perception, phonology, and articulation; (b) understanding of spoken language; and (c) complexity of expressive language to determine whether these improved after a period of intensive SLT that included weekly visits and homework. Reading skills were also assessed, since changes in listening and spoken language skills may benefit a child's literacy development. It was, however, hypothesized that a longer period of SLT intervention or literacy-focused intervention would be needed to see gains in reading. This was an exploratory study, in preparation for a future controlled study, to determine the range of baseline abilities among school-aged children and to determine whether it is possible to measure improvements in speech, language, reading, and speech perception after a relatively short period of SLT intervention.

Methodology

Participants

The 7 participants (2 boys and 5 girls) were a convenience sample of children with moderate-to-profound hearing losses and ages ranging from 5 to 17 years old at the start of the study who lived at home with their parents. All participants were native New Zealanders. An additional eighth participant recruited into the study was withdrawn when the extent of her speech perception and language difficulties was revealed by baseline testing. She was referred for a cochlear implant (CI) assessment and was excluded from the study (by the CI program) when she was accepted as a CI candidate. Participant recruitment occurred via advertisement to otolaryngologists, audiologists, and teachers of the deaf. Five of the 7 participants had a bilateral profound hearing loss and used at least one CI. Six of the participants used an FM system consistently in the classroom. Table 1 lists participant ages and amplification type. Table 2 shows aided thresholds. No secondary disorder was identified for any of the children. All had nonverbal IQs in the average range, and none had attention deficit hyperactivity disorder. The family status of participants varied widely, from being a single child in a single-parent family to being one of six siblings in a two-parent family. Three of the children lived in single-parent families. All attended mainstream local schools.

The age at which hearing loss was diagnosed ranged from 1 to 4 years. Universal newborn hearing screening was not mandatory at the time the participants were born, which would have contributed to the late diagnoses. For the 5 participants with CIs, activation of the first CI ranged from 1 to 10 years. With the exception of 1 participant, all children were born to parents who had typical hearing. One child (participant 2) had meningitis at age 8 months.

Table 1. Amplification characteristics, age at diagnosis of hearing loss (years: months), age when first CI was received (years:months), and age at entry to the study (years:months). All participants received hearing aids within a few weeks of diagnosis

<i>Participant</i>	<i>Hearing aid (HA)/</i>		<i>Age at diagnosis</i>	<i>Age at first CI</i>	<i>Age at test</i>
	<i>Cochlear implant (CI)</i>	<i>Gender</i>			
P1	Right CI	F	1:1	5:1	17:7
P2	Right CI	M	0:11	1:0	8:7
P3	Bilateral HA	F	3:0	n/a	5:5
P4	Bilateral HA	F	3:2	n/a	10:4
P5	Right CI	M	2:6	4:4	10:9
P6	Bilateral CI	F	1:6	2:2	9:5
P7	Bilateral CI	F	1:5	1:7	6:8

F = female, M = male

Table 2. Aided hearing thresholds (dB HL) for the 7 participants

Frequency (Hz)	P1 <i>Right CI</i>	P2 <i>Right CI</i>	P3 <i>Bilat HA</i>	P4 <i>Bilat HA</i>	P5 <i>Right CI</i>	P6 <i>Right CI</i>	P7 <i>Bilat CI</i>
500	30	25	25	20	25	30	15
1000	35	30	20	15	30	25	20
2000	30	35	30	50	20	25	15
4000	35	30	20	40	25	25	20
8000	30			35	15		15

CI = cochlear implant; HA = hearing aid; Bilat = bilateral

He subsequently lost his hearing and developed kidney failure. At age 6, after 2 years of dialysis, he received a kidney transplant. The health difficulties this child experienced further exacerbated the speech and language delays associated with his bilateral profound hearing loss. This participant had severe language deficits at the outset of the study. His poor school attendance in previous years and lack of therapy due to the severity of his illness are likely to have contributed to his language delay.

Assessments

Each participant was brought to the University of Auckland Listening and Language Clinic for weekly SLT sessions provided by the first author, a speech-language pathologist with 25 years of experience who is also certified as a Listening and Spoken Language Specialist in Auditory-Verbal Therapy (LSLS Cert. AVT). The children were given weekly follow-up activities for home practice (approximately 20 minutes per day). Baseline assessment occurred over several sessions at the beginning of the participant’s AVT program, and follow-up assessment occurred immediately after 20 individual 1-hour appointments. Language, phonology, articulation, and reading were assessed using the Australian version of the CELF-4 (Clinical Evaluation of Language Fundamentals, 4th Edition), HAPP-3 (Hodson Assessment of Phonological Patterns Edition, 3rd Edition), NZAT (New Zealand Articulation Test), and WIAT-II (Wechsler Individual Achievement Test, 2nd Edition), respectively. Pre- and posttherapy assessments were performed in the same order over three or four assessment sessions.

Speech recognition in noise was evaluated using words from the Lexical Neighborhood Test (LNT) (Eisenberg, Martinez, Holowecky, & Pogorelsky, 2002; Kirk, Pisoni, & Osberger, 1995), re-recorded with a native New Zealand female speaker. LNT words were presented via a loudspeaker at zero degrees azimuth. Multi-talker speech babble was presented simultaneously via a custom-built mixer through the three other speakers, located at 90, 180, and 270 degrees azimuth, to simulate classroom listening. The LNT includes eight lists altogether: four “easy” lists (e.g., *give, monkey, juice, ducks, myself, pocket*)

and four “hard” lists (e.g., *taught, us, trick, worm, were, rain, cups, cats*) of 15 words per list. Easy words are words frequently heard with few lexical neighbors. Hard words are infrequently heard with many lexical neighbors (Eisenberg, et al., 2002). Fifteen easy words (one list) and 15 hard words (one list) were presented in the sound field at 70 dB SPL at a +5 dB signal-to-noise ratio (SNR). List order was counterbalanced across test sessions and children.

Language, Speech, and Reading Assessments

Language

Language was assessed using the Australian version of the CELF-4 (Semel, Wiig, & Secord, 2003). This is standardized on children who have hearing within typical limits. The CELF-4 has 19 subtests, but not all are applicable across the age range of participants in this study. There are two test forms, one for ages 5–8 years and one for ages 9–21 years. Results are reported here for norm-referenced subtests with age norms for all or most participants. They include three receptive language subtests (Concepts and Following Directions, Understanding Spoken Paragraphs, Word Classes-Receptive) and three expressive language subtests (Word Classes-Expressive, Formulated Sentences, Recalling Sentences). Core Language scores are based on four subtests that the test authors describe as “most discriminating [of disorder]” at each age level.

Phonology

Phonological development was assessed using the HAPP-3 (Hodson, 2004). This test is standardized for children with no hearing difficulties and measures developmental simplification processes present in typical speech development. All the sounds of spoken English are categorized into groups such as fricatives (s, z, ʃ, f, v), plosive stops (p, b, t, d, k, g), and nasals (m, n). Concern arises when developmental processes persist beyond the typical age range, or when there is an occurrence of phonological processes not seen in children with typical speech development. One example is the simplification process of “stopping” in which long, continuous fricative sounds are replaced by short, stopped sounds.

Articulation

Articulation was assessed using the NZAT, which is a picture-based test with 82 individual colored pictures illustrating vocabulary items familiar to New Zealand children (Moyle, 2005). The therapist phonetically transcribes each word as the child produces it. Words are scored as correct or incorrect, producing error scores that can be converted to standard scores using New

Zealand normative data for children ages less than 8 years old, the upper limit of the normative sample age range. The complete test assesses each consonant phoneme at the beginning, middle, and end of words, such as the /b/ in *bath*, *table*, and *web*. Consonant clusters are also assessed with words such as *bread*, *flower*, and *scratch*.

Speech Perception in Noise

Each participant was assessed before and after the block of therapy on his/her ability to listen to and then repeat a set of words in a background noise at +5 dB SNR. Recorded words were used to ensure that the stimuli were consistently delivered and responses were transcribed. During the assessment and intervention time, none of the participants had a change in CI or hearing aid settings that could affect speech scores.

Reading

Reading was assessed using the Word Reading, Pseudoword Decoding, and Reading Comprehension subtests of the WIAT-II, developed by the Psychological Corporation in 2002. Standardized scores were derived using the Australian norms for this test (Pearson PsychCorp, 2007). Skillful decoding of words can mask poor reading comprehension; hence it is important to assess a range of literacy skills. Reading comprehension is more closely related to listening comprehension, as both require understanding of vocabulary, verb tenses, and inferential statements.

Therapy Approach

Each week, the participant and a caregiver arrived at The Listening and Language Clinic at the University of Auckland. In general, one parent brought the participant, but sometimes an aunt or grandparent came instead. The intervention occurred at the same time each week and lasted for 1 hour. Unless a specific activity required a variation, the participant and therapist sat beside each other so that speech reading was minimized and a small distance was maintained between the speaker and listener. The caregiver was required to practice the activities and goals for the week at home. The caregiver and therapist took turns working with the participant on an activity. As part of this caregiver involvement, discussion occurred between the therapist and caregiver so that an activity could be extended to the home environment. For example, for the word *open*, the therapist explained why the word *open* had been selected. Possible reasons include production of the “p” sound in the middle of a word; practice of verb-noun phrases (e.g., “Open the door” or “Open the box.”); practice of adverbs (e.g., “Open it... quietly, slowly, later.”); and practice of past tense (e.g., “He **opened** the drawer.”).

A rapid review of the Ling Sounds (Ling, 1976) was conducted at the beginning of the session to verify CI and/or hearing aid function. Occasionally for participants using bilateral CIs, this check resulted in the discovery that one CI had a flat battery. On two occasions, a participant had audible feedback from her hearing aid and was referred to an audiologist for ear mold review.

Sessions followed a particular theme, divided into goals for expressive language, receptive language, articulation, and an age-appropriate cognitive activity that would reinforce the language concepts. Each session utilized toys, pictures, photos, books, and role play. For participant 2, one real challenge was selecting cognitively appropriate activities that could also be used in the context of extreme language delays. By following a topic-based approach each week, vocabulary was targeted systematically within different semantic areas. In conjunction with new vocabulary, appropriate concepts and a variety of verb forms were taught and practiced during the sessions. Session goals were selected as appropriate for the individual child. An example of this strategy is as follows:

- **Topic:** seasons, with an emphasis on autumn.
- **Nouns:** Wind, gusts, leaf, leaves, tree, bush, bushes, house, houses, child, children, weather, season.
- **Verbs:** blowing, blew, has blown, falling, fell, have fallen, running, ran, have run, shine, shone, has shone.
- **Adjectives:** hot, cold, wet, cool, warm, moist, humid, tropical.
- **Prepositions:** in, on, under, between, opposite.

To prevent confusion at home, target goals for home practice were written into the participant's book each week. With the younger participants, the target goals could be reinforced through picture activities, which resembled the toys and activities used during the session. The picture activities generally included one themed picture from the Verb Activity Sheets developed by van Asch Deaf Education Centre (n.d.). For the older participants, the format of the homework tended to follow a write-read-practice pattern rather than looking at pictures or engaging in adult-led activities to practice the goals. The intervention was also individualized by the use of Duplo and Playmobil® for younger participants, photos and pictures for older participants, and authentic pamphlets, brochures, and articles for the teenage participant.

Family involvement in therapy practice at home was monitored by documenting what the parent had achieved with the participant during the week. This was done at the end of the session so that therapy occurred first to optimize the participant's engagement in the therapy. All families were able to comply with the home activities, but every family had the occasional week (no more than 2 weeks per family) when other commitments took precedence and home-based activities did not occur. Due to New Zealand's paucity of SLT

services for school-aged children with hearing loss, families were highly motivated to engage in therapy practice at home.

Inclusion of AVT Principles in Therapy Approach

Principles One and Two: Focused Audiological Management and Immediate Fitting of Appropriate Hearing Technology

As part of the baseline measurement, the third author completed an audiological evaluation for each participant to check that participants' hearing instruments were giving them access to the speech spectrum. The 5 participants wearing CIs were confirmed to have bilateral profound hearing loss. One participant (originally recruited as an eighth participant) had profound hearing loss in the higher frequencies (2000–6000 Hz) and, despite the use of powerful hearing aids, her thresholds could not be lifted into the speech spectrum. Consequently, she was referred to the CI program to assess her candidacy for a CI. The CI assessment team stipulated that she discontinue her participation in the research while this evaluation took place, and hence she was excluded from further participation in the study.

For the remaining 7 participants, a brief Ling Six-Sound Test (Ling, 1976) was conducted at the start of each therapy session to check hearing instruments. On two occasions, 1 participant with hearing aids had a problem with acoustic feedback, so she was immediately seen by an audiologist. On another occasion, new ear mold impressions were taken.

Principle Three: Present the Auditory Stimulus as the Main Sensory Input

Listening and spoken language contrasts with methodologies that highlight visual aspects of language, which may focus on lip-reading or tactile cues. In New Zealand, New Zealand Sign Language was adopted as an official third language in 2002 and other auditory approaches that highlight visual components of spoken language are widely used. In line with international trends, however, AVT was used in the current study. Acoustic highlighting was extensively utilized. Therapy strategies used to improve the auditory stimulus included emphasis of suprasegmental speech characteristics, slightly slower speech patterns, reduced distance, and modification of background noise levels. For participants with CIs, the implanted ear was directed toward the therapist.

Principle Four: Provide Early, Intense (Re)habilitation

In New Zealand, it has been difficult to adhere to this principle as universal newborn hearing screening is only now being introduced. Some hearing loss is detected early through high-risk registers; however, in general, the age of hearing loss detection has been high. Because this AVT principle has

not consistently been met in New Zealand, the current investigation aimed to determine whether therapy based on listening and spoken language techniques could positively affect the outcomes of children diagnosed at an older age with hearing loss.

Principle Five: Adopt a Family-Centered Approach

Parents/guardians were involved in every session. Parents or guardians observed, wrote notes, took turns with the activities, asked questions, and listened to comments from the therapist. The sessions were tailored to suit the specific needs of each family and parent constellation. Cultural variations were also considered. A variety of family groupings existed among participants, and the therapist worked closely with each family to ensure key family members were involved.

Principle Six: Teach Language and Speech through Listening

By teaching language and speech through listening, the therapist utilizes the potential of the hearing instruments to maximum effect. The powerful hearing aid of 2 participants and the CIs of the other 5 participants all enabled auditory access across the speech spectrum. As noted above, short distance, low background noise, pitch variation, and slowed speech were used to optimize the auditory signal provided by the therapist. In addition, auditory information was repeated several times to aid recall of sounds, words, phrases, and sentences.

Principle Seven: Integrate Listening into Every Aspect of Life

All except one of the participants used listening and spoken language as their sole means of communication. The oldest participant is also competent in New Zealand Sign Language, which she uses when socializing with other individuals who are deaf and use sign language. The integration of listening into everyday life was reinforced by participants' involvement in one or more of the following activities: jazz, ballet, Polynesian cultural music activities, piano, school choir, and Māori language learning.

Principle Eight: Promote Placement in Mainstream Classrooms

All participants attend school in their local district school and are fully integrated into mainstream classes.

Assessment Areas Targeted by the Therapy Approach

Skills specific to each participant were targeted within each session, using a designated topic for context. Phonological, auditory, and language goals were

specified. For receptive language, these ranged from one-item, auditory memory closed-set language goals to open-set conversational turns for the more advanced participants. The complexity of tasks was based on participants' ability to retain and process auditory information. Actual items could be "Get the bowl," advancing to "Find the container that is used for mixing." Within expressive language goals there were targets for syntax, vocabulary, and verb forms. For example, using the same general topic of baking, target goals could range from simple to complex syntactical structures: "The girl is baking," to "The youngest girl is baking small chocolate muffins." Vocabulary items could range from "knife" to "muffin tins." Verb forms included variations from "She baked the cakes," to "Although she hasn't baked the cakes yet, she will soon."

Phonological development and the continued inappropriate use of specific processes were targeted for each participant. For Participant 1, the process of nasalization significantly reduced overall intelligibility. The nasalization affected the fricatives /s/, /z/, and /sh/ as well as the glide /l/. Many vowels and diphthongs were produced with a nasal overlay. Techniques included reinforcement of diaphragmatic breathing, improved breath control, and increasing the amount of intra-oral air. Articulation was also targeted. Each participant exhibited a number of unexpected idiosyncratic articulation errors that did not fit into the pattern of a phonological process.

Reading was not a targeted activity during sessions, although reading was evaluated to determine whether therapy affected literacy skill development. Speech perception in noise was also not specifically addressed in individual sessions. Since therapy was focused on listening, however, it was hypothesized that therapy would generalize to listening in noise.

Results

Tables 3 and 4 summarize pre- and posttherapy scores. Table 5 shows individual participants' pre- and posttherapy results for speech perception and production, language, and reading measures. There was considerable individual variability across all measures. CELF-4 language and WIAT-II reading scores are presented as standard scores (mean is either 100 [SD 15], or 10 [SD 3] for some subtests). For these standardized measures, each child is compared to similar-aged peers who have typical hearing. Other tests have percent error (HAPP-3, NZAT) or percent correct scores (speech recognition). For HAPP-3 and NZAT, reductions in error scores reflect improvement over time; for other tests, increased scores reflect improvement. Pre- versus posttherapy scores were compared using nonparametric Wilcoxon Matched Pairs Tests. Nonparametric testing was performed to determine statistical significance because there were a small number of participants. Because of the small sample size and large standard deviations, negative findings may reflect a lack of statistical power. There were a number of statistically significant findings that are of interest, however.

Table 3. Means and standard deviations (in parentheses) for measures with scores for most (N = 6 or 7) participants. Scores that improved significantly ($p < .05$) after therapy are shown in bold. CELF-4 standard scores have a mean of 10 and SD of 3. WIATT-II standard scores have a mean of 100 and SD of 15

	Type of measure	Subtest/score	N	Pre-therapy	Post-therapy
CELF-4	Receptive language	Concepts & Following Directions	6	7.17 (6.05)	9.33 (3.98)
		Understanding Spoken Paragraphs	7	7.29 (3.64)	10.14 (4.88)
		Word Classes-Receptive	7	8.57 (4.31)	9.00 (2.94)
	Expressive language	Word Classes-Expressive	7	8.57 (4.12)	8.86 (4.38)
		Formulated Sentences	7	7.57 (4.43)	7.86 (4.41)
		Recalling Sentences	7	7.29 (3.77)	7.43 (4.04)
	Core language		7	85.57 (28.37)	90.43 (25.32)
HAPP-3	Phonological processing	Error score	7	37.29 (27.95)	15.88 (12.21)
NZAT	Articulation	Word error score	7	22.14 (13.43)	11.14 (6.53)
WIAT-II	Reading	Word Reading	7	83.29 (20.80)	85.29 (24.12)
		Pseudoword Decoding	7	83.71 (18.09)	87.43 (17.24)
		Reading	7	91.71 (24.34)	94.58 (39.69)
		Comprehension			
Word recognition	Speech perception	Easy words	7	37.5% (16.3)	45.7% (13.0)
		Hard words	7	20.8% (14.9)	47.6% (13.0)

Table 4. Percentage of participants with scores that are more than 1 standard deviation below the mean (> 1 SD) for measures with standard scores. A reduction in the percentage indicates that more participants fell within the typical range after therapy. Scores that improved significantly ($p < .05$) after therapy are shown in bold

	Type of measure	Subtest/score	N	Pre-therapy	Post-therapy
CELF-4	Receptive language	Concepts & Following Directions	6	67%	17%
		Understanding Spoken Paragraphs	7	43%	14%
		Word Classes-Receptive	7	29%	14%
	Expressive language	Word Classes-Expressive	7	14%	29%
		Formulated Sentences	7	29%	29%
		Recalling Sentences	7	43%	29%
	Core language		7	29%	29%
WIAT-II	Reading	Word Reading	7	29%	14%
		Pseudoword Decoding	7	43%	14%
		Reading	7	29%	29%
		Comprehension			

Table 5. Individual pre- (bolded) and posttherapy scores for the 7 participants for each assessment. Speech perception scores for easy and hard words and NZAT scores are percent correct scores. HAPP-3 speech productions scores are percent error scores. There were three receptive language subtests (Concepts and Following Directions, Understanding Spoken Paragraphs, Word Classes-Receptive) and three expressive language subtests (Word Classes-Expressive, Formulated Sentences, Recalling Sentences). Core Language scores are a combination of expressive and receptive results for the four subtests described by the test authors as “most discriminating” at each age level. The final column contains Reading Composite scores, which are the sum of Word Reading, Reading Comprehension, and Pseudoword Decoding WIATT-II reading subtest scores

#	Age (yrs)	Easy words (%)	Hard words (%)	HAPP-3 error score (%)	NZAT (%)	Concept & follow direc	Under spoken para	Word classes Rec	Word classes Exp	Form sent	Rec sent	Core lang	Word read	Read compr	Pseud decod	Read comp
1	17.6	46.7	20	74.1	35	NA	6	6	9	7	9	99	82	113	90	285
		53.3	53.3	10.3	19	NA	7	7	10	7	9	103	86	123	81	290
2	8.8	26.7	13.3	44.5	27	1	1	1	1	1	1	41	45	56	65	143
		33.3	33.3	26	21	8	1	11	1	1	0	53	40	40	62	142
3	5.4	40.0	26.7	51.2	13	4	7	12	8	8	9	79	116	62	68	246
		46.7	46.7	37.2	7	6	13	10	9	10	10	94	86	100	89	275
4	10.3	13.3	20.0	63.8	37	4	6	7	8	4	5	70	84	87	68	239
		40.1	33.3	19	20	8	12	4	6	5	6	68	89	99	82	270
5	10.8	33.3	6.7	7.2	10	12	12	10	10	8	6	91	77	97	86	260
		60	66.7	9.3	7	10	12	8	10	10	7	96	86	104	88	278
6	9.4	46.7	46.7	10.6	12	5	8	10	9	10	8	85	86	77	75	238
		26.7	60	3.73	7	7	10	10	11	7	7	87	87	53	98	238
7	6.7	66.7	33.3	9.6	21	17	11	14	15	15	13	134	114	130	119	363
		60	40	5.6	13	17	16	13	15	15	13	132	123	156	118	397

NA = not applicable

Individual pretherapy CELF-4 language scores displayed in Tables 5 and 6 show variation across participants and across subtests. "Recalling Sentences" was difficult for participants 2, 4, and 5. "Understanding Spoken Passages" was difficult for participants 1 and 2 in both pre- and posttherapy. Participants 3 and 4 also had difficulty on the "Understanding Spoken Paragraphs" pretherapy, but their scores were within the normal range during posttherapy. Participant 2, who has very severe language difficulties, fell below the 1st percentile for all CELF-4 measures. The Core Language score is a measure of overall language performance indicative of the presence or absence of a language delay/disorder. Four of the 7 participants had a language delay at the start of therapy based on this measure (score 1 standard deviation or more below the normative mean).

Pre- versus posttherapy scores were compared statistically for all subtests that were age appropriate for all participants, and hence yielded complete data. There was a significant improvement in scores for CELF-4 "Understanding Spoken Paragraphs" subtest ($p = .043$). As Table 3 shows, scores improved on average by just over two standard score units. Table 4 lists the percentage of children falling more than 1 standard deviation (SD) below the mean on the assessments with standard scores. This percentage drops from 43% to 14% for "Understanding Spoken Paragraphs," the subtest that showed a significant improvement in mean scores. One other receptive language measure,

Table 6. Individual pre- and posttherapy CELF-4 subtest standard scores for the 7 participants, for the subtests producing standard scores for all participants. Scores for Recalling Sentences and Understanding Spoken Paragraphs fell outside the normative range (mean standard score of 10 minus 1 SD of 3) for several children. Participant 2 performed very poorly across all subtests

		<i>Recalling sentences</i>	<i>Formulated sentences</i>	<i>Word Classes- Receptive</i>	<i>Word Classes- Expressive</i>	<i>Word Classes- Total</i>	<i>Understanding Spoken Paragraphs</i>	<i>CORE</i>
1	Pre	9	7	6	9	14	6	99
	Post	9	7	7	10	16	7	103
2	Pre	1	1	1	1	1	1	41
	Post	0	1	1	1	1	1	53
3	Pre	9	8	12	8	14	7	79
	Post	10	10	10	9	15	13	94
4	Pre	5	4	7	8	15	6	70
	Post	6	5	4	6	5	12	68
5	Pre	6	8	10	10	9	12	91
	Post	7	10	8	10	10	12	96
6	Pre	8	10	10	9	7	8	85
	Post	7	7	10	11	10	10	87
7	Pre	13	15	14	15	19	11	134
	Post	13	15	13	15	15	16	132

"Concepts and Following Directions," also showed a substantial reduction in the number of children falling below the mean by more than 1 SD after therapy (from 67% to 17%), but the improvement in mean scores was not statistically significant. This may be due to the substantial variability in pretherapy scores.

Phonology

Pretherapy HAPP-3 phonological error scores were particularly variable, ranging from 7% to 74% (mean = 37%; SD = 28%). All participants showed a pattern of sound substitutions or delay in developing mature phonological patterns on the HAPP-3. These were analyzed as phonological deviations, sound substitutions, or consonant category deficiencies. HAPP-3 error scores improved significantly after therapy ($p = .028$). As Table 3 shows, error scores more than halved compared with baseline scores.

Articulation

Articulation errors measured using the NZAT also dropped significantly after therapy ($p = .018$). Although a significant improvement was found, the NZAT was not an optimal test for the study participants since it does not adequately describe changes in intelligibility that occurred. For example, one participant had a number of errors due to her process of gliding /r/ → /w/. This has a minimal impact on intelligibility; however, the NZAT has multiple words containing /r/ (*green, brush, frog, crab*, etc.), and therefore her error score was high. Other participants with similar scores had a variety of errors. Another limitation of the NZAT is the use of single words rather than connected speech. Participant 2, whose connected speech was very difficult to understand, had relatively few errors on the NZAT. The test also does not measure nasal resonance, which affected intelligibility of some participants.

Reading

Pretherapy pseudoword reading standard scores ranged from 65 to 119 (mean = 84; SD = 18), and word reading standard scores were 45–114 (mean = 83; SD = 21). Thus, there was a very wide range of reading abilities among participants. Pretherapy reading comprehension scores fell well outside the normal range at both extremes, with participant 2 performing very poorly and participant 7 well above the normal range (56–130; mean = 92; SD = 24). All reading scores improved after therapy, on average, by a few standard score units, but there was still a wide range of scores and no statistically significant changes were seen for the group.

Speech Perception in Noise

Table 3 includes average pre- and posttherapy speech recognition percent correct scores measured using the "easy" and "hard" words taken from the LNT.

Figure. Pretherapy versus posttherapy mean present correct speech perception (word) scores for the “easy” and “hard” words taken from the LNT. Scores show the anticipated advantage for easy words pretherapy but this difference disappears after therapy due to a significant improvement in scores for the lexically hard words. Error bars indicate 95% confidence intervals for the means.

Speech recognition scores improved an average of 15%, from 31.4% (SD 13.3%, median 33.4%) to 46.7% (SD 10.2%, median 46.7%). This difference was statistically significant ($p = .046$). As the Figure shows, the greatest improvements in speech scores occurred for the hard words (planned comparisons, $p = .018$).

Profiles of Individual Participants

Participant 1 was 17 years old and has a congenital bilateral profound hearing loss. She received a CI at the age of 5. Posttest measures for both receptive and expressive skills on the CELF-4 show small gains over pretest measures for word definitions and understanding spoken paragraphs. This demonstrates a link between receptive and expressive language skills. This participant progressed from a moderate phonological disorder based on the HAPP-3 in the pretest measures, to a mild disorder in the posttest. This was primarily due to eliminating the processes of nasalization, consonant reduction, and prevocalic

voicing, which had significantly impaired her intelligibility. The NZAT also showed gains in articulation, primarily as a result of correcting /s/ omission and /l/ substituted as /n/ speech errors. There was also a small change in reading. Overall, her reading skills were at an acceptable level prior to therapy, and she had completed high school with satisfactory results. Speech perception scores also improved considerably.

Participant 2 was 10 years old and received a CI at age 1 after contracting meningitis at the age of 8 months. He experienced kidney failure, resulting in several years of hospitalization, dialysis, and an organ transplant. Receptive and expressive skills on the CELF-4 show no gains over the pretest measures. His chronological age is well above his language age, so the gains made were not observable on age-appropriate standardized assessments. A language sample shows an increase in vocabulary, verbs, and a small number of two-word utterances. His family and teachers report that he is speaking much more than in the past. HAPP-3 and NZAT scores show no gains in phonological development or articulation, although speech perception scores improved. This participant speaks clearly at the single-word level. However, minor speech errors, such as /th/ replaced by /f/, reduced his scores significantly. Reading scores were very poor and did not improve.

Participant 3 was 6 years old and received bilateral hearing aids at the age of 2. Posttest CELF-4 expressive skills showed gains over pretest measures, particularly for grammatical structures such as plurals, possessives, contractions, and the perfect present tense. Receptive scores did not show gains, which is consistent with the focus of therapy on expressive language goals. However, HAPP-3 phonological scores did show gains. This participant progressed from a moderate phonological delay to a mild delay. The processes of cluster reduction and stopping no longer occur at the single-word level during testing and are generally absent in spontaneous speech. NZAT articulation scores improved owing to appropriate production of /s/ in the final position of words and in consonant clusters. WIATT-II reading scores were unchanged overall. Speech perception scores improved, particularly for hard words.

Participant 4 was 11 years old and received bilateral hearing aids at the age of 3. Both receptive and expressive CELF-4 language scores showed gains. Receptive scores improved for the "Concepts & Following Directions" and "Understanding Spoken Paragraphs" subtests. Expressive scores improved for "Recalling and Formulating Sentences." Phonological development progressed from a moderate delay to a mild delay after therapy. After therapy, the processes of cluster reduction and final consonant deletion occurred only rarely at the single-word level and were generally absent from spontaneous speech. NZAT articulation scores improved owing to appropriate production of /s/ in word final position and in consonant clusters. Reading and speech perception also improved, but there were only modest gains in word reading. After completing the therapy intervention, this participant was re-referred to the CI program and subsequently received a CI.

Participant 5 was 10 years old and has a congenital, bilateral profound hearing loss. He received a CI at the age of 4. Receptive and expressive language scores improved for recalling sentences, receptive word classes, understanding spoken paragraphs, and formulated sentences. Although his HAPP-3 phonological scores indicated mild delays both pre- and posttherapy, there was a reduction in errors. He had acceptable speech clarity pretherapy and hence there was no change in his NZAT articulation score. Reading improved slightly and speech perception scores improved substantially.

Participant 6 was 10 years old and received her first CI at age 2 and her second CI at age 8. Receptive and expressive language showed some gains for “Concepts & Following Directions” and “Word Definitions” subtests. Her HAPP-3 scores indicated a mild delay both pre- and posttherapy, but there were some improvements. She had occasional errors with triple blends but these did not significantly impact intelligibility. In spontaneous conversation, she was easy to understand. NZAT showed some articulation gains but minor errors still occurred, with /th/ produced as /f/ and /str/ produced as /shtr/. Overall, her reading scores were unchanged and speech perception scores improved, but only slightly.

Participant 7 was 8 years old and received her first CI at age 2 and her second CI at age 4. Receptive and expressive skills showed gains in all areas. She was in an accelerated class at her school and her CELF-4 results indicated strong language learning abilities, consistent with her other academic abilities. It was easy for listeners to understand her in spontaneous conversation. HAPP-3 phonological scores indicated a mild delay pre- and posttherapy. Specific error scores showed some improvement. NZAT scores improved, owing mainly to the correct production of /th/, produced as /f/ in the pretest condition. The remaining error of /r/ produced as /w/ resulted in a number of errors posttherapy in blends such as /gr/ as well as the singleton /r/. Scores improved for word reading and reading comprehension subtests. This participant did not show improvements in speech perception.

Discussion

Therapy Outcomes

Overall, the results of this exploratory study support the notion that SLT with an emphasis on auditory-verbal techniques does improve outcomes for children with hearing loss in the areas of receptive language, phonological development, articulation, and listening in noise. Twenty sessions of SLT were associated with an improvement in participants’ speech production and understanding of language. For the assessments that showed improvements after therapy, the greatest changes did not occur for the youngest children and hence there is no clear link to age of intervention for this small sample. This is not surprising given that the participants were

generally identified late and were all school-aged when they participated in the study.

Studies of children whose hearing loss was identified early by newborn hearing screening show that the gap between typical language development and the language development of a child with hearing loss can widen over time, particularly for children with greater degrees of hearing loss (Yoshinaga-Itano, 2006) as language skills become increasingly sophisticated in children who have typical hearing. It is important, therefore, that standardized measures be repeated to determine whether children with hearing loss are keeping up with their peers. A study by Kiese-Himmel (2008) showed that children with profound hearing loss slipped behind their peers who have typical hearing for receptive vocabulary as they were followed longitudinally. With universal newborn hearing screening and early intervention, we anticipate that language outcomes will improve for all children with hearing loss. However, as noted by Jiménez, Pino, and Herruzo (2009), it is not yet clear how early-identified children with hearing loss will fare as they progress through school and require more advanced language, such as the metalinguistic skills that enable us to understand metaphors.

There were statistically significant gains in the understanding of spoken language. This appears to be a result of the therapy sessions that focused on the development of vocabulary (nouns, verbs, prepositions) and the understanding of synonyms, such as “cold-chilly,” “moist-wet,” and “big-large.” Language that the participants had previously understood only in its contextual situation was really only understood when supported by visual and environmental information. One major component of therapy sessions was to develop a genuine understanding of the language used to describe concepts such as “before,” “after,” and “between.” Significant gains in expressive language were not seen on standardized testing, suggesting a need to modify the therapy approach and/or assessment tools if improved expressive language is a therapy goal. Alternatively, this negative finding may have been caused by a lack of statistical power due to the small sample size and large standard deviations, or the time frame for therapy and assessment may have been too short. Interestingly, Law, Garrett, and Nye’s (2003) systematic review of the literature indicate that, for children without hearing loss who have speech and language delays, gains in expressive language are more common than gains in receptive language after intervention.

Vocabulary

Vocabulary was not formally measured as part of the study, but anecdotally it appeared that vocabulary improved with therapy for all participants. Changes in vocabulary were most evident in participants 2 and 7 whose pre-therapy measurements were at the two extremes of performance (below the 1st and above the 99th percentile, respectively). Participant 7 made gains in

her vocabulary scores by working on sophisticated synonyms such as “fragile-delicate,” “palace-castle,” and “ornament-decoration.” Figurative language was also used with expressions such as “A clean bill of health” and “A one dollar bill.” Participant 2, whose CELF-4 language scores were below the 1st percentile both pre- and posttherapy, had language goals incorporating both understanding and then use of words such as “tree, grass, sun” and “up, down, in,” and verbs such as “fall, blow, shine” (for the autumn theme). Participant 2 received his CI relatively early, with activation at age 12 months. After therapy, participant 2 showed progress on informal language measures; these gains were not detected by the CELF-4. His expressive vocabulary increased from approximately 100 to 300 words, measured informally using the MacArthur-Bates Communicative Developmental Inventories (CDI) (Fenson et al., 1993). Thal, DesJardin, and Eisenberg (2007) reported excellent validity of the words and gestures, and words and sentences forms of the CDI for children with CIs. Participant 2 also made gains from using two-word phrases such as “Boy up” and “Me go,” to “Boy up tree” and “Me going home.” In receptive language areas, he made gains understanding simple phrases such as “What’s the name of your sister/brother?” versus “What’s your name?” and “How are you?” versus “How old are you?”. Language sampling that can capture these gains may be a useful addition to future studies.

Participant Selection

Although not linked to the study goals, an important, unanticipated outcome of this research was that one potential participant, initially recruited into the study, performed so poorly on the baseline assessments with her hearing aids that she was referred to the CI program and subsequently received a CI. She performed poorly for all the subtests of receptive and expressive language, phonological development, speech perception in noise, and reading comprehension. This was despite her consistent use of hearing aids, good academic input, and significant support from her mother. In the previous 5 years, this child had been referred twice for CI evaluation. On both occasions, an implant was not approved because she was “performing too well.” The CI team did not detect this child’s difficulties, in part because of her use of a variety of compensatory strategies, strong cognitive abilities, skilled speech reading, and determination to appear “just like everybody else.” This case highlights one of the benefits of performing standardized tests in a wide range of different areas and continuing to follow children with hearing loss who are in mainstream schools.

Choice of Assessment Materials

This exploratory study highlighted the difficulty in finding assessment materials that are applicable across all participants with a range of abilities.

Few measures are both sensitive enough to use with participants who have severe delays in their language development and also broad enough to measure more sophisticated and mature aspects of language, such as double meanings, inference, and implication. In the current study, pre- and posttherapy language assessments incorporated subtests that depended on the chronological age of each participant. Only a small number of CELF-4 subtests occurred across the age bands so only those subtests could be statistically analyzed. Several areas of language showed improvements for individuals but could not be included in the statistical analysis because data was not available for all participants. The challenge is to identify language assessment tools suitable for a wide range of abilities. One option may be to use a standardized test such as the CELF-4 to determine the level of language delay relative to typical development, and to use other tools, such as language sampling and vocabulary measures, to assess progress in therapy. However, the variation in performance across CELF-4 subtests highlights the importance of looking comprehensively at language abilities, and hence it is likely that several measures are needed to determine language outcomes.

Achieving Cognitive Potential

The language of children who have worn a CI for 5 to 6 years may appear to be age appropriate, particularly if they have good speech intelligibility. However, if the language is assessed in depth, it is clear that the appearance of adequate language is superficial in some cases. With the exception of 1 child, all participants showed some degree of language delay. Participant 7 had language levels significantly above those expected for her chronological age, but this child has been described as “intellectually gifted” by the school. The challenge for this participant and her family will be to ensure that language levels remain commensurate with overall cognitive skills. Negative effects on quality of life have been reported for children with hearing loss (Wake, Hughes, Poulakis, Collins, & Rickards, 2004). Presumably, these negative effects would be greater for children who are not able to reach their cognitive potential because of their hearing loss.

Articulation and Phonology

The articulation skills of all participants improved during the intervention. This was particularly encouraging as parents often report “clear speech” that is easy to understand to be the most important goal of intervention. Therapy sessions targeted phonological processes that yielded the highest impact. The targets varied among the participants, although for several participants the focus was on production of /s/. Although the CI allows the wearer to access the /s/ sound acoustically, the phoneme /s/

was often omitted or inaccurate in the spontaneous speech of several participants. Intelligibility improved with systematic therapy targeting errors in speech production. By selecting “high-impact” processes that were most destructive of speech intelligibility, the effect of remediation was probably more dramatic than if other processes had been selected. This approach is based on Grunwell’s (1992) principles of treatment planning for phonological therapy.

Using AVT techniques, participants’ speech production improved. Therapy moved in progressive steps from isolated sounds to word and phrase level. Although individual participants made gains, generalization to spontaneous conversation had not occurred for all children at the end of 20 weeks of therapy. Hypernasal speech and nasalization of many consonants was a significant problem for the oldest participant (participant 1). This participant used sign language as a young child and did not receive a CI until the age of 5. At that time, she was only able to produce four recognizable words and a series of nasalized vowels. Minimizing the process of nasalization during therapy resulted in a significant improvement in phoneme accuracy and intelligibility for this participant.

Voicing and nasalization errors were noted in two of the older participants (participants 1 and 6), including hypernasal resonance, prevocalic voicing of some plosive consonants, especially /p/→/b/ and /t/→/d/, and nasalization of the /l/ phoneme. Nasal resonance as an overlay on all vowels was particularly noticeable for participant 1. Therapy for this participant was highly focused on increasing oral resonance. Techniques included teaching the participant to understand how the soft palate functions to shut off the nose, and direct observation of the therapist’s soft palate moving up and backward during production of “ah” breathing activities to improve diaphragmatic breathing and reduce shallow clavicular breathing. This led to progressively less intense physical occlusion of the nose during production of elongated vowel sounds such as /ah/. In these activities, both the therapist and the participant would occlude their own nostrils to start the sound, such as /ah/ or /s/. The fingertip pressure was progressively reduced while the sound was maintained with the same level of oral resonance, and this was transferred from sounds to words and then phrases. Another technique was tactile awareness of intra-oral air pressure. By placing the back of the hand in front of the mouth during production of the voiceless plosive sounds such as /p/ and /t/, participants 1 and 6 built up an awareness of intra-oral air pressure. This awareness was then linked to listening activities so that they could hear the difference in sounds produced with appropriate intra-oral air pressure and those that lacked the appropriate levels. The speech errors observed in the current study are consistent with Peng, Weiss, Cheung, and Lin’s (2004) report that 6- to 12-year-old CI users with approximately 2 to 6 years of CI experience had most errors for nasals, affricates, fricatives, and the lateral approximant /l/, and fewest errors for plosives.

Speech Perception in Noise

One unexpected finding was that 6 out of 7 participants showed improvements on the speech perception in noise test. This can probably be accounted for as an effect of improved auditory discrimination. One focus of AVT is listening to speech, and all the participants were aware that “concentrating” when they listened improved their ability to understand what others were saying. When working on audition and receptive language goals in the weekly sessions, auditory discrimination was taught by having participants listen for words such as “path/bath” or “multiple/multiply,” or pointing to pictures that may illustrate “I will catch it” versus “I will cash it.”

Speech recognition was evaluated in noise at a +5 dB SNR. This SNR is better than what can occur in classrooms containing approximately 30 children and other noise sources. In New Zealand classrooms, the SNR is typically less than 0 dB (i.e., noise is louder than speech signal) (Valentine et al., 2000). Although speech perception scores were better posttherapy, they were still quite poor, highlighting the need for these children to use additional technology to support listening in the classroom, and also highlighting the need to assess speech perception in noise as well as quiet. The words from the LNT were re-recorded using a New Zealand speaker, and hence it is not possible to compare results directly with published data for the original North American version of the test. Also, the LNT is typically administered in quiet settings rather than in noisy settings. Results in noise have greater relevance to classroom listening, since busy classrooms are typically noisy (and reverberant).

Reading

Although reading did not change significantly, the scores for pseudoword recognition showed some improvement. This is likely due to the following phonics-based activities that occurred during some sessions:

- Visual reinforcement of acoustic differences in different letters, e.g., “When you see the letter ‘b’ in the word ‘book,’ the ‘b’ letter is telling you to make the sound /b/. Here are some other words with the same letter ‘b’ at the beginning of the word. They are *bag*, *ball*, *bath*. Now look at these words that are almost the same. They have a different letter at the beginning. They start with the letter ‘s.’ The letter ‘s’ makes the sound /s/.”
- Establishing a strong sound-letter association for consonants, e.g., “I have a box of toys and a box of labels. I will say a word and you match the word to the toy: *rope*, *soap*, *ring*, *sing*, *run*, *sun*. Now, you’re the teacher. You tell me the word. I have to listen really carefully and then you check to see if I am right.”

- Establishing sound-letter associations for some vowels as they incidentally occurred during activities with another focus. A similar role-play was used, but words varied by vowels such as *book, bake, look, lake*.

The ability to manipulate speech sounds in words (phonological awareness) is regarded as a key ability underlying reading success. Hence, reading difficulties in children with hearing loss are not surprising if they cannot easily discriminate speech sounds in words. Easterbrooks, Lederberg, Miller, Bergeron, and Connor (2008) found that generally, 5-year-old children with hearing loss lagged behind their peers in phonological awareness, and that phonological awareness correlated with literacy skill development. Spencer and Oleson (2008) examined the link between early speaking and listening in 72 CI users and found that a large proportion of the variance (59%) in reading ability was accounted for by speech perception and speech production abilities measured approximately 4 years earlier. This highlights the need to look comprehensively at speech perception (ideally in listening conditions that simulate real classroom conditions), speech production, and literacy, as well as receptive and expressive language, when evaluating outcomes for children with hearing loss.

Limitations and Social Validity

This pilot study does not examine the social validity of AVT for school-aged children with hearing loss. As noted by Eriks-Brophy (2004), the social development of such children has been largely ignored. Future research should examine the relationship between communicative competence, speech intelligibility, and academic achievement and the social acceptance of children with hearing loss in mainstream settings. Concepts such as same gender and opposite gender peer acceptance would be valuable areas for consideration. Improvements in communication outcomes should ultimately lead to a reduction in the high incidence of unemployment and underemployment currently experienced by individuals with severe-to-profound hearing loss.

There is a general lack of clear research findings to support the efficacy of listening and spoken language therapies. This is due to a number of factors, many of which are described by Eriks-Brophy (2004). Participant groups tend to be very small and to be heterogeneous with regard to the age of diagnosis and the type and length of intervention, which reduces the ability to generalize outcomes from one group to another. Often reports in the literature are anecdotal or retrospective and do not use comparison groups. In addition, there is diversity between the therapies and therapy settings (hospital, school, or home setting).

Participants in this study were older than the typical age at which children with hearing loss first enter an AVT program. This is due to several factors. First, New Zealand's average age of detection for hearing loss is high

due to the delayed implementation of universal newborn hearing screening. Additionally, due to funding and other restrictions, there is a cohort of New Zealand children who did not meet the very strict criteria for cochlear implantation in place prior to 2005. At that time, no more than 25 CI surgeries were funded each year for the entire population, including adults with acquired hearing loss. The acceptance criteria for CIs have broadened and there has been some increase in funding for devices, but government-funded (re)habilitation for school-aged children is still not widely available.

Conclusion

Overall, this exploratory study has shown that SLT with a listening and spoken language focus shows promise as a successful form of intervention for children with hearing loss. The results highlight the need for children with hearing loss, including those with CIs, to have ongoing (re)habilitation, not just assistance in the first few years. Although the initial few years of a child's life are crucial for establishing the fundamentals of language, subsequent years are crucial for establishing the sophisticated, subtle nuances of expressive and receptive language. Ongoing support is required to avoid a widening gap between the listening and spoken language skills of children with typical hearing and those with hearing loss, and to ensure that children with hearing loss do not fall behind in literacy skill acquisition. How to achieve this in an efficient, cost-effective way for all children with hearing loss remains a challenge.

Future Research

The pretherapy results were very variable, and hence a "one-size-fits-all approach" is not appropriate for children with hearing loss. This highlights a problem for randomized controlled trial research designs in this area. Although not as robust in terms of level of evidence, a case study control design, in which the participants serve as their own control prior to implementation of therapy, is preferable due to the difficulty in obtaining matched groups of participants. For language, the greatest improvements were seen in two areas of receptive language, which is consistent with the goals of listening and spoken language therapy. Anecdotally, there appeared to be gains in other aspects of language that were not identified using the CELF-4 assessment tool. Wider ranging and more sensitive outcome measures may be required. Hence, future studies should include formalized language sampling and standardized receptive and expressive vocabulary measures in addition to the CELF-4. Observation of the participants' speech production during therapy highlighted the need for assessment and therapy for voice and nasalization errors. This may be particularly helpful for children who receive their CIs at a later-than-optimum age. Future studies should

include assessment of these speech sound errors as an additional outcome measure.

References

- Caleffe-Schenck, N. (1992). The auditory-verbal method: Description of a training program for audiologists, speech language pathologists, and teachers of children with hearing loss. *The Volta Review*, 94, 65–68.
- Connor, C.M., Craig, H.K., Raudenbush, S.W., Heavner, K., & Zwolan, T.A. (2006). The age at which young deaf children receive cochlear implants and their vocabulary and speech-production growth: Is there an added value for early implantation? *Ear and Hearing*, 27, 628–644.
- Estabrooks, W. (n.d.). *COMENIUS 2.1 Action. Qualification of educational staff working with hearing-impaired children (QESWHIC). Study letter 7 auditory-verbal practice*. Retrieved October 18, 2010, from <http://www.qeswhic.eu/downloads/letter07en.pdf>.
- Easterbrooks, S.R., Lederberg, A.R., Miller, E.M., Bergeron, J.P., & Connor, C.M. (2008). Emergent literacy skills during early childhood in children with hearing loss: Strengths and weaknesses. *The Volta Review*, 108, 91–114.
- Eisenberg, L.S., Martinez, A.S., Holowecky, S.R., & Pogorelsky, S. (2002). Recognition of lexically controlled words and sentences in children with normal hearing and children with cochlear implants. *Ear and Hearing*, 23, 450–462.
- Eriks-Brophy, A. (2004). Outcomes of auditory-verbal therapy: A review of evidence and a call for action. *The Volta Review*, 104, 21–35.
- Fenson, L., Dale, P.S., Reznick, J.S., Thal, D., Bates, E., Hartung, J.P., et al. (1993). *The MacArthur-Bates communicative development inventories: User's guide and technical manual*. Baltimore, MD: Brookes.
- Goldberg, D.M., & Flexer, C. (1993). Outcome survey of auditory-verbal graduates. *Journal of the American Academy of Audiology*, 4, 189–200.
- Grunwell, P. (1992). Processes of phonological change in developmental speech disorders. *Clinical Linguistics & Phonetics*, 6, 101–122.
- Hodson, B.W. (2004). *HAPP-3 Hodson Assessment of Phonological Patterns* (3rd Ed.). Austin, TX: PRO-ED.
- Jiménez, M.S., Pino, M.J., & Herruzo, J. (2009). A comparative study of speech development between deaf children with cochlear implants who have been educated with spoken or spoken+sign language. *International Journal of Pediatric Otorhinolaryngology*, 73, 109–114.
- Kiese-Himmel, C. (2008). Receptive (aural) vocabulary development in children with permanent bilateral sensorineural hearing impairment. *The Journal of Laryngology & Otology*, 122, 458–465.
- Kirk, K., Pisoni, D., & Osberger, M. (1995). Lexical effects on spoken word recognition by pediatric cochlear implant users. *Ear and Hearing*, 16, 470–481.

- Law, J., Garrett, Z., & Nye, C. (2003). Speech and language therapy interventions for children with primary speech and language delay or disorder. *The Cochrane Database of Systematic Reviews*, 3, Art. No.: CD004110. doi: 10.1002/14651858.CD004110.
- Lim, S.Y., & Simser, J. (2005). Auditory-verbal therapy for children with hearing impairment. *Annals, Academy of Medicine, Singapore*, 34, 307–312.
- Ling, D. (1976). *Speech and the hearing-impaired child: Theory and practice*. Washington, DC: Alexander Graham Bell Association for the Deaf and Hard of Hearing.
- Moyle, J. (2005). The New Zealand Articulation Test Norms Project. *New Zealand Journal of Speech-Language Therapy*, 60, 61–75.
- Pearson PsychCorp. (2007). Wechsler Individual Achievement Test Second Edition Australian Standardised Edition (WIAT-II) Australian. Pearson Clinical and Talent Assessment, Sydney, 2002 Australian Adaptation; 2007 NCS Pearson Ltd.
- Peng, S.C., Weiss, A.L., Cheung, H., & Lin, Y.S. (2004). Consonant production and language skills in Mandarin-speaking children with cochlear implants. *Archives of Otolaryngology—Head & Neck Surgery*, 130, 592–597.
- Rhoades, E.A. (1982). Early intervention and development of communicative skills for deaf children using an auditory-verbal approach. *Topics in Language Disorders*, 2, 8–16.
- Rhoades, E.A. (2006). Research outcomes of auditory-verbal intervention: Is the approach justified? *Deafness & Education International*, 8, 125–143.
- Rhoades, E.A., & Chisholm T.H. (2000). Global language progress with an auditory-verbal approach for children who are deaf or hard of hearing. *The Volta Review*, 102, 5–25.
- Semel, E.M., Wiig, E.H., & Secord, W. (2003). *Clinical Evaluation of Language Fundamentals (CELF-4)*. San Antonio, TX: The Psychological Corporation.
- Spencer, L.J., & Oleson, J.J. (2008). Early listening and speaking skills predict later reading proficiency in pediatric cochlear implant users. *Ear and Hearing*, 29, 270–280.
- Thal, D., DesJardin J.L., & Eisenberg L.S. (2007). Validity of the MacArthur-Bates Communicative Development Inventories for measuring language abilities in children with cochlear implants. *American Journal of Speech-Language Pathology*, 16, 54–64.
- Valentine, J., Wilson, O., Halstead, M., Dodd, G., McGunnigle, K., Hellier, A., & Wood, J. (2000). Classroom acoustics – A New Zealand perspective. *Proceedings of the 15th Conference of New Zealand Acoustical Society*, pp. 127–142.
- van Asch Deaf Education Centre. (n.d.). *Verb activity sheets*. Retrieved November 1, 2010, from http://www.vanasch.school.nz/catalog.php?cat_id=222.
- Wake, M., Hughes, E.K., Poulakis, Z., Collins, C., & Rickards, F.W. (2004). Outcomes of children with mild-profound congenital hearing loss at 7 to 8 years: A population study. *Ear and Hearing*, 25, 1–8.

- Wray, D., Flexer, C., & Vaccaro, V. (1997). Classroom performance of children who are hearing impaired and who learned spoken communication through the auditory-verbal approach: An evaluation of treatment efficacy. *The Volta Review*, 99, 107–120.
- Yoshinaga-Itano, C. (2006). Early identification, communication modality, and the development of speech and spoken language skills: Patterns and considerations. In P.E. Spencer & M. Marschark (Eds.), *Advances in the spoken language development of deaf and hard-of-hearing children* (pp. 298–327), Oxford, UK: Oxford University Press.

Use of Differential Reinforcement to Increase Hearing Aid Compliance: A Preliminary Investigation

Sandie M. Bass-Ringdahl, Ph.D., CCC-A; Joel E. Ringdahl, Ph.D.; and Eric W. Boelter, Ph.D.

Compliance with hearing aid use can be difficult to achieve with children. This difficulty can be increased when a child presents with other disabilities, such as developmental delays. Behavioral treatments, including differential reinforcement, might be one strategy for increasing compliance by these children. In the clinical scenario discussed, the hearing aid compliance of a young child with Sanfilippo syndrome was increased by systematic application of differential reinforcement and escape extinction. Treatment appeared to be successful both in the clinic and during a 1-month follow-up visit to the child's educational placement.

Introduction

It is currently estimated that 3 to 4 per 1,000 children are born with permanent, congenital hearing loss each year (National Center for Hearing Assessment and Management, 2008), with the highest incidence in populations of children who are medically fragile. All of these children are considered potential hearing amplification/assistive device candidates, with the decision to use such devices based on family choice and progress made in communication development. Because hearing aids increase audibility or access to the speech signal, they are considered an essential component of the intervention process when the goal is to maximize listening and spoken language development (American Academy of Audiology, 2004; Ontario Ministry of Children

Sandie M. Bass-Ringdahl, Ph.D., CCC-A, is an Assistant Professor in the Department of Communication Sciences and Disorders at the University of Iowa. Joel E. Ringdahl, Ph.D., is an Assistant Professor at the University of Iowa. Eric W. Boelter, Ph.D., is a Clinical Psychologist at Seattle Children's Hospital in Seattle, WA. Correspondence concerning this article can be directed to Dr. Bass-Ringdahl at sandie-bass-ringdahl@uiowa.edu.

and Youth Services, 2007; The Pediatric Working Group, 1996). Audition and communication are intimately linked in terms of their effect on one another. At the earliest stages of speech development, lack of audibility has been linked to the delayed onset of canonical babble (a precursor to first word production) in children with all degrees of hearing loss (Eilers & Oller, 1994; Moeller, Tomblin, Yoshinaga-Itano, Connor, & Jerger, 2007; Nathani, Neal, Olds, Brill, & Oller, 2001; Oller & Eilers, 1988). Bass-Ringdahl (2010) provided additional evidence regarding the role of audibility as well as the role of amplification device use on the onset of canonical babble. Specifically, the participants in the Bass-Ringdahl study began canonical babbling only after achieving a minimum amount of listening experience with an audible speech signal.

Lack of audibility has also been linked to a delay in listening and spoken language development, decreased academic achievement, and behavioral concerns. Moeller et al. (2007) reviewed the current literature regarding delays typically noted for children who are deaf or hard of hearing. The results of their review indicated that children with hearing loss were at risk for delays in vocabulary, syntax, morphology, and social communication development. In addition, Davis, Elfenbein, Schum, and Bentler (1986) and Wake, Hughes, Poulakis, Collins, and Rikards (2004) provided evidence that the academic performance of children with hearing loss was well below that of children who have typical hearing. Davis et al. also reported on the psychosocial characteristics of 40 children with mild to moderate hearing loss using the Child Behavior Checklist (Achenback & Edelbrock, 1983). As a group, the children from this study were described by their parents as having behavior problems, including aggression, impulsivity, immaturity, and resistance to discipline and structure. Additional studies have investigated the root of increased behavior problems in children with hearing loss. Marshall (1997) hypothesized that receptive communication skills would be highly related to increased behavioral problems. One significant find from this study was that mothers' perception of poor mother-child communication accurately predicted children with increased behavior problems.

The impact of hearing loss and reduced audibility on speech, language, academic, and behavioral outcomes highlights the importance of hearing aid compliance for children who rely on hearing aids for communication development. One of the most comprehensive investigations of hearing aid compliance is from an ongoing, longitudinal study conducted by Moeller, Hoover, Peterson, and Stelmachowicz (2009). Families enrolled in this study were interviewed every 3 months concerning the consistency of amplification device use. Preliminary results suggested that few infants in the study actually wore their hearing aids during all waking hours. Moeller et al. (2009) speculated that factors including infant temperament, parenting skills, and parental understanding of the importance of amplification contributed to the inconsistency. The importance of achieving good hearing aid compliance is heightened for children who have concomitant developmental delays. Achieving hearing aid

compliance can be more difficult for these children because they might exhibit behavioral problems beyond those caused by the communication difficulties associated with the hearing loss, and likely also have receptive language delays that serve as obstacles. Given the (a) impact that the lack of audibility has on speech and language development, academic performance, and behavior; (b) apparent difficulty achieving compliance with consistent hearing aid use; and (c) additional obstacles faced by developmental disabilities, strategies are needed to achieve compliance with hearing aid use for children who are deaf or hard of hearing and have additional developmental disabilities.

One behavioral strategy that has been used to increase compliance with various academic, daily living, and other tasks is differential reinforcement (DR) of compliance (e.g., Ringdahl, et al., 2002). DR is a procedure in which the target response of compliance (e.g., completing the instruction in the absence of the target problem behavior) results in access to a reinforcer. Disruptive behavior, such as the removal of a hearing aid, results in either withholding the reinforcer (if the individual is not accessing the reinforcer at the time of the disruptive response) or removing the reinforcer (if the individual is accessing the reinforcer at the time of the disruptive response). In the current case study, a DR procedure was used to increase hearing aid compliance by a young child diagnosed with Sanfilippo syndrome. The treatment program was initially assessed in a clinic setting. Follow-up data were then obtained in the child's education placement.

Method

Participant

One individual took part in the evaluation. Tony was a 5-year-old boy diagnosed with bilateral, moderate sensorineural hearing loss, attention deficit hyperactivity disorder, and developmental delays secondary to Sanfilippo syndrome (MPS III). Sanfilippo syndrome is a disorder that affects long chains of sugar molecules used in the building of connective tissues in the body known as mucopolysaccharides. Children with this inherited recessive disorder are missing an enzyme essential to the breakdown of used mucopolysaccharides. The incompletely broken down mucopolysaccharides remain stored in the cells of the body, causing progressive cell damage (National MPS Society, 2008). The disease tends to progress through stages. The first stage occurs during the child's preschool years. The preschool-aged child with Sanfilippo syndrome tends to lag behind in development and displays overactive and difficult-to-manage behavior. As the disease progresses, the child shows extreme activity, restlessness, and difficult-to-manage behavior. Eventually the child's activity level slows and the child regresses in skill development. Ultimately, individuals with Sanfilippo syndrome typically live into their teenage years (National MPS Society, 2008).

Tony was referred to an intensive behavioral treatment program to address problem behavior that included attempts to elope (i.e., leave rooms) and flopping to the ground when required to complete a nonpreferred task. Of particular relevance to the current study was Tony's noncompliance with wearing his hearing aids. Parental reports, at the point of referral, indicated that Tony would not tolerate wearing his hearing aids for any appreciable amount of time. Attempts to have Tony wear his hearing aids had stopped in both the home and school settings for fear that the hearing aids would be damaged.

Tony was followed on a routine basis by a hospital-based pediatric audiologist as well as an educational audiologist to monitor his hearing sensitivity and hearing aid function and to remake his ear molds. The settings of Tony's hearing aids were monitored through the use of real ear measures to ensure that his hearing aid fit was appropriate for his degree of hearing loss.

Setting and Materials

Sessions were conducted in the therapy room of a clinic-based day treatment program that specialized in conducting behavioral assessment and treatment evaluations of challenging behavior exhibited by individuals with developmental disabilities. The session room was approximately 4 meters by 5 meters and contained a table and chairs, a workstation (desk and chair), and an array of preferred stimuli. Preferred stimuli were identified via a free operant preference assessment (Roane, Vollmer, Ringdahl, & Marcus, 1998) in which Tony had free access to an array of stimuli chosen by parental and staff reports of potential reinforcement items. Data were collected to determine which stimuli/toys he preferred based on his allocation of interaction among the stimulus array. A follow-up observation was conducted in Tony's classroom, a special education program placement at his local elementary school. His classroom was a resource room where he and other children with disabilities received educational services.

Response Definition

The dependent variable for the evaluation was "time in," defined as Tony keeping the relevant component (e.g., ear mold only or full unit, depending on the condition requirement) inserted in his ear. This response was recorded using a duration recording method (i.e., number of seconds) and was expressed as a percentage of total session time.

Data-Collection and Interobserver Agreement

During the clinic-based observations, data were recorded using laptop computers and customized behavioral data-collection software. The software allowed for data to be collected on the duration of the dependent variable

(i.e., "time in," in seconds) and summarized as a percentage of session time. Sessions were 10 minutes across all clinic-based conditions.

A second, independent observer collected data during 15% of the clinic-based sessions. Interobserver agreement (IOA) scores were calculated using an interval-by-interval comparison. Specifically, each session was divided into a series of 10-second intervals and the observers' records were compared. An agreement was scored if both observers indicated the hearing aid was in for any portion of the 10-second interval or if both observers indicated that the hearing aid was out for the entire 10-second interval. A disagreement was scored if one observer indicated the hearing aid was in for any portion of the interval, while the other indicated the hearing aid was out for the entire 10-second interval. Agreements were then summed and divided by the total number of intervals for an observation (i.e., 60). The mean agreement score was 97% (range: 87–100%).

During the follow-up observation in the school setting, data were collected using a pencil and paper, 6-second partial interval recording system. Compliance with wearing the hearing aid was scored if Tony kept the hearing aid in his ear for the entire 6-second interval. This method of data collection allowed for data to be summarized as a percentage of 6-second intervals. The alternative method was used in the school setting because the computer-based data collection was not portable (i.e., the computers had to stay in the clinical setting for use with the ongoing clinical services). Sessions were 10 minutes across all conditions. Due to personnel limitations, IOA could not be collected during the school visit.

Either doctoral-level graduate students or clinical therapists collected all data. Each observer had been trained on specific criteria in the use of computer-based and paper-based behavioral scoring. Criteria used to determine that a graduate student or clinical therapist was proficient in data collection included achieving IOA scores of greater than 90% for three consecutive sessions across two different patients relative to previously trained observers. Each observer had several years of experience in observing and scoring child and adult behavior relative to disruptive behavior, appropriate behavior, and compliance and noncompliance, and had met a predetermined criterion regarding their proficiency as behavioral data collectors as part of their orientation to the clinical service for which they were working.

Procedures

The treatment evaluation began with the collection of baseline data on the percentage of session time Tony would keep both hearing aids in his ears. A treatment was designed that consisted of DR and escape extinction (EE) (Iwata, Pace, Kalsher, Cowdery, & Cataldo, 1990). Recall that DR is a procedure in which the target response (e.g., compliance with wearing the hearing aid or ear mold in the absence of the target problem behavior) results in access

to a reinforcer whereas responses consisting of disruptive behavior (e.g., the removal of a hearing aid or ear mold) result in withholding the reinforcer or removal of the reinforcer. EE refers to the inability to escape the stimulus (e.g., the immediate reinsertion of the hearing aid or ear mold).

The combined treatment of DR and EE was pursued for two reasons. First, reinforcement-based procedures are often more effective than treatment strategies that do not include a reinforcement component (e.g., timeout or response cost), and are the typical first line of treatment when attempting to address behavioral concerns (Pelios, Morren, Tesch, & Axelrod, 1999). Second, it was hypothesized that hearing aids were a negative reinforcer. Thus, noncompliance with wearing the hearing aid was likely maintained by escape from the hearing aids. An EE component (described in further detail below) was implemented to address this potential response-reinforcer relationship.

Baseline. Both hearing aids were inserted and turned on. If Tony engaged in noncompliance (i.e., pulled the hearing aids out), he was allowed a 60-second break from the hearing aids (i.e., they were not replaced for 60 seconds). After the 60-second break, the aids were reinserted with the same contingency in place (i.e., 60-second break for noncompliance).

Differential Reinforcement + Escape Extinction (DR + EE). Prior to beginning treatment sessions, a preference assessment based on Roane et al. (1998) was conducted. At the outset of each treatment session, Tony was given access to preferred toys and attention. Tony maintained access to these stimuli contingent on compliance with wearing either the ear mold or the whole hearing aid unit(s) (depending on the phase of treatment implementation). Noncompliance resulted in immediate removal of preferred toys and the cessation of verbal attention from parent/staff. In addition, the hearing aid (or ear mold, depending on the condition) was immediately reinserted. Once the hearing aid or ear mold was back in the ear, access to the toys and attention was allowed. Thus, Tony could maintain access to the preferred toys and attention through ongoing compliance with wearing the hearing aids (or components thereof).

The evaluation was conducted in two settings: clinic and school. In the clinic setting, the treatment strategy was sequentially applied to compliance with the ear mold or hearing aid in the left ear only, then in the right ear, and, finally, in both ears. In addition, when treatment was conducted with the left ear, the treatment strategy was sequentially applied to wearing the ear mold, followed by wearing the entire unit, and, finally, wearing the unit with the unit switched on. When treatment was conducted with compliance with the right ear, the treatment strategy was applied only to wearing the ear mold before probing compliance with wearing both hearing aids together (entire units and switched on). The treatment sequence was initially meant to be a fading program: starting with the presumed least stimulus (i.e., ear mold), building tolerance, and working up to the entire unit (i.e., ear mold plus hearing aid).

The decision to deviate from the treatment sequence used in the left ear was a clinical decision due to the degenerative nature of Tony's disorder. The team felt a time pressure to increase hearing aid compliance as soon as possible in order to maximize Tony's quality of life. Treatment implementation, therefore, continued with both units in place and switched on. Following implementation in the clinic, a 1-month maintenance follow-up was conducted at Tony's school. Tony was observed across a variety of work (e.g., speech session, one-on-one work time) and free (e.g., snack, play) times, and his compliance with wearing both hearing aids (switched on) was recorded. No attempts were made to manipulate the implementation of the treatment procedures. It should be noted that the teacher did replace hearing aids when pulled out. However, no DR was delivered. Thus, this observation provides a measure of the maintenance of treatment effects (i.e., continued behavior change in the absence of treatment; see Catania, 1999).

Results

The top panel of the Figure (next page) displays the whole-phase mean for each phase of the evaluation. During baseline, Tony was compliant with wearing his hearing aids during an average of 26.6% of the time (range: 20–38%). When the treatment procedures were put in place for compliance with wearing the device in his left ear, compliance averaged 98% of session time (range: 85–100%). When the same treatment was implemented for compliance with the right hearing-aid ear mold, compliance averaged 80% of session time (range: 62–90%). When treatment was implemented for both units in and turned on, compliance averaged 72.7% of session time (range: 48–87%). When Tony was later observed in his classroom, compliance with wearing both units, switched on, averaged 84.7% of the 6-second intervals across a wide variety of activities (range: 52–100%).

The bottom panel of the Figure displays the same results in a session-by-session manner. That is, each data point represents Tony's compliance for a given 10-minute observation. These data are included to demonstrate that compliance during each treatment phase either was stable (e.g., left ear, right ear, and follow-up) or revealed an upward trend (e.g., both units). In addition, compliance during the maintenance observation conducted at school was stable across time and activity.

Discussion

In this evaluation, a DR procedure with EE was used to increase the percentage of time a young child with Sanfilippo syndrome complied with wearing hearing aids. The treatment procedures were implemented in a clinic setting, and maintenance data were collected in the school setting. Results suggested the program was effective for increasing hearing aid compliance.

Figure. Mean percentage with hearing aid (or ear mold) successfully worn across pre-treatment baseline, left ear treatment application, right ear treatment application, treatment application for both ears, and maintenance follow-up (upper panel). Percentage of session time with hearing aid (or its components) successfully worn across each session (lower panel).

This evaluation makes several modest contributions to the existing literature. First, a review of the behavioral and audiology literature did not yield any studies that have used behavioral procedures to improve compliance with hearing aid use. Thus, this evaluation represents a unique demonstration of DR and EE for this specific behavioral concern. Second, the evaluation was conducted with a child diagnosed with Sanfilippo syndrome. This syndrome is a rare, degenerative condition that includes progressive hearing loss over the life span. Despite the degenerative nature of Tony's condition, his behavior appeared to be sensitive to the contingencies and treatment results were maintained at follow-up. Third, researchers (e.g., Palmer, Adams, Bourgeois, Durrant, & Rossi, 1999) have established that hearing aid compliance can reduce problem behavior in some populations. Though not the specific focus of the current investigation, it is plausible that identifying behavioral strategies to increase hearing aid compliance could have the complementary effect of reducing problem behavior.

This evaluation also has some limitations. Most of these limitations are derived from the abbreviated nature of the evaluation, the clinical setting in which it was conducted, and the fact that only one individual participated. First, experimental control of the response was not established. That is, no specific single-subject design was employed. However, given the clinical nature of the intervention and the degenerative nature of Tony's disorder, the clinical team decided not to remove the contingencies once the desired response was being observed. Second, no IOA was collected during the follow-up observation in the classroom. The follow-up observation was conducted by a member of the clinical team that initially worked with Tony. However, the clinical team could not afford to send a second member, given its ongoing clinical responsibilities. Third, longer term follow-up data were not obtained. Thus, while the treatment seemed to have produced maintained effects at 1 month, it is unknown for how long these effects were maintained and whether or not reexposure to the treatment would be needed to reestablish desired response, given any decrease in compliance. Finally, as mentioned earlier, only 1 individual participated in the study, which could limit its external validity.

Given these limitations, the current data set should be considered preliminary. However, each of the limitations also establishes a potential area of future clinical research. Thus, these data can be considered pilot data for a variety of future studies related to the use of behavioral treatments to increase the use of hearing aids and other sensory enhancement devices. Of particular interest would be the generalizability of these procedures to other individuals with and without developmental disabilities. The behavioral literature is replete with single-subject design examples of effective treatment. External validity for behavioral approaches to treatment based on single-subject design is strengthened through replication; in particular, replication across a wide range of participant characteristics. Additionally, it would be interesting to

parse out the components of treatment that were responsible for the observed effect. Future studies could be designed to evaluate the contribution of the sequential application of the hearing aid components and the relative contributions of DR and EE.

References

- Achenback, T., & Edelbrock, C. (1983). *Manual for the child behavior checklist and revised child behavior profile*. Burlington, VT: Queen City Printers.
- American Academy of Audiology. (2004). Pediatric amplification guideline. *Audiology Today*, 16, 46–53.
- Bass-Ringdahl, S.M. (2010). The relationship of audibility and the development of canonical babbling in young children with hearing impairment. *Journal of Deaf Studies and Deaf Education*, 15(3), 287–310.
- Catania, A.C. (1999). *Learning*. Cranbury, NJ: Prentice Hall.
- Davis, J.M., Elfenbein, J., Schum, R., & Bentler, R. (1986). Effects of mild and moderate hearing impairments on language, educational, and psychosocial behavior of children. *Journal of Speech and Hearing Disorders*, 51, 53–62.
- Eilers, R.E., & Oller, D.K. (1994). Infant vocalizations and the early diagnosis of severe hearing impairment. *Journal of Pediatrics*, 124, 199–203.
- Iwata, B.A., Pace, G.M., Kalsher, M.J., Cowdery, G.E., & Cataldo, M.F. (1990). Experimental analysis and extinction of self-injurious escape behavior. *Journal of Applied Behavior Analysis*, 23, 11–27.
- Marshall, L.A. (1997). *Communication accessibility, behavioral ratings, and childhood deafness: Investigation of three modes of communication* (Unpublished doctoral dissertation). Washington, DC: Gallaudet University.
- Moeller, M.P., Hoover, B., Peterson, B., & Stelmachowicz, P. (2009). Consistency of hearing aid use in infants with early-identified hearing loss. *American Journal of Audiology*, 18, 14–23.
- Moeller, M.P., Tomblin, J.B., Yoshinaga-Itano, C., Connor, C.M., & Jerger, S. (2007). Current state of knowledge: Language and literacy of children with hearing impairment. *Ear & Hearing*, 28, 740–753.
- Nathani, S., Neal, A.R., Olds, H., Brill, J., & Oller, D.K. (2001, April). *Canonical babbling and volubility in infants with moderate to severe hearing impairment*. Paper presented at the International Child Phonology Conference, Boston, MA.
- National Center for Hearing Assessment and Management (NCHAM). (2008). Retrieved January 4, 2010, from <http://www.infanthearing.org>.
- National MPS Society. (2008). Retrieved November 4, 2010, from <http://www.mpsociety.org>.
- Oller, D.K., & Eilers, R.E. (1988). The role of audition in infant babbling. *Child Development*, 59, 441–449.

- Ontario Ministry of Children and Youth Services. (2007). *Ontario infant hearing program protocol for the provision of amplification, version 3.1*. Retrieved December 28, 2009, from www.mountsinai.on.ca/care/infant-hearingprogram/documents/amplification_revision_2007_006.pdf.
- Palmer, C.V., Adams, S.W., Bourgeois, M., Durrant, J., & Rossi, M. (1999). Reduction in caregiver-identified problem behaviors in patients with Alzheimer disease post-hearing aid fitting. *Journal of Speech, Language, and Hearing Research*, 42, 312–328.
- The Pediatric Working Group. (1996). Amplification for infants and children with hearing loss. *American Journal of Audiology*, 5, 53–68.
- Pelios, L., Morren, J., Tesch, D., & Axelrod, S. (1999). The impact of functional analysis methodology on treatment choice for self-injurious and aggressive behavior. *Journal of Applied Behavior Analysis*, 32, 185–195.
- Ringdahl, J.E., Kitsukawa, K., Andelman, M.S., Call, N., Winborn, L., Barretto, A., & Reed, G. K. (2002). Differential reinforcement with and without instructional fading. *Journal of Applied Behavior Analysis*, 35, 291–294.
- Roane, H.S., Vollmer, T.R., Ringdahl, J.E., & Marcus, B.A. (1998). Evaluation of a brief stimulus preference assessment. *Journal of Applied Behavior Analysis*, 31, 605–620.
- Wake, M., Hughes, E.K., Poulakis, Z., Collins, C., & Rikards, F.W. (2004). Outcomes of children with mild-profound hearing loss at 7 to 8 years: A population study. *Ear and Hearing*, 25, 1–8.

Literature Review

Concerns Regarding Direct-to-Consumer Hearing Aid Purchasing

Suzanne H. Kimball, Au.D.

An individual over age 18 can purchase a hearing aid online or through mail order if they sign a waiver declining a medical evaluation, while those under 18 are required to be seen by a physician to obtain medical consent. However, in many states there is nothing to prevent a parent or caregiver from purchasing hearing aids for their child from a direct-to-consumer distributor. Online purchasing of hearing aids may be particularly tempting for parents and caregivers due to the cost savings involved. This manuscript summarizes the results of three previously published studies that evaluated the effectiveness of online hearing aid distribution utilizing adult populations. These results are consistent with other investigations that suggest purchasing hearing aids through mail order or the Internet may carry potential health and amplification risks. Based on the outcomes of these three studies, implications of online purchases for children and adolescents are discussed.

Introduction

According to CNNMoney.com (2007), e-commerce product sales represent about 6% of total retail product sales in the United States. Similar trends are seen in other countries, such as England, where in 2009 the Office for National Statistics indicated that approximately 5% of product sales were online (Office for National Statistics, Great Britain, 2009). The advantages to purchasing products online or through mail order (a similar business model) include convenience, access, and travel time. Consumers can search for products much faster than at a traditional bricks-and-mortar retailer. Products can be shipped

Suzanne H. Kimball, Au.D., is an Assistant Professor in the Department of Communication Sciences and Disorders at the University of Oklahoma Health Sciences Center. Correspondence concerning this article should be directed to Dr. Kimball at suzanne-kimball@ouhsc.edu.

directly to a home or business, and cost comparison can take place rapidly without the hassle of driving from store to store or checking local advertisements. Products can also often be found cheaper through online and mail order distributors than in a retail establishment.

It may not be surprising, then, that purchases of hearing aids through the Internet and mail order have increased over the past several years. Estimates from 2008 suggest that direct-to-consumer hearing aid purchases, which include mail order and Internet, accounted for 4–5% of all hearing aid purchases in the United States (Kochkin, 2009). According to Sweetow (2009), there are three different ways consumers can obtain hearing aids through online/mail order sources. Two business models require consumers to work with a licensed hearing health care professional for hearing testing and hearing aid fitting services. For the first, “consumer/patient referral sources,” the consumer is referred to a hearing professional, either an audiologist or hearing instrument specialist (HIS), from a website interaction or phone call. All services take place through the local professional, including the purchase of the hearing aids. For the “marketing/point of sale with face-to-face fitting” model, the consumer responds to a website and is referred to a local hearing professional (audiologist or HIS) for services. The hearing aid purchase takes place between the web-based company and the consumer. The local professional is paid separately by the web-based company for the fitting and follow-up services. The last hearing aid service delivery model is referred to as “direct-to-consumer” (DTC), “mail order,” or “Internet sale.” Businesses in this model may accept (a) hearing test results that can be mailed or faxed to the company, (b) an online hearing test, or (c) possibly no hearing test results at all. These companies sell either “one-size-fits-most” in-the-ear (ITE) or open-fit behind-the-ear (BTE) nonprogrammable analog or digital hearing aids, or custom-fitted, programmable digital hearing aids supplied by various manufacturers. The hearing aids are sent directly to the consumer. Generally, no face-to-face fitting takes place with this business model (Sweetow, 2009). If the hearing aids need to be modified or reprogrammed, the consumer must either return them via mail to the company to be adjusted (or returned for credit), or seek the services of a local hearing health care professional at the consumer’s own expense.

To sell hearing aids directly to consumers, a company is required only to post a statement by the Food and Drug Administration advising consumers to seek medical clearance before purchasing the aid(s). This statement advises that it is in the consumer’s best interest to see a physician, preferably an otolaryngologist, to determine if there are any contraindications to the use of a hearing aid. Consumers over the age of 18 can sign a waiver declining the medical evaluation, while those under the age of 18 are required to be seen by a physician to obtain medical consent before being fitted with a hearing aid. While the medical consent is necessary for those under 18 years old, in many states there is no law to prevent a parent or caregiver from purchasing hearing

aids for their child directly through an online or mail order distributor. While it might be considered “best practice” for a child to be seen by a licensed and clinically certified audiologist, it is not required by law consistently throughout the United States. Illinois, for example, does not require a physician’s consent. The online mode of distribution may be particularly tempting for parents and caregivers as hearing aids purchased online are often less expensive than those purchased through a traditional method. That being said, several studies have shown that mail-order, over-the-counter (OTC), and online hearing aids may not be suitable for everyone, including children and adolescents (Callaway & Punch, 2008; Cheng & McPherson, 1999; Kasper, Spitzer, & Rodriguez, 1999; Sweetow, 2001; Zimmerman, 2004). In general, these studies have concluded that hearing aids purchased via a DTC method pose potential health risks and may provide inappropriate or insufficient amplification for individuals with hearing loss.

Much debate has arisen about DTC hearing aid sales that are conducted without a face-to-face meeting and fitting by a licensed audiologist or HIS. While the regulation of hearing aid dispensing is generally determined by each individual state, the Medical Device Amendments to the Food, Drug and Cosmetic Act provides that a state law can be preempted if it differs from the federal law. State laws in Missouri, Texas, and Florida have been preempted by a federal ruling, resulting in the states’ inability to regulate online and mail order sales. California currently requires that, before a DTC hearing aid purchase takes place, the Internet seller receive a statement signed by a California-licensed physician, audiologist, or hearing aid dispenser to verify that a direct observation of the ear canal has been performed and no contraindications to hearing aid use exist. Individuals who are diagnosed with certain medical conditions are precluded under the California law from purchasing hearing aids directly from an online or mail order source (Sweetow, 2009). While this law has yet to be challenged in California, the state law could likely be preempted due to the legal precedence set in the other states.

In order to purchase a hearing aid online, consumers are asked to do three things: (a) send a copy of hearing test results obtained from a local hearing professional directly to the company or take an online hearing test; (b) agree to purchase aids from the selection offered by the online dispenser; and (c) for custom-made products, send in ear mold impressions made by a local professional, or self-make ear mold impressions with instructions and materials sent by mail from the online company. The consumer then sends the impression(s) back to the company in order for the hearing aid(s) to be manufactured. Some online dispensers do not require hearing test results or an ear mold impression, depending on the type of aid the consumer selects. Hearing aids obtained through mail order generally do not require test results or ear mold impressions. As mentioned previously, studies have indicated that hearing aids obtained in this manner may not provide adequate amplification and may

pose potential health risks (Callaway & Punch, 2008; Cheng & McPherson, 1999; Kasper, et al., 1999).

Method

To determine the effectiveness of the online hearing aid dispensing model, three separate experiments were conducted at the Illinois State University Eckelmann-Taylor Speech and Hearing Clinic. The purpose was to determine potential health risks or hearing aid fitting concerns specifically resulting from online dispensation. The first experiment assessed the quality of one online hearing test (Kimball, 2008a). The second assessed how well individuals were able to “self-make” ear mold impressions, as is required for some online purchases (Kimball, 2008b). The third chronicled 2 individuals throughout the entire process of online hearing aid purchasing (Kimball & Yopchick, 2009). The results of these studies have been previously reported in *The Hearing Journal* and are reprinted with the permission of *The Hearing Journal* and its publisher, Lippincott, Williams & Wilkins.

While each of the three studies was published independently in *The Hearing Journal* in 2008 and 2009, the target audience for this journal is specifically hearing health care providers. Parents of children with hearing loss should be aware of all of the various hearing aid distribution models and any potential dangers thereof in order to make an informed decision regarding their child’s hearing health care needs. Therefore, while all three of these experiments were conducted on adult subjects, similar results would likely be obtained for pediatric and adolescent populations, or, at a minimum, negative findings might possibly create a more deleterious outcome. Each study is summarized below.

Experiment One

The purpose of the first experiment was to determine the accuracy of one online hearing test. For this online company, the results of the online hearing test can be used to program hearing aids purchased. A total of 81 subjects in three age groups took part in the study (Kimball, 2008a).

All subjects took the online hearing test as well as a standard hearing test in a sound suite at the university’s clinic. For the online test, subjects were each seated in front of one of four computers (two Dell desktops and two Dell laptops) and were provided Aiwa HP-A091 on-the-ear headphones. These headphones are the same style issued by airlines for in-flight listening to music or movies and are similar to those provided by the online company, if requested. It should be noted that for the online testing, consumers can use any privately owned, commercially available headphones, but each subject used the same type for this experiment.

Subjects also received a hearing test in a sound-treated room (IAC) using standard audiometric test procedures and calibrated audiometric equipment

(Grason-Stadler GSI-61 Audiometer). The recommended Hughson-Westlake down 10 dB-up 5 dB ascending pure-tone testing method was utilized as compared with the descending only method that was used in the online test. All sound room testing was completed by a certified audiologist or an audiology doctoral student.

Results for Experiment One

Results indicated statistically significant differences, using paired T-tests at the .05 confidence level, between the online and professional testing conditions across all groups and by individual age groups for the frequency range 250–4000 Hz. Some subjects were identified as having mild hearing losses via the computer test, whereas the booth test indicated typical hearing sensitivity. In addition, the computer test underestimated hearing loss in some subjects with known hearing losses. It may also be reasonable to assume that differences occurred for the online test due to a testing effect because the subjects initiated the test signals themselves.

Experiment Two

The purpose of the second experiment was to determine how well individuals were able to “self-make” ear mold impressions compared with those made by a hearing professional. Data for this study were collected from 83 subjects (Kimball, 2008b).

First, the author anonymously contacted an online company and obtained its kit for the purpose of making an ear mold impression at home. The kit included (a) a polyethylene syringe, (b) one pack of silicone ear mold impression material (to make two impressions), (c) a medium cotton otoblock, (d) a cotton swab (which purchasers were to obtain at home, but was included in the test kits), and (e) two sets of instructions. The contents of the kit were replicated for use by the subjects in the project.

Each subject was asked to pair with a partner (a friend, family member, or spouse, as was suggested in the kit instructions) and each pair was given an impression-making kit. While most likely in reality only one individual would require an impression for the purposes of obtaining a hearing aid, for this experiment both partners made ear impressions on each other. Each pair was asked to follow the included instructions and make an impression for one of their partner’s ears, and then switch places and have the partner make one on them. When the subject pair finished, they then had an impression made on the same ear by a member of the research team using traditional, professional hearing aid mold supplies and equipment. Once the impressions were all completed, both the subject-made and professional-made molds were sent to the online hearing aid manufacturer for evaluation.

Results for Experiment Two

The impressions were rated by two anonymous audiologists employed by the online manufacturer. Ratings values (0–5) were averaged for each impression (A [amateur] and B [professional]). A score of 0 indicated that the impression contained none of the criteria necessary to build a hearing aid from the impression, and a score of 5 indicated that the impression had all of the necessary criteria. The results suggested that having average consumers make ear mold impressions without professional assistance was a difficult task at best. Approximately 50% of the participants' self-produced ear mold impressions received a score of 2 or lower, and almost 80% received a score of 3 or lower. Only 20.5% of the "amateur" impressions were rated with an average score of 3.5 or higher. In contrast, more than 97% of the impressions produced by the research team received a score of 3.5 or higher, with less than 3% of the "professional" impressions scoring a 3. Almost 93% of the impressions made by the research team rated 4.0 or higher, with 25.3% scoring an average rating of a perfect 5. No impressions produced by the research team received a score lower than 3. While it was not surprising that trained professionals could produce ear mold impressions superior to those who were untrained, it does speak to how well average consumers can self-make their own impressions.

Experiment Three: Individual Case Studies

Two adult subjects were recruited for the study and were asked to purchase hearing aids of their choice through an online hearing aid distributor. Subject recruitment was limited to two case studies due to the expense of purchasing hearing aids. The 2 subjects were then asked to follow the online purchasing process as detailed through the website, which included taking the online hearing test, making an ear mold impression (for subject 1), and navigating through the general purchasing process. No professional advice was given by anyone on the research team. For comparison purposes, after the online purchase, both subjects were subsequently fitted with hearing aids at the university clinic by a licensed and certified audiologist (Kimball & Yopchick, 2009).

Results for Experiment Three

Using real-ear verification set to the NAL-NL1 prescriptive method, an analysis of the hearing aids ordered from the online company by the first subject indicated that the aids did not reach targets on either of the two "pre-set" programs, even after repeated reprogramming. The second subject selected one noncustom device from the online company. Again with this subject, real-ear verification using the NAL-NL1 prescription method indicated insufficient hearing aid gain as was indicated through the verified

hearing loss. After receiving the aids in the mail, subject 2 initially perceived all sounds as too loud. Both subjects returned their aids to the online company for credit.

Both subjects were subsequently fitted at the university clinic with mixed success, even though a recommended fitting protocol (Valente, Valente, & Mispagel, 2007) and pre- and postoutcome measures were used to determine appropriate amplification needs. Subject 1 returned the university-obtained hearing aids for credit. Subject 2 kept the two recommended aids purchased from the university clinic but reportedly does not use them on a continual basis.

Discussion

The combined results of these and prior studies (Callaway & Punch, 2008; Cheng & McPherson, 1999; Kasper, et al., 1999; Sweetow, 2001; Zimmerman, 2004) indicate that the DTC purchasing of hearing aids via the Internet or mail order is risky at best. There are several downfalls to purchasing hearing aids without face-to-face interaction with a hearing professional. The first is the online hearing test. As seen in experiments one and three, hearing thresholds can easily be under- or overestimated in the absence of standard test procedures performed utilizing calibrated test equipment and a sound-treated testing room. Additionally, only pure-tone air conduction results can be obtained from online hearing testing, and in the case of experiment one, only through 4000 Hz with no interoctave frequencies. No speech threshold, loudness measures (most comfortable or loudness discomfort), or bone conduction thresholds are obtained.

While the use of speech testing has not necessarily been shown to predict hearing aid satisfaction (Killion & Gudmundsen, 2005), the use of loudness discomfort levels has been shown to be useful in successful hearing aid outcomes (Mueller & Bentler, 2005). In addition, bone conduction measurements are recommended for use by the American Academy of Audiology's Pediatric Amplification Protocol (2003). Also, while not all studies agree, the use of threshold information above 4000 Hz may be important for hearing aid benefit, particularly in children (Ching, Dillon, & Katsch, 2001; Stelmachowicz, 2001). Many hearing aids currently on the market have frequency response capabilities that reach beyond the 4000 Hz range. While online hearing tests may post a caution that the results from the test should not be considered professional advice, hearing aids can be purchased and programmed based on the online results. Additionally, hearing aid recommendations can be based on the online test, as in the case of subject 2 in experiment three, which can also be of concern.

Another potential downfall with DTC fittings is the lack of real-ear verification techniques. According to Valente (2006), verification through verified protocols such as real-ear measurements (conducted with the hearing aid user

present) is the most important aspect of a hearing aid fitting. Valente states that when a hearing aid is programmed to "first fit" and no hearing aid verification is utilized, as in DTC distribution, there will be a high probability for error due to the "one-size-fits-all approach" that cannot account for individual differences. Mueller (2005) and Killion (2004) support this notion and indicate that often hearing aid manufacturers' proprietary formulae for calculating amplification requirements result in a hearing aid user not being able to hear quiet sounds. Mueller suggests that when a patient is fitted for a hearing aid that is set to "first fit," real-ear verification should be performed and gain settings should be adjusted to meet specified targets as indicated by verified prescriptive methods, such as NAL-N11/2 (Dillon, Ching, Keidser, & Katsch, 2008) or Desired Sensation Level (Seewald, Moodie, Scollie, & Bagatto, 2005). There is ample evidence to demonstrate the effectiveness of real-ear verification and its importance as a clinical tool in hearing aid fittings with children, adolescents, and adults (Mueller, 2005; Mueller & Bentler, 2005; Valente, 2006; Westwood & Bamford, 1995). Additional fitting errors may be caused by DTC fittings due to the lack of appropriate ear acoustic measures, such as the real-ear to coupler difference, which can help to account for any individual or age differences (Moodie, Seewald, & Sinclair, 1994). Overall, the lack of all of these measures may result in inappropriate amplification. For children and adolescents in particular, appropriate amplification is imperative to address their social and educational needs. It is unknown if dispensers who distribute DTC products test the purchased hearing instruments in a 2cc coupler to ensure that the aids are functioning according to the manufacturer's specifications prior to shipping. Failure to do so may result in additional fitting and use concerns.

The overall success of a hearing aid fitting cannot rely solely on the prescribed gain settings, but in the case of custom products or BTE hearing aids, must also take into account the quality and fit of the ear mold impression and the manufactured ear mold. Hoover, Stelmachowicz, and Lewis (2000) concluded that for at least a limited range of conditions found in clinical practice, ill-fitting ear molds may influence clinical decisions about the type of hearing aid fitted and the amount of gain provided. Poorly fitted ear molds can result in reduced hearing aid benefit.

Experiment two clearly showed that self-made ear mold impressions could potentially result in ill-fitting ear molds or custom-fit hearing products. Many of the amateur ear mold impressions in the experiment had short ear canals because the subjects were uncomfortable placing the otoblock deeply into their research partner's ear canal. Pirzanski (2006) stated that if an ear mold impression has a shallow otoblock position, the impression material will not stretch the cartilage within the seal area and the resulting ear mold may fit loosely, have retention problems, and be susceptible to acoustic feedback. Realistically, it may be difficult to get an appropriate hearing aid fit if the ear mold impression is not professionally made. Additionally, potential health risks arise if injury occurs as the result of the process. Parents or caregivers who opt to

make ear mold impressions on their child may encounter some or all of these issues.

Another area that is not addressed in DTC fittings is the need for counseling on various topics such as instructions on the care, use, and maintenance of the hearing aids; realistic expectations; and any communication strategies available to wearers to enhance their listening experience. An important counseling topic for parents of children with hearing loss is options for hearing aid insurance and techniques, and suggestions for keeping the hearing aid on the child's ear. According to Kochkin (2005), hearing aid dispensers take an average of about 45 minutes to explain this information to hearing aid patients. It is likely that pediatric/adolescent evaluations require even more time. Humes (2006, as cited in Desjardins & Doherty, 2009) has shown that individuals who have greater difficulty managing and manipulating their hearing aids within the first 2 weeks of use are not as satisfied, perceive less benefit, and reportedly wear their hearing aids less than those that have fewer problems within the first 2 weeks. This may indicate that if consumers do not have a full understanding of the use and capabilities of the hearing aids after purchasing them online, they may ultimately reject their usage. The same could be said for a parent or caregiver of a child with a hearing loss. During experiment three, the research team had an opportunity to inspect the product information that accompanied the hearing aids purchased online. Instructions covered the care, use, and maintenance of the aids; how to cut the ear mold tubing to fit the hearing aids; and a booklet on hearing with the aids. While the information was not particularly difficult for the trained professional to follow, it is unknown how well a typical consumer can digest this type of information. Parents could potentially have a more difficult time with the information provided by an online distributor, as they may or may not have a full understanding of their child's hearing aids or hearing loss. Consumer-purchased hearing aids also lack the opportunity to utilize the data logging feature found on many current hearing aids, as consumers do not have the hearing aid software readily available to them for monitoring purposes. For children or adolescents, this may hamper progress for future reprogramming needs or for cochlear implant candidacy.

Conclusion

The potential risks involved with DTC hearing aid distribution outweigh any cost savings involved in the purchase. Parents and caregivers may still be tempted to purchase products online, particularly when a licensed hearing professional has performed the hearing testing and made the ear mold impressions. While these professional services may have been utilized for the child, it still does not necessarily mean that a hearing aid purchased online or through mail order will result in an adequate fit or provide adequate gain for successful hearing aid usage, particularly in an academic setting. The results

of this series and prior studies suggest that parents are best served by having their children see a licensed and certified hearing health care professional, particularly one who has experience with the pediatric and adolescent population and uses standardized diagnostic test procedures and real-ear verification measures to determine appropriate amplification needs.

References

- American Academy of Audiology. (2003). *Pediatric amplification protocol*. Retrieved October 20, 2010, from <http://www.audiology.org/resources/documentlibrary>.
- Callaway, S., & Punch, J. (2008). An electroacoustic analysis of over-the-counter hearing aids. *American Journal of Audiology*, 17(1), 14–24.
- Cheng, C., & McPherson, B. (1999). Over-the-counter hearing aids: Electroacoustic characteristics and possible target client groups. *Audiology*, 39, 110–116.
- Ching, T., Dillon, H., & Katsch, R. (2001). Do children require more high-frequency audibility than adults with similar losses? *A Sound Foundation through Early Amplification: Proceedings of the Second International Conference* (pp. 141–152). Chicago: Phonak, Inc.
- CNNMoney.com. (2007). *Online sales spike 19 percent*. Retrieved May 14, 2007, from http://money.cnn.com/2007/05/14/news/economy/online_retailng.
- Desjardins, J., & Doherty, K. (2009). Do experienced hearing aid users know how to use their hearing aids correctly? *American Journal of Audiology*, 18(1), 71–76.
- Dillon, H., Ching T., Keidser, G., & Katsch, R. (2008). *Derivation of the NAL-NI2 prescription formula*. Paper presented at the AudiologyNow 2008 Convention, Charlotte, NC.
- Hoover, B., Stelmachowicz, P., & Lewis, D. (2000). Effect of earmold fit on predicted real ear SPL using a real ear to coupler difference procedure. *Ear and Hearing*, 21(4), 310–317.
- Humes, L. (2006, November). Hearing-aid outcome measures in older adults. *Proceedings of the International Conference, A Sound Foundation Through Early Amplification*. Chicago: Phonak, Inc.
- Kasper, C., Spitzer, J., & Rodriguez, H. (1999). Mail-order hearing aids and patient safety. *Hearing Journal*, 52(7), 41–44.
- Killion, M. (2004). Myths about hearing aid benefit and satisfaction. *Hearing Review*, 11(9), 14–20.
- Killion, M., & Gudmundsen, G. (2005). Fitting hearing aids using clinical pre-fitting speech measures: An evidence-based review. *Journal of the American Academy of Audiology*, 16, 439–447.
- Kimball, S. (2008a). Inquiry into online hearing test raises doubt about its validity. *Hearing Journal*, 61(3), 38–46.

- Kimball, S. (2008b). Making earmold impressions at home: How well can untrained consumers do it? *Hearing Journal*, 61(4), 26–30.
- Kimball, S., & Yopchick, S. (2009). Study compares hearing aids fitted online with clinical findings. *Hearing Journal*, 62(3), 44–47.
- Kochkin, S. (2005). MarkeTrak VII: Customer satisfaction with hearing aids in the digital age. *Hearing Journal*, 58(9), 30–37.
- Kochkin, S. (2009). MarkeTrak VIII: 25-year trends in the hearing health market. *Hearing Review*, 16(10), 12–31.
- Moodie, K., Seewald, R., & Sinclair, S. (1994). An approach for ensuring accuracy in pediatric hearing instrument fitting. *American Journal of Audiology*, 3, 23–31.
- Mueller, H. (2005). Fitting hearing aids to adults using prescriptive methods: An evidence-based review of effectiveness. *Journal of the American Academy of Audiology*, 16(7), 448–460.
- Mueller, H., & Bentler, R. (2005). Fitting hearing aids using clinical measures of LDL: An evidence-based review of effectiveness. *Journal of the American Academy of Audiology*, 16(7), 461–472.
- Office for National Statistics, Great Britain. (2009, December). *Statistical bulletin: Retail sales*. Retrieved January 2010 from <http://www.statistics.gov.uk/pdfdir/rs0110.pdf>.
- Pirzanski, C. (2006). Earmolds and hearing aid shells: A tutorial part 2: Impression-taking techniques that result in fewer remakes. *Hearing Review*, 13(5), 39–46.
- Seewald, R., Moodie, S., Scollie, S., & Bagatto, M. (2005). The DSL method for pediatric hearing instrument fitting: Historical perspectives and current issues. *Trends in Amplification*, 9(4), 145–157.
- Stelmachowicz, P. (2001). The importance of high frequency amplification in young children. *A Sound Foundation through Early Amplification: Proceedings of the Second International Conference*. Chicago: Phonak, Inc.
- Sweetow, R. (2001). An analysis of entry-level, disposable, instant-fit, and implantable hearing aids. *Hearing Journal*, 54(2), 28–42.
- Sweetow, R. (2009). Hearing aid delivery models, part I. *Audiology Today*, 21(5), 49–58.
- Valente, M. (2006). Valente leads development of national adult hearing-aid fitting guidelines. *Record*, Washington University, St. Louis, 30(32).
- Valente, M., Valente, M., & Mispagel, K. (2007). Hearing aid selection and fitting. In J. Katz (Ed.), *Handbook of clinical audiology* (5th ed., pp. 707–728). Baltimore, MD: Lippincott, Williams and Wilkins.
- Westwood, G., & Bamford, J. (1995). Probe-tube microphone measures with very young infants: Real ear to coupler differences and longitudinal changes in real ear unaided response. *Ear and Hearing*, 16, 263–273.
- Zimmerman, A. (November 20, 2004). Cheaper hearing aids crop up online; as prescription devices get costlier, consumers turn to web; skipping the medical exam. *Wall Street Journal*, D1.

Book Review

Auditory-Verbal Practice: Toward a Family Centered Approach

Auditory-Verbal Practice: Toward a Family Centered Approach

Ellen Rhoades, Ed.S., LSLS Cert. AVT, and

Jill Duncan, Ph.D., LSLS Cert. AVT, Editors

Charles C. Thomas Publishers

Hard Cover, 2010, \$79.95, 420 pages

This is an important, comprehensive work that will not only support the growth of family support skills in listening and spoken language professionals, but of all rehabilitation practitioners. In that sense, it is somewhat unfortunately titled. It would be a loss if the readership of this book were confined to listening and spoken language professionals.

Ellen Rhoades and Jill Duncan divide the book into three separate sections: "Auditory-Verbal Practice," "Systemic Family Perspective," and finally, "Family-Based Auditory-Verbal Intervention." The chapters and sections flow well and are very well integrated, and the writing is clear and engaging throughout. The research bases of the discussions presented are exhaustive and current. This book would be worth reading for the literature reviews alone, but it offers much more than that. The editors and contributing authors provide not only the basic information to enable the novice reader to better evaluate evidence and understand practice, but extensive discussions that will challenge even lifelong students of family-centered intervention and listening and spoken language development.

The first section, "Auditory-Verbal Practice," provides the theoretical, historical, and evidentiary bases for auditory-verbal practice (AVP). The authors' use of the term "practice" versus "therapy" is intentional, as the term "therapy" immediately connotes a non-normative process and puts the emphasis on the professional rather than on collaborating with the families to best support listening and spoken language development for children with hearing loss. The third chapter in this section provides a discussion of ethical considerations related to AVP that will both challenge and inform. Indeed, that is the case throughout this book. The editors and authors do not shy away from potentially contentious issues, but meet them head on with a balanced, thorough discussion. Issues such as the locations in which AVP is carried out, the potential dangers of communicating a "more is better" philosophy to families, a narrow focus on parents (usually mothers), and the typically higher socioeconomic status and majority culture membership of both traditional families and practitioners are examined in a fearless, even-handed way within the theoretical framework of family-centered practice.

The second section, "Systemic Family Perspective," lays the groundwork for a family-systems approach to intervention. This section provides a breadth

of information that will support the development of systemic thinking in new practitioners, and depth that will enhance the understanding of more experienced readers. Chapters include discussions of enablement and environment, circles of influence (based on the work of Urie Bronfenbrenner), and an introduction to systemic family therapy and core constructs of family therapy. If there is a danger in this book, it might be in the temptation of newer practitioners to take on the role of a family therapist. The editors and authors are clear that this is not their intent, but argue, convincingly, that the theoretical bases of this discipline can provide insight regarding how practitioners might more effectively engage family systems versus individuals in supporting optimal listening and spoken language development.

The final section, "Family Based Auditory-Verbal Intervention," provides a rich array of "how to" advice to enable readers to become more family centered in their practice. Chapters address progressive steps toward more family-centered practice, socio-emotional considerations, ways to support families, family-centered assessments, application of family therapy constructs to listening and spoken language specialist practice, a family intervention framework, and an excellent discussion by Mary McGinnis of appropriate goals to support providers. The section closes with two family retrospectives that highlight the need for and benefit of an equal collaboration between family systems and practitioners in relation to achieving optimal listening and spoken language outcomes for children who are deaf or hard of hearing.

This book has an impressive international list of contributors including certified listening and spoken language specialists (LSLSs), educators, family therapists, and parents of children who are deaf or hard of hearing. The editors state that it is intended as a text for graduate students in educational audiology, deaf education, speech-language pathology and early childhood special education. I would extend that to all of rehabilitation medicine, including physicians, occupational and physical therapists, psychologists, administrators, and policy-makers. Many organizations and practitioners espouse adherence to family-centered practice, but the practical application of this philosophy varies tremendously. This book will, I hope, serve as a means to clarify not only what we mean when we say we practice family-centered intervention, but how that philosophy might look in actual practice. I have ordered this book for both my practice and academic libraries, and plan to make several of the chapters required reading for my annual LSLS seminars with speech-language pathology graduate students. I strongly encourage others to do the same.

Kathryn Ritter, Ph.D., LSLS Cert. AVT, is an Adjunct Associate Professor in the Faculty of Rehabilitation Medicine at the University of Alberta and a Listening and Spoken Language Specialist in the Department of Communication Disorders at Glenrose Rehabilitation Hospital in Edmonton, Alberta. She has no personal or professional affiliation with the editors of this book.

Book Review

Developmental Language Disorders: Learning, Language, and the Brain

Developmental Language Disorders: Learning, Language, and the Brain

Diane L. Williams, Ph.D.

Plural Publishing, San Diego, CA

Soft Cover, 2010, \$69.95, 336 pages

Numerous books and articles have been written about developmental language disorders. In “Developmental Language Disorders: Learning, Language, and the Brain,” Diane Williams, a Ph.D. with 18 years of clinical experience, takes a neurological perspective to the topic. She notes in the preface that most of the relevant research in this area is in highly specialized publications that are not readily available outside research and academic settings. The target audience of the book is graduate students, practicing speech-language pathologists (SLPs), and special educators.

The book is divided into three sections, with three chapters in each section. The first section, “Brain Development for Learning,” has separate chapters explaining how the brain is organized for learning language, the cortical basis of learning language, and measuring the brain-behavior relationship. The first two chapters provide a good review of neuroanatomy and brain-language relationships. The third chapter discusses the various neuro-imaging techniques used to evaluate brain function (e.g., PET scans, fMRI, and ERP). Readers will find the information about these techniques useful.

The second section, “Neurobiological Research on Developmental Language Disorders,” contains chapters on specific language impairment (SLI) and dyslexia, autism, Downs Syndrome, Williams Syndrome, and Fragile X. The chapter on autism is the strongest one, in part because it is the longest of the three and only covers one disorder. The other chapters provide a nice review of neurobiological studies of the other disorders.

The final section of the book is entitled “Brain-Based Intervention.” There are separate chapters on brain-based learning, brain-based assessment and intervention with young children, and brain-based assessment and intervention with older children and adolescents. Although there is certainly useful information in these final chapters, there is a fundamental problem with the notion of brain-based approaches to assessment and intervention. All of the biological and cognitive mechanisms for learning are brain-based. Learning occurs in the brain, not in other organs of the body such as the heart, lungs, or kidneys. There is no such thing as non-brain-based language learning. Language assessments and interventions are thus all brain-based, though

some assessments (e.g., neuroimaging) directly measure neurological structures and activation patterns whereas behavioral measures more directly measure language functions. Both forms of assessment are discussed in these final chapters.

In chapter 7, Williams presents findings from neuroimaging research on individuals with dyslexia, showing that their “brains are somewhat plastic and responsive to remediation” (p. 214). She fails to mention, however, that neuroimaging studies are simply confirming what the traditional measures of reading have shown. Indeed, when fMRIs or other measures (e.g., ERPs) are used to assess language abilities or language change, the findings from these neurological measures are often compared to behavioral assessments with proven reliability and validity.

Williams recognizes the usefulness of behavioral measures of language, but curiously refers to them as measures of brain function as opposed to standard measures of language function. Readers will thus be surprised to see that Table 8.1, entitled “Behaviors to Assess that Relate to Brain Function” (p. 246), contains the following questions:

- How does the child attend to and respond to language?
- How does the child respond to auditory/verbal stimuli in the environment?
- Does the child imitate a word when it is modeled?
- Does the child initiate turn-taking?

Readers will be similarly surprised to read the following suggestions for brain-based intervention:

- Give the child time to respond (p. 250).
- Use a reduced amount of language (single words or short phrases) with a young child (pp. 251-252).
- Pair a spoken word with a gesture like pointing (p. 254).
- Use contextual cues to teach language (p. 260).

These are all useful and appropriate suggestions for assessment and intervention, which makes it particularly puzzling why Williams felt the need to dress them up as brain based. I find the notion of the brain as a disembodied entity that can be studied, evaluated, and treated somewhat troubling. I prefer the focus be on the child or adolescent rather than their brains. Our brains may be responsible for our thoughts, feelings, emotions, and cognitive abilities and skills, but I would like to keep the illusion that we are treating *children* with language disorders, not simply their brains.

In summary, this book provides an overview of developmental language disorders from a neurological perspective. Although there is information in the book that can be applied to the deaf or hard of hearing, there is no mention of deaf or hard of hearing populations in the book. Clinicians who have

diverse caseloads of children with developmental language disorders may find some useful information in the book. I would not envision this book being widely read by clinicians and educators who work exclusively with the deaf and hard of hearing.

Alan G. Kamhi, Ph.D., is a Professor in the Department of Communication Sciences and Disorders at the University of North Carolina in Greensboro.

2010 AG Bell Research Symposium Proceedings

Re-Modeling the Deafened Cochlea for Auditory Sensation: Advances and Obstacles

These are the proceedings of 2010 AG Bell Research Symposium, presented June 27, 2010, as part of the AG Bell 2010 Biennial Convention. The session was moderated by Carol Flexer, Ph.D., CCC-A, LSLS Cert. AVT. For more information about the symposium, visit www.agbell2010convention.org.

What Stops the Inner Ear from Regenerating? By Andy Groves, Ph.D.

Hearing loss is one of the most common disabilities in the United States. The most common form of hearing loss is caused by the death of cochlear hair cells in the organ of Corti and once lost, hair cells in humans and other mammals do not regenerate. In contrast, non-mammalian vertebrates, such as birds, can functionally recover from deafening injury by mobilizing supporting cells in the cochlea to divide and differentiate, replacing lost hair cells. Since the discovery of hair cell regeneration in birds in the 1980s, research has focused on trying to understand the cellular and molecular mechanisms underlying regeneration and why these processes do not occur in mammals.

To address this question, we have developed methods to isolate pure populations of supporting cells from newborn mice and to grow these cells in culture. Surprisingly, we observe that supporting cells from young mice are able to start dividing and can generate hair cells in a manner reminiscent of that seen in birds. In particular, we observed that a negative regulator of cell division – a gene called *p27kip1* – is switched off in supporting cells as they start dividing. However, when we repeat these experiments in older mice that are able to hear, we find that the older supporting cells do not divide and do not switch off the *p27* gene. We are currently investigating the basis for these age-dependent changes in the cochlea. We are also trying to understand how

new hair cells are generated from supporting cells. We have found that an evolutionarily ancient system of signaling – the Notch signaling pathway – is deployed during cochlear development to regulate the correct numbers of hair cells and supporting cells. We have found that manipulating the Notch signaling pathway leads to the production of extra hair cells, and we are currently trying to understand if such manipulations might allow restoration of hair cells in a therapeutic context.

Introduction

One in 500 children is born deaf, with many having some form of hereditary hearing loss. One in 200 3-year-olds have acquired hearing loss. It is also estimated that half of Americans will have lost at least some of their hearing by the time they retire. By the time we reach age 75, about half of us will suffer from balance problems too. Although the causes of hearing loss and balance disorders can be wide-ranging, a frequent common denominator can be found in the cells within our inner ear that mediate our senses of hearing and balance – sensory hair cells.

When we hear sound waves, they are captured by our external ear, passed across the middle ear by the three smallest bones in our body – the hammer, anvil, and stirrup – and conveyed to the cochlea. The cochlea contains about 15,000 hair cells – so-called because of the tiny hair-like projections that stick up from each cell’s surface like organ pipes (Figure 1). As sound waves travel

Figure 1. The top surface of a single hair cell in the cochlea. The hair-like projections sticking up from the surface of this cell form a characteristic “W” shape. It is the movement of these tiny projections that cause the hair cell to become electrically active. *Image courtesy of the House Ear Institute, Los Angeles.*

through the fluid within the cochlea, they stimulate these projections, which cause the hair cells to become electrically active. Each hair cell is connected by a nerve cell to the brain, which then interprets these electrical signals as sound. Hair cells are incredibly mechanically sensitive – it has been estimated that their hair-like projections only have to be moved by a diameter of a few atoms in order for the cell to become stimulated. To put this on a human scale, this amount of movement can be compared to moving the very tip of the Empire State Building by a few inches. This sensitivity allows us to hear extremely soft sounds, such as a whisper, the sound of a pin dropping, or an orchestral pianissimo.

This exquisite sensitivity of hair cells comes at a price – they are extremely vulnerable to damage. In addition to the aging process, a variety of insults and injuries can damage hair cells – principally loud noises, but also chemicals such as certain kinds of antibiotics or chemotherapy drugs. Certain hereditary conditions can also make individuals more prone to losing their hair cells. One of the first signs of damage to hair cells can be perceived as a ringing in the ears, which may progress to full-blown tinnitus. The damage to hair cells can ultimately lead to their death. Hearing loss that results from the death of hair cells can be gradual, as is commonly seen in the wear and tear of old age, or immediate and catastrophic, such as deafness resulting from explosions or gunshots. In either case, the hearing loss is permanent, as the lost hair cells are never replaced (Figure 2). Although humans cannot replace their lost hair cells, this is not true for other animals. It has been known for over 20 years that birds, frogs, and fish can regenerate their hair cells naturally. A bird that has been deafened by exposure to loud noises or drugs that kill hair cells is able to grow back almost its entire complement of hair cells and hear again almost perfectly

Figure 2. The surface of a mammalian cochlea (left), showing three rows of outer hair cells, and a single row of inner hair cells. On the right, a mammalian cochlea that has been damaged by sound. Many of the hair cells have been destroyed and will never regenerate, leading to permanent hearing loss. *Image courtesy of the House Ear Institute, Los Angeles.*

in a matter of weeks. However, mammals apparently lost the ability to regenerate hair cells at some point over the last 300 million years when the ancestors of mammals separated from the ancestors of modern birds and reptiles.

Hair Cell Regeneration

How are birds able to regenerate their hair cells? Every bird hair cell is surrounded by four to seven “supporting cells” to form a repeating mosaic pattern. The death of a bird hair cell somehow triggers one of its neighboring supporting cells to divide and produce two daughter cells. One of the daughter cells then turns into a hair cell, and this process of division followed by transformation into a hair cell restore the mosaic of hair cells and supporting cells back to its original state. This replenishment of cells by division and transformation happens all the time in many parts of the mammalian body, such as our skin, the bone marrow (which produces blood cells), and our digestive system (for example, we lose about 10 trillion cells from the lining of our guts every day, and will continue to do so until the day we die). Intriguingly, this process of division and transformation even occurs in parts of the adult mammalian brain thought to play a role in short-term memory. However, although mammals also have supporting cells surrounding their hair cells like birds, there is almost no regeneration of hair cells after damage (Figure 3).

Why are mammals unable to do something that birds do so well? There are a number of possible explanations. For example, mammalian supporting cells might simply have lost the ability to divide and make hair cells at some point during evolution. Alternatively, they might still be able to regenerate hair cells, but the signal to trigger regeneration might be blocked or missing. To test these possibilities, my lab collaborated with the lab of Dr. Neil Segil

Figure 3. When bird hair cells (HC) are killed, some of the supporting cells (SC) are triggered to divide (grey cells). One of the two daughter cells turns back into a hair cell, restoring the system back to normal. In damaged mammals, the supporting cells do not divide or make new hair cells.

at the House Ear Institute in Los Angeles, Calif., to carry out a conceptually simple but technically difficult experiment. We developed ways of purifying supporting cells from the ears of newborn mice, an age where all the cell types of the cochlea are formed and in place. We also devised ways of keeping the supporting cells alive in a culture dish to study their ability to divide and make hair cells. After several years of trial and error, we were able to show that newborn mouse supporting cells could behave like those of birds in a culture dish – they were able to divide, and their daughters were able to turn into hair cells.

We then started to ask why the mouse supporting cells were able to divide in a dish, but not in the intact mouse ear. Cells in our bodies have a whole battery of genes whose function is to stop cells from dividing when they are not meant to. These genes serve an extremely important role, as their malfunction can lead to inappropriate growth of cells, which can lead to cancer. Mouse supporting cells express one such growth-blocking gene with the rather unimaginative name of p27. We found that when we take supporting cells out of the mouse ear and grow them in a dish, about half the supporting cells switch off the p27 gene within 12 hours, and it is these supporting cells that start dividing. So far, so good – these experiments showed that mammalian supporting cells do have the capacity to divide and make new hair cells, just like those in birds, provided we put them in the right environment. However, all our experiments were done in newborn mice, and unlike humans, mice do not start to hear until about two weeks after birth. We decided to repeat our experiments in two week old mice that can hear and to our surprise, the purified supporting cells were now unable to switch off the p27 gene and unable to divide. Our results suggested that the ability to switch off the p27 gene seemed to correlate with the ability to divide. We settled the issue by genetically inactivating the p27 gene in 2 week old mice – and once again, their supporting cells were able to divide. Learning how to switch the p27 gene off in the right place at the right time might help our supporting cells divide and transform into hair cells. However, our experiments also tell us that the potential of the inner ear to regenerate hair cells changes quite quickly with age, and so understanding the biological basis of these changes will be very important for understanding how to regenerate hair cells.

Notch Signaling Pathways

If p27 is a gene that can control how supporting cells divide, are there genes that affect their ability to transform into hair cells? A number of labs, including our own, have been looking at an evolutionarily conserved way in which cells can communicate with each other, called the Notch signaling pathway. Notch signaling is frequently used to create mosaic patterns of different cell types, such as the mosaic repeating pattern of hair cells and supporting cells. Under this scheme, supporting cells make a protein on their cell surfaces – the Notch

receptor. Hair cells make proteins on their cell surfaces that bind to the Notch receptor, like a key fitting into a lock. Binding to the Notch receptor is believed to actively inhibit supporting cells from turning into hair cells. As a result, the pattern of hair cells and supporting cells remains stable in the cochlea.

If this model is correct, it suggests that blocking Notch signaling between hair cells and supporting cells would remove the barriers that normally prevent supporting cells from turning into hair cells. We and others have used both genetic and pharmacological approaches to block Notch signaling in the cochlea (Figure 4). We find that supporting cells can indeed actively transform

Figure 4. Hair cells make proteins on their surfaces that bind the Notch receptor on supporting cells. Activation of the Notch receptor blocks hair cells from becoming supporting cells. When Notch signaling is blocked, the supporting cells transform into new hair cells.

into hair cells when Notch signaling is blocked. In some experiments, we can turn at least 50% of supporting cells into hair cells. What is especially intriguing is that recent evidence from birds suggests that the Notch pathway is also employed when birds regenerate their hair cells, suggesting that manipulating this signaling pathway with drugs could conceivably be used to make new hair cells in mammals. However, as with the story of p27, the process is far from simple. We have recently shown that the ability to make new hair cells by blocking Notch signaling only works in very young mice, and not older animals. Once again, the mammalian cochlea seems to start out with an intrinsic capacity to repair itself, but this capacity disappears as the cochlea matures.

Conclusion

What has all this told us? We now believe that mammals may have the capacity for hair cell regeneration, but that this capacity is typically blocked. Our experiments have raised two possible targets – the p27 gene and the Notch signaling pathway – in our quest to trick the mammalian ear into behaving more like a bird's. However, we are a very long way from attempting to try such experiments in humans. We need to find answers to many more basic questions, such as how to trigger just the right amount of supporting cell division and transformation into hair cells. The cochlea is an extremely mechanically sensitive structure, and it is likely that producing too many hair cells may do more harm than good. Finally, our work tells us that the mammalian cochlea becomes less and less amenable to repair as it gets older. Understanding the basis of this maturation will help us design new targets for possible therapies. Despite these challenges, it's clear that birds are showing us the way ahead, and only time (and a lot more research) will tell if we will be able to follow their cue.

Andy Groves, Ph.D., is an associate professor in the Department of Neuroscience, Department of Molecular and Human Genetics and Program in Developmental Biology at the Baylor College of Medicine in Houston, Texas. He received his undergraduate degree from Sidney Sussex College in Cambridge and his Ph.D. from the University College of London. Before joining Baylor in 2008, Dr. Groves was a group leader at the House Ear Institute in Los Angeles, Calif. Prior to researching ear system development, Dr. Groves' work focused on the glial and neural progenitors in the central nervous system.

What We Can Do with Stem Cells and What We Cannot Do By Stefan Heller, M.S., Ph.D.

Millions of patients are diagnosed with hearing loss, which is mainly caused by degeneration of sensory hair cells in the cochlea. The underlying reasons for hair cell loss are highly diverse, ranging from genetic disposition, drug side effects, and traumatic

noise exposure to the effects of aging. Whereas modern hearing aids offer some relief for individuals with mild hearing loss, the only viable option for patients who have a severe to profound hearing loss is the cochlear implant. Despite their successes, hearing aids and cochlear implants are not perfect. Particularly, frequency discrimination, performance in noisy environments, and general efficacy of the devices vary among individual patients. The advent of regenerative medicine, the publicity of stem cells and gene therapy, and recent scientific achievements in inner ear cell regeneration have created an emerging spirit of optimism among scientists, health care practitioners, and patients. Here, I introduce the different types of stem cells that are being utilized by scientists in the field of hearing research. I will explain some of their most important features and attempt to illustrate the limitations of stem cell technology and the state of current research as well as future directions.

Introduction

In the time that it takes you to read this sentence, your body has lost about 1,000,000 cells, which are already replaced by now. So, nothing to worry about.

Your body's remarkable regenerative capacity is the result of a highly controlled population of stem cells that bestow most of your organs with the striking ability of continuous regeneration. The organs maintain shape during this process, which is even more remarkable. Unfortunately, our regenerative capacity is not perfect – it slows down as we get older and, as a result, our bodily functions decline with increasing age. To make matters worse, some of our organs have only very limited or no regenerative capacity. New neurons in the human brain, for example, are only rarely regenerated, and sensory hair cells in the cochlea or retinal photoreceptors do not regenerate at all.

Responsible for this regenerative potential are somatic stem cells, which are the workforce keeping us going. Each organ with regenerative capacity maintains its own specific type of stem cells. Placing the stem cells into a virtual cage, a so-called niche, allows the organ to control the powerful growth potential (Li and Clevers, 2010). Uncontrolled stem cell growth can result in tumors. Therefore, it is very important to keep the cells in check throughout our long lives. Regeneration is not a novel concept, but the dream for the fountain of youth has been fueled in the last decade by numerous claims that stem cells will provide novel therapies for many ailments and that they can fix many diseases for which no cures exist.

The ongoing public debate characterized by misinformation as well as political and religious bias, combined with public and private funding frenzies, has not helped in providing the public with independent resources about stem cells. Even for the informed layperson, it is often difficult to assess what we as scientists and clinicians are able to do with these cells. In the following paragraphs, I would like to introduce several different types of stem cells and explain what

makes them so distinctively special and different from each other. Toward the end of this article, I will try to put into the context of hearing loss what we know about stem cells and how this knowledge could culminate in the development of novel therapies. I will discuss some of the roadblocks that exist with regard to stem cell applications to the inner ear, but I will also attempt to predict how future generations might benefit from current research.

Different Kinds of Stem Cells

Probably the most well-known type of stem cells are *embryonic stem (ES) cells* that can be isolated from the inner cell mass of blastocysts, which are one of the very early stages of embryonic development. Mouse or human ES cells are pluripotent, which means that they are able to become any tissue in the body. In the case of mouse ES cells, it is routine practice to genetically modify the cells. Mice generated from ES cells with mutations that are comparable to corresponding human mutations often show a similar hearing loss phenotype to human patients. This strategy allows researchers to use transgenic mice as animal models to study particular forms of hearing loss.

Because of their ability to generate every cell type of the body, ES cells have raised a tremendous amount of hope that they can be used to generate replacement cells for degenerative diseases. However, only a single clinical trial has emerged from nearly a decade of research on human ES cells (Alper, 2009). In 2009, the FDA approved the use of high-purity human ES cell-derived glial cells for the treatment of spinal cord injuries. Glia is the connective tissue of the nervous system, and ES cell-derived glia has been shown to improve function (leg movement) in paralyzed rats. The reason why this clinical trial was approved by the FDA is that human ES cells can be differentiated in the culture dish into glia cells with high efficiency, and that the glia cells can be purified so that no pluripotent ES cells are being transplanted. This is very important because human or mouse ES cells, when transplanted, are highly tumorigenic (Figure 5). To suppress this potentially “evil” side of ES cells, it is important that stringent safety measures are employed before we prematurely jump to human trials, no matter which disease we are dealing with.

A potentially safer variant than ES cells are *adult stem cells*, which are also called *somatic stem cells*. These cells can be isolated from a regenerating tissue or organ, further purified, and then transplanted into the damaged organ of a sick patient. Blood-forming stem cells are routinely used for human bone marrow transplants and they have saved the lives of many leukemia patients. Ligament and tendon injuries or chronic diseases, such as arthritis in animals, are more and more commonly treated with somatic stem cells. In these still experimental treatments, somatic stem cells are isolated from the animals’ fat. Fat stem cell populations are not pure, but they do not have tumorigenic potential, which allows them to be safely used in veterinary medicine. The stem cells are then reintroduced into the same animal at the site of the injury.

Figure 5. Embryonic stem (ES) cells are pluripotent and can give rise to any tissue in the body. They are also tumorigenic, which means that they grow into cell masses called teratoma when transplanted into a recipient. The population of cells shown in the center form when ES cells are subjected to specific guidance protocols. If these protocols are efficient, most ES cells have differentiated along a certain pathway and are committed to this (i.e. they are unable to fall back into the pluripotent state). This stage is useful for laboratory experiments, but for clinical settings the cells need to be further purified and tumorigenic and ES cell remnants need to be 100% removed. Current translational research with such cells often focuses on improving efficiency and ensuring/improving safety.

Because the cells came from the recipient, they have virtually no risk of rejection and there is no need to suppress the function of the recipient's immune system. In transplantation terms, this procedure is called an allograft. Adult stem cells, such as bone marrow and fat-derived stem cells, are much less controversial than ES cells because they can be obtained without the destruction of an embryo.

A few years ago, a Kyoto University research team developed a method to convert simple connective tissue cells from mice and humans into pluripotent stem cells. These *induced pluripotent stem (iPS) cells* are combining the benefits from ES cells and patient-derived adult stem cells. Human iPS cells can be generated from a patient's skin cells, allowing for the generation of allografts – the recipient is receiving her or his own cells without need for immunosuppression. The generation of iPS cells initially required the use of viruses and the introduction of the c-Myc gene that can cause cancer. Safety concerns arising from the use of these tools were confirmed when researchers found that mice generated from mouse iPS cells showed an increased tendency to develop tumors when compared to mice generated from ES cells. Nevertheless, several new approaches are being developed by numerous laboratories around the world to create iPS cells without using viruses and introduction of potentially harmful genes. It appears that this hurdle will be taken in 2010, which will open the door for bringing iPS cell technology into the clinical practice. I will introduce a use for iPS and ES cells in the context of inner ear cell regeneration in the last section of this article.

An interesting line of inner ear research focuses on *inner ear stem cells*. The phrase “inner ear stem cells” is somewhat paradoxical because the human cochlea is not able to regenerate lost hair cells. Nevertheless, birds and fish, and presumably all other non-mammalian vertebrates, are able to regenerate hair cells throughout life. When the inner ears of these animals were examined closer, it turned out that the supporting cells left behind after hair cells die have the ability to divide into new hair cells and supporting cells. This regenerative capability is preserved throughout life, and even repeated deafening is no challenge for this powerful regeneration mechanism. The cells in the inner ear of animals that have this regenerative capacity are the so-called supporting cells that remain quiescent (i.e. not dividing) unless the loss of a hair cell triggers them to come to life. When stem cell terminology is applied to the inner ears of non-mammalian vertebrates, it becomes very obvious that the supporting cells are the stem cells and that the surrounding hair cells are providing the niche that controls the supporting cells’ regenerative abilities (Figure 6).

It turns out that evolution has, for some unknown reason, sacrificed the regenerative capacity of mammalian supporting cells. We describe this circumstance as a loss of stemness because the supporting cells of the mouse or human inner ear are still present but they lack regenerative potential. Interestingly, the loss of the regenerative capability of the mammalian inner ear is not complete. In the adult balance organs of rodents, for example, a small number of supporting cells still have the ability to divide and to regenerate a few hair cells after drug-induced hair cell loss (Forge et al., 1993; Warchol et al., 1993). Based on this original observation, our laboratory went on to isolate proliferating supporting cells from adult mouse balance epithelia, and we demonstrated in 2003 that the isolated cells have *bona fide* stem cell characteristics (Li et al., 2003) (Figure 6). Several research groups, including our own, subsequently showed that the supporting cells of the cochlea of newborn mice still have proliferative potential, or stemness (Oshima et al., 2007;

Figure 6. The defining feature of a stem cell is that it is able to self-renew. This means that when the stem cell is activated, it divides and generates a replacement cell as well as an identical copy of itself. Applied to the inner ear of non-mammalian vertebrates, such a mechanism exists when a supporting cell asymmetrically divides into a replacement hair cell and a new supporting cell. The inner ears of non-mammalian vertebrates as well as the vestibular sensory epithelia of mammals, therefore, harbor *bona fide* stem cells.

White et al., 2006). Unlocking this feature, however, required completely destroying the mouse cochlea and isolating the cells away from their surrounding cells. Again, applying stem cell terminology to this situation suggests that the niche exerts a strong suppression of the stemness of cochlear supporting cells. Unfortunately, this stemness appears to be completely gone in animals older than three weeks, suggesting not only a strong suppressing function exerted by the stem cell niche, but also that the supporting cells lose their stem cell features when they mature after birth.

Current State of Research

Ongoing research in several laboratories is focusing on two main strategies that utilize stem cell technology. The first one aims to explore the potential of stem cells *in vitro* with the goal of generating precursor cells, for example from ES or iPS cells that show commitment to become only inner ear cells but that lack the pluripotency and the tumorigenicity of plain ES and iPS cells. Transplantation studies only make sense with cells that are safe as well as cells that have been proven to generate appropriate inner ear cell types. Research is just beginning to advance in this direction and in 2010, for the first time, it has been shown that ES and iPS cell-generated hair cells are indeed functional. These *in vitro*-generated hair cells are mechanosensitive and they have the appropriate morphology that unequivocally identifies them as inner ear hair cells (Oshima et al., 2010).

The second strategy is highly hypothetical but it involves the fact that ES and iPS cell generated inner ear cell types and inner ear tissues display many of the characteristics of mouse or human inner ears. For example, we assume that stem cell-generated mammalian supporting cells will behave similarly as their native counterparts, which means that these cells will not be able to regenerate lost hair cells. Arguing that mammalian supporting cells evolved from supporting cells that back in time had stem cell characteristics leads us to further presume that some of the cellular pathways that confer stemness might still be present. ES and iPS cells are an unlimited source for *in vitro* generation of supporting cells and we predict that such cell populations will be used in future drug discovery to identify novel drugs with the ability to confer regenerative capacity. These drugs could be candidates for further studies to regenerate cochlear hair cells and to ameliorate hearing loss.

Finally, we are hopeful that transplantation experiments could become more and more successful – at least in appropriate animal models. Methods now exist to guide ES and iPS cells toward inner ear cell fates and isolated inner ear stem cells have been shown to generate hair cell-like cells after transplantation into embryonic inner ears. We are quite certain that the transplanted cells are able to become hair cells. At the same time, the accumulated knowledge of cochlear function, the complex micromechanical organization, and the discovery that almost every part of the cochlea is essential for proper

function presents a plethora of new challenges for cochlear hair cell regeneration (Brigande and Heller, 2009). It is not clear whether transplanted cells will be able to find the correct place to integrate and restore function. What happens, for example, when some of the cells become situated at locations where their presence interferes with cochlear mechanics? Will new hair cells be fully integrated with the nervous system? How long will new hair cells be able to survive?

At the moment, many challenges remain but it is promising to note that we have at least identified and defined many of the persisting issues, which allows us to systematically and specifically address roadblocks to the development of potential treatment options. All this requires three things that we all have so little of: groundbreaking ideas, time, and money – but nobody said that it would be easy.

References

- Alper, J. (2009). Geron gets green light for human trial of ES cell-derived product. *Nat Biotechnol*, 27, 213–214.
- Brigande, J.V., & Heller, S. (2009). Quo vadis hair cell regeneration? *Nat Neurosci*, 12, 679–685.
- Forge, A., Li, L., Corwin, J.T., & Nevill, G. (1993). Ultrastructural evidence for hair cell regeneration in the mammalian inner ear. *Science*, 259, 1616–1619.
- Li, H., Liu, H., & Heller, S. (2003). Pluripotent stem cells from the adult mouse inner ear. *Nat Med*, 9, 1293–1299.
- Li, L., & Clevers, H. (2010). Coexistence of quiescent and active adult stem cells in mammals. *Science*, 327, 542–545.
- Oshima, K., Grimm, C.M., Corrales, C.E., Senn, P., Martinez Monedero, R., Geleoc, G.S., et al. (2007). Differential distribution of stem cells in the auditory and vestibular organs of the inner ear. *J Assoc Res Otolaryngol*, 8, 18–31.
- Oshima, K., Shin, K., Diensthuber, M., Peng, A.W., Ricci, A.J., & Heller, S. (2010). Mechanosensitive hair cell-like cells from embryonic and induced pluripotent stem cells. *Cell*, in press.
- Warchol, M.E., Lambert, P.R., Goldstein, B.J., Forge, A., & Corwin, J.T. (1993). Regenerative proliferation in inner ear sensory epithelia from adult guinea pigs and humans. *Science*, 259, 1619–1622.
- White, P.M., Doetzlhofer, A., Lee, Y.S., Groves, A.K., & Segil, N. (2006). Mammalian cochlear supporting cells can divide and trans-differentiate into hair cells. *Nature*, 441, 984–987.

Stefan Heller, M.S., Ph.D., is a professor and director of research in the Departments of Otolaryngology – Head & Neck Surgery and Molecular & Cellular Physiology at Stanford University School of Medicine in Stanford, Calif. He received his Ph.D. in genetics from the Johannes Gutenberg University in Mainz, Germany, and completed his post-doctoral training at the Rockefeller University in New York, N.Y.

Initiating Cell Regeneration in the Mammalian Cochlea

By Jian Zuo, Ph.D.

Exposure to chemotherapy, antibiotics, and loud noise often causes loss of sensory hair cells in the cochlea of the inner ear, leading to permanent hearing loss in humans. While mammals cannot replace damaged sensory hair cells, chickens, fish, and amphibians can by division of neighboring supporting cells which then adopt sensory hair cell characteristics. Is it possible to give a mammal the chicken's ability to regenerate auditory hair cells? We propose a three-step working model for regeneration of mammalian cochlear hair cells in vivo that involves hair cell damage, supporting cell division, and cell fate change (or transdifferentiation) to hair cells. Recent advances in mouse genetics provide an unprecedented avenue to test this model. We first focused on the neonatal mouse cochlea because neonatal tissue is generally more plastic and regenerative than adult tissue. Since it is difficult to damage hair cells in the neonatal mouse cochlea with ototoxic drugs or noise, we have instead adopted a novel genetic method to kill hair cells by expressing a toxin specifically in hair cells beginning at birth. To further induce proliferation of supporting cells, we have deleted one of two key cell cycle regulators specifically in neonatal supporting cells, the retinoblastoma (Rb) protein and p27^{Kip1}. Finally, to make new hair cells, we are manipulating several transcription factors (i.e., Atoh1) in neonatal supporting cells. Our studies complement other strategies for hair cell regeneration in the mammalian cochlea and provide promising therapeutic avenues to restore hearing.

Hearing Loss and Causes

Cochlear hair cells (HCs) of the inner ear are sensory receptors that transduce sound into electrical signals. Many genetic and environmental factors can cause irreversible damage to auditory HCs. Children are particularly prone to such damage. Hearing loss afflicts approximately 2 to 3 in 1,000 infants, and 1 in 1,000 infants are born deaf. This can occur as a consequence of developmental abnormalities or exposure to antibiotics, noise, or chemotherapy.

Cisplatin, in particular, is a potent chemotherapy drug that is used to treat a variety of tumors. In addition to its anti-tumor effects, cisplatin causes many adverse effects that may limit the amount of doses that one can safely take, thus preventing the patient from completing therapy. This toxicity results in destruction of HCs, leading to permanent hearing loss. This acute dose-limiting toxicity may also have a long-term adverse effect on the ability of the child to perform in school. The mechanisms underlying cisplatin and antibiotic-induced hearing loss are unclear (Zhang et al., 2003; Cheng et al., 2005), although otoprotective measures can be effective in preventing hearing loss, but only to a certain extent.

Similarly, noise-induced hearing loss is becoming more and more prominent in the digital age, especially among adolescents. There are also many sources of potentially damaging noise in military settings, including weapons systems, vehicle engines, aircraft, ships, and explosions. Even the best hearing protection devices are only partially effective.

Because many genetic and environmental causes of hearing loss are uncontrollable and unavoidable, it would be useful to regenerate functional HCs after damage as an alternative to protective measures and cochlear implants.

HC Regeneration in Mammals and Nonmammals

In mammals, including humans, auditory HCs cannot regenerate spontaneously. However, other vertebrates such as birds, fish, and amphibians can regenerate their lost sensory HCs by proliferation and transdifferentiation of neighboring supporting cells (SCs) (Ryals & Rubel, 1988; Corwin & Oberholtzer, 1997). In chickens, the loss of HCs evokes SC proliferation within approximately 200 μm of the site of damage after roughly 16 hours, and SC division further give rise to both HCs and SCs (Warchol & Corwin, 1996). Resembling embryonic development of HCs and SCs, HC regeneration in mature non-mammalian vertebrates requires three distinct types of molecules: 1) signaling molecules released from damaged HCs or immune cells attracted to the damage site, 2) proliferating factors that trigger SCs to come out of the quiescent state to enter the cell cycle, and 3) differentiating factors that allow the new cells to transdifferentiate into HCs.

Several differences between mammals and non-mammals could explain their differing regenerative capacities. First, young and mature HCs may have different responses to HC damage. Second, SCs may have differential expression of cell cycle regulatory genes in mammals and non-mammals. And third, differentiation factors and pathways may be different in mammals and non-mammals. Recently, considerable progress has been made in inducing mammalian HC regeneration. Ectopic expression of *Atoh1* in a subset of SCs outside the organ of Corti produced HCs in postnatal rat cochlear culture (Zheng & Gao, 2000). In addition, virus-mediated transduction of *Atoh1* in adult guinea pigs with damaged HCs remarkably induced partially functional HCs, although the mechanisms for HC regeneration have yet to be determined (Izumikawa et al., 2005). Recently, isolated SCs from postnatal mouse cochleas were found to divide and transdifferentiate into HCs in vitro (White et al., 2006).

Based on how Mother Nature regenerates HCs in non-mammals, it is reasonable to propose a three-step working model for HC regeneration in mammals: first, HCs are damaged; second, SCs divide; and third, SCs are subsequently converted into HCs (Figure 7). In this model, the proliferative responsiveness of postmitotic, quiescent SCs holds the key for HC regeneration in the mammalian cochlea.

Figure 7. A working model for hair cell regeneration in the mammalian cochlea. In the neonatal or adult mouse cochlea, both HCs (dark grey) and SCs (light grey) are post-mitotic and quiescent; they are believed to share same progenitors during embryonic development. When HCs are damaged (black cross) by ototoxic drugs or noise, SCs will be stimulated to divide and proliferate. Appropriate factors will convert the proliferating SCs into HCs.

Mouse Models of Hair Cell Regeneration

The study of HC regeneration in mouse models is promising. The mouse inner ear is anatomically and physiologically similar to the human inner ear. Unlike other mammalian species (such as cat, rat, guinea pig, and chinchilla), the genetics of the mouse can be easily manipulated – deleting a gene in a specific cell type at a particular time during the life-span of the mouse is feasible. In particular, recently developed Cre-loxP technology (Figure 8) allows the precise deletion of genes in HCs or SCs at neonatal and adult ages (Weber et al., 2008; Yu et al., 2010). For example, the use of *Atoh1*-CreER allows the removal of genes with nearly 100% efficiency in neonatal cochlear HCs (Weber et al., 2008). *Prox1*-CreER allows deletion of genes specifically in two types of SCs within the organ of Corti: pillar and Deiters' cells that are located directly underneath cochlear HCs (Yu et al., 2010). Compared to viral transduction of cochlear cells (Minoda et al., 2007), the use of Cre-loxP restricts gene alteration in a reproducible manner to specific cell types, some of which are not accessible to the virus. This ability to genetically manipulate genes in cochlear cells in a spatially and temporally controlled manner has great potential to help us understand the genetic pathways that are important for HC regeneration in mammals.

However, the study of HC regeneration using mouse models also remains a challenge. By approximation, the cochlea of a new-born mouse resembles that of a human fetus in the third trimester; the cochlea of a newborn human is similar to that of a three-week old mouse. Therefore, HC regeneration should be best studied in mouse models at adult ages. However, the capacity of isolated SCs to proliferate and change into HCs significantly decreases with age (White et al., 2006). It is therefore sensible to first study the neonatal cochlea, which is likely more plastic and regenerative than the adult cochlea. If HC regeneration is successful in mouse models at neonatal ages, we can then

Figure 8. Cre-loxP technologies for gene manipulation in HCs and SCs of the neonatal mouse cochlea. Top panel: The HC-specific, inducible Cre mouse in which HCs are labeled with the lacZ reporter. Modified from (Weber et al., 2008). Bottom panel: The SC-specific, inducible Cre mouse where SCs are labeled with a fluorescent protein. HCs are labeled with a myosin VIIa antibody; nuclei are labeled with Hoechst. Right panel: cross section showing HCs and two types of SCs: pillar and Deiters' cells (labeled with arrows and arrowheads). Modified from (Yu et al., 2010).

expand what we've learned to adult ages. It is still possible that additional or different mechanisms are required for HC regeneration in the adult mammalian cochlea.

Damaging HCs in the Neonatal Mouse Cochlea In Vivo

Damage to HCs in adult mice by ototoxic drugs or noise is relatively easy (Hirose et al., 2005; Oesterle & Campbell, 2009). In the neonatal mouse, however, this is extremely difficult, if not impossible, because ototoxic drugs are lethal to neonatal mice and because mice do not hear until two weeks after birth. To overcome such technical difficulties, we have adopted a novel technology that allows expression of a toxin (DTA) specifically in neonatal mouse cochlear HCs. Expression of DTA induces cell death that is similar to HC death caused by ototoxic drugs. In DTA mice, we observe rapid HC loss within two weeks. It is our hope that such induced HC death in the neonatal mouse cochlea would stimulate SC proliferation and possibly regeneration of HCs.

SC Proliferation In Vivo by Inactivating Cell Cycle Inhibitors

Neonatal mouse cochlear SCs and HCs do not normally divide. Can we induce proliferation of SCs *in vivo* by removing genes that block cell cycle entry? We have focused on inactivating several key cell cycle inhibitors, the retinoblastoma (*Rb*) protein and p27^{Kip1}.

Rb Inactivation in SCs

Rb was the first tumor suppressor identified (Friend et al., 1986; MacPherson et al., 2004). The deregulation of the *Rb* pathway is a hallmark of most, if not all, human cancers. Many inactivating mutations and deletions within *Rb* are known. *Rb* is expressed prominently in embryonic HC and SC precursors, postnatal HCs, and moderately postnatal SCs (Mantela et al., 2005; Sage et al., 2005; Sage et al., 2006). Conditional deletion of *Rb* in HC and SC precursors resulted in proliferation of cochlear HCs and SCs followed by subsequent HC and SC death (Mantela et al., 2005; Sage et al., 2005; Sage et al., 2006). We have acutely deleted *Rb* in postnatal HCs and observed similar HC cell cycle reentry, followed by HC death (Weber et al., 2008). These studies together provide evidence that postmitotic HCs can reenter the cell cycle by inactivating *Rb*, although subsequent cell death is inevitable. Thus inactivating *Rb* in residual HCs in partially damaged cochleae may not represent a promising avenue to regenerate HCs. On the other hand, deletion of *Rb* in two subtypes of neonatal SCs (pillar and Deiters' cells) causes both SC types to reenter the cell cycle (Figure 9). However, only pillar cells can complete the cell cycle and proliferate for multiple divisions (Yu et al., 2010).

The fact that *Rb*-negative HCs and Deiters' cells cannot complete the cell cycle and proliferate, whereas *Rb*-negative pillar cells can, may indicate that *Rb* plays essential roles in other phases of the cell cycle in postmitotic HCs

Figure 9. Proliferation of SCs after *Rb* deletion in the neonatal mouse cochlea at postnatal day 4. Two dividing SC nuclei are labeled with a cell cycle marker (phosphohistone 3) antibody; HCs (three rows of outer hair cells and one row of inner hair cells) are labeled with myosin VIIa antibody. Modified from (Yu et al., 2010).

and Deiters' cells but not in postmitotic pillar cells. Alternatively, pillar cells may be less mature or differentiated, and thus more able to proliferate than HCs and Deiters' cells at this postnatal age. In this regard, it is tempting to hypothesize that pillar cells are more "stem cell-like" and thus can more easily reenter the cell cycle and proliferate. Such striking heterogeneity among various types of SCs is also supported by in vitro studies of isolated SCs (White et al., 2006) and the different pathways for differentiation and maintenance of Deiters' cells versus pillar cells (Doetzlhofer et al., 2009). In addition, a recent study found differential expression of a cell cycle protein, cyclin D1, between neonatal pillar and Deiters' cells, which may contribute to the ability of pillar cells to proliferate (Laine et al., 2009).

Despite proliferation of SCs in vivo in our model, SCs eventually died followed by subsequent HC loss (Yu et al., 2010). It is unknown whether death of proliferating SCs in this model is primarily caused by *Rb* deletion or by the lack of space and nutrition needed for rapidly proliferating cells in the precise architecture of the organ of Corti. It remains to be further determined whether an intrinsic mechanism exists for proliferating *Rb*-negative pillar cells to stop dividing. It is also known that SCs play key roles in providing trophic and mechanical support for HCs, which may be the reason that HCs died in this model. Moreover, during postnatal development, efferent fibers also initiate their transition from inner to outer HC areas through SCs.

p27^{Kip1} Inactivation in SCs

p27^{Kip1} is a cell cycle regulator which prevents proliferation. Isolated SCs from the postnatal mouse cochlea can down-regulate the cell cycle inhibitor, p27^{Kip1}, proliferate, and transdifferentiate into HCs in vitro (White et al., 2006). In the adult guinea pig cochlea, forced expression of Skp2, an enzyme that enhances the degradation of p27^{Kip1}, led to proliferation of non-sensory cells outside the organ of Corti (Minoda et al., 2007). Moreover, in p27^{Kip1} germline knockout mice, postnatal pillar cells, but not Deiters' cells, were found to reenter the cell cycle at P6 (Chen & Segil, 1999). It would be therefore interesting to determine whether deletion of p27^{Kip1} in neonatal mouse SCs within the organ of Corti would lead to proliferation and transdifferentiation into HCs in vivo.

Similarity of SC Proliferation Between Mammals and Non-Mammals

An interesting feature of our findings is that a majority of dividing SCs were observed near and in parallel to the luminal surface of the sensory epithelium. These proliferating SCs in the models appear to undergo cell cycle and migratory changes similar to those in the regenerating SCs in non-mammals (Stone & Cotanche, 2007). Our findings, therefore, suggest that mammalian SCs follow similar mechanisms to avian SCs to induce cell division and nuclear migration, which is the first stage of HC regeneration.

Transdifferentiation of SCs into HCs in Mammals

Unfortunately, no new HCs were detected in proliferating SCs in our *Rb*-negative mouse models (Weber et al., 2008; Yu et al., 2010). Failure of newly generated SCs to transdifferentiate or change cell fate into HCs may be due to their inability to reestablish a quiescent state since these cells now lack *Rb*; therefore, transient deletion of *Rb* in SCs may represent a better strategy for HC regeneration. In addition, several potential factors may facilitate transdifferentiation into HCs. *Atoh1* is a transcription factor that is both necessary and sufficient for the differentiation of HCs in the mammalian cochlea (Bermingham et al., 1999; Zheng & Gao, 2000; Chen et al., 2002; Shou et al., 2003; Izumikawa et al., 2005; Gubbels et al., 2008). Inhibition of other genes, such as *Ids*, *Prox1*, and *Sox2*, may up-regulate *Atoh1* in *Rb*-negative proliferating SCs (Jones et al., 2006; Khidr & Chen, 2006; Kirjavainen et al., 2008). Therefore, a combination of transient *Rb* deletion and *Atoh1* activation or inactivation of *Ids*, *Prox1*, or *Sox2* may represent an effective measure to regenerate cochlear HCs. Future development of small molecules or drugs that can remove the block for SC proliferation and/or enhance the transdifferentiation into HCs will further translate discoveries in mouse models into therapeutic applications for human patients.

Acknowledgement

The author acknowledges the contribution of the members in the Zuo lab and support from the National Institutes of Health (DC06471, DC05168, DC008800, 1F31DC009393, 1F32DC010310, and CA21765), Office of Naval Research (N000140911014), Sir Henry Wellcome Postdoctoral Fellowship, National Organization for Hearing Research Foundation (NOHR), and the American Lebanese Syrian Associated Charities (ALSAC) of St. Jude Children's Research Hospital. Dr. Zuo is a recipient of The Hartwell Individual Biomedical Research Award.

References

- Bermingham N.A., Hassan B.A., Price S.D., Vollrath M.A., Ben-Arie N., Eatock R.A., et al. (1999). Math1: an essential gene for the generation of inner ear hair cells. *Science*, 284, 1837–1841.
- Chen P., & Segil N. (1999). p27(Kip1) links cell proliferation to morphogenesis in the developing organ of Corti. *Development*, 126, 1581–1590.
- Chen P., Johnson J.E., Zoghbi H.Y., & Segil N. (2002). The role of Math1 in inner ear development: Uncoupling the establishment of the sensory primordium from hair cell fate determination. *Development*, 129, 2495–2505.
- Cheng A.G., Cunningham L.L., & Rubel E.W. (2005). Mechanisms of hair cell death and protection. *Curr Opin Otolaryngol Head Neck Surg*, 13, 343–348.

- Corwin J.T., & Oberholtzer J.C. (1997) Fish n' chicks: Model recipes for hair-cell regeneration? *Neuron*, 19, 951–954.
- Doetzlhofer A., Basch M.L., Ohyama T., Gessler M., Groves A.K., & Segil N. (2009). Hey2 regulation by FGF provides a Notch-independent mechanism for maintaining pillar cell fate in the organ of Corti. *Developmental Cell*, 16, 58–69.
- Friend S.H., Bernards R., Rogelj S., Weinberg R.A., Rapaport J.M., et al. (1986). A human DNA segment with properties of the gene that predisposes to retinoblastoma and osteosarcoma. *Nature*, 323, 643–646.
- Gubbels S.P., Woessner D.W., Mitchell J.C., Ricci A.J., & Brigande J.V. (2008). Functional auditory hair cells produced in the mammalian cochlea by in utero gene transfer. *Nature*, 455, 537–541.
- Hirose K., Discolo C.M., Keasler J.R., & Ransohoff R. (2005). Mononuclear phagocytes migrate into the murine cochlea after acoustic trauma. *J Comp Neurol*, 489, 180–194.
- Izumikawa M., Minoda R., Kawamoto K., Abrashkin K.A., Swiderski D.L., Dolan D.F., et al. (2005). Auditory hair cell replacement and hearing improvement by Atoh1 gene therapy in deaf mammals. *Nat Med*, 11, 271–276.
- Jones J.M., Montcouquiol M., Dabdoub A., Woods C., & Kelley M.W. (2006). Inhibitors of differentiation and DNA binding (Ids) regulate Math1 and hair cell formation during the development of the organ of Corti. *J Neurosci*, 26, 550–558.
- Khidr L., & Chen P.L. (2006). RB, the conductor that orchestrates life, death and differentiation. *Oncogene*, 25, 5210–5219.
- Kirjavainen A., Sulg M., Heyd F., Alitalo K., Yla-Herttuala S., Moroy T., et al. (2008). Prox1 interacts with Atoh1 and Gfi1, and regulates cellular differentiation in the inner ear sensory epithelia. *Developmental Biology*, 322, 33–45.
- Laine H., Sulg M., Kirjavainen A., & Pirvola U. (2010). Cell cycle regulation in the inner ear sensory epithelia: Role of cyclin D1 and cyclin-dependent kinase inhibitors. *Developmental Biology*, 337(1), 134–46.
- MacPherson D., Sage J., Kim T., Ho D., McLaughlin M.E., & Jacks T. (2004). Cell type-specific effects of Rb deletion in the murine retina. *Genes Dev*, 18, 1681–1694.
- Mantela J., Jiang Z., Ylikoski J., Fritzsche B., Zacksenhaus E., & Pirvola U. (2005). The retinoblastoma gene pathway regulates the postmitotic state of hair cells of the mouse inner ear. *Development*, 132, 2377–2388.
- Minoda R., Izumikawa M., Kawamoto K., Zhang H., & Raphael Y. (2007). Manipulating cell cycle regulation in the mature cochlea. *Hearing Research*, 232, 44–51.
- Oesterle E.C., & Campbell S. (2009). Supporting cell characteristics in long-deafened aged mouse ears. *J Assoc Res Otolaryngol*, 10, 525–544.
- Raphael Y. (1992). Evidence for supporting cell mitosis in response to acoustic trauma in the avian inner ear. *Journal of Neurocytology*, 21, 663–671.

- Ryals B.M., & Rubel E.W. (1988). Hair cell regeneration after acoustic trauma in adult *Coturnix* quail. *Science*, 240, 1774–1776.
- Sage C., Huang M., Vollrath M.A., Brown M.C., Hinds P.W., Corey DP, et al. (2006). Essential role of retinoblastoma protein in mammalian hair cell development and hearing. *Proc Natl Acad Sci*, 103, 7345–7350.
- Sage C., Huang M., Karimi K., Gutierrez G., Vollrath M.A., Zhang D.S., et al. (2005). Proliferation of functional hair cells in vivo in the absence of the retinoblastoma protein. *Science*, 307, 1114–1118.
- Shou J., Zheng J.L., & Gao W.Q. (2003). Robust generation of new hair cells in the mature mammalian inner ear by adenoviral expression of *Hath1*. *Molecular and Cellular Neurosciences*, 23, 169–179.
- Stone J.S., & Cotanche D.A. (2007). Hair cell regeneration in the avian auditory epithelium. *International Journal of Developmental Biology*, 51, 633–647.
- Tsue T.T., Watling D.L., Weisleder P., Coltrera M.D., & Rubel E.W. (1994). Identification of hair cell progenitors and intermitotic migration of their nuclei in the normal and regenerating avian inner ear. *J Neurosci*, 14, 140–152.
- Warchol M.E., & Corwin J.T. (1996) Regenerative proliferation in organ cultures of the avian cochlea: Identification of the initial progenitors and determination of the latency of the proliferative response. *J Neurosci*, 16, 5466–5477.
- Weber T., Corbett M.K., Chow L.M., Valentine M.B., Baker S.J., & Zuo J. (2008). Rapid cell-cycle reentry and cell death after acute inactivation of the retinoblastoma gene product in postnatal cochlear hair cells. *Proc Natl Acad Sci*, 105, 781–785.
- White P.M., Doetzlhofer A., Lee Y.S., Groves A.K., & Segil N. (2006). Mammalian cochlear supporting cells can divide and trans-differentiate into hair cells. *Nature*, 441, 984–987.
- Yu Y., Weber T., Yamashita T., Liu Z., Valentine M.B., et al. (2010). In vivo proliferation of postmitotic cochlear supporting cells by acute ablation of the retinoblastoma protein in neonatal mice. *J Neurosci*, 30(17), 5927–5936.
- Zhang M., Liu W., Ding D., & Salvi R. (2003). Pifithrin- α suppresses p53 and protects cochlear and vestibular hair cells from cisplatin-induced apoptosis. *Neuroscience*, 120, 191–205.
- Zheng J.L., & Gao W.Q. (2000). Overexpression of *Math1* induces robust production of extra hair cells in postnatal rat inner ears. *Nat Neurosci*, 3, 580–586.

Jian Zuo, Ph.D., is a member/professor of the Department of Developmental Neurobiology at St. Jude Children's Research Hospital in Memphis, Tenn. He received his undergraduate and graduate degrees in biomedical engineering from Huazhong University of Science and Technology in China, and his Ph.D. in physiology from the University of California, San Francisco. Dr. Zuo is a recipient of The Hartwell Individual Biomedical Research Award.

Directory of Professional Programs

The Alexander Graham Bell Association for the Deaf and Hard of Hearing is not responsible for verifying the credentials of the service providers below. Listings do not constitute endorsements of establishments or individuals, nor do they guarantee quality. If you are interested in listing your program, please contact Gary Yates, manager of advertising, exhibit sales and sponsorships, at gyates@agbell.org or (202) 204-4683.

California

Sprint Relay, 2455 Naglee Road, Suite 179, Tracy, CA 95304 • Phone: 866-540-4657 • E-mail: Chameen.r.stratton@sprint.com • Websites: www.sprintrelay.com/800i; www.sprintcaptel.com.

Conversations flow freely with Sprint's Captioned Telephone Services! Read conversations real-time while speaking and listening to callers!

CapTel 800i

Telephones with large displays that, when connected to high speed internet & phone line, display captions of calls. www.sprintrelay.com/800i.

Sprint WebCapTel

No specialized phone equipment needed. If you have a computer with high speed internet and any telephone, you are ready to read captions on calls! www.sprintcaptel.com.

Illinois

Expanding Children's Hearing Opportunities (ECHO) Program at Carle Foundation Hospital, 611 West Park Street, Urbana, IL 61801 • Phone: (217) 383-4375 • E-mail: echo@carle.com • Website: www.carle.org/ECHO • The Expanding Children's Hearing Opportunities (ECHO) Program was established in 1989 to serve children with hearing loss and their families. ECHO has grown to encompass two programs: the Pediatric Hearing Center (PHC) and Carle Auditory Oral School (CAOS). PHC provides audiologic, speech language and early intervention services as well as an experienced pediatric cochlear implant team. CAOS supports children with hearing loss in developing their spoken language and listening skills from preschool through second grade.

New York

Nazareth College Deafness Specialty Preparation Program – Rochester, NY (joint program with National Technical Institute for the Deaf) • Point of Contact: Paula Brown, PhD, CCC-SLP • Email: pbrown8@naz.edu • Phone: 585-389-2796 • Website: <http://www.rit.edu/ntid/spslp/>. The Nazareth College Deafness Specialty Preparation Program in Rochester, NY, prepares graduate students in Speech-Language Pathology for work with children who are deaf or hard of hearing, especially those with cochlear implants. This joint program with the National Technical Institute for the Deaf provides specialized coursework and practica in spoken language and auditory assessment and intervention.

North Carolina

FIRST YEARS – Kathryn Wilson, Program Director • Phone: 919-966-0103 • Email: kathryn_wilson@med.unc.edu • Administered by the University of North Carolina-Chapel Hill, **FIRST YEARS** is an in-service, certificate program committed to enhancing the knowledge and skills of professionals practicing in deaf education, speech-language pathology, audiology, and early intervention. The **FIRST YEARS** courses in listening and spoken language development combine convenient distance education with a hands-on mentorship experience to meet the needs of practicing professionals.

Australia

University of Newcastle – GradSchool, GradSchool Services Building, University of Newcastle, Callaghan, NSW 2308, Australia • Phone: 61249217373 • Fax: 61249218636 • Master of Special Education, distance education through the University of Newcastle. The program provides pathways through specialisations in:

- Generic special education
- Emotional disturbance/behaviour problems
- Sensory disability
- Early childhood special education

The Master of Special Education (Sensory Disability specialisation) is available through the Renwick Centre, which is administered by the Australian Royal Institute for Deaf and Blind Children. Program information and application is via GradSchool: www.gradschool.com.au, +61249218856, or email gs@newcastle.edu.au.