Journal of Experimental Biology and Agricultural Sciences

<http://www.jebas.org>

ISSN No. 2320 – 8694

EVALUATION OF THE EFFECT OF ELEVATED CO₂ ON BIOEFFICACY OF BUPROFEZIN INSECTICIDE AGAINST BROWN PLANT HOPPER, *Nilaparvata lugens* (STÅL)

Srinivasa N^{1,2}, Subhash Chander^{2*}, Padala Vinod Kumar³

¹Department of Entomology and Agricultural Zoology, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh-221005

²Division of Entomology, Indian Agricultural Research Institute, New Delhi-110012

³ICAR-Research Complex for Eastern Region, Research Centre for Makhana, Darbhanga Bihar-846007

Received – October 22, 2020; Revision – December 21, 2020; Accepted – February 03, 2021

Available Online – February 27, 2021

DOI: [http://dx.doi.org/10.18006/2021.9\(1\).5.11](http://dx.doi.org/10.18006/2021.9(1).5.11)

KEYWORDS

Brown plant hopper

Buprofezin

Elevated CO₂

Rice

Spray volume

climate change

ABSTRACT

The effect of elevated CO₂ (570±25ppm) on the brown plant hopper (BPH) population, rice yield parameters, and efficacy of buprofezin (0.05%) in terms of spray volume was studied in an open top chamber (OTCs) during rainy season 2017 and 2018. The pest population was observed to be higher during 2017 compared to the rainy season of 2018. Under elevated CO₂, rice plants had more vegetative tillers (18%) and reproductive tillers (22.1%), but there was a decrease in 1000-seed weight (11.2%), seed number per panicle (3.91%), and grain yield (18.8%) in comparison to ambient CO₂ grown rice plants. The spray volumes of 700, 600, 500, and 400 l/ha each caused higher BPH mortality under ambient CO₂ compared to elevated CO₂. A spray volume of 500 l/ha did not prove as effective under elevated CO₂ as under ambient CO₂. Lower efficacy of spray volume of 500 l/ha under elevated CO₂ could be ascribed to higher canopy size under elevated CO₂ due to higher tillering. Increased crop canopy size under elevated CO₂ may thus require higher spray volume to ensure proper coverage. Results of the study suggested a need to revise spray volume recommendations to facilitate effective management of BPH under climate change.

* Corresponding author

E-mail: schanderthakur@gmail.com (Subhash Chander)

Peer review under responsibility of Journal of Experimental Biology and Agricultural Sciences.

Production and Hosting by Horizon Publisher India [HPI]
(<http://www.horizonpublisherindia.in/>).
All rights reserved.

All the articles published by [Journal of Experimental Biology and Agricultural Sciences](http://www.jebas.org) are licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by/4.0/) Based on a work at www.jebas.org.

1 Introduction

Rice (*Oryza sativa* L.) is one of the important cereals of the world particularly in Asian countries and forms a staple diet for more than 50 percent of the population (Khush, 2004). It is grown on 33.3% of the area with 41% of total food grain production in India and with the productivity of 2404 kg/ha (DAC, 2016-17). Among important insect pests of rice, brown plant hopper, (BPH) *Nilaparvata lugens* (Stål) is a serious pest, wherein both nymphs and adults suck the phloem sap from the plants causes yellowing and drying of plants results “hopper burn”(Figure 1a). Besides, it also causes harm indirectly as a vector for rice grassy stunt and ragged stunt diseases of rice (Anjaneyalu, 1974; Zheng et al., 2014). Before the green revolution, BPH was a minor pest, but, post 1970 it attained major pest status due to the growth of susceptible high yielding varieties (HYVs) of rice and indiscriminate use of insecticides that killed natural enemies of BPH in Asia (Dale and Kenneth, 2012). In the northern part of India also, it caused several outbreaks since 2008 (Chander & Hussain, 2018). Further, many studies indicate that under elevated CO₂ the BPH problem will be aggravated (Pandi et al., 2017 and 2019).

According to the Intergovernmental panel on climate change (IPCC) the earth's temperature has already increased by 0.74°C between 1906 and 2005 and CO₂ is expected to increase up to 445–640 ppm by 2050 due to an increase in anthropogenic emissions of greenhouse gases (IPCC, 2007). Climate change will have an impact on insects through temperature, which affects the insect's pests directly while CO₂ affects them indirectly through host plants and natural enemies (Netherer & Schopf, 2010). Under elevated CO₂, an increase in carbon: nitrogen (C:N) ratio in plants resulted in the accumulation of carbon that forced herbivores to feed more to acquire the required amount of amino acids (Coviella & Trumble, 2000; Rao et al., 2009), which in turn inflicted more crop damage as compared to ambient CO₂. Although under elevated CO₂ and a combination of elevated CO₂ and temperature, rice plants produced more vegetative and reproductive tillers and grains per panicle but suffered higher yield loss due to a higher BPH population (Prasannakumar et al., 2012; Pandi et al., 2018). Besides affecting insect population, climate change is expected to have a significant influence on pest management tactics such as host plant resistance (Guo et al., 2012 and 2017), biological control (Coll & Hughes, 2008; Yin et al., 2009) and insecticide efficacy. Many studies have been mostly concentrated on the impact of climate change on insect pests (Sudderth et al., 2005; Reddy et al., 2010; Prasannakumar et al., 2012) while the information on the impact of climate change on the efficacy of insecticides is meager. Delcour et al. (2015) opined that increased pesticide use is expected in the form of higher doses, frequency, and different products under climate change scenarios. Pandi et al. (2019) reported that the spray of 400 and 500 l/ha was not sufficient to

manage the BPH under elevated temperature and CO₂ with imidacloprid 17.8SL @ 0.006%. Insecticides that are effective against insect pests but relatively innocuous to non-target organisms including biological control agents are known as biorationals. Among these, an insect growth regulator (IGR) buprofezin is a promising one against BPH compared to neonicotinoids (Uchida et al., 1985; Hegde & Nidagundi, 2009; Shivashankar & Gowda, 2015). Keeping this in view, this biorational pesticide was evaluated in the present study against BPH under elevated CO₂ conditions to determine future requirements with regards to its spray volume.

2 Materials and Methods

2.1 Maintenance of Rice plants

The effect of different spray volumes of buprofezin 25SC(0.05%) against BPH under elevated CO₂ (570±25ppm) *vis-à-vis* ambient CO₂ (400±25ppm) was investigated on potted rice plants kept in open top chambers (OTCs) during rainy season 2017 and 2018 at the Division of Plant Physiology, ICAR-Indian Agricultural Research Institute, New Delhi (28.6377° N, 77.1571° E). Twenty-two days old seedlings of scented rice Pusa Basmati 1121 were transplanted in 100 plastic pots (22.5 X 15 cm²) on 20th July and 18th July during 2017 and 2018, respectively, and raised further by following recommended agronomic practices.

2.2 Plant exposure to CO₂, BPH infestation and insecticide

The study was undertaken in two OTCs one having ambient CO₂ (400±25ppm) and the other had elevated CO₂ (570±25ppm). Seedlings were transplanted in 100 pots and 15 days after transplanting (DAT) 50 pots were kept in each OTCs. The CO₂ was released in OTC during day time only between 9.30 AM to 4.30 PM until the maturation of paddy grains as only during daytime plants utilize CO₂ for photosynthesis. Under elevated CO₂, potted rice plants were exposed to elevated CO₂ (570±25ppm) for 15 days before the release of 5 pairs of laboratory reared BPH adults in each potted plants to acquire infestation under both ambient and elevated CO₂ condition. The BPH population was monitored regularly in the OTCs to ensure the insecticide application at peak incidence of the pest. The BPH population was found to peak after 45 days of its release in OTCs and thus treated at this stage. The insecticidal treatments comprised four spray volumes *viz.*, 400, 500, 600, and 700 l/ha along with untreated control (spray with plain water), each treatment have 10 replicates each. The different spray volumes of buprofezin (0.05%) were applied using hand held mist sprayer and observations on BPH mortality were recorded on three, six, and ten days after the application as it's being an IGR acts only at the nymphal stage of the pest (Uchida et al., 1985; Alam & Das 2017). The plant growth and yield parameters such as vegetative tiller number, reproductive

Figure 1(a) Hopper burn symptoms on rice caused by rice planthoppers in Delhi: 1(b) Difference rice yield parameters under elevated and ambient CO₂

tiller number, 1000-seed weight, and seeds per panicle were recorded across the treatments. BPH mortality was compared among different spray volumes both under ambient and elevated CO₂. However, yield parameters were pooled across the spray volumes and compared broadly between ambient and elevated CO₂.

2.3 Statistical Analysis

Pre spray BPH populations across different treatments were averaged. The pre-spray BPH populations and BPH mortality with different spray volumes of buprofezin (0.05%) under ambient and elevated CO₂ were analysed using one-way analysis of variance (ANOVA) and Tukey's honest significant difference (HSD), while the effect of elevated CO₂ on growth and yield parameters of BPH infested plants was analysed through t-test.

3 Results

3.1 Effect of elevated CO₂ on BPH population

The BPH population at peak infestation (45 days after infestation) was found to be higher in different treatments during rainy season 2017 both under elevated (135.3±25.9 to 239±28.68/hill) and ambient CO₂ (115.6±12.3 to 278.2±34.02/hill) compared to rainy season 2018 with respective peak populations ranging from 75.3±9.9 to 93.8±12.5/hill and 76.5±9.20 to 91.6±11.98/hill. The pest population was thus non-significantly higher under elevated CO₂ than the ambient CO₂ during 2017 but did not differ between two CO₂ levels during 2018 (Table 1).

3.2 Effect of elevated CO₂ on rice yield parameters

Different yield parameters of BPH-infested rice plants differed significantly under ambient and elevated CO₂ during both the years of study (Table 2). Pooled data of two years indicated that rice plants grown under elevated CO₂ had more number vegetative tillers (18%) and reproductive tillers (22.1%) than ambient CO₂ (Table 2). However, grain yield (18.8%), seed number per panicle (3.91%), and 1000-seed weight (11.29%) were observed to be lower under elevated CO₂ than ambient CO₂ (Table 2; Figure 1b).

3.3 Effect of elevated CO₂ on bio-efficacy of buprofezin insecticide

During 2017, the BPH mortality data revealed that buprofezin (0.05%) at 3 days after spraying (DAS) resulted in very low BPH mortality under both ambient and elevated CO₂ that remained around 18% under both the CO₂ conditions with all the spray volumes (Table 3). Likewise, during the rainy season, 2018 buprofezin (0.05%) resulted in poor mortality (≤13%) at 3DAS under both ambient and elevated CO₂ with all the spray volume (Table 4)

At 6 DAS also, the mortality was low and was observed to be 31 and 26% under ambient and elevated CO₂ conditions respectively with 700 l/ha spray volume during 2017 (Table 3). Similarly, the mortality remained low at 6 DAS during 2018 and was recorded to be 27 and 24% under ambient and elevated CO₂ condition, respectively with 700 l/ha spray volume (Table 4)

However, at 10 DAS, buprofezin (0.05%) caused the highest mortality with a spray volume of 700 l/ha (45.6±2.5%) followed by 600 (42.2±1.99%), 500 (38.7±4.64%), and 400 (31.8±2.71%) l/ha under ambient CO₂ compared to corresponding mortalities under elevated CO₂ as 40.61±3.50, 36.92±2.49, 32.65±1.9 and 28.44±2.16 % during the rainy season 2017 respectively (Table 3). The same trend was observed during 2018 and buprofezin (0.05%) at 10 DAS inflicted the highest mortality with a spray volume of 700 l/ha (44.2±2.4%) followed by 600 (37.93±2.5%), 500 (33.50±2.29%), and 400 (21.75±1.5%) l/ha under ambient CO₂ compared to corresponding mortalities under elevated CO₂ as 41.25±1, 33.51±1.9, 25.04±1.7 and 20.68±.59 during rainy season 2018 respectively (Table 4)

Table 1 Brown planthopper (BPH) peak population (Number/hill) prior to buprofezin (0.05%) application

Treatment	No. of BPH/hill during 2017		No. of BPH/ hill during 2018	
	Ambient CO ₂ (400 ± 25 ppm)*	Elevated CO ₂ (570 ± 25 ppm)*	Ambient CO ₂ (400 ± 25 ppm)*	Elevated CO ₂ (570 ± 25 ppm)*
400 l/ha	191.9±23.04 ^{ab}	194.8±25.33 ^{ab}	79.3± 5.7 ^a	88.3±10.21 ^a
500 l/ha	278.2±34.02 ^a	239±28.68 ^a	76.5±9.20 ^b	91.8±5.51 ^a
600 l/ha	115.6±12.36 ^b	220.8±21.72 ^{ab}	91.6±11.98 ^a	79.5±4.1 ^a
700 l/ha	220.2±16.57 ^a	208.1±19.37 ^{ab}	81.1±9.83 ^a	75.3±9.9 ^a
Control	198.3±20.69 ^{ab}	135.3±25.59 ^b	80.8±11.36 ^a	93.8±12.5 ^a
Average	200.8±23.4	199.6±15.7	81.8±2.2	85.74±3.20
F value	6.070	2.688	.227	1.42
P value	0.001	0.043	0.889 ^{NS}	0.264 ^{NS}

Brown planthopper counts with different lowercase letter in the same column differ significantly at P=0.05%, NS:Nonsignificant. * mean of ten replicates.

Table 2 Rice growth and yield parameters under elevated and ambient CO₂

Parameter	Rainy season 2017			Rainy season 2018			Average	
	Ambient CO ₂ * (400 ± 25 ppm)	Elevated CO ₂ * (570 ± 25 ppm)	t-Statistics	Ambient CO ₂ * (400 ± 25 ppm)	Elevated CO ₂ * (570 ± 25 ppm)	t-Statistics	Ambient CO ₂ (400 ± 25 ppm)	Elevated CO ₂ (570 ± 25 ppm)
No. of tillers per hill	20.3±0.53	24.1±0.79	t= -3.7(p=0.001)	17.2±0.72	19.7±1.11	t= -1.7(p=0.103) ^{NS}	18.79±1.5	22.18±1.9
No. of reproductive tillers per hill	18.3±0.42	22.5±0.55	t= -5.7(p<.0001)	16±0.61	18.8±1.10	t= -2.1 (p=0.050)	17.15±1.1	20.95±1.5
1000 seed weight (g)	19.7±0.28	16.9±0.37	t= 5.7(p<.0001)	17.4±0.38	15.8±0.38	t= 2.9 (p=0.009)	18.51±1.8	16.42±0.5
Seeds per panicle	79.12±0.85	73.27±1.36	t= 3.18(p=0.005)	80.8±2.75	75.9±2.50	t=1.04(p=0.310) ^{NS}	77.50±3.3	74.58±3.9
Yield per hill (g)	27.2±0.30	23.1±0.62	t= 5.6(p<.0001)	25.2±0.38	20.8±0.52	t= -6.40 (p<0.001)	26.21±0.9	22.05±1

Yield parameters were mean±SE, significant at 5% level of significance *: average of ten replicates. NS, Non-significant

Table 3 Mean BPH mortality (%) at different intervals after buprofezin (0.05%) application during rainy season 2017

Treatment	3 DAS		6 DAS		10 DAS	
	Ambient CO ₂ * (400 ± 25 ppm)	Elevated CO ₂ * (570 ± 25 ppm)	Ambient CO ₂ * (400 ± 25 ppm)	Elevated CO ₂ * (570 ± 25 ppm)	Ambient CO ₂ * (400 ± 25 ppm)	Elevated CO ₂ * (570 ± 25 ppm)
400 l/ha	13.59±1.14 (21.45 ^a)	11.15±2.58 (18.57 ^b)	19.41±1.8 (25.84 ^b)	18.50±2.35 (25.08 ^b)	31.79±2.71 (34.03 ^b)	28.44±2.16 (32.05 ^b)
500 l/ha	14.91±1.78 (22.33 ^a)	11.83±1.19 (19.89 ^{ab})	24.03±2.87 (28.96 ^{ab})	20.90±1.65 (27.02 ^{ab})	38.72±4.64 (38.17 ^{ab})	32.65±1.95 (34.75 ^{ab})
600 l/ha	17.64±1.80 (24.59 ^a)	14.88±1.37 (22.45 ^{ab})	27.88±1.6 (31.76 ^a)	24.07±1.92 (29.18 ^{ab})	42.20±1.99 (40.46 ^{ab})	36.92±2.49 (37.27 ^{ab})
700 l/ha	18.61±1.63 (25.36 ^a)	17.15±1.01 (24.37 ^a)	31.36±2.38 (33.70 ^a)	26.18±2.09 (30.61 ^a)	45.62±2.53 (42.44 ^a)	40.61±3.50 (39.46 ^a)
Control	0.72±0.22 (3.67 ^b)	1.78±0.37 (7.27 ^c)	2.46±0.38 (8.73 ^c)	3.53±0.56 (10.52 ^c)	3.51±0.53 (10.45 ^c)	5.61±1.02 (13.18 ^c)
F value	50.91	25.31	47.12	37.24	50.08	45.69
P Value	P<0.0001	P<0.0001	P<0.0001	P<0.0001	P<0.0001	P<0.0001

DAS: Days After Spray. *= average of ten replicates (mean±SE) mortality data with different lowercase letters in same column differ significantly. Data in parenthesis are arc sin transformed values. Arc sign transformed values were used in ANOVA followed by Tukey HSD test at 5% significance.

Table 4 Mean BPH mortality (%) at different intervals after buprofezin (0.05%) application during rainy season 2018

Treatment	3DAS		6DAS		10DAS	
	Ambient CO ₂ (400 ± 25 ppm)	Elevated CO ₂ (570 ± 25 ppm)	Ambient CO ₂ (400 ± 25 ppm)	Elevated CO ₂ (570 ± 25 ppm)	Ambient CO ₂ (400 ± 25 ppm)	Elevated CO ₂ (570 ± 25 ppm)
400 l/ha	7.10±0.85 (15.33 ^b)	7.10±0.59 (15.37 ^b)	13.92±1.38 (21.76 ^b)	12.62±1.10 (20.70 ^c)	21.75±1.54 (27.72 ^c)	20.68±0.59 (27.03 ^c)
500 l/ha	8.08±0.65 (16.44 ^{ab})	9.04±0.87 (17.36 ^{ab})	18.68±1.08 (25.55 ^{ab})	15.34±0.47 (23.04 ^{bc})	33.50±2.29 (35.30 ^b)	25.04±1.71 (29.96 ^{bc})
600 l/ha	10.80±1.26 (19.01 ^{ab})	10.78±1.02 (19.08 ^{ab})	23.73±3.13 (28.91 ^a)	20.45±1.98 (26.71 ^{ab})	37.93±2.59 (37.96 ^{ab})	33.51±1.96 (35.31 ^{ab})
700 l/ha	13.08±1.44 (21.07 ^a)	12.31±0.45 (20.51 ^a)	27.03±2.03 (31.24 ^a)	24.53±0.91 (29.65 ^a)	44.2±2.42 (41.67 ^a)	41.25±1.50 (39.93 ^a)
control	0.93±0.40 (4.20 ^c)	1.396±0.35 (5.98 ^c)	3.18±0.58 (10.01 ^c)	2.023±0.15 (8.14 ^d)	4.56±0.72 (12.13 ^d)	2.76±0.40 (9.44 ^d)
F value	25.16	28.80	29.92	67.14	66.10	51.85
P Value	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001

DAS: Days After Spray. * = average of ten replicates (mean±SE) mortality data with different lowercase letters in same column differ significantly. Data in parenthesis are arc sin transformed values. Arc sign transformed values were used in ANOVA followed by Tukey HSD test at 5% significance.

4 Discussion and Conclusions

Increased pesticide use is being projected in the form of higher doses and frequencies to effectively combat crop pests under climate change scenarios (Chen & McCarl, 2001). An increase in temperature, ultraviolet radiation (UV), and decrease in relative humidity may render plant protection techniques like natural plant products, entomopathogenic viruses, bacteria, fungi, nematodes, and synthetic pesticides less effective because these are highly sensitive to environmental variation (Sharma, 2014) and such effects would be more pronounced on biopesticides (Isman, 1997). In the present study, elevated CO₂ increased the number of tillers and reproductive tillers that resulted in increased rice crop canopy, and the pest population was recorded to be non-significantly higher under elevated CO₂ than ambient CO₂ during 2017. Earlier studies have reported a higher BPH population under elevated CO₂ (Prasannakumar et al. 2012) and under the combination of elevated CO₂ and temperature (Pandi et al., 2019) than ambient CO₂, because of a more favourable microenvironment for the pest multiplication. In the present study, the pest-induced yield loss was observed to be higher under elevated CO₂ than ambient CO₂ mainly due to the higher sucking rate of the pest under elevated CO₂ as the pest population did not differ significantly under two CO₂ levels. In previous studies, higher yield loss due to BPH under elevated CO₂ was ascribed to both higher pest population and sucking rate than under ambient CO₂ (Prasannakumar et al., 2012; Pandi et al., 2019). Buprofezin (0.05%) proved to be most effective against the pest at a spray volume of 700 l/ha followed by 600, 500, 400 l/ha under both ambient and elevated CO₂ conditions. However, BPH mortality always happened to be higher under ambient than elevated CO₂ at respective spray volumes. The buprofezin application did not produce significant mortality under both the

CO₂ conditions at 3 and 6 DAS which could be perhaps ascribed to insect growth regulatory (IGR) activity of buprofezin, which is slower compared to neurotoxic insecticides, which act faster resulting in the instant kill of the pest. Similarly, slow action of buprofezin against plant hoppers of rice has also been reported earlier (Uchida et al., 1985; Shivashankar & Gowda, 2015; Alam & Das, 2017). A spray volume of 500 l/ha, which is currently recommended for the management of rice insect pests did not prove to be effective under elevated CO₂. Ineffectiveness of spray volume of 500 l/ha could be ascribed to increased canopy size due to increased tillering under elevated CO₂ that resulted in increased canopy circumference. Increased crop canopy might thus require higher spray volume to ensure proper coverage against the pest. In this context, earlier also pesticides are more effective against insect pests at higher spray volumes under elevated CO₂ and temperature in the case of imidacloprid (0.006%) against BPH (Pandi et al., 2019) and at ambient conditions in the case of *Beauveria bassiana* against western flower thrips, *Frankliniella occidentalis* (Ugine et al., 2007), methoxyfenozide or fenpropathrin against grape berry moth, *Paralobesia viteana* (Wise et al., 2010) and buprofezin (15%) + acephate (35%) against rice plant hoppers (Ghoshal et al., 2018). Another study also indicated the reduced efficacy of triazophos under elevated CO₂ against BPH (Ge et al., 2015). The current study revealed that elevated CO₂ may have a nutritive effect on rice yield through increased tillering and greater canopy size thereby turning the microenvironment more favourable for the pest. Consequently, the presently recommended spray volume may not suffice to ensure proper spray coverage of foliage to safeguard the crop against insect pests. There may thus be a need to revise spray volume recommendations to facilitate effective management of BPH under changing climatic conditions.

Acknowledgements

We are thankful to Dr. Madan Pal Singh, Division of Crop Physiology, ICAR-Indian agricultural research Institute, New Delhi-1100012 for providing OTCs facility to conduct research and field workers Naresh Ji, and Ashok Ji for help during studies.

Conflict of Interest

Authors declare that they have no conflict of interest

References

- Alam MJ, Das G (2017) Laboratory evaluation of buprofezin on the mortality of brown planthopper, *Nilaparvata lugens* (Delphacidae: Hemiptera). *Journal of Entomology and Zoology Studies* 5(6):2172-2175.
- Anjaneyalu A (1974) Identification of grassy stunt, a new virus disease of rice in India. *Current Science* 43:416-417.
- Chander S, Husain M (2018) Brown plant hopper (BPH) outbreak in rice: analysis of weather parameters during outbreak and non-outbreak years. *Indian Journal of Entomology* 3: 677-682.
- Chen CC, McCarl BA (2001) An investigation of the relationship between pesticide usage and climate change. *Climate Change* 61:475-487.
- Coll M, Hughes L (2008) Effects of elevated CO₂ on an insect omnivore: a test for nutritional effects mediated by host plants and prey. *Agriculture, Ecosystems & Environment* 123: 271-279.
- Coviella CE, Trumble JT (2000) Effect of elevated atmospheric carbon dioxide on the use of foliar application of *Bacillus thuringiensis*. *Biocontrol* 45 (3):325-336.
- Department of Agriculture, Cooperation and Farmers welfare (2016) Annual Report 2016-17. Ministry of Agriculture and Farmer's welfare GOI, Pp.3
- Dale GB, Kenneth GS (2012) Resurrecting the ghost of green revolutions past: The brown planthopper as a recurring threat to high-yielding rice production in tropical Asia. *Journal of Asia-Pacific Entomology* 15(1): 122-140.
- Delcours I, Spanoghe P, Uyttendaele M (2015) Literature review: impact of climate change on pesticide use. *Food Research International* 68: 7-15.
- Ge LQ, Wu JC, Sun YC, Ouyang F, Ge F (2013) Effects of triazophos on biochemical substances of transgenic Bt rice and its nontarget pest *Nilaparvata lugens* (Stål) under elevated CO₂. *Pesticide Biochemistry and Physiology* 107: 188-199.
- Ghoshal A, Dolai AK, Chatterjee ML (2018) Evaluation of a new ready mixed insecticide (Buprofezin 15% + Acephate 35% WP) against BPH (*Nilaparvata lugens* (Stal)) and WBPH (*Sogatella furcifera* (Horvath)) of rice in West Bengal. *Journal of Crop and Weed* 14(1): 205-211.
- Guo H, Han S, Ge F (2017) Effect of elevated CO₂ and O₃ on phytohormone-mediated plant resistance to vector insects and insect-borne plant viruses. *Science China Life Sciences* 60 (8): 816-825.
- Guo H, Sun Y, Ren Q, Zhu-Salzman K, Kang L, et al. (2012) Elevated CO₂ reduces the resistance and tolerance of tomato plants to *Helicoverpa armigera* by suppressing the JA signaling pathway. *PLoS ONE* 7(7) :e41426. doi: 10.1371/journal.pone.0041426.
- Hegde M, Nidagundi J (2009) Effect of newer chemicals on planthoppers and their mirid predator in rice. *Karnataka Journal of Agricultural Sciences* 22(3): 511-515.
- IPCC (2007) Summary for policy makers. In: Solomon et al. (eds) *Climate Change 2007: the Physical Science Basis. Contribution of Working Group I to the IV Assessment Report of the Intergovernmental Panel on climate Change*, Cambridge University Press Pp.1-8
- Isman MB (1997) Neem and other botanical pesticides: Barriers to commercialization. *Phytoparasitica* 25: 339-344
- Khush GS (2004) Harnessing science and technology for sustainable rice-based production systems. In *FAO Rice conference*, 12-13 February, Rome Italy, Pp13. Available at <http://www.fao.org/rice2004/en/pdf/khush.pdf>.
- Netherer S, Schopf A (2010) Potential effects of climate change on insect herbivores general aspects and a specific example (Pine processionary moth, *Thaumetopoea pityocampa*). *Forest ecology and Management* 259:831-838.
- Pandi GGP, Chander S, Singh MP, Pathak H (2018) Impact of Elevated CO₂ and temperature on Brown Planthopper Population in Rice Ecosystem. *Proceedings of national academy of sciences India, Section B Biological Sciences* 88:57-64.
- Pandi GGP, Chander S, Pal M (2017) Impact of elevated CO₂ on rice brown planthopper, *Nilaparvata lugens* (stal.). *Indian Journal of Entomology* 79 (1): 82-85.
- Pandi GGP, Chander S, Pal MP, Soumia PS and Sujithra M (2019) Imidacloprid efficacy against brown planthopper, *Nilaparvata lugens* under elevated carbon dioxide and temperature. *Current Science* 117(7) : 1199-1206. doi: 10.18520/cs/v117/i7

- Prasannakumar N R, Chander S, Singh M P (2012) Assessment of impact of climate change with reference to elevated CO₂ on rice brown planthopper, *Nilaparvata lugens* (Stål.) and crop yield. *Current Science* 103:1201-1205.
- Rao MS, Srinivas K, Vanaja M et al. (2009) Host plant (*Ricinus communis* Linn.) mediated effects of elevated CO₂ on growth performance of two insect folivores. *Current Science* 97:1047–1054.
- Reddy AR, Rasineni GK, Raghavendra AS (2010) The impact of global elevated CO₂ concentration on photosynthesis and plant productivity. *Current Science* 99: 46–57.
- Sharma HC (2014) Climate change effect on insects: Implications for crop protection and food security. *Journal of Crop Improvement* 28:229-259.
- Shivashankar T, Gowda DKS (2015) Evaluation of efficacy of insecticides against brown planthopper (BPH), *Nilaparvata lugens* (Stål) in rice ecosystem. *Environment and Ecology* 33(1):228-231.
- Sudderth EA, Stinson KA, Bazzaz FA (2005) Host-specific aphid population responses to elevated CO₂ and increased N availability. *Global Change Biology* 11: 1997–2008.
- Uchida M, Asai T, Sugimoto T (1985) Inhibition of cuticle deposition and chitin biosynthesis by a new insect growth regulator, buprofezin, in *Nilaparvata lugens* (Stål). *Agricultural and Biological Chemistry* 49(4):1233-1234.
- Ugine TA, Wright SP, Sanderson JP (2007) Effects of manipulating spray-application parameters on efficacy of the entomopathogenic fungus *Beauveria bassiana* against western flower thrips, *Frankliniella occidentalis*, infesting greenhouse impatiens crops. *Biocontrol Science and Technology* 17:193-219.
- Wise JC, Jenkins PE, Schilder AMC, Vandervoort C, Isaacs R (2010) Sprayer type and water volume influence pesticide deposition and control of insect pests and diseases in juice grapes. *Crop Protection* 29: 378–385.
- Yin J, Su NY, Wu G, Parajulee MN, Ge F (2009) Effects of elevated CO₂ on the population relationship between cotton bollworm, *Helicoverpa armigera* Hübner (Lepidoptera: Noctuidae), and its parasitoid, *Microplitis mediator* Haliday (Hymenoptera: Braconidae). *Agriculture Ecosystem and Environment* 132 (3): 267-275.
- Zheng L, Mao Q, Xie L, Wei T (2014) Infection route of rice grassy stunt virus, a tenuivirus, in the body of its brown planthopper vector, *Nilaparvata lugens* (Hemiptera: Delphacidae) after ingestion of virus. *Virus Research* 8:170-3. doi: 10.1016/j.virusres.2014.04.008.