AREAL GEOLOGY AND PETROLOGY OF THE IGNEOUS ROCKS OF THE SANTA ANA REGION, NORTHWEST SONORA

Guillermo A. Salas *

RESUMEN

Rocas metamórficas del Precámbrico fueron encontradas durante un estudio geológico de reconocimiento en el área de Santa Ana, en el norponiente de Sonora. Estas rocas pertenecen a las facies de esquistos verdes, de almandita-anfibolita y de granulita de metamorfismo regional, y en algunas localidades a las facies de corneana de hornblenda de metamorfismo de contacto. Las rocas de metamorfismo regional son las más comunes en el área estudiada y presenciaron, por lo menos, dos eventos metamórficos; uno durante el Precámbrico "Anterior" y otro durante el Precámbrico "Posterior" (grupo Coyotillo). Tanto los gneises como los esquistos forman las sierras con orientación regional de norte-sur y están yuxtapuestas a bloques fallados durante el Terciario temprano. El dinamometamorfismo relacionado con este fallamiento reconstituyó localmente los gneises y los esquistos en cataclasitas, milonitas y filonitas.

Rocas paleozoicas no afloran en el área. Carbonatos, grauvacas, lutitas y conglomerados fueron depositados en cuencas eugeosinclinales durante el Cretácico Temprano. La extravasación de andesitas, algunas de ellas probablemente submarinas, seguidas por levantamientos locales, erosión, plegamiento y fallamiento intensos, acompañados por la intrusión de rocas graníticas y volcánicas asociadas tuvieron lugar durante el Cretácico Tardío.

Las rocas terciarias de la Sierra de La Lumbre fueron cartografiadas en detalle. Estudios de campo, petrográfico y químico atestiguan tres períodos de actividad magmática (extravasación de andesitas, basaltos y material piroclástico félsico). Estas rocas no son comagmáticas pues el vulcanismo ha sido intermitente con etapas largas de intemperismo y erosión. Las rocas más abundantes son la andesita basáltica, brecha andesítica, andesita de hornblenda, basalto de olivino y toba litica. Sin embargo, propilita y ceniza volcánica localmente abundan.

El evento volcánico basáltico final, que consistió en la extravasación por fisuras durante el Plioceno tardío o el Pleistoceno y está representado por la Formación Comedores, cubre extensos depósitos terrígenos conglomeráticos terciarios (?) de la Formación Báucarit.

ABSTRACT

Reconnaissance mapping in the Santa Ana area, northwest Sonora, revealed the presence of Precambrian metamorphic rocks belonging to the greenschist, almandine-amphibolite,

* *Present address of the author: Sierra Gorda 12, México 10, D F.*

and granulite facies of regional metamorphism and local development of hornblende-hornfels facies of contact metamorphism. These rocks are the most abundant in the mapped area and represent, at least, two metamorphic events; one during the "Older" Precambrian and the other during the "Younger" Precambrian (Coyotillo group). All metamorphic rocks form north-south trending sierras bounded by Early Tertiary fault blocks. Dynamic metamorphism related to this faulting reconstituted the "Older" Precambrian gneisses and schists into locally developed cataclasites, mylonites, and phyllonites.

No Paleozoic rocks are exposed. Carbonates, graywackes, shales, and conglomerates were deposited in shallow eugeosynclinal basins during the Early Cretaceous. Eruption of andesites, some possibly under water, and later uplift, erosion, and intense folding and faulting, accompanied by emplacement of granites and associated volcanic rocks followed during the Late Cretaceous.

The Tertiary rocks of the Sierra de La Lumbre were mapped in detail. Three episodes of magmatic activity (eruption of andesitic lava, extrusion of basaltic magma, and extrusion of felsic pyroclastic material) were revealed by field, petrographic, and chemical studies. The rocks are not comagmatic and volcanism was intermittent with long periods of weathering and erosion. Basaltic andesite, andesite breccia, hornblende andesite, olivine basalt, and lithic-crystal tuff are most common in the Sierra de La Lumbre. However, propylite and volcanic ash are locally abundant.

A final basaltic volcanic event, consisting of fissure eruption during late Pliocene or Pleistocene time, represented by the Comedores Formation, produced a cover that overlies the extensive terrigenous conglomeratic deposits of the Tertiary (?) Báuçarit Formation.

INTRODUCTION

The mapped area lies in the Santa Ana Quadrangle, which takes its name from the town of Santa Ana in the southeastern part of the map. The area of approximately 1800 km² is limited by parallels 30°30' and 31°00' north latitude and by meridians 111°00' and 11°20' west longitude (Figure 1). Field work was done during the summer of 1966 and July of 1967. The geology was plotted on a topographic base map (scale 1:55,000, contour interval 50 m) with the aid of vertical aerial photographs.

The purpose of the present study is threefold: first, to present a general reconnaissance of the geology of the quadrangle; second, to describe and interpret the extrusive rocks of the Sierra de La Lumbre; and third, to compare the structural and petrogenic history of the quadrangle with nearby parts of north-west Mexico and southern Arizona.

ACCESSIBILITY.—The Santa Ana region is readily accessible. Federal Highway 45, which extends from Mexico City to Nogales, transects the southeastern part of the area, and Federal Highway 2, which extends from Santa Ana to Tijuana, crosses the entire southern portion of the quadrangle. The Sud-Pacífico Guadalajara-Nogales Railroad connects Santa Ana and Magdalena (Figure 1).

An unimproved rural road, which joins Highway 2 west of Santa Ana and passes through the villages of Coyotillo and La Cebolla, provides access to the north-central part of the area. A similar road, which connects Magdalena and Tubutama, gives easy access to the central and northwestern portions. Most of the region of the small ranches and ejidos are well served by all-weather dirt roads.

FIG. 1. Index map, showing the location of the mapped area.

PHYSIOGRAPHY.—The Santa Ana Quadrangle is located near the extreme eastern limit of the Sonoran Desert which corresponds to the Physiographic Province termed *Sierras Sepultadas* (Raisz, 1959), a southern extension of the Basin and Range Province.

The main orographic units, with summits as high as 1,400 m, are the Sierra del Rincón and its southern prolongations, the Sierra de Guacomca and the Sierra de Magdalena, which together constitute a north-south trending positive unit that marks the eastern limit of the area. To the north and west the Sierra de La Tortuga, the Sierra del Potrero, the Sierra de Las Jarillas, and the Sierra del Picacho also forming a massive positive unit trend in a north-south direction with peaks that reach more than 1,200 m. West of the Sierra de La Tortuga a relatively flat area surfaced by basalt has an average elevation of 750 m.

The large valley of the region is broader and higher to the north where the average elevation is 900 m, and descends southward toward the southern limit of the map where the average elevation is 635 m.

There are three main drainage systems: that of the Magdalena River which flows southwestward and drains the extreme southeastern part of the quadrangle; that of the Arroyo del Coyotillo, which describes a rough north-south arc convex to the west and drains the central portion of the area; and that of the Arroyo de Búsani which flows southwestward and drains the northern half of the region. All drainage is intermittent and carries water only after the severe thunderstorms which are characteristic of the area. The Magdalena River flows steadily during the rainy season but rarely carries water in winter.

CLIMATE, VEGETATION, AND LAND USE.—Thornthwaite (1943) described the region as having a very dry semi-arid climate with a deficiency of rainfall in all seasons. According to Koeppen, the climate is arid because the mean annual temperature is above 64°F, the winter is the dry season, and 70 percent of the precipitation occurs in the summer. Xerophytic brush, cacti, and a wide variety of desert grasses are the major vegetation groups.

The most important population center is the town of Santa Ana which has about 10,000 inhabitants. Its principal economic activity is trading whereas that of the region as a whole is cattle raising and agriculture. The latter is developed in the relatively wide floodplains of the Magdalena River and of the Arroyo del Coyotillo.

ACKNOWLEDGMENTS.—The author wishes to thank Dr. George T. Stone for his helpful suggestions and guidance in the field and in the office, and Dr. Clifford A. Merritt for his patience in revising and correcting thin section studies. Thanks are also due to Dr. William E. Ham who made illuminating comments on the limestone petrography, and to Dr. Charles J. Mankin who critically reviewed the manuscript. Ing. Jerjes Pantoja-Alor, of the Instituto de Geología of Universidad Nacional Autónoma de México helped the author in the field in many ways.

The field work was carried out with a four-wheel drive vehicle, placed at

the author's disposal by the Instituto de Geología, where also all illustrations have been prepared.

GENERAL GEOLOGY

PREVIOUS INVESTIGATIONS.—The only published work that treats any part of the Santa Ana Quadrangle is that of Flores (1927), who mapped and studied the rocks that crop out along the Sud-Pacifico Railroad between Santa Ana and Magdalena. However, more recently, a number of workers have published work which is relevant to the studied area.

Garfias and Chapin (1949), Eardley (1951), and de Cserna (1960 and 1961) studied the regional tectonic aspects of north-central Sonora. Stratigraphic and paleontologic work has been done by Gloria Alencáster (1961) who also gave a summary of the Triassic paleogeography; by Silva (1961) and Miller (1961); and by Giletti and Damon (1961) and Damon, Livingston, and Giletti (1961). The latter workers established a geochronological sequence of events for the Precambrian of Arizona, Sonora, and the remainder of northwestern Mexico.

Regional mapping has been done by Aguilera (1896), Dumble (1900a, 1900b), Burckhardt (1923), Keller (1929), Taliaferro (1933), King (1934, 1939), Santillán (1936), Valentine (1936), Imlay (1939), Cooper and Arellano (1952) and others. Their work is important because it constitutes a framework within which parts of the present study have been made. An excellent summary of the economic geology of northwest Mexico was given by Wisser (1966).

NATURE AND DISTRIBUTION OF ROCK TYPES.—Many rock types collectively ranging in age from Precambrian to Recent (excluding Paleozoic) with a total thickness of more than 2,000 m of metamorphic, sedimentary, and igneous rocks are exposed in the area (Figure 2).

The prominent sierras contain well exposed sections of Precambrian metamorphic rocks that are characterized by distinctive facies changes.

The sedimentary rocks are of Early Cretaceous age and are exposed only in the southeastern and southwestern parts of the quadrangle. The volcanic rocks, which range in age from Oligocene to Pleistocene are found exposed in the northeastern, southeastern and northwestern portions of the map. The Baucarit Formation (upper Tertiary) is the only unit which has been formally named.

In describing the metamorphic and sedimentary units, the general lithology, distribution, geometry and thickness, age, and probable mode of origin are discussed in that order. The igneous rocks, exclusive of those of the Sierra de La Lumbre, are treated in the same manner. The latter group is considered separately.

FIG. 2. Composite stratigraphic column.

METAMORPHIC ROCKS

The metamorphic rocks are divided according to the rank or grade of their mineral assemblages and according to their inferred relative ages. Of the many classifications proposed for metamorphic mineral assemblages (for example, Eskola, 1939; Turner, 1948; Fyfe and others, 1958; Ramberg, 1952; Barth, 1962; and Turner and Verhoogen, 1960), the one outlined by Turner and Verhoogen is used here.

The metamorphic rocks of the Santa Ana Quadrangle generally represent facies of regional metamorphism of former sedimentary and igneous rocks including the greenschist facies, the almandine-amphibolite facies, and the granulite facies.

Rocks tentatively assigned to the hornblende-hornfels facies (containing andalusite) of contact metamorphism are locally associated with gneiss and intrusive bodies of granitic composition. Mylonites, cataclasites, and phyllonites are found along sheared and faulted zones. The principal mineral assemblages indicative of each facies are shown in Table 1.

ROCKS OF THE GREENSCHIST FACIES.—The rocks of this facies include the common products of low-grade regional metamorphism: marble, phyllite, metaconglomerate schist, quartzite, semi-schist and metagraywacke.

Because of their distinctive lithologic characteristics and age relationships, the metamorphic rocks of the south-central part of the quadrangle are described first. They are a thick sequence of thin-to medium-bedded metasediments which are cut by hornblende andesite dikes and quartz veins (Figure 2). They are divided into 8 units which collectively comprise what is here termed the Coyotillo group:

- Unit A: Cordierite-bearing quartzite (oldest)
- Unit B: Sericite-chlorite schist
- Unit C: Magnetite-rich sericite-chlorite phyllite
- Unit D: Metaconglomerate with intercalations of thin sandy layers.
- Unit E: Sericite-chlorite phyllite
- Unit F: Metagraywacke
- Unit H: Impure marble (youngest)

Detailed megascopic and petrographic descriptions of each unit appear in Appendix A.

The Coyotillo group crops out in an unnamed homoclinal sierra of roughly circular outline. Because of their isolated position and distinctive lithologies, these metamorphic rocks are considered to represent a different environment and age of metamorphism than the remainder of the metamorphic rocks in the area.

Rocks of the greenschist facies are exposed in the eastern front of the Sierra de La Tortuga, the southern parts of the Sierra de Las Jarillas and the Sierra del Picacho, and parts of the Sierra de Magdalena. In eastern Sierra de La Tortuga the most common types are phyllites in which epidote, biotite, and

TABLE 1.—Metamorphic facies; Santa Ana Quadrangle

SAMPLE	FACIES	PRESSURE-TEMP. CONDITIONS		TYPE OF METAM.	MINERAL		ASSEMBLAGES			
SP		Heinrich, 1956			Carbonate rocks	Aluminous (pelitic) rocks	Quartz Feldspar rocks	Basic rocks		
9-1, 9-2 9-3, 9-4 9-5, 4-1 22-3 22-6 24-2 38, 39 45, 48 55, 62, 63 154, 202	Greenschist	P	moderate to low	Regional	Calcite Quartz Epidote	Chlorite Biotite Muscovite Almandine moderate T and P	Albite Quartz Epidote Biotite	Quartz Epidote Albite Chlorite Actinolite		
		T	low							
		+ stress							Cordierite Albite Epidote Microcline	
23-3 25, 50 51, 64?	Almadine- amphibolite	P	moderate to high	Regional	Calcite Quartz Diopside Zoisite Grossularite Epidote?	Quartz Muscovite Staurolite				
		T	moderate to high rather strong stress							
1, 5, 4-1 58, 59-2 153	Granulite	P	very high	Regional		Quartz Cordierite? Oligoclase	Quartz Microcline Almandine Pyroxene? Amphibole?			
		T	very high							
		+ stress								
24-1, 52	Hornblende- hornfels	P	moderate	Contact		Andalusite Cordierite Quartz Oligoclase				
		T	moderate							
		— stress								
		Metasomatism								
151				Contact			Tourmaline Garnet			
42A, 43 44, 47A 53, 501B 814		Extreme dis- location, cataclasis, granulation		Kinematic (grades locally with regional)		Andalusite Quartz Plagioclase Epidote Cordierite?	Quartz Oligoclase Cordierite Almandine Epidote

Fig. 1. Photomicrograph of tourmaline-bearing gneiss. Semi-idioblastic schorl (black and dark gray) embedded in a quartz-mosaic matrix. Note characteristic sieve texture of the tourmaline. Crossed polars, 96 X. Sample Sp-151.

Fig. 2. Photomicrograph of epidote-bearing quartzo-feldspathic schist. Some epidote (high-relief, left-center) shows sieve texture and evidence of slight rotation during deformation. Finer grained material is quartz and chlorite. Crossed polars, 96 X. Sample SP-55.

PETROGRAPHIC DETAILS OF METAMORPHIC ROCKS

Fig. 1. Typical outcrop of medium to thin-bedded metamorphosed limestones of the southern Sierra de Las Jarillas. Note contortion of individual beds.

Fig. 2. Photomicrograph of chlorite-epidote bearing calc-schist. Alternating thin bands of epidote-chlorite (dark) and calcite (light) are characteristic. Plain light, 96 X Sample SP-39.

LIMESTONE FROM SIERRA DE LAS JARILLAS AND PETROGRAPHIC DETAILS
OF CALC SCHIST

chlorite are abundant. The rocks have a yellowish weathered surface, and a fine-grained lepidoblastic texture; locally abundant poikiloblastic quartz is found. Locally interbedded with the phyllites, schistose rocks of quartzo-feldspathic composition are found.

Pegmatite dikes up to 10 m in width are common. These bodies resist more than the country rock and stand out as isolated hills easily recognizable in the field. The most common effect of contact metamorphism near the pegmatites is the presence of tourmaline (Plate 1, figure 1), probably formed by gases escaping from the dikes. The pegmatite dikes are simple pegmatites. Microcline, microperthite, and quartz are the essential constituents. The central portions of these bodies contain the larger crystals, progressively diminishing in size towards the margins.

In the country rocks near the intrusive contacts tourmaline occurs in almost euhedral stout prisms which range in length from 5 mm to 1 cm, and most of which are perpendicular to the wall of the dike. These crystals diminish in size away from the intrusive body, and they are embedded in a quartz mosaic producing a black and white rock of distinctive appearance (Plate 1, figure 1).

The southern portions of the Sierra de Las Jarillas consist of a series of quartzo-feldspathic schists and associated phyllites intercalated with calc-schists. North of the Cerro del Picacho, the sequence consists predominantly of dark greenish-gray quartzo-feldspathic schists. They are thin-bedded and contain numerous fractures which have an average strike of N 8° E. Their texture is generally schistose and fine-grained, but locally somewhat gneissic. Epidote is a major constituent (up to 30 percent) of some layers (Plate 1, figure 2).

Interbedded with these rocks are metamorphosed limestones, generally in beds of medium thickness, thinly laminated, and which present a weathered surface of a distinctive yellowish-brown color. They are present in isolated small outcrops. The rock is highly fractured and contorted (Plate 2, figure 1), and has an approximate strike of N 13° W. Dikes of milky quartz and local concentrations of micaceous gneiss are present.

Tightly folded and highly fractured calc-schists (Turner, 1948) and phyllites are increasingly abundant to the west and form a broad north-south trending valley. The calc-schists southeast of the Rancho El Rincón represent the quartz-albite-muscovite-chlorite subfacies of the greenschist facies. They are very fine-grained dark to light-green, slightly cataclastic chlorite and epidote-bearing rocks (Plate 2, figure 2), containing contorted schistose bands of chlorite and calcite alternating with bands of chlorite, calcite, quartz, epidote, and magnetite. The calcite is hetero-granular, consisting of either minute interlocking anhedral (less than 0.05 mm in maximum diameter) or coarser grains from 0.6 mm up to 4 mm in maximum diameter. Bent polysynthetic twinning is a common feature of the coarser calcite and has been interpreted by Heinrich (1965) as evidence of cataclastic deformation.

West of the Rancho El Rincón, the calc-schists are richer in biotite, magnetite, sericite, and depleted in quartz. The presence of minute lime-garnet crystals suggests a slightly higher grade of metamorphism. In contrast with the calc-schists to the east, these metasediments are medium-to thick-bedded and have a brownish color on both fresh and weathered surface due to oxidation. The

calcite is in layers of equigranular crystals. The finer crystals commonly have complex denticulate to sutured margins. Phyllites near the Rancho El Rincón are commonly dark gray, fine-grained, in medium-bedded layers intruded by quartz dikes, and are characterized by abundant biotite sometimes up to 10 percent (Plate 3, figures 1 and 2).

The southern Sierra del Picacho is composed of marbles interbedded with phyllites and schists to the east, and predominantly muscovite schists to the west near the southern end of the Sierra del Potrero. The fine-grained marbles are impure and presumably were originally quartzose calcilutites and calcisiltites. The average thickness of these strata is 1 m.

In one peculiar bed (SP 64 on Plate 4) the marble has "xenoliths" of micaceous schist. Pantoja (personal communication) believes that a viscous paste of carbonate rock at depth incorporated other rock fragments as a magma would acquire country rock.

At least three different marble lithologies are recognizable in the field:

Unit A: Epidote-bearing calcite marble. The essential constituents are very minute, interlocking, anhedral prisms of calcite (60 percent), locally forming bands of fine to almost cryptocrystalline material; and epidote (10 percent) commonly showing twinning and parting and inclusions of magnetite and zircon.

Unit B: Banded hematite-bearing, fine-grained calcite marble. The unusual banding of this rock consists of thin laminae of magnetite altered to hematite (15 percent) alternating with bands of polysynthetically twinned calcite elongated in the direction of foliation. The crystals are commonly bent by strong mechanical deformation.

Unit C: White marble with localized iron oxide stain. The rock is almost all calcite (98 percent) with minor quartz, magnetite, hematite, and sphene.

These units strike approximately due east and are separated by layers of micaceous schists and phyllites containing secondary epidote on bedding planes.

The southwestern Sierra del Potrero comprises north-south trending hills of metasedimentary rocks cut by pegmatitic and quartz dikes which vary in width from less than 2.5 cm to 50 cm. This sequence is generally quartzitic with little or no mica but chlorite and muscovite are progressively more abundant upward in the section (Plate 5, figure 1). The texture is schistose to granoblastic and the rocks are properly termed muscovite-chlorite semi-schists (Turner, 1948).

Immediately south of Rancho el Sauz in the southern Sierra del Potrero, a few dikes of basic composition were observed. Petrographic examination revealed albite (10 percent), epidote (25 percent), and actinolite (25 percent), with minor magnetite, sphene, quartz, hematite, and brown chlorite (Plate 5, figure 2).

Calc-schists and garnetiferous muscovite schists are well exposed in the Sierra de Magdalena near the Rancho Las Cuevitas. Phyllites are abundant to the south. To the west, the sequence is predominantly thin to medium-bedded muscovite schists, with large (up to 1 cm in maximum diameter) idioblastic

Fig. 1. Thin-bedded phyllites west of Rancho El Rincón in the Sierra de las Jarillas.

Fig. 2. Thin- to medium-bedded phyllites near Rancho El Rincón. Light-colored rock on top is a pegmatite dike, one of many in this area. Note a slight synclinal structure.

PHYLLITES FROM SIERRA DE LAS JARILLAS

Fig. 1. Photomicrograph of muscovit-chlorite semi-schist; southwestern Sierra del Potrero. Crossed polars, 96 X. Sample SP-48.

Fig. 2. Photomicrograph of albite-epidote-actinolite schist; southern Sierra del Potrero. The rock is a metamorphosed mafic dike in the metasedimentary sequence. Abundant actinolite (bottom) denotes the albite-epidote-biotite subfacies of the greenschist facies. Plain light, 96 X. Sample SP-4-1.

PETROGRAPHIC DETAILS OF SCHISTS FROM SIERRA DEL POTRERO

porphyroblasts of red garnet. The garnet is largely altered to iron oxide giving the entire rock a reddish appearance.

The rocks east of Rancho Las Cuevitas consist of thin-bedded gray muscovite schists which lack garnet. These metasedimentary rocks are interbedded with a series of low-grade calc-schists and lenses of green shale. Locally, some units are *finely laminated and show contorted bedding*.

North of the Rancho El Alamito in the Sierra de Magdalena, the calc-schists are associated with dark grayish-green epidote-bearing phyllites. Their texture is extremely fine-grained and strongly lepidoblastic. Epidote is an abundant constituent (23 percent) as individual crystals and granular aggregates with inclusions of quartz. This sequence is also intruded by quartz dikes with local concentrations of andalusite in the country rock.

ROCKS OF THE ALMANDINE-AMPHIBOLITE FACIES.—The central Sierra de Magdalena and the central and west-central Sierra del Potrero are the only areas known to the author to have rocks of this facies. The diagnostic mineral assemblages (Table 1) are locally associated with rocks of the hornblende-hornfels facies of contact metamorphism. A similar occurrence in the Caborca-Altar region of northwest Sonora is described by Damon and others (1962).

Near pegmatite dikes in the Santa Ana Quadrangle, there is a greater concentration of andalusite and tourmaline, and an increase in abundance of schist and some varieties of gneiss at the expense of phyllite.

Southwest of the Mina Dolores in the Sierra del Potrero, the rocks are strongly schistose and consist of a series of brownish to light gray muscovite-staurolite schists (Plate 6, figure 1). The texture is porphyroblastic-poikiloblastic (staurolite) and lepidoblastic (muscovite). The staurolite (15 percent) attains *a maximum diameter of 3 mm*, but the muscovite flakes are considerably larger and highly deformed, bending around staurolite porphyroblasts. To the north, the schists are richer in tourmaline (30 percent; Plate 6, figure 2) due to a greater abundance of pegmatitic intrusions, and staurolite (3 percent) gives way to cordierite.

A few kilometers north of Rancho El Sauz, the schists are subordinate to andalusite and magnetite-bearing gneiss (Plate 7, figure 1) with foliated, nematoblastic, and locally cataclastic texture. Numerous open pits have been dug in this area because the gneisses locally contain abundant malachite.

About 500 m south of Rancho El Sauz the dominant lithology is quartzofeldspathic gneiss (true granite gneiss) of the granulite facies. This suggests a rough zonation: a central portion of granulite facies rocks surrounded by rocks of lower metamorphic grade. A similar zonation is present in the central and southern Sierra de Magdalena where rocks of the greenschist facies to the west are flanked by staurolite-muscovite semischists and calc-schists of the almandine-amphibolite facies on the east.

North of Rancho El Alamito in the Sierra de Magdalena, a sequence of purplish-pink contorted calc-schists (Plate 7, figure 2) with crystalloblastic, fine-grained granoblastic texture contain calcite (60 percent) in highly contorted layers. Minor diopside (5 percent) is slightly altered to zoisite. It is porphyro-

blastic and highly fractured with abundant inclusions of calcite, quartz, and opaque minerals. Locally abundant streaks of xenoblastic magnetite are oxidized and give the rock its distinctive color.

To the east, light reddish-brown, thin-bedded, highly deformed staurolite-sericite schists are interbedded with calc-schists and are locally deep orange-red on weathered surfaces, due to hematite. Rocks of the granulite facies are present farther east.

ROCKS OF THE GRANULITE FACIES.—The distribution of rocks of the granulite facies is apparently related to metamorphosed granitic intrusions. The most common lithologies are true granite gneiss which progressively change to schistose rocks away from the intrusive cores (Plate 8, figure 1). Alternation of light sialic and dark mafic bands is common but not universal. It is emphasized that boundaries between metamorphic facies are gradational and that, therefore, no definite contact lines appear on the map. Field information is incomplete and more work is needed to define lithologic boundaries in more detail.

The texture of the gneiss and granulites is largely dependent on the texture of the original rock. The granites and granodiorites have retained their textures to a greater extent (Plate 8, figure 2) than the original quartzo-feldspathic and pelitic rocks, except where they have been intensely sheared by fault movements. However, as seen in thin section, most granite gneiss is cataclastically deformed.

At the Rancho El Sauz in the Sierra del Potrero, the gneiss is quartzo-feldspathic. The feldspars are plagioclase (An_{12}), myrmekite, perthite, orthoclase, and microcline. As much as 5 percent porphyroblastic tourmaline was recorded in some samples. In western Sierra de Las Jarillas, the granite gneiss displays a characteristic flaser texture (Plate 9, figure 1). Lepidoblastic muscovite and garnet are locally abundant.

A chemical analysis of a gneiss sample from 3 km east of the Mina Magdalena in the Sierra de Magdalena shows a granodioritic composition (Table 2). Other similar analyses are included for comparison. It is noted that this analysis and those to follow represent only a minute fraction of the lithology. Care was taken to select the most abundant lithologies for chemical analyses.

The gneiss of the Sierra de Magdalena was extensively studied by the author because of the excellent exposures. In the extreme southern portions of this sierra white to gray cordierite-bearing gneiss is cataclastically deformed (Plate 9, figure 2) and shows incipient augen structure in the outcrop. "Eyes" of porphyroblastic albite (30 percent) and cordierite (30 percent) are conspicuous. Biotite and muscovite commonly form less than 3 percent of the rock but they are locally segregated into thin bands giving the rock a more orthodox gneissic fabric.

In the core of the Sierra de Magdalena, the gneiss is probably formed as a result of regional metamorphism of granitic intrusive rocks. The rocks are lithologically similar to those near the Rancho El Sauz. The gneiss locally grades into schistose rocks and the compromise term "schistose granite gneiss" is used to describe them (Plate 8, figure 1).

PRODUCTS OF DYNAMIC METAMORPHISM.—Dynamic metamorphism in the Santa Ana Quadrangle resulted from faulting and granulation. The common products of this type of metamorphism are cataclasites, mylonites, and phyllonites

Fig. 1. Photomicrograph of muscovite-staurolite schist; south-central Sierra del Potrero. Rotated semiisoblastic porphyroblasts of staurolite (left) displaying a strong sieve texture are conspicuous. Muscovite flakes bend around these porphyroblasts suggesting deformation. Crossed polars, 96 X. Sample SP-50.

Fig. 2. Photomicrograph of tourmaline-muscovite schist; south-central Sierra del Potrero. This rock was collected a few hundred meters north of SP-50 (Fig. 15). Tourmaline (dark gray and black) is the characteristic mineral. The rock does not show evidence of deformation. Crossed polars, 96 X. Sample SP-51.

PETROGRAPHIC DETAILS OF SCHISTS FROM SIERRA DEL POTRERO

Fig. 1. Photomicrograph of andalusite-magnetite gneiss; central Sierra del Potrero. Note size variation and sieve texture of andalusite (bottom and top of microphotograph). Picture shows part of an andalusite-rich layer. Magnetite-rich layer not shown. Plain light, 96 X. Sample SP-52.

Fig. 2. Highly contorted calc-schist; Sierra de Magdalena. Natural size, Sample SP-25.

DETAILS OF METAMORPHIC ROCKS FROM SIERRA DEL POTRERO AND SIERRA DE MAGDALENA

TABLE 2.—Chemical composition of granodiorites and gneiss

	SP-2	I	II	III	IV
SiO ₂	66.80%	67.86%	66.88%	63.9%	70.7
TiO ₂	0.14	0.46	0.57	0.8	0.5
Al ₂ O ₃	19.10	16.62	15.16	17.0	14.5
Fe ₂ O ₃	0.54	0.81	1.33	2.4	1.6
FeO	0.41	2.16	2.59	3.0	2.0
MnO	0.02	—	0.07	—	0.1
MgO	0.40	1.52	1.57	1.8	1.2
CaO	3.79	2.13	3.56	4.1	2.2
Na ₂ O	5.80	5.08	3.84	3.7	3.2
K ₂ O	2.47	2.54	3.07	3.1	3.8
P ₂ O ₅	0.16	—	0.21	—	0.2
SO ₃	0.00	—	—	—	—
CO ₂	0.19	—	—	—	—
H ₂ O+	0.25	0.95	0.65	—	—
H ₂ O—	0.05	—	—	—	—
S	—	—	—	—	—
	100.12%	100.19%		99.80%	100.0

SP-2* — Gneiss, 3 km east of the Mina Magdalena, Sierra de Magdalena, Santa Ana Quadrangle, Sonora, Mexico.

I — Granodiorite (sodic type), Glen Benvie, Scotland; N. Holgate, Quar. Journ. Geol. Soc.; v. 106, p. 438, 1951.

II — Average granodiorite; S. R. Nockolds, Bull. Geol. Soc. America, v. 65, 1954.

III — Average Precambrian, Canada, composed of gneiss, gneissic granites, and granites; F. F. Grout, J. Geol., v. 46, p. 502, 1938.

IV — Average of 250 quartzofeldspathic gneiss; E. Poldervaart, Geol. Soc. America Spec. Paper 62, p. 135, 1955.

* Analyst: Ing. Quim. Alberto Obregón, Instituto de Geología, Universidad Nacional Autónoma de México.

according to the terminology of Turner (1948). The mylonites (Plate 10, figure 1) are restricted to the major fault zones and are best exposed in the western front of the Sierra de Magdalena and the eastern front of the Sierra del Picacho and Las Jarillas. The main constituents are characteristically quartz and feldspar which generally are granulated and even pulverized, although some complete relict crystals survive. Mylonites consisting almost entirely of quartz crop out south of the San Miguel Mine in the Sierra de Magdalena.

Phyllonites also reflect reduction in grain size, and recrystallization of some minerals (for example, mica, chlorite, epidote) is evident in thin section (Plate 10, figure 2). Locally, as in parts of the western edge of the Sierra de Magdalena "mylonitic phyllonites" are present. These rocks retain coarser relict minerals such as poikiloblastic quartz, in association with abundant epidote, micas, and pyrite.

The cataclasites were deformed by shattering without chemical reconstitution. Southwest of the Rancho El Sauz, they are rather coarse-grained, porphyroclastic, and highly gneissic. Quartz (45 percent) is very fine-grained, elongate parallel to foliation, and strongly deformed. Sodic oligoclase (17 percent), and perthite (23 percent) both porphyroclastic, are bent, broken, and displaced by micro-faults. Myrmekite, cordierite, garnet, and muscovite are other constituents.

AGE AND PETROGENESIS.—As early as 1927, Flores (p. 202) considered the metamorphic rocks of the Santa Ana Quadrangle to be of Precambrian age because of the presence of high grade metasediments in the sequence even though they are overlain by Tertiary volcanic and sedimentary rocks. Santillán (1936), Imlay (1939), and Arellano (1956) described basement rocks of Precambrian age elsewhere in Sonora.

Terminology used in Canadian Shield	Terminology used in SW United States & this paper	Terminology used for Minnesota — U. S.			
		ERA	SYSTEM	OROGENY	AGE (10 ⁶ yr)
ERA	ERA				
Proterozoic	"Younger" Precambrian	Late Precambrian	Keweenawan	Grenvillian—1000	
	Mazatzal				
	Orogeny			Penokean —1700	
		Middle Precambrian	"Huronian"		
	"Older" Precambrian			Algoman —2500	
Archeozoic		Early Precambrian	Timiskamian		
			Ontarian	Laurentian— ? —	

FIG. 3.—Geochronologic scale for the Precambrian Era; after Goldich and others (1961).

Fig. 1. Photomicrograph of "schistose" granite gneiss; Sierra de Magdalena. The texture is between schistose and gneissic. Granulate quartz (light) and slightly bent muscovite flakes indicate cataclastic deformation. Traces of euhedral garnet (lower right) are present. Crossed polars, 96 X. Sample SP-58.

Fig. 2. Granite gneiss; Sierra de Magdalena. Note incipient development of augen structures. Natural size.

GRANITE GNEISS FROM SIERRA DE MAGDALENA

Fig. 1. Photomicrograph of granite gneiss; western Sierra de Las Jarillas. Note characteristic flaser texture. Crossed polars, 96 X. Sample SP-5.

Fig. 2. Granite gneiss; southern Sierra de Magdalena. Texture almost mylonitic. Large crystals of albite (center) and cordierite (not shown) stand out as porphyroblastic "eyes". Crossed polars, 96 X. Sample SP-1.

PETROGRAPHIC DETAILS OF GRANITE GNEISS FROM SIERRA DE LAS JARILLAS
AND SIERRA DE MAGDALENA

More recently, Damon and others (1962) successfully correlated the "Older" and "Younger" Precambrian of Arizona with metamorphic rocks in Sonora. Their conclusions establish an approximate minimum age of at least 1680 m. y. for the time of deposition and subsequent regional metamorphism of the "Older" Precambrian. The metamorphic rocks of the main sierras of the Santa Ana Quadrangle are tentatively assigned to the "Older" Precambrian.

It is recalled that the Coyotillo group represents low grade metamorphism and forms a homoclinal sierra. In Arizona, as well as in other parts of Sonora (for example, the northern portion of the Benjamin Hill Quadrangle), the "Younger" Precambrian rocks are of low metamorphic grade (greenschist facies) and show little or no structural deformation. In the absence of radiometric age determinations, it is suggested that the Coyotillo group is of "Younger" Precambrian age (Figure 3). However, the Altar Schist of northwestern Sonora is lithologically similar to the Coyotillo group, but Damon and others (1962) assign it a radiometric age of 57 ± 3 m. y. For this reason, the author does not rule out the possibility of a Laramide metamorphic event in the Santa Ana Quadrangle although the sedimentary rocks could be of Precambrian age.

The intimate association of granite and gneissic terranes in the area suggests that regional metamorphism was at least locally accompanied by granitic intrusions. The relatively widespread occurrence of pegmatitic material in schists, phyllites, marbles, and gneiss supports this contention. The association with igneous rocks implies that, at least locally, heat was influential to some extent in controlling the formation and distribution of metamorphic facies.

However, the presence of metasedimentary rocks, such as the Coyotillo group, not visibly related to igneous intrusions cannot be explained by thermal activity alone. The influence of load and directed pressure must also be considered. The wide range of physical conditions reflected in the various mineral assemblages developed during metamorphism not only in the rocks of the Coyotillo group but also in the remainder of the metasedimentary rocks, suggests a depth of formation equivalent to the mesozone of Buddington (1959).

Thin sections of several rocks show evidence of mild to strong mechanical deformation accompanied by recrystallization under the influence of strong directional stresses. It is suggested that rock deformation in itself played a significant role in the regional metamorphism, perhaps by permitting chemical reconstitution of the rocks being deformed.

In summary, the metamorphic rocks of the Santa Ana Quadrangle reflect gradations from localized thermal metamorphic effects of granites and other intrusive rocks to what may properly be termed regional metamorphism in its broadest sense. Table 3 is a summary of the author's interpretation of development of the metamorphic rocks from the original materials.

TABLE 3.—Metamorphic series

ORIGINAL MATERIAL	ROCKS OF INCREASING METAMORPHIC GRADE				
	GREENSCHIST	AMPHIBOLITE	GRANULITE		
Clay-shale (argillite)	Phyllite	mica schist	actinolite schist	gneiss	
Sandstone		Quartzite			
Conglomerate		Metaconglomerate			
Graywacke		Metagraywacke			
Limestone	Marble	marble		marble	
Psephites		Quartzo-feldspathic schist			
Impure calcisil- tite and calcilutite		Calc-schist	calc-schist		
Acid igneous rock				Gneiss	

SEDIMENTARY ROCKS

The sedimentary rocks of the Santa Ana Quadrangle are divided into three general units (Figure 2): (1) graywackes, shales, and limestones of Lower Cretaceous age; (2) red conglomerates of Tertiary age with intercalations of thin sandy layers; and (3) stratified valley-fill deposits mainly gravels and conglomerates of the upper Pliocene Báucarit Formation.

ROCKS OF CRETACEOUS AGE.—In 1927, Flores mapped and described a thick sedimentary sequence which crops out immediately southeast of Santa Ana. The same sequence is also exposed as isolated but extensive outcrops in the southwestern part of the quadrangle, west and south of Magdalena (Sierra de La Ventana), and in the central portion of the Sierra de Magdalena. In the latter two localities, the sediments, mostly limestones, are found together with extrusive rocks. These limestone are wholly recrystallized, highly fractured, and locally mineralized suggesting incipient metamorphism (Ham, personal communication).

El Represo Section.—The thickest exposure (approximately 485 m) of undisturbed Lower Cretaceous rocks is the section found 500 m northeast of the Rancho El Represo north of the Sierra de La Lumbre (Appendix B). Flores (1927) assigned an Early Cretaceous (Neocomian) age to this sequence based on the presence of the following fossils: *Trigonia* aff. *hertzogi* Hausmann (Goldfuss), *Trigonia* cfr. *subventricosa* Stanton, *Ctenostrion* sp., *Exogira* sp., and *Ostrea* sp. Upon examination of several thin sections the author found the foraminifer *Orbitolina texana*, recrystallized fragments of crinoid stems, indivi-

Fig. 1. Photomicrograph of mylonite; western Sierra de Magdalena. Granulated and pulverized quartz and feldspar (light) in a "dusty" matrix of unidentifiable material (dark). Note abundance of microfractures. Plain light, 96 X. Sample SP-44.

Fig. 2. Photomicrograph of quartz-mica phyllonite; central Sierra de Magdalena. Plain light, 96 X. Sample SP-42A.

PHOTOMICROGRAPHIC DETAILS OF MYLONITE AND PHYLLONITE FROM SIERRA DE MAGDALENA

Fig. 1. Medium to thick-bedded limestone of el Represo section, north of the Rancho El Represo. Note abundance of *Ostrea* and *Pecten*.

Fig. 2. Photomicrograph of quartzose calcisiltite; north of the Rancho El Represo. Note large pelecypod showing partial recrystallization at edges of shell and some microfractures. Plain light, 96 X. Sample SP-26.

LIMESTONES FROM RANCHO EL REPRESO

dual sponge spicules and spicular meshworks, and fragments of gastropods and pelecypods.

Limestone of the El Represo Section.—The limestones in this section are invariably massive and thick-bedded (Appendix C). Locally they are completely silicified and nodular when in shales. Most beds give a strong fetid odor when struck with the hammer, probably due to the presence of large amounts of organic matter.

The silicified units have a characteristic purple color. Gastropod shells are relatively abundant, locally forming 10 to 15 percent of the rock. The allogenic constituents, forming less than 10 percent of the rock, are quartz, zircon, magnetite, tourmaline (dravite), ilmenite, leucosene, and plagioclase (An_{22}). The main constituent is silica in the form of quartz and chalcedony (65 percent) which replaced calcite silt. The quartz is anhedral where replacing fossil shells and euhedral and aligned perpendicular to the wall where filling pores or fractures. Some hematite and sparry calcite, which constitute about 15 percent of the rock postdate the euhedral quartz where filling open spaces. The author interprets the beds to have originally been fine-grained bioclastic limestones.

The "fresh" limestones at this locality are skeletal calcite mudstones and locally, spicular calcite mudstones. The textures are detrital comprising calcite mud supporting silt-size quartz and organic remains (Plate 11, figure 1). Allogenic constituents are quartz (locally as much as 10 percent), magnetite, sericite, zircon, and clay minerals. The micro-skeletal remains are generally not well preserved, being partly or completely recrystallized to sparry calcite and commonly broken and even displaced by microfaults. A few incomplete pelecypod shells have retained their original fibrous carbonate texture. Calcite mud is probably the chief authigenic constituent. Patches of sparry calcite, recrystallized from the mudstone are minor. The calcite mud is locally intraclastic and associated with small euhedral pyrite crystals. Hematite is common in fractures staining the adjacent rock.

Quartzose calcisiltites in the section (Plate 11, figure 2) contain subrounded quartz (15-20 percent), plagioclase, clay, iron oxides (10 percent), and silt-size calcite with inclusions of minute grains of magnetite. Partly silicified fossils are rare. Sparry calcite, as a recrystallization product of the original carbonate, and pyrite and silica are locally abundant.

Limestone nodules are abundant in some of the shales of the El Represo Section. They are gray, oval-shaped, from 5 to 30 cm long, and have a characteristic very fine-grained texture (Plate 12, figure 1). Calcite mud is the essential constituent with minor accessory sparry calcite, pyrite, and hematite.

Limestone from other localities.—To the southwest and southeast of Santa Ana, and west of the Rancho Los Federicos in the southwestern portion of the quadrangle, isolated limestone hills crop out showing variable trends. Lithologically and paleontologically they are similar to those of the El Represo Section although part of the sequence west of the Rancho Los Federicos are predominantly micritic calcarenites with abundant mudstone intraclasts (Plate 12, figure 2), and coated grains (pseudo-oolites; Ham, personal communication), and breccia

with sharp angular pieces of calcite mudstone, commonly about 5 cm in maximum diameter, cemented by hematite and red clay. The breccia forms lenses about half a meter thick and 2 to 3 m long in the calcarenites.

The limestone associated with extrusive rocks are markedly different. The presence of sericite, coarse (1-4 mm), recrystallized calcite, and hematite filling fractures suggests mild metamorphism. The hematite could be a result of hydrothermal oxidation related to the formation of the volcanic rocks. A small amount of dolomite is present northeast of the Ejido San Lorenzo. According to Ham (personal communication), the dolomite could have been originally a coarse-grained calcarenite.

Southeast of the Rancho Las Cuevitas, in the Sierra de Magdalena, is a sequence, 10 m in thickness, of thinly laminated calcisiltites which have non-clastic, microcrystalline, and granular textures. Thin section examination revealed minute sericite-magnetite-rich layers alternating with layers of anhedral calcite. Although they are mapped as sedimentary rocks by the author and by geologists of the Consejo de Recursos Naturales no Renovables, they may be low-grade metamorphic limestone. No association with igneous rocks is evident.

Conglomerates, graywackes, and siltstones.—The Lower Cretaceous limestones are interbedded with graywackes, conglomerates and siltstones.

A sequence 1 to 2 m thick of grayish-green, fine-grained lithic conglomerates is ubiquitous interbedded with the Lower Cretaceous limestones (see Appendix B). The conglomerates consist of rounded to sub-rounded fragments of chert, spherulitic chalcedony, phyllite, quartzite, igneous rocks (probably andesite porphyry and basalt with diabasic texture), shale, quartz, and plagioclase (An_{12}), in a matrix of hematite and cemented by silica. This unit is interpreted to represent a hiatus in the deposition of the normal Lower Cretaceous section exposed in the area.

The essential characteristic of the graywackes of the Santa Ana Quadrangle is that they contain numerous rock fragments, and therefore, are called here lithic graywackes (Turner and others, 1954). The particle size distribution is variable from fine-to medium-grained gravel to fine silt. Some units show well developed graded bedding.

The graywackes are dark reddish-brown, generally well consolidated, and invariably underlain by dark shales. "Intraclasts" of the shales are incorporated within the graywacke beds locally indicating considerable "cannibalism." Soft mud stirred up by currents or waves on bottom of the depositional basin may have been mixed with depositing graywacke. A representative sequence about 4 km southeast of Santa Ana that shows this phenomenon consists of the following gross lithologies:

- Unit 1: reddish-brown shale (oldest)
- Unit 2: conglomeratic graywacke
- Unit 3: reddish-brown shale
- Unit 4: graywacke with "intraclasts" of underlying shale
- Unit 5: graywacke (youngest)

Fig. 1. Photomicrograph of partly recrystallized calcilutite in limestone nodule; north of the Rancho El Represo. Anhydrous sparry calcite (center) has recrystallized at the original carbonate. Plain light, 96 X. Sample SP-506.

Fig. 2. Photomicrograph of micritic calcarenite; northwest of the Cerro del Alamito, extreme southwestern corner of the Quadrangle. Abundant mudstone intraclasts (euhedral and elongated grains) and coated grains (lower right) are supported by a partly recrystallized mudstone matrix. Plain light, 96 X. Sample SP-6.

PHOTOGRAPHIC DETAILS OF LIMESTONES FROM RANCHO EL REPRESO AND CERRO DEL ALAMITO

Fig. 1. Protomicrograph of lithic graywacke; west of Cerro de La Lumbra (from specimen shown in Fig. 33). A few lithic fragments, plagioclase (twinned) and quartz (light colored) in matrix of sericite and chlorite. Crossed polars. 90X. Sample SP-15.

Fig. 2. Outcrop of thick-bedded, fine to medium-grained, lithic red conglomerated of questionable Upper Pliocene age; 5 km northeast of the Ejido San Lorenzo.

GREYWACKE FROM CERRO DE LA LUMBRERA AND CONGLOMERATE
SAN LORENZO

Under the microscope (Plate 13, figure 1), the graywackes are characterized by angular to sub-rounded quartz, plagioclase (albite and oligoclase), orthoclase, and rock particles. The lithic constituents are fragments of altered volcanic rocks, quartzite, chert, shale, and metamorphic rocks, probably altered schists and phyllites. The matrix is a mixture of minute aggregates of silica, sericite, chlorite, and hematite which is commonly indistinguishable from the cement of calcite (ankerite?) and hematite. Accessory minerals include apatite, epidote, zircon, biotite, and tourmaline.

The finer-grained graywackes grade into calcareous sandy siltstones and siltstones. The percentage of rock fragments decreases from approximately 35 percent in the graywackes to about 15 percent in the siltstones, but the percentage of calcite as cement increases to about 25 percent.

ROCKS OF TERTIARY AGE.—Red conglomerates.—During his pioneering trip through Sonora, Flores (1927) mapped local discontinuous outcrops of red conglomerates of questionable late Pliocene age (Plate 13, figure 2). These coarse-grained conglomerates are exposed in topographic depressions, generally overlain by the Báucarit Formation and resting unconformably on Mesozoic rocks. The main constituents are rounded fragments of gneiss, quartzite, and volcanic rocks in a matrix of red sand and clay cemented by argillaceous, siliceous, and/or calcareous material. As shown on the map, this unit has a very restricted areal distribution.

Báucarit Formation.—The Báucarit Formation in the Santa Ana Quadrangle consists of poorly consolidated sandstones and conglomerates which display rather poor stratification. This unit of intermontane valley-fill deposits has the largest area of outcrop of all the rock units mapped in the quadrangle.

The Báucarit Formation was originally called the "Báucarit division" by Dumble (1900a). The name was changed to Báucarit Formation by King (1934; 1939) who designated as type locality the Cedros River basin in central Sonora. Flores (1927) considered this formation to be of Pleistocene age in the Santa Ana Quadrangle; however, in accordance with Pantoja (personal communication) and others, the author believes that the Báucarit Formation may, at least in part, be of Tertiary age. It is roughly equivalent to the Gila Conglomerate of Arizona (Pye, 1959).

The composition of the Báucarit Formation varies sympathetically with that of exposed preexisting rock masses. Thus, it consists largely of gneiss fragments and volcanic rock fragments around sierras of metamorphic and volcanic rocks, respectively.

A 30 m section of the Báucarit Formation was measured south of Ejido San Lorenzo on the south bank of the Magdalena River. There it consists of well sorted horizontal conglomeratic beds in which about 80 percent of the particles are sub-rounded; the remainder are sub-angular. The particles range from boulders 80 to 90 cm in maximum diameter to pebbles less than 3 cm in maximum diameter. About 98 percent of the fragments are andesite and andesite porphyry; the remainder are boulders of gneiss and schist. The matrix is

generally angular to subrounded hematite-stained sand and silt grains. This unit characteristically forms steep bluffs or cliffs.

Alluvial deposits.—Unconsolidated alluvial sand, silt, and mud of Pleistocene and Recent age were largely derived from the Báucarit Formation but also from the other preexisting rocks in the area. Concentrations of magnetite, hornblende, and tourmaline sands are localized at the intersection of some small arroyos. Most agricultural activity of the region takes place in the sands and silts of the alluvium.

PETROGENESIS OF THE SEDIMENTARY ROCKS.—Table 4 suggests the provenance of the major types of Cretaceous sedimentary rocks of the Santa Ana Quadrangle. Low to high-grade metamorphic rocks and sialic to perhaps intermediate igneous rocks contributed to the clastic fractions of the graywackes, siltstones, and limestones. Pre-Cretaceous mafic igneous rocks probably did not contribute to these clastic fractions of the exposed sedimentary rocks except for small dikes in the granitic bodies.

The lithologic associations of the Lower Cretaceous sediments suggest a eugeosynclinal environment of deposition. The author considers this sequence to represent the "Graywacke Association of Shallow Burial" of Krumbein and Sloss (1963, p. 530). The sediments may have accumulated in a downwarped trough with an uplifted area on either or both sides providing immature clastics. Temporary interruptions of uplift and erosion in the source area and/or failure of the clastic sediments to reach the geosyncline may have allowed accumulation of limestone and shale.

Post-depositional effects such as cementation, quartz enlargement, silicification, and authigenesis of feldspars were minor in the graywackes. However, recrystallization of fossils and calcite mud to sparry calcite is common in some of the carbonate rocks.

Flores (1927) postulated that the red conglomerates were deposited in lakes (as a result of erosion of fragments from the surrounding higher areas) because they occupy topographic depressions. However, a deltaic environment of deposition is not ruled out by the author. For lack of evidence to support either postulate the matter is open for future research.

The Báucarit Formation was probably deposited by streams as a series of coalescing *bajadas* which covered intermontane pediment surfaces. Diagenetic processes have operated little. Poor consolidation of beds and ill-defined stratification are characteristic.

IGNEOUS ROCKS

The granites and associated rocks, the volcanic rocks (andesites) associated with the sedimentary sequence, and the Comedores Basalt are considered here. The igneous geology of the Sierra de La Lumbre is described separately.

TABLE 4.—Inferred sources of detrital minerals of Lower Cretaceous sedimentary rocks in the Santa Ana Quadrangle

<i>From reworked sediments source</i>	
Chert (in lithic graywacke, calcareous siltstone, conglomerate)	
Zircon (in silicified fossiliferous limestone, calcareous siltstone)	
Tourmaline (in silicified fossiliferous limestone, calcareous siltstone)	
Quartz (in lithic graywacke)	
<i>From low-grade metamorphic source</i>	
Quartzite fragments (in lithic graywacke)	
Quartz (in lithic graywacke, calcite mudstone, silicified fossiliferous limestone, calcisiltite, conglomerate)	
Chlorite (in calcareous siltstone, lithic graywacke)	
Biotite (in lithic graywacke)	
<i>From high-grade metamorphic source</i>	
Sodic plagioclase (in lithic graywacke, calcareous siltstone, conglomerate)	
Magnetite (in calcite mudstone, silicified fossiliferous limestone, calcareous siltstone).	
Epidote (in calcareous siltstone)	
Quartz (in calcisiltite)	
Biotite (in graywacke?)	
Andalusite (in calcisiltite?)	
<i>From sialic and intermediate igneous source</i>	
Volcanic rock fragments (in lithic graywacke)	
Apatite (in lithic graywacke)	
Magnetite (in lithic graywacke, calcite mudstone?, calcareous siltstone)	
Muscovite (in calcareous siltstone)	
Microcline (in lithic graywackes)	
<i>From mafic igneous source</i>	
Intermediate plagioclase (in silicified fossiliferous limestone?)	
Leucoxene (in silicified fossiliferous limestone)	
Ilmenite (in silicified fossiliferous limestone)	
<i>From pegmatite source</i>	
Muscovite (in calcareous siltstone, calcite mudstone)	
Microcline (in lithic graywacke)	
Myrmekite (in lithic graywacke)	

GRANITES AND ASSOCIATED ROCKS.—Two main classes of granitic rocks are exposed in the area: (1) granites and dikes associated with the metamorphic sequence, and (2) the granitic complex exposed in the extreme northeastern corner of the quadrangle which includes rhyolites and rhyolite porphyries.

The western fronts of the Sierra de La Tortuga and the Sierra del Rincón, the northwestern part of the Sierra de Guacomea, and parts of the core of the Sierra de Magdalena comprise granitic rocks of Precambrian age which are, in general, highly altered and metamorphosed. The author was able to find outcrops of fresh, unaltered granite in one locality; northeast of the Rancho San Antonio in northwestern Sierra de Guacomea, in the deepest parts of ravines and characteristically associated with and overlain by granite gneiss.

The granites are pink, generally medium-grained, holocrystalline, and hypidiomorphic granular. The modal constituents in volume percent are quartz (25), oligoclase (35), microcline (5), perthite (5), and myrmekite (5), plus rare local concentrations of almandite (up to 5 percent).

The quartz is anhedral with sutured contacts between some of the grains. Some crystals are fractured and even crushed exhibiting a "mosaic" appearance and locally giving the rock a cataclastic texture. Rows of minute inclusions of magnetite and plagioclase are common in the quartz. Microcline and perthite are characteristically anhedral and display poikilitic texture with inclusions of quartz and some alteration to sericite. Oligoclase crystals are commonly broken or bent. Selective sericitization is localized along twinning planes and in the cores of some crystals. Almandite up to 2 mm in maximum diameter is euhedral; inclusions of magnetite and highly irregular fractures are common. Magnetite, biotite, and zircon are accessory minerals in the granites of metamorphic terranes. Magnetite is subhedral as inclusions or intergranular crystals; some is clearly an oxidation product after biotite. Biotite is pleochroic in shades of brown and occurs as small tabular flakes locally associated with magnetite. Some is altered to chlorite.

Aplitic dikes a few centimeters thick are abundant in the parent granite and commonly penetrate the adjacent metamorphic rocks. The aplites are white or pink, and fine-grained. The pink varieties are mineralogically similar to the parent granite except that almandite is less abundant. A chemical analysis of the parent granite is shown in Table 5. Also see Appendix C for normative and modal compositions. Notice the striking similarity to Nockolds' average alkali granite.

The white aplites contain biotite and have a remarkably even, fine-grained saccharoidal texture, which, under the microscope, is seen to be holocrystalline, allotriomorphic-granular. The principal minerals in volume percent are quartz (35), oligoclase (25), microcline (10), and biotite (10). Accessories include magnetite, zircon, myrmekite, perthite, sericite, sphene, and garnet. Although a chemical analysis of this rock is not available, according to Hatch and others (1961) rocks of this type are characterized by high contents of silica and alkalis with minor iron and magnesia.

The sialic rocks described above are the "normal granites" of Flores (1927) rather than "binary granites" (essentially all quartz-orthoclase) which were not found in the area.

TABLE 5.—Chemical composition of granites

	SP-152	I	II
SiO ₂	73.94%	72.08%	73.86%
TiO ₂	0.10	0.37	0.20
Al ₂ O ₃	13.96	13.86	13.75
Fe ₂ O ₃	1.20	0.86	0.78
FeO	0.06	1.67	1.13
MnO	0.10	0.06	0.05
MgO—	0.25	0.52	0.26
CaO	0.87	1.33	0.72
Na ₂ O	3.98	3.08	3.51
K ₂ O	5.35	5.46	5.13
P ₂ O ₅	—	0.18	0.14
SO ₃	—	—	—
CO ₂	—	—	—
H ₂ O+	0.16	0.53	0.47
H ₂ O—	0.03	—	—
S	—	—	—
	100.00%		

SP-152* — Granite, northwest Sierra de Guacameza, Santa Ana Quadrangle, Sonora, Mexico.

I — Average calc-alkali granite; S. R. Nockolds, Bull. Geol. Soc. America, v. 65, 154.

II — Average alkali granite; S. R. Nockolds, Bull. Geol. Soc. América, v. 65, 1954.

* Analyst: Ing. Quim. Alberto Obregón, Instituto de Geología, Universidad Nacional Autónoma de México.

East of the Magdalena Mine in the Sierra de Magdalena, quartz diorite (?) dikes intrude the granites (Plate 14, figure 1). The quartz diorite (?) is highly altered; the original ferromagnesian minerals have been completely replaced by pseudo-radial clusters of chlorite and euhedra of green hornblende (up to 40 percent). Quartz (5 percent) appears to be interstitial between plagioclase (40 percent by volume) grains. Other minerals in the rock are opaques (13 percent), hematite, biotite, muscovite (traces, probably after chlorite), zircon, and calcite. The latter is found filling micro-fractures and as an alteration product of plagioclase, and associated with traces of sericite.

All rocks exposed in the northeastern corner of the quadrangle are included under the name "Igneous Complex." These rocks were considered by Flores (1927) to be rhyolite and rhyolite porphyry but granite and small andesite bodies are also present. The granites and rhyolites are grouped together because they are associated in the field. According to Flores (1927), the rhyolites are

intrusive and have the form of either dikes or volcanic necks of limited areal extent. Flores' "porphyritic rhyolites," the most abundant rock type of this "Igneous Complex," form large intrusive bodies and are composed mainly of large orthoclase and quartz phenocrysts set in a fine-grained phanecritic groundmass of the same minerals. This mineralogy is the same as the "binary granites" of Flores (1927). Flores also observed minor bodies of andesite of probable Late Cretaceous or early Tertiary age (Pantoja, personal communication) in the vicinity of the Rancho El Pozo.

ANDESITES.—South of the Rancho La Jojoba (Sierra de Magdalena) and northeast of the Ejido San Lorenzo, the Cretaceous limestones are interbedded with pinkish-gray, slightly altered hornblende andesites (Plate 14, figure 2). The andesites are holocrystalline, aphanitic, and porphyritic containing megaphenocrysts which range from 1 to 2.5 mm in maximum diameter. The groundmass texture varies from intergranular to intersertal and pilotaxitic.

Plagioclase (35 percent) phenocrysts are zoned and locally grouped in glomeroporphyritic clusters. The cores of the zoned plagioclase crystals are altered to a combination of calcite and sericite. Unaltered biotite (4 percent) in small tabular crystals approximately 0.5 mm in maximum diameter is intimately associated with magnetite. Hornblende (25 percent) exhibits all degrees of deuteric oxidation to magnetite (Plate 14, figure 2) from early stage replacement by magnetite granules to the extreme in which only the shape of the original hornblende phenocrysts has survived. Intergranular primary magnetite is ubiquitous as small euhedral to subhedral crystals. Accessory minerals include green serpentine, altered pyroxene (?), and traces of quartz. The gray matrix consists of opaque minerals, plagioclase microlites, and minor quartz, and makes up about 30 percent of the rock.

Because of their interbedding with Cretaceous rocks, these extrusive rocks have been considered to be of Cretaceous age by several geologists (Pantoja, Lucarelli, personal communication). Moreover, the andesites of the Sierra de La Ventana, although not associated with limestones, are lithologically similar to those just described. For this reason, Pantoja and others believe the Sierra de La Ventana volcanic rocks to be of Cretaceous age also. Clearly more field relations must be determined to establish the relative ages of these volcanic rocks as well as the Tertiary volcanic rocks of the Sierra de La Lumbre.

The andesites that intrude rocks of the Coyotillo group as hypabissal dikes ranging from 0.5 to 1.0 m in thickness are pinkish-gray, aphanitic, porphyritic, and glomeroporphyritic. The principal modal constituents are megaphenocrysts of oxyhornblende, andesine, and pyroxene in a vitrophyric, slightly pilotaxitic groundmass of glass, plagioclase microlites, sericite, and opaque minerals. The oxyhornblende (10 percent), which is green in the hand specimen, is strongly twinned and contains inclusions of magnetite, apatite, and zircon. The centers of some crystals are corroded and replaced by a fine-grained (possibly chlorite) material which could not be identified under the microscope. The andesine (30 percent) is slightly zoned, and locally altered to calcite and sericite. Only the crystal outline is left of the original pyroxene (10 percent) which has altered

Fig. 1. Highly altered quartz-diorite (?) dike intruding granites of the Sierra de Magdalena. Dikes of this type are common in the central Sierra de Magdalena.

Fig. 2. Photomicrograph of hornblende andesite; south of the Rancho La Jojoba, Sierra de Magdalena. Phenocrysts of hornblende set in a fine-grained holocrystalline groundmass. Note the high degree of oxidation of the hornblende to magnetite. Plain light, 240 X. Sample SP-42.

DIKE AND HORNBLENDE ANDESITE FROM SIERRA DE MAGDALENA

SP-507

SP-503

SP-510.

SP-511.

Photomicrographs of olivine basalts of the Comedores Formation; extreme northwest corner of the Quadrangle. Olivine phenocrysts (dark and medium gray) and large labradorite laths in a semi-pilotaxitic matrix. Note alteration of olivine to iddingsite. Crossed polars, 80 X; Samples SP-507 and SP-511. Plain light, 80 X; Samples SP-503 and SP-510.

PETROGRAPHIC DETAILS OF OLIVINE BASALTS OF THE COMEDORES FORMATION

to calcite, magnetite, chlorite, and locally even to hornblende. The accessory minerals include apatite, zircon, epidote and *green radial chlorite in the matrix*. A chemical analysis of this andesite given in Table 6 is similar to that of andesite of the Tertiary Pachuca formation of central Mexico, but it is higher in potash and lower in lime and total iron than the others shown in Table 6.

COMEDORES FORMATION.—The youngest igneous rocks of the region form a series of flat-topped hills of dark-gray olivine lava flows termed the Comedores Formation. These lava rocks crop out west of the Sierra de La Tortuga and have considerable areal extent. Locally, the flows dip slightly to the west, but they are generally horizontal. Their thickness ranges from about 50 m near the centers of the flows, where widely spaced, well developed columnar jointing was observed in one locality, to 2 m or even less at the edges. Ropy lava surfaces are locally preserved on the youngest of several flows. Stone (personal communication) pointed out pillow bases near the edges of one of the flows.

Megascopically, the Comedores basalts are aphanitic and contain *anhedral*, dark-green, glassy-looking phenocrysts, up to 4 cm in maximum diameter which, upon microscopic examination, are found to be crystals of olivine. Microscopically, the basalts are holocrystalline, aphanitic, porphyritic, and generally somewhat vesicular. Flow surfaces are more vesicular and even wholly amygdaloidal. The groundmass is *intergranular to intersertal and felty* (Plate 15).

The modal mineralogy is that of picrite basalts: olivine (27 percent), labradorite (30 percent), opaque minerals (8 percent), clinopyroxene (probably titanaugite judging by TiO_2 content shown in the chemical analysis of Table 7, and Appendix C), and a yellowish matrix (25 percent) in which serpentine (?) and/or *montmorillonite* (?) is relatively abundant. Calcite and minor quartz are secondary minerals. Olivine occurs as phenocrysts and as small subhedral crystals in the groundmass, where its maximum diameter is generally less than 0.2 mm. Olivine is altered to iddingsite along fractures. Some crystals have abundant inclusions of magnetite and locally are rimmed by it. Labradorite laths attain a maximum length of 0.3 mm. They are locally broken and displaced by micro-faults, and connected by calcite veins. Magnetite inclusions are also common in *labradorite and augite*. A few vesicles are partly filled with calcite.

Prof. Eduardo Schmitter, of the Instituto de Geología de la Universidad Nacional Autónoma de México, studied three samples of the Comedores basalts and reported that the clinopyroxene is augite, and that some cavities contain zeolites.

AGE AND PETROGENESIS OF THE "IGNEOUS COMPLEX" AND COMEDORES FORMATION.—The age of rocks elsewhere in Sonora which are similar mineralogically and genetically to those of the "Igneous Complex" has been the subject of some speculation and has aroused the interest of several geologists. Damon and others (1962) have determined a radiometric age of 85 ± 6 million years for sialic intrusive bodies around El Arpa, Sonora. Independently, Pantoja (personal communication) had suggested a Late Cretaceous or early Tertiary age for these rocks.

TABLE 6.—Chemical composition of andesites

	SP22-5	I	II	III	IV
SiO ₂	60.69%	61.12%	61.49%	60.8%	54.20%
TiO ₂	0.74	0.42	0.96	0.84	1.31
Al ₂ O ₃	17.67	17.65	14.98	15.80	17.17
Fe ₂ O ₃	1.92	2.89	1.51	2.00	3.48
FeO	2.14	2.40	4.05	2.90	5.49
MnO	0.08	—	—	0.08	0.15
MgO	3.15	2.44	3.22	3.50	4.36
CaO	3.87	5.80	4.56	5.70	7.92
Na ₂ O	4.08	3.83	3.59	3.30	3.67
K ₂ O	3.25	1.72	2.80	2.20	1.11
P ₂ O ₅	0.33	—	—	0.24	0.28
SO ₃	—	—	—	—	—
CO ₂	0.54	—	—	0.71	—
H ₂ O+	1.60	1.43	1.91	2.20	0.86
H ₂ O-	0.10	—	—	—	—
S	—	—	—	—	—
	100.16%	100.00%	100.45%	100.3%	

SP22-5* — Hornblende andesite, Coyotillo group, Santa Ana Quadrangle, Sonora, Mexico.

I — Hornblende andesite, av. 20 analyses; Daly, 1933.

II — Hornblende andesite, Glencoe, Scotland; Mem. Geol. Survey, p. 183, 1916.

III — Andesite, Pachuca Formation, Pachuca — Real del Monte Mining District, Hidalgo, Mexico; A. R. Geyne *et al.*, Consejo de Recursos Naturales No Renovables, Publ. 5E, p. 70, 1963.

IV — Average effusive andesite; S. R. Nockolds, Bull. Geol. Soc. America, v. 65, 1954.

* Analyst: Ing. Quim. Alberto Obregón, Instituto de Geología, Universidad Nacional Autónoma de México.

Two independent lines of evidence indicate that the Comedores Formation is probably of Plio-Pleistocene age and younger than the Báucarit Formation. First, the Báucarit Formation contains no clasts of basalt in the Santa Ana Quadrangle, suggesting pre-Comedores deposition. Second, well marked angular unconformities within the Báucarit Formation were observed in a few localities. The basalts have not been subjected to strong tectonic movements and are therefore younger than at least part of the Báucarit Formation. The angular unconformities within the Báucarit Formation also support the view that the Báucarit Formation, at least in part, is of different Tertiary age.

TABLE 7.—Chemical composition of basalts

	SP-155	I	II	III
SiO ₂	47.53%	43.69%	46.77%	50.83%
TiO ₂	2.16	2.12	3.00	2.03
Al ₂ O ₃	15.60	9.06	14.65	14.07
Fe ₂ O ₃	4.90	3.46	3.71	2.88
FeO	5.10	9.43	7.94	9.06
MnO	0.19	0.16	0.15	0.18
MgO	7.46	19.68	6.82	6.34
CaO	9.84	9.18	12.42	10.42
Na ₂ O	3.00	1.49	2.59	2.33
K ₂ O	1.28	0.69	1.07	0.82
P ₂ O ₅	0.65	0.30	0.37	0.23
SO ₃	—	—	—	—
CO ₂	—	—	—	—
H ₂ O +	2.38	0.74	0.51	0.91
H ₂ O -	0.19	—	—	—
S	—	—	—	—
	100.28%			

SP-155* — Olivine basalt, Comedores Formation, Santa Ana Quadrangle, Sonora, Mexico.

I — Average olivine rich basalts; S. R. Nockolds, Bull. Geol. Soc. América, v. 65, 1954.

II — Average alkali basalts; S. R. Nockolds, Bull. Geol. Soc. América, v. 65, 1954.

III — Average tholeiite basalts; S. R. Nockolds, Bull. Geol. Soc. America, v. 65, 1964.

* Analyst: Ing. Quím. Alberto Obregón, Instituto de Geología, Universidad Nacional Autónoma de México.

The genesis of both Precambrian and Cretaceous plutonic rocks was probably magmatic. Granites are associated with rhyolites and andesites in the "Igneous Complex," and the Precambrian granites are intruded by basic and intermediate dikes in the Santa Ana Quadrangle as well as by dikes of lamprophyre and diabase in El Arpa and Bátori in northwestern Sonora (Damon and others, 1962). Moreover, dikes and apophyses from these major granitic bodies cut the enclosing country rocks. The author disagrees with de Cserna (1961) who stated that Cretaceous plutonic rocks in northwest Mexico, are "anatexitic rocks" formed by high temperature metamorphic processes.

Although one may consider the basalts of the Comedores Formation as representative of the "alkaline-olivine basalt association on the continents" of Turner and Verhoogen (1960) who, incidentally, postulated a primary olivine basalt magma for this kindred, it is more difficult to assign the granites and the andesites to definite igneous rock associations. Until more regional petrologic

studies are made, it is probably not a meaningful pursuit. However, as is described subsequently, de Cserna (1961) classified the igneous rock associations of northwest Mexico on the basis of tectonic relationships.

GEOLOGY OF THE SIERRA DE LA LUMBRERA

The volcanic rocks that crop out in the Sierra de La Lumbra (Plate 16), east of Santa Ana, were mapped in detail in the present study. In his reconnaissance mapping of 1927, Flores divided the volcanic rocks into three units (hornblende andesite, basalt, and tuff) but five units were recognized in the present study: basaltic andesite, andesite breccia, hornblende andesite, olivine basalt, and lithic-crystal tuff. In addition, a sedimentary sequence was mapped and described (Plate 22).

SEDIMENTARY ROCKS.—A sequence of sedimentary rocks approximately 90 m in thickness is exposed in the northwestern part of the Sierra de La Lumbra as an inlier in younger volcanic rocks which overlie it with angular unconformity. A paucity of fossils prevents the assignment of an age to this sequence. However, its regional strike suggests possible correlation with the Lower Cretaceous rocks exposed to the west.

Three limestone units were mapped:

Unit K1-1 (oldest): Reddish-brown oolitic limestone. Oolites (80 percent) of calcite mud, from 0.5 to 1 mm in diameter. In most oolites there appears to be no nucleus and radial structures are well preserved. Quartz (?) (10 percent) locally replaces the calcite wholly or in part, without destroying the radial structures. Inclusions of magnetite and hematite are common. Calcite mud (5 percent) fills the interstices between oolites. Traces of detrital quartz are present.

Unit K1-2: White, granoblastic, medium-grained limestone in which euhedral celestite (96 percent) has replaced the original calcite. Only 3 to 5 percent sparry calcite remains with minor magnetite, biotite, and altered rock fragments.

Unit K1-3 (youngest): Pinkish-gray, fine-grained calcisiltite. Silt-size calcite (95 percent) is the main constituent although some sparry calcite has formed as a recrystallization product of the original carbonate. Pellets, magnetite, and quartz are minor.

A few thin shale and siltstone beds are intercalated with the limestones. Partial silicification of the limestone units was observed locally. This is interpreted to be post-diagenetic, perhaps due to descending silica-rich waters from the overlying igneous rocks.

IGNEOUS ROCKS.—*Basaltic andesite.*—The oldest unit of the extrusive sequence consists of basaltic andesites up to 10 m in thickness. They are exposed

VERTICAL AERIAL VIEW OF SIERRA DE LA LUMBRERA

SP —11A

SP —67-6

SP —67-6

PETROGRAPHIC DETAILS OF HORNBLENDE ANDESITES FROM SIERRA DE LA LUMBERA

only near the sedimentary outcrops, and thus they have limited areal distribution. This highly altered, yellowish-green rock is holocrystalline, aphanitic, and porphyritic; its groundmass is intergranular and pilotaxitic. The principal modal minerals are plagioclase (60 percent), magnetite (10 percent), olivine (5 percent), secondary calcite (15 percent), altered pyroxene, and hematite. Plagioclase microlites (averaging in length 0.5 mm) are unaltered whereas the phenocrysts are altered completely to calcite. Olivine and altered pyroxene attain a maximum diameter of 0.5 mm and are euhedral. Most calcite is after plagioclase but it locally fills cavities. Hematite is an oxidation product of magnetite.

Hornblende-biotite andesite dikes which vary in width from 10 cm to 1 m intrude the basaltic andesites. These dikes have a distinctive reddish-brown color and are aphanitic and porphyritic. Megaphenocrysts of biotite, oxyhornblende, and plagioclase (An_{44}) are embedded in a fine, holocrystalline, pilotaxitic groundmass. Magnetite, secondary calcite and hematite are minor.

Andesite breccia.—The southern portion of the Sierra de La Lumbre is composed of andesite breccia exhibiting various degrees of alteration. In a few places where the alteration has been intense (propylitization) all traces of the original texture and much of the original mineralogy have been obliterated. The unaltered rock is light pink and contains angular to subangular fragments of biotite-hornblende andesite, ranging from a few centimeters to 50 cm in maximum diameter, in a red matrix rich in iron oxides (20 percent) and clay minerals.

The rock fragments consist of plagioclase (An_{68}), biotite, hornblende, calcite, magnetite hematite, altered pyroxene, and traces of quartz and chalcedony (?). Plagioclase is present as euhedral to subhedral phenocrysts (1-2 mm averaging in length, and microlites (less than 0.05 mm in length). Oscillatory zoning is common and most of the crystals are either broken or fractured. The cores, rims, or entire crystals of plagioclase are altered to calcite, the distribution of which is controlled largely by compositional zoning and twinning.

Biotite is present as tabular phenocrysts of variable thickness up to 1 mm in length and as euhedral inclusions in the plagioclase. Partially resorbed crystals intimately associated with magnetite are not uncommon and some crystals are bent indicating internal deformation. Hornblende and pyroxene are extensively altered to calcite and/or magnetite and commonly only the crystal outline is preserved. Most quartz and silica (?) are probably secondary but some may be primary. Anhedral quartz "phenocrysts" are corroded and show a slight undulatory extinction.

Although the breccia is the most common lithology, it is by no means the only type found in southern Sierra de La Lumbre. In the extreme southwestern portion, mineralogically similar breccia is characteristically reddish-brown and has a blocky weathered surface. The angular blocks are also hornblende-biotite andesite, but the matrix is more relatively abundant (30 percent) and contains fragments of plagioclase (An_{44}), altered biotite and hornblende, magnetite, and red clay. The rock is commonly propylitized and cut by numerous thin dikes

(up to 10 cm in thickness) of a reddish opaque material which is unidentifiable under the microscope.

Hornblende andesite.—Hornblende andesite is one of Flores' original units and is limited to the eastern margin of the area. The rock is distinctively white to gray and locally exhibits a characteristic banding. Banding was also observed in several localities where the andesite breccia is exposed.

The texture is aphanitic porphyritic and commonly glomero-porphyritic (Plate 17). Prismatic phenocrysts (up to 5 mm or more in length) are oxyhornblende and oscillatory zoning plagioclase (An_{10}) in a felty, hypocrySTALLINE groundmass. The oxyhornblende (15-20 percent) is generally zoned, rimmed by magnetite, and slightly altered to clay minerals and calcite. Andesine (20 percent), diopside (less than 5 percent), and plagioclase microlites (50 percent) are the other principal constituents. Magnetite, hematite, calcite, apatite, sericite, biotite, ilmenite? and leucocene are minor.

At Rancho El Represo the andesites differ somewhat in composition. The plagioclase phenocrysts are more calcic (An_{18}) and more abundant (as much as 40 percent; Plate 18, figure 1). Pyroxene (aegerine-augite or hypersthene?) makes up about 10 percent of the rock whereas hornblende constitutes only 5 percent. A trace of olivine is present.

Olivine basalt.—The northern portion of the Sierra de La Lumbre is characterized by a series of northwest-southeast trending hills of olivine-rich basalt which varies from dark gray to green on fresh surfaces to deep brown on the weathered surfaces. Their texture is holocrystalline, aphanitic, and slightly pilotaxitic and the principal constituents are labradorite (55 percent), magnetite (7 percent), olivine (15-20 percent), and pyroxene (5-10 percent; Plate 18, figure 2). Plagioclase is slightly altered to clays and calcite, and inclusions of apatite and magnetite are common. Radiating semi-circular clusters of plagioclase laths were observed in some thin sections. A trace of chlorite (?) is also present.

Field evidence indicates that there was only one episode of basaltic extrusion. Examination of several thin sections from widely separated localities revealed a range of olivine content from less than 5 percent to more than 20 percent, the larger amounts concentrated in the lower parts of the flow suggesting some gravitational settling.

Lithic-crystal tuff.—Lithic-crystal tuffs are the youngest igneous rocks of the Sierra de La Lumbre. They are exposed in the southeastern portion of the area as long, narrow hills that generally trend northwest and reach a maximum length of approximately 1,000 m and have average width of 20 m and height of 3 m. The main constituents are plagioclase (An_{17} ; 8 percent), altered igneous rock fragments (4 percent), fine-grained silica (quartz?, 4 percent), and partly devitrified glass (75 percent). Biotite, olivine, chlorite, hematite, magnetite, and calcite are present in minor amounts. Plagioclase and rock fragments range from 0.1 mm to 1 mm in maximum diameter. Most plagioclase crystals are subhedral but a few are broken and highly fractured, and some corroded out-

Fig. 1. Photomicrograph of pyroxene-bearing andesite; Rancho El Represo, Sierra de La Lumbre. Crossed polars, 96 X. Sample SP-28.

Fig. 2. Photomicrograph of olivine basalt; Sierra de La Lumbre. Note olivine crystals oxidized to magnetite and hematite (black) in felty groundmass. Plain light, 96 X. Sample SP-67A.

PHOTOGRAPHIC DETAILS OF ANDESITE AND BASALT FROM SIERRA DE LA LUMBRERA

Metaconglomerate unit of the Coyotillo group ("Younger" Precambrian). Main constituents are clasts of quartzite, chert, altered igneous rock fragments, and quartz. Minor amount of magnetite, chlorite, plagioclase, and microcline are present. Sample SP-22-6; natural size.

lines and alteration to sericite were observed. Brown glass and glass shards constitute the groundmass material.

A single outcrop of thin-bedded volcanic ash was found northeast of the northernmost tuff outcrops. The ash is overlain with angular unconformity by the Bâucarit Formation and was probably deposited in a lake as shown by its even, thin stratification.

Chemical composition.—Analyses of igneous rocks from the Sierra de La Lumbra are given in Table 8. Analyses of basalts from different localities throughout the world are given, for comparison, in Tables 9 and 10. CIPW norms are presented in Appendix C.

TABLE 8.—Chemical composition of rocks of the Sierra de La Lumbra

	SP 32* I	SP 29* II	SP 3J* III	SP 12* IV	SP14-2* V	SP67A-1* VI
SiO ₂	66.33%	58.31%	56.60%	64.02%	48.39%	45.03%
TiO ₂	1.85	0.68	0.56	0.41	2.07	1.95
Al ₂ O ₃	13.43	12.44	14.51	13.84	16.18	15.04
Fe ₂ O ₃	4.00	4.55	3.36	0.00	3.30	7.81
FeO	0.08	0.09	0.00	1.33	5.92	1.32
MnO	0.18	0.42	0.27	0.08	0.18	0.15
MgO	1.10	0.08	0.05	1.21	6.75	4.10
CaO	1.65	7.21	7.71	3.14	7.51	11.27
Na ₂ O	1.92	0.60	0.78	0.96	3.61	3.42
K ₂ O	6.07	9.20	9.85	3.17	2.73	2.47
P ₂ O ₅	0.22	0.12	0.37	0.05	1.24	0.80
SO ₃	0.00	0.00	0.00	0.00	0.00	0.00
CO ₂	0.05	5.08	5.59	0.00	0.09	2.94
H ₂ O+	2.68	1.20	0.48	10.55	2.13	3.40
H ₂ O—	0.40	0.16	0.04	1.20	0.11	0.28
S	—	—	—	—	—	—
	99.95%	100.14%	100.17%	99.96%	100.20%	99.98%

I — Altered biotite-hornblende andesite.

II — Biotite-hornblende andesite.

III — Andesite breccia.

IV — Lithic-crystal tuff.

V — Olivine basalt.

VI — Amygdaloidal olivine basalt.

* Analyst: Ing. Quím. Alberto Obregón, Instituto de Geología, Universidad Nacional Autónoma de México.

The andesites (SP-29 and SP-3J) are similar in chemical composition. (Also see Table 6.) Sample SP-32 (a brecciated propylite) contains more silica than SP-3J and SP-29 but less potash and lime. This abundance of silica suggests dacitic composition, but petrographic study reveals an altered andesite in which silicification has occurred. Comparison with the average effusive andesite of Nockolds (1954; Table 6) reveals that the three samples are extraordinarily rich in potash, and, together with the fact that relatively high contents of ferric iron and CO_2 were also recorded, suggests a high degree of alteration. However, if the potash is not the result of alteration, the andesites could have been derived from an alkali-rich magma (normative quartz does not appear in the norm).

Sample SP-14-2 (Table 8) has higher contents of potash and P_2O_5 than those reported for most olivine basalts throughout the world (Table 9) and is similar in this respect to the average alkali basalt of Nockolds (1954; Table

TABLE 9.—Chemical composition of olivine basalts

	SP 14-2	I	II	III	IV
SiO_2	48.38%	48.35%	48.44%	52.9 %	45. %
TiO_2	2.07	2.77	4.29	1.86	
Al_2O_3	16.18	13.18	13.27	18.20	15.
Fe_2O_3	3.30	2.35	4.06	1.80	
FeO	5.92	9.08	8.27	7.30	13.
MnO	0.18	0.14	0.15	0.14	
MgO	6.75	9.72	8.21	4.60	8.
CaO	7.51	10.34	7.72	7.60	9.
Na_2O	3.61	2.42	3.47	3.80	2.5
K_2O	2.73	0.58	1.55	1.00	0.5
P_2O_5	1.24	0.34	0.57	0.12	
SO_3	0.00	—	—	—	
CO_2	0.09	—	—	0.00	
H_2O^+	2.13	—	—	0.60	
H_2O^-	0.11	—	—	—	
Other constituents	—	0.73	—	—	
	100.20%	100.00%	100.00%		93%

I — Average of 53 olivine basalts, Hawaii; G. A. Macdonald, Bull. Geol. Soc. América, v. 60, 1949.

II — Average of 5 olivine basalts, Tutuila, Samoa; R. A. Daly, Am. Acad. Arts Sci. Proc., v. 62, p. 71, 1927. Water free compositions.

III — Olivine basalt, Pachuquilla, Hidalgo, Mexico; A. R. Geyne and others, Consejo de Recursos Nat. no Renovables, Publ. 5E, 1963. Basalt somewhat rich in silica and alumina.

IV — Olivine basalt magma-type; W. Q. Kennedy, Am. Jour. Sci., v. 25, p. 241, 1933.

10). Sample SP-67A1, which is an amygdaloidal olivine basalt, has larger amounts of CO_2 and H_2O than SP-14-2 and the others due to the presence of calcite and zeolites filling cavities.

The lithic-crystals tuffs approach a rhyodacitic composition, except for less soda and more H_2O (Table 10). The absence of ferric iron is noteworthy because rock of a rhyodacitic composition have at least 2 percent ferric iron (Nockolds, 1954). The magma responsible for its formation may have been a residual, relatively silica-rich liquid with low oxygen fugacity.

TABLE 10.—Average chemical composition of alkali basalts and rhyodacites

	SP-67A1	I	SP-12	II
SiO_2	45.03%	46.77%	64.02%	66.27
TiO_2	1.85	3.00	0.41	0.66
Al_2O_3	15.04	14.65	13.84	15.39
Fe_2O_3	7.81	3.71	0.00	2.14
FeO	1.32	7.94	1.33	2.23
MnO	0.15	0.15	0.08	0.07
MgO	4.10	6.82	1.21	1.57
CaO	11.27	12.42	3.14	3.68
Na_2O	3.42	2.59	0.96	4.13
K_2O	2.47	1.07	3.17	3.01
P_2O_5	0.80	0.37	0.05	0.17
SO_3	0.00	—	0.00	—
CO_2	2.94	—	0.00	—
H_2O^+	3.40	0.51	10.55	0.68
H_2O^-	0.28	—	1.20	—
	99.88%		99.96%	

I — Average alkali basalts; S. R. Nockolds, Bull. Geol. Soc. América, v. 65, 1954.

II — Average rhyodacite; S. R. Nockolds, Bull. Geol. Soc. América, v. 65, 1954.

Petrogenesis of the igneous rocks.—Surface eruption of andesites, basalts, and tuffs, probably during and following orogenic activity in Oligocene to Pliocene time, were major events in the area of the Santa Ana Quadrangle. This volcanic phase followed long after Cretaceous folding and plutonic activity which affected the sedimentary rocks and formed the "Igneous Complex" in the Santa Ana Quadrangle. The "alkaline olivine-basalt association on the continents" of Turner and Verhoogen (1960, p. 165) seems appropriate for the basaltic rocks of the Sierra de La Lumbre. A Harker variation diagram for the rocks of the Sierra de la Lumbre is shown in Figure 3. The lack of smoothness of the curves suggests discontinuities in the trends of magmatic evolution and is also

a reflection of the altered nature of the rocks studied. It also suggests that the extrusive rocks of the Sierra de La Lumbre are not comagmatic.

FIG. 3. Harker variation diagram for volcanic rocks of the Sierra de La Lumbre. Total Iron oxide plotted as FeO.

The basalts are much less abundant than the andesites in the Sierra de La Lumbre. Moreover, in Early Cretaceous time great volumes of andesite were erupted with no accompanying basalt. The same is true of other provinces in North America. Thus, on quantitative grounds alone, it seems unlikely that the andesitic magma could have been derived from basaltic magma by fractional crystallization. The restriction of the andesites to an orogenic region in the Sierra de La Lumbre suggests possible mixing of basaltic magma with sialic material as a source of andesitic magma.

Three episodes of magmatic activity are suggested to account for the present distribution of rock types: first, eruption of andesitic lava which presumably formed as a result of mingling of acidic and basic magmas; second, extrusion of basaltic magma; third, extrusion of pyroclastic rocks of rhyodacitic composition.

Volcanism was intermittent and probably interrupted by relatively long periods of weathering and erosion as reflected by the areal distribution and stratigraphic position of the rocks.

Fig. 1. Lineation in metamorphic rocks defined by axes of small crenulations in schist; Sierra de Magdalena. Sample SP-36. Natural size.

Fig. 2. Steep normal fault in metasedimentary rocks; southern Sierra del Potrero.

STRUCTURAL DETAILS OF SIERRA DE MAGDALENA AND SIERRA DEL POTRERO

STRUCTURAL GEOLOGY

Large structural features, their geomorphic effects, and some structural characteristics of the metamorphic and igneous rocks are considered in this chapter. Only the major folds, fracture systems, and faults are shown on the geologic map of the Santa Ana region. Some of the major faulting and jointing was originally suggested in an unpublished report by several geologists of the Consejo de Recursos Naturales no Renovables of Mexico who were doing geochemical sampling at the time of the author's field work.

LINEATION.—Examples of macroscopic lineation in the Santa Ana Quadrangle are of three main classes: (1) elongated or "stretched" pebbles, for example, in the metaconglomerate unit of the Coyotillo group (Plate 19); (2) preferred orientation of elongate crystals, for example, tourmaline in the schists of the Sierra del Potrero (Plate 6, figure 2); (3) axes of small crenulations in schist (Plate 20, figure 1).

Several forms of microscopic lineation are also conspicuous: (1) rotation axes of andalusite porphyroblasts in schists, (2) strings of muscovite in gneiss, and (3) elongate grains of calcite in marbles calc-schists.

The most commonly observed lineation present is that defined by the preferred orientation of mineral grains (Plate 4). The average regional strike of the lineation seems to be northeast.

FAULTS.—The faults and fractures in the Santa Ana Quadrangle may be subdivided into three main groups: (1) a north-south striking fault group, (2) a northeast-southwest striking group, and (3) a northwest-southeast striking group. The geologic map shows examples of groups 2 and 3 but not group 1. However, north-central Sonora is regionally characterized by a series of north-south to slightly northwest trending mountain ranges. These ranges are generally bordered by faults (Eardley, 1962), and, in the Santa Ana Quadrangle, the high metamorphic sierras are probably the geomorphic expression of normal block-faulting. The stress orientation responsible for this deformation was, probably, Sigma 1 vertical, and Sigma 3, horizontal. For any secondary normal faults, such as the one shown in Plate 20, figure 2, a similar stress orientation is proposed.

Mr. L. J. Luccarelli (personal communication) believes that on a regional scale, no major thrust faulting has occurred in the area of the Santa Ana Quadrangle; this view is supported by de Cserna (1961). However, minor thrust faulting was observed in the Santa Ana Quadrangle.

FRACTURES AND JOINTS.—The large-scale fracturing and jointing is shown on the accompanying geologic map. Field observation indicates that the major fractures have vertical or near-vertical dips and a northeast strike. The northwest trending group may be complementary to the main northeast trending group. These fractures commonly control the courses of small streams and

TABLE 11.—Tectonic history of the Santa Ana Quadrangle.

TIME	EVENTS
Quaternary	Basaltic volcanism Faulting
Upper Tertiary	Faulting Báucarit Fm; Red cgl. Faulting Volcanism
Lower Tertiary	Volcanism Block faulting Volcanism
Upper Cretaceous	erosion Folding and faulting Intrusions and volcanism
Lower Cretaceous	Graywackes, limestones, andesites
Jurassic and Triassic	Deposition followed by erosion
Paleozoic	
Precambrian	Present; Coyotillo Group; Younger Precambrian? Igneous intrusions

TABLE 12.—The Mexican Tectonic Cycle, in Sonora.

	STRUCTURAL EVOLUTION	PETROGENIC EVOLUTION
Third Phase	10 — Fissure eruptions probably due to tensional effects in the crust	Eruption of olivine basalt lava flows.
	9 — <i>Taphrogeosynclinal</i> conditions	Deposition of continental debris.
	8 — Eruption of volcanics like those of the <i>Sierra de La Lumbre</i>	Deposition of andesites, andesite breccia, basalts, and tuffs.
Second Phase	7 — Block faulting	<i>Formation of cataclasites in metamorphic rocks.</i>
	6 — Intense folding and faulting	Emplacement of granites and associated rocks.
	5 — Volcanism and/or fissure eruptions	Deposition of andesites and some limestones.
First Phase	4 — Presence of eugeosynclinal conditions	Deposition of thick sequence of graywackes, limestones and shales.
	3 — Minor structural pulses. Shape and location of Coyotillo group dependent on 2	Metamorphism to greenschist facies of the Coyotillo group.
	2 —	Metamorphism to granulite facies with partial melting of pelitic and semi-pelitic rocks. Emplacement of igneous rocks.
	1 — Possible earlier structural episodes not recognized	Depositions of thick succession of pelites, semipelites, quartzite, and minor limestone. Igneous Rocks?

arroyos, and a trellis drainage pattern can be seen in several aerial photographs. Fracture azimuth frequency diagrams compiled from aerial photographs are included in the geologic map for some parts of the area (Plate 4).

FOLDS.—No regional pattern of folding in the metasedimentary rocks is apparent, perhaps in part because of a lack of detailed field information, but also because the rocks are highly deformed (Plate 21). Several dip reversals were observed in the central and eastern parts of the Sierra de La Tortuga, southern Sierra de Guacamea, Sierra de Las Jarillas, Sierra del Picacho, and Sierra del Potrero, suggesting several sets of broad anticlines and synclines.

The Lower Cretaceous sedimentary rocks have variable strikes; although the regional strikes seems to be east-west, southwest of Santa Ana the trend is north, and to the northeast the regional strike is northeast. However, the outcrops are too small and widely separated for meaningful structural correlation and interpretation.

TECTONIC HISTORY.—Table 11 summarizes the tectonic history of the Santa Ana Quadrangle. For comparison, the tectonic and igneous events suggested by de Cserna (1961) as "the Mexican Tectonic Cycle" for all Sonora are included. A summary of the structural and petrogenic events is proposed in Table 12.

ECONOMIC GEOLOGY

No commercial mining is taking place in the area at present. However, the region witnessed intensive activity during the period 1900-1940; the principal products being gold, silver, copper, and manganese, and minor tungsten and iron.

Wisser (1965) assigned the Santa Ana Quadrangle to the "Epithermal Precious Metal Province of Northwest Mexico." A list of the most common features of epithermal deposits which are applicable to the Santa Ana Quadrangle follows:

1. The mineral deposits are typically associated with volcanic rocks of the andesite-family.
2. They were emplaced during Middle to Late Tertiary.
3. They occur mainly as veins occupying fractures.
4. Fluids effecting "hydrothermal alteration" are responsible for the mineralization. The earliest fluids caused propylitization in andesitic rocks, with the formation of chlorite and carbonates. Later fluids caused sericitization and perhaps silicification. Stress behaved in a brittle manner; they were brecciated or split into complex fracture patterns.
5. At relatively shallow depths (from a few hundred to perhaps 600 m) the rocks under patterns. Mineralizing solutions could have taken advantage of this situation.
6. The principal gangue mineral is quartz.

Schmitt (1950) suggested that metallic elements of epithermal ores may have been transferred to their place of deposition by volcanic gas-vapor emanations. This would explain the association of ore deposits with volcanism.

EXPLANATION

MESOZOIC		CENOZOIC	
Tb	Balear formation	Tu	Quaternary
K1-3	Green andesite	Tho	Miocene andesite and conglomerate
K1-2	Green andesite	Tob	Andesite (porphyry and andesite breccia)
K1-1	Limestone		
			PLIOCENE TO OLILOCENE
			LOWER CRETACEOUS?

- SYMBOLS
- Strike and dip of beds
 - Fault, indicates downthrown side dashed where concealed or inferred
 - Fracture
 - Contact line
 - Stream
 - Old road
 - Anticline

0 100 200 300 400 500 600
M E T E R S
GEOLOGY BY GUILLERMO A. SALAS, JULY 1967

GEOLOGIC MAP OF THE SIERRA DE LA LUMBRERA, MUNICIPALITY OF SANTA ANA, SONORA

Minor copper mineralization was observed northwest of the Rancho El Sauz in the Sierra de Las Jarillas. Examination of small abandoned pits revealed abundant localized pockets of malachite which coats the gneissic and schistose country rock in which it is found. In Ejido San Lorenzo, manganese deposits were mined as recently as the 1950's. The ore mineral, psilomelane(?), is restricted to the Lower Cretaceous andesites and limestones.

Flores (1927) subdivided the mineral deposits of the Santa Ana-Carbó region into three types:

1. Those associated with veins; in which the principal ore minerals are: native gold and silver, cerargyrite, wulfenite, malachite, azurite, hematite, limonite, chalcopryrite, pyrite and galena.
2. Ores that are the result of contact phenomena. These are most abundant in the association andesite-limestone (for example, the Mina Jojoba).
3. Replacement ores. The least common of all. These deposits are typically found in limestone. Examples of this occurrence were not found in the Santa Ana Quadrangle.

The reader is referred to Flores' paper for a more complete discussion.

SUMMARY OF GEOLOGIC HISTORY

The following is a brief summary of the geologic events that have occurred in the Santa Ana region.

Sometime approximately 1,600 m. y ago (Damon and others, 1962) a thick sequence of interstratified shales, sandstones, calcilutites, and calcarenites was deposited in a north-northwest trending basin (Eardley, 1962). After sedimentation, these strata were metamorphosed ("Mazatzal Revolution") to various degrees; locally up to *granulite facies*. The metamorphism was accompanied by intrusions of granitic stocks, and followed by the emplacement of mafic dikes.

Erosion of these "Older" Precambrian rocks followed, and, probably during early "Younger" Precambrian time, a coastal plain of low-relief was formed. A marine transgression followed and marine sedimentation became dominant. Orogenic movements and metamorphism occurred during late "Younger" Precambrian time. It is recalled that Damon and others (1962) and Pye (1959) considered that the Altar Schist of northwest Sonora, similar in many respects to those of the Coyotillo group, was metamorphosed during the Laramide Orogeny. However, the present writer believes that the Coyotillo group is of "Younger" Precambrian age for reasons that were stated previously.

Paleozoic rocks do not crop out in the Santa Ana region. However, compilations of regional studies made by Eardley (1962) indicate that sediments accumulated in a north-south trending basin from a source area to the west during Cambrian and Ordovician times. Uplift, folding, and subsequent erosion took place during the Silurian Period, after which uninterrupted sedimentation occurred during the rest of the Paleozoic. Thicknesses of up to 2,000 m may have accumulated locally.

In Early Triassic, sedimentation persisted in a miogeosynclinal (?) basin bordered by land masses to the east and west. Late Triassic and Early Jurassic times witnessed a period of orogenic activity (Eardley, 1962) followed by a long erosional interval.

The Cordilleran Geanticline (Eardley, 1962) and what is now known as the Mexican Geosyncline (P. B. King, 1949) laid to the west and east respectively of the Santa Ana region in Early Cretaceous time. At this time, all of the area covered by the quadrangle was inundated by a shallow sea in which fossiliferous limestones, graywackes, and extrusive rocks (probably erupted under water), were deposited. Folding and some faulting took place from Late Cretaceous to early Tertiary time. Today, the few outcrops of limestone and volcanic rocks in the area of metamorphic rocks are thought to be remnants preserved in down-faulted blocks (Plate 4).

An erosional interval was followed by the late early Tertiary eruption of andesites and basalts. Block faulting was concurrent with the late stages of this magmatic activity. The faulting caused taphrogeosynclinal conditions (de Cserna, 1960). Thick accumulations of clastic material (terrigenous) such as the Báucarit Formation and the "red conglomerates," were deposited in the topographic depressions adjacent to the raised mountain ranges.

Intermittent tectonic movements continued during the rest of late Tertiary and early Pleistocene times, folding fracturing, and faulting the Báucarit Formation.

Recent tensional stresses in the crust facilitated the extrusion of basaltic lava, such as those forming the Comedores Formation. Judging from the thickness of flows, their original areal extent probably covered hundreds of square miles.

REFERENCES CITED

- AGUILERA, J. C., 1896. *Bosquejo geológico de México*: Inst. Geol. México, Bol. 4-6.
- ALENCASTER, GLORIA, 1961. *Estratigrafía del Triásico Superior de la parte central del Estado de Sonora*: Univ. Nal. Autón. México, Inst. Geología, Paleontología Mexicana 11, pt. 1, 18 p.
- ARELLANO, A. R. V., 1956. *Relaciones del Cámbrico de Caborca, especialmente con la base del Paleozoico*. In: *El Sistema Cámbrico, su paleogeografía y el problema de su base*. Congr. Geol. Internat., 20a. Sesión, México, v. 2, p. 509-527.
- BARTH, T. F. W., 1962. *Theoretical petrology*: John Wiley, New York, 416 p.
- BUDDINGTON, A. F., 1959. *Granite emplacement, with special reference to North America*: Bull. Geol. Soc. America, v. 70, p. 671-747.
- BURCKHARDT, KARL, 1923. *Quelques remarques critiques sur l'ouvrage de M. W. Fruedenberg "Geologie von Mexico"*: Mém. Soc. Cient. Alzate, v. 41, 185 p. (Discovery of large Triassic faunas in Sonora).

- COOPER, G. A., and others, 1952. *Cambrian stratigraphy and paleontology near Caborca, northwest Sonora, Mexico*: Smithsonian Miscell. Coll., v. 119, v. 1, 184 p.
- CSERNA, ZOLTAN DE, 1960. *Orogenesis in time and space in Mexico*: Geol., Rundschau, v. 50, p. 595-605.
- , 1961. *Tectonic map of Mexico*: Geol. Soc. America, New York, scale 1:2,500,000.
- DAMON, P. E., LIVINGSTON, D. E., and GILETTI, B. J., 1962. *Extension of the Older Precambrian of the Southwest into Sonora, Mexico*: Geol. Soc. America, Spec. Paper 68, p. 158-159.
- , and others, 1962. *Estudios geocronológicos de rocas mexicanas, parte II, Edad del Precámbrico "Anterior" y de otras rocas del zócalo de la región de Caborca-Altar, de la parte noroccidental del Estado de Sonora*: Univ. Nal. Autón. México, Inst. Geología, Bol. 64, p. 11-44.
- DUMBLE, E. T., 1900a. *Notes on the geology of Sonora, Mexico*: Am. Inst. Min. Eng., Tr., v. 29, p. 122-152.
- , 1900b. *Triassic coal and coke in Sonora*: Bull. Geol. Soc. America, v. 11, p. 10-14.
- EARDLEY, A. J., 1962. *Structural geology of North America*: Harper and Row, New York, 743 p.
- ESKOLA, PENNTI, 1939. *Die metamorphen Gesteine*: In: *Entstehung der Gesteine*, Borntraeger, Berlin, p. 263-407.
- FLORES, TEODORO, 1927. *Reconocimientos geológicos en la región central del Estado de Sonora*: Inst. Geol. México, Bol. 49, 267 p.
- FYFE, W. S., and others, 1958. *Metamorphic reactions and metamorphic facies*: Geol. Soc. America, Mem. 73, p. 1-259.
- GARFIAS, V. R., and CHAPIN, T. C., 1949. *Geología de México*: Editorial Jus, México, 202 p.
- GILETTI, J. B., and DAMON, P. E., 1962. *Rubidium strontium ages of some basement rocks from Arizona and northwestern Mexico*: Bull. Geol. Soc. America, v. 72, p. 639-644.
- GOLDICH, S. S., and others, 1961. *The Precambrian geology and geochronology of Minnesota*: Univ. Minn. Press, Minneapolis, 193 p.
- HATCH, F. H., and others 1961. *Petrology of the igneous rocks*: Thomas Murby, London, 515 p.
- HEINRICH, E. W., 1965. *Microscopic identification of minerals*: McGraw-Hill, New York, 414 p.
- IMLAY, R. W., 1939. *Paleogeographic studies in northwestern Sonora*: Bull. Geol. Soc. America, v. 50, p. 1723-1744.
- KELLER, W. T., 1928. *Stratigraphische Beobachtungen in Sonora, Nordwest Mexico*: Eclogae Geol. Helv., v. 21, p. 327-355.

- KING, P. B., 1949. *Pre-Tertiary history of Sierra Madre Occidental of Sonora and Chihuahua and some adjacent parts of central Sonora*: Proc. Sixth Pacific Sci. Congr., v. 1, p. 217-222.
- KING, R. E., 1934. *Geological reconnaissance in central Sonora*: Am. Jour. Sci. 5th ser., v. 28, p. 81-101.
- , 1939. *Geological reconnaissance in northern Sierra Madre Occidental of Mexico*: Bull. Geol. Soc. America, v. 50, p. 1625-1722.
- KOEPFEN, W., 1931. *Grundriss der Klimakunde*: Grwyter, Berlin, 388 p.
- KRUMBEIN, W. C., and SLOSS, L. L., 1963. *Stratigraphy and sedimentation*: W. H. Freeman, San Francisco, 660 p.
- MILLER, H. W., 1961. *Belemnoides del Triásico Superior del Estado de Sonora*: Univ. Nal. Autón. México, Inst. Geología, Paleontología Mexicana 11, pt. 4, 21 p.
- NOCKOLDS, S. R., 1954. *Average chemical composition of some igneous rocks*: Bull. Geol. Soc. America, v. 65, p. 1007-1032.
- PYE, W. D., 1959. *Catalog of principal sedimentary formation names in southern Arizona and northern Sonora Mexico*: Ariz. Geol. Soc., Guidebook II, p. 274-281.
- RAISZ, ERWIN, 1959. *Landforms of Mexico*: Cambridge, Mass., map with text, scale 1:3,000,000.
- RAMBERG, HANS, 1952. *The origin of metamorphic and metasomatic rocks*: Univ. Chicago Press, Chicago, 317 p.
- SANTILLÁN, MANUEL, 1936. *Synopsis of the geology of Mexico*: Bull. Am. Assoc. Petrol. Geol. v. 20, p. 394-402.
- SCHMITT, H. A., 1950. *Origin of the "epithermal" deposits*: Econ. Geology, v. 45, p. 191-200.
- SILVA, ALICIA, 1961. *Flora fósil de la Formación Santa Clara (Cárnico) del Estado de Sonora*: Univ. Nal. Autón. México, Inst. Geología, Paleontología Mexicana 11, pt. 2, 48 p.
- TALIAFERRO, N. L., 1933. *An occurrence of Upper Cretaceous sediments in northern Sonora, Mexico*: Jour. Geol., v. 41, p. 12-37.
- THORNTWAITE, C. W., 1943. *Problems in the classification of climates*: Geog. Review, v. 33, p. 233-255.
- TURNER, F. J., 1948. *Mineralogical and structural evolution of the metamorphic rocks*: Geol. Soc. America, Mem. 30, 342 p.
- TURNER, F. J., and VERHOOGEN, J., 1960. *Igneous and metamorphic petrology*: McGraw-Hill, New York, 694 p.
- TURNER, F. J., WILLIAMS, H., and GILBERT, C. M., 1954. *Petrography: an introduction of the study of rocks in thin sections*: W. H. Freeman, San Francisco, 406 p.

VALENTINE, W. G., 1936. *Geology of the Cananea Mountains, Sonora, Mexico*: Bull. Geol. Soc. America, v. 47, p. 53-86.

WISSER, EDWARD, 1966. *The epithermal precious-metal province of northwest Mexico*: Nevada Bur. Mines, Report 13, pt. C, p. 63-92.

APPENDIX A

ROCKS OF THE COYOTILLO GROUP ("Younger" Precambrian) OF SONORA.

Unit A: (oldest): Cordierite-bearing quartzite. Quartz (60 percent), cordierite (10 percent), albite (5 percent), chlorite (5 percent), magnetite (4 percent), hematite (4 percent), sericite (3 percent), apatite, zircon, sphene.

Quartz is xenoblastic with numerous inclusions of sericite, apatite, and opaques. Some grains show sutured margins and others are granulated indicating cataclastic deformation. Cordierite is characteristically twinned and altered to pinitic. Chlorite flakes are arranged in semi-parallel manner giving the rock a crude schistose appearance.

Texture of rock: granoblastic, fine-grained, slightly schistose.

Sample SP-22-3.

Unit B: Sericite-chlorite schist. Sericite (60 percent), chlorite (17 percent), quartz (14 percent), epidote (6 percent), hematite (1 percent), apatite, albite, magnetite, and calcite.

Sericite is present in flaky crystals commonly less than 1 mm in length; some crystals bend around larger quartz grains. Minor sericite is alteration of chlorite and albite. Chlorite is either porphyroblastic (up to 2.0 cm in maximum diameter) or minute (less than 0.1 mm in length) flaky crystals. It is characteristically green pleochroic. The cores of some of the porphyroblastic chlorite are highly altered to calcite. Magnetite and apatite inclusions are locally abundant in the chlorite. Quartz is xenoblastic commonly with inclusions of apatite, rutile (?), and chlorite.

Texture of rock: porphyroblastic, fine-grained, schistose.

Rock belongs to the quartz-albite-muscovite-chlorite subfacies of the greenschist facies of regional metamorphism of Turner and Verhoogen (1960, p. 534).

Unit C: Magnetite-rich sericite-chlorite phyllite. Sericite (30 percent), chlorite (28 percent), quartz (25 percent), magnetite (10 percent), biotite, albite, garnet, epidote, and calcite.

Sericite in minute (less than 0.1 mm in length), slender, tabular crystals commonly but not universally aligned parallel to foliation. Chlorite (less than 0.2 mm in length) is brown and green pleochroic and contains abundant inclusions of magnetite. Slightly coarser brown chlorite fills micro-fractures. Chlorite is locally associated with biotite. Quartz (less than 0.2 mm in maximum diameter) is crudely aligned parallel to foliation. Few crystals show poikiloblastic texture with inclusions of chlorite. Magnetite is xenoblastic and minute (less than 0.2 mm in maximum diameter). In hand specimen calcite is filling fractures.

Texture of rock: Fine-grained, phyllitic.

Rock belongs to the quartz-albite-muscovite-chlorite subfacies of the greenschist facies of regional metamorphism of Turner and Verhoogen (1960, p. 534).

Sample SP-9-3

Unit D: Metaconglomerate. Quartz (37 percent), chert (20 percent), igneous rocks fragments (13 percent), chlorite (5 percent), clay minerals (5 percent), albite (4 percent), hematite (4 percent), magnetite (3 percent), zircon, calcite, microcline, and perthite.

Two types of quartz are present: (a) individual grains (up to 1 mm in maximum diameter: 7 percent which are xenoblastic, with sutured margins, and highly fractured, (b) equigranular mosaics of interlocking quartz grains (30 percent) with abundant inclusions of chlorite and idioblastic magnetite. The latter quartz grains show some evidence of cataclastic deformation (slight to moderate undulatory extinction). Chert fragments contain numerous inclusions of quartz, magnetite, chlorite, and microcline. Igneous rocks fragments are highly altered (chloritized). Most fragments retained their original "diabasic" texture. Some of the plagioclase in these fragments is albite. Chlorite is present in minute (less than 0.5 mm in length) flaky crystals; but some chlorite forms radial fibrous aggregates. Chlorite found in altered igneous rocks fragments is oriented parallel to the "lineation" of the original rock. Magnetite and hematite are concentrated in the igneous rocks fragments.

Texture of rock: Crystalloblastic, fine-grained metaconglomeratic with fragments rounded and elongate in direction of foliation, slightly schistose, relict textures in igneous rocks fragments are present.

Sample SP-22-6

Unit E: Sericite-chlorite phyllite. Quartz (40 percent), chlorite (20 percent), sericite (13 percent), magnetite (10 percent), epidote (8 percent), calcite (4 percent), hematite, albite (?), and zircon.

Size of quartz grains ranges from 0.1 mm to 0.4 mm in maximum diameter. Quartz grains are commonly xenoblastic. In the coarser grains some overgrowth is marked by inclusions (magnetite, chlorite). Few quartz grains show poikiloblastic texture. Smaller quartz grains are aligned with schistosity. Chlorite and sericite (0.1 to 0.2 mm in maximum length) are commonly aligned parallel to schistosity. Size of magnetite grains varies from less than 0.1 mm to 0.5 mm in maximum diameter. Smaller crystals are xenoblastic whereas larger ones are strongly idioblastic. Magnetite alters slightly to hematite. Epidote (less than 0.2 mm in maximum diameter) occurs in aggregates and in individual xenoblastic to semi-idioblastic crystals with abundant inclusions of magnetite.

Texture of rock: Fine-grained, phyllitic, slightly schistose.

Rock belongs to the quartz-albite-muscovite-chlorite subfacies of the greenschist facies of regional metamorphism of Turner and Verhoogen (1960, p. 534).

Sample SP-9-5

Unit F: Metagraywacke. Quartz (28 percent), calcite (18 percent), chlorite (15 percent), cordierite (7 percent), albite (7 percent), microcline (5 percent), sericite (5 percent), hematite (after pyrite, 5 percent), magnetite (4 percent), zircon, epidote, and apatite.

Most quartz grains are xenoblastic (less than 0.1 mm to 0.6 mm in maximum diameter): a few are interlocking grains with idioblastic outlines. Some quartz has indented margins against sericite and/or calcite and some show poikiloblastic texture with inclusions of sericite, magnetite, calcite, and chlorite. Calcite is xenoblastic and probably represents part of the original cement which has recrystallized. Inclusions of magnetite and hematite stain are common. Parts of hand specimen and thin section lack calcite; this is interpreted to be the result of weathering (leaching). Chlorite and sericite are minute, slender, tabular to flaky crystals (less than 0.2 mm in length). Cordierite shows characteristic poikiloblastic texture (inclusions of pinites) and has a brownish color without crossed polars, probably due to hematite stain. Albite and microcline are relatively free of alteration. Large (up to 1 cm in maximum diameter), idioblastic crystals of hematite-pyrite were observed in the hand specimen.

Texture or rock: Granoblastic, fine to medium-grained, poikiloblastic.

Sample SP-9-4

Unit G: (youngest): Impure marble. Calcite (60 percent), quartz (25 percent), hematite (10 percent), magnetite (3 percent), sericite (2 percent), zircon, epidote, apatite, albite, and biotite.

Calcite is xenoblastic in minute (less than 0.1 mm in maximum diameter) crystals with numerous inclusions of magnetite and hematite. Localized patches of slightly coarser calcite were observed. Calcite filling fractures is much finer-grained. Distribution of most hematite and magnetite is fracture-controlled. Amount of hematite (stain) decreases with increasing distance from fractures. Minute (less than 0.1 mm in length; locally up to 1 mm in maximum length) crystals of sericite are partially aligned with the foliation giving the rock a semi-schistose character.

Texture of rock: Granoblastic, fine to medium-grained, schistose; fractures very abundant.

Sample SP-202

APPENDIX B

MEASURED SECTION OF LOWER CRETACEOUS ROCKS AT RANCHO EL REPRESO

Measured July 25, 1966; revised July 4, 1967.

	Thickness (meters)
37. Shale; light to yellow on both fresh and weathered surface, blocky fracture; Strike N57°E, dip 54°SE; valley former	29.9
36. Covered; limestone float; ridge former	29.6
35. Limestone; gray on both fresh and weathered surface, fine-grained (calcite mudstone), massive, thick-bedded; <i>Ostrea</i> sp. abundant; highly fractured, strike of fractures N23°W, vertical dip; ridge former; fetid odor	56.2
34. Siltstone and shale; light yellow on both fresh and weathered surface. Siltstone is fine-grained, has blocky fracture. Unit is intercalated with thin layers of fossiliferous limestone; valley former	35.0
33. Limestone; light brown of weathered surface, light gray of fresh surface; fine-	

	Thickness (meters)
grained (calcsiltite); massive, thick-bedded; <i>Pecten</i> sp, <i>Ostrea</i> sp, and crinoid stems abundant; reddish patches due to hematite stain; strike N58°E, dip 40°SE; ridge former	2.2
32. Siltstone; light brown on both fresh and weathered surface, coarser-grained, thin-bedded; intercalated with lenses of very thin-bedded, light brown to gray fossiliferous limestone	32.4
31. Covered interval, probably siltstone or shale	10.6
30. Siltstone and shale; light brown to deep reddish-brown on both fresh and weathered surface, thin-bedded, blocky fracture. Sequence is an alternation of siltstones and shales	40.8
29. Covered interval, probably siltstone or shale	23.8
28. Siltstone and shale; light brown to deep reddish-brown on both fresh and weathered surface in lower part of unit. Conglomerate in upper 5 meters, gray to green on fresh surface; rounded to sub-rounded fragments of chert, quartzite, metamorphic and igneous rock fragments; hematite and silica cement. Average size of fragments is 0.5 cm	16.0
27. Limestone; same as Unit 23	0.6
26. Covered interval, probably limestone	6.0
25. Limestone, same as Unit 23	0.3
24. Covered interval, probably limestone	3.9
23. Limestone; light gray on both fresh and weathered surface; fine-grained (calcsiltite and calcite mudstone), massive, thick-bedded; <i>Ostrea</i> sp, crinoid stems abundant, <i>Pecten</i> sp. absent; highly fractured, strike of fractures N16°E, almost vertical dip. Attitude of beds: strike N45°E, dip 38°SE; ridge former.	1.5
22. Limestone; light gray on both fresh and weathered surface; fine-grained (calcsiltite and calcite mudstone), massive, thick-bedded; <i>Pecten</i> sp. <i>Ostrea</i> sp. crinoid stems abundant; ridge former. This unit is in places intercalated with thin redbrown limestones	4.3
21. Limestone; dark red-brown on both fresh and weathered surface; fine-grained (calcsiltite), massive, thick-bedded; <i>Ostrea</i> sp. common; weathered surface is rough; ridge former	6.9
20. Siltstone; dark brown on weathered surface, light brown on fresh surface; fine-grained, thin-bedded, blocky fracture, ridge former.	3.3
19. Calcareous siltstone; light brown on weathered surface, pinkish on fresh surface; fine-grained, medium-bedded, blocky fracture, fossils absent, abundant calcite cement, ridge former	7.2
18. Covered interval, probably siltstone	18.8
17. Limestone; light on weathered surface, brown on fresh surface; <i>Ostrea</i> sp. abundant; massive, medium to thick-bedded	0.5
16. Covered interval, probably limestone	3.3
15. Limestone; dark red-brown on both fresh and weathered surface; massive, thick-bedded; <i>Pecten</i> sp. and <i>Ostrea</i> sp. common; rough weathered surface, ridge former	2.3
14. Covered interval, probably limestone, ridge former	48.5
13. Limestone (largely silicified calcarenite); gray to brown on both fresh and weathered surface; massive, thick-bedded; <i>Pecten</i> sp, <i>Ostrea</i> sp, and crinoid stems abundant; highly fractured, most fractures filled with calcite; attitude: strike N47°E, dip 49°SE	1.9
12. Siltstone and shale; tan on both fresh and weathered surface; fine-grained, blocky fracture; intercalated with thin limestone beds	7.0
11. Limestone; gray on both fresh and weathered surface; fine to medium-grained	

	Thickness (meters)
<i>calcsiltite to calcarenite</i>), massive, thick-bedded; <i>Ostrea</i> sp. very abundant; ridge former	3.8
10. Shale; alternating tan and pink on both fresh and weathered surface; with thin (15 cm) intercalations of shaly limestone	7.3
9. Limestone; gray on both fresh and weathered surface; fine to medium-grained (<i>calcsiltite to calcarenite</i>), massive, thick-bedded, weathered surface is very rough; <i>Pecten</i> sp, <i>Ostrea</i> sp, and crinoid stems abundant; highly fractured; locally oxidized to a deep red-brown color, ridge former; strike N61°E, dip 29°SE	26.0
8. Limestone; gray on both fresh and weathered surface; fine-grained, (<i>calcsil- tite and calcite mudstone</i>), masive, thick-bedded, weathered surface is very rough; <i>Pecten</i> sp. and <i>Ostrea</i> shells abundant; jointed, some fracture open, others filled with calcite; ridge former	18.7
7. Siltstone; tan to light brown on both fresh and weathered surface; fine grained, calcite cemented; blocky fracture; fossils absent	6.1
6. Covered interval, shale float on top	1.9
5. Limestone; sandy; light tan on both fresh and weathered surface; fine-grained, thin-bedded; some mollusk shells present; numerous vertical fractures trending N80°W were observed	0.9
4. Shale; brownish-red on both fresh and weathered surface fine-grained (<i>calcsil- tite to calcite mudstone</i>); massive thick-bedded; blocky fractures; with thin (less than 3 cm) intercalations of limestone	8.2
3. Limestone; reddish-brown on both fresh and weathered surface; fine-grained (<i>calcsiltite to calcite mudstone</i>); massive, thick-bedded; contains <i>Pecten</i> sp. shells and corals (?); ledge former; parts are shaly and sandy	1.6
2. Shale; brownish-red on both fresh and weathered surface; fine-grained, thin- bedded, blocky fracture, valley former	4.4
1. Limestone; light gray to tan on both fresh and weathered surface; fine-grained (<i>calcsiltite to calcite mudstone</i>); massive, thick-bedded; abundant <i>Ostrea</i> sp. shells, other mollusk shells less prominent; ridge former	4.7
0. Siltstone and shale; light brown on fresh surface, tan on weathered surface; siltstone is fine-grained, thin-bedded, blocky fracture; valley former	12.6
Measured thickness of El Represo Section	485.0

APPENDIX C

CHEMICAL ANALYSES, NORMS, AND MODES OF THE IGNEOUS ROCKS, SANTA ANA QUADRANGLE, SONORA.

LOCALITY	EXTREME NW OF QUAD.	COYOUILLO GROUP	SIERRA DE GUACOMEA	SIERRA DE LA LUMBRERA	
				SP-67AI Amygdaloidal olivine basalt Chemical analysis*	SP-142 Olivine basalt Chemical analysis*
Sample number	SP-155	SP-152	SP-22-5		
Rock name	Olivine basalt Chemical analysis*	Fine-grained leucogranite Chemical analysis*	Hornblende andesite Chemical analysis*		
SiO ₂	47.53	60.69	73.94	45.03	48.38
TiO ₂	2.16	0.74	0.10	1.85	2.07
Al ₂ O ₃	15.60	17.67	13.96	15.04	16.18
Fe ₂ O ₃	4.90	1.92	1.20	7.81	3.30
FeO	5.10	2.14	0.06	1.32	5.92
MnO	0.19	0.08	0.10	0.15	0.18
MgO	7.46	3.15	0.25	4.10	6.75
CaO	9.84	3.87	0.87	11.27	7.51
Na ₂ O	3.00	4.08	3.98	3.42	3.61
K ₂ O	1.28	3.25	5.35	2.47	2.73
P ₂ O ₅	0.65	0.33	0.00	0.80	1.24
CO ₂	0.00	0.54	0.00	2.94	0.09
H ₂ O ¹	2.38	1.60	0.16	3.40	2.13
H ₂ O ²	0.19	0.10	0.03	0.28	0.11
Total	100.28	100.16	100.00	99.88	100.20

¹Mainly clay and mica.

²Laboratorio de Química, Instituto de Geología, Universidad Nacional Autónoma de México.

**Mainly magnetite; probably some ilmenite.

***Modal analysis denotes a vitric tuff.

APPENDIX C

CHEMICAL ANALYSES, NORMS, AND MODES OF THE IGNEOUS ROCKS, SANTA ANA QUADRANGLE, SONORA.

(Continuation)

	Norm (C. I. P. W. Classification)	Norm (C. I. P. W. Classification)	Norm (C. I. P. W. Classification)
Quartz	7.78	28.40	12.78
Orthoclase	7.78	31.14	19.46
Albite	25.15	33.54	34.58
Anorthite	25.29	4.17	14.45
Corundum		0.20	
Diopside	15.80		
Hypersthene	7.92		9.32
Magnetite	7.10	0.23	2.78
Hematite		1.12	4.84
Ilmenite	4.10	0.15	3.04
Apatite	1.34		2.01
Rutile			0.32
Calcite			6.70
Olivine	2.92		2.00
Total	95.27	99.55	13.72
Normative plagioclase	An 50	An 11	102.60
Rock Classification	35.44	1413	An 40 2534

¹Mainly clay and mica.

*Laboratorio de Quimica, Instituto de Geología, Universidad Nacional Autónoma de México.

**Mainly magnetite; probably some ilmenite.

***Modal analysis denotes a vitric tuff.

APPENDIX C
CHEMICAL ANALYSES, NORMS, AND MODES OF THE IGNEOUS ROCKS, SANTA ANA QUADRANGLE, SONORA.
(Continuation)

	Mode	Mode	Mode	Modes
Quartz	trace			
Microcline-perthite		trace	25.6	
Myrmekite			20.7	
Plagioclase			2.8	
Hornblende	30 (An 47)	10 (andesine)	35.3 (An 14)	30 55 (labradorite)
Biotite		10.5		
Chlorite		1	0.8	
Pyroxene	?	10.4	0.1	trace
Opaque minerals**		5.1		11
Apatite	8.5	trace	1.5	10.5
Zircon		trace		trace
Sphene			trace	
Ilmenite	3			
Epidote		1		4
Calcite	trace	1.5		5
Ilmenite		trace		
Olivine	27		trace	10
Garnet				20
Zeolites			4.5	
Serpentine?	23			20
Igneous rock fragments				4
Volcanic glass				
Feldspar alteration ¹		20	5.3	
Groudmass, undifferentiated		7		

¹Mainly clay and mica.

*Laboratorio de Química, Instituto de Geología, Universidad Nacional Autónoma de México.

**Mainly magnetite; probably some ilmenite.

***Modal analysis denotes a vitric tuff.

APPENDIX C
CHEMICAL ANALYSES, NORM, AND MODES OF THE IGNEOUS ROCKS, SANTA ANA QUADRANGLE, SONORA.
(Continuation)

LOCALITY		SIERRA DE LA LUMBRERA			
	Sample number Rock name	SP-31 Andesite breccia	SP-29 Propylite Chemical analysis*	SP-32 Brecciated propylite (Dacite?)	SP-12 Lithic crystal tuff***
		56.60 0.56 14.51 3.36 0.00 0.27 0.05 7.71 0.78 9.85 0.37 5.59 0.48 0.04 100.17	58.31 0.68 12.44 4.55 0.09 0.42 0.08 7.21 0.60 9.20 0.12 1.20 5.08 0.16 100.14	66.32 1.85 13.43 4.00 0.08 0.18 1.10 1.65 1.92 6.07 0.22 2.68 0.05 0.40 99.95	64.02 0.41 13.84 0.00 1.33 0.08 1.21 3.14 0.96 3.17 0.05 0.00 10.55 1.20 99.96
	SiO ₂				
	TiO ₂				
	Al ₂ O ₃				
	Fe ₂ O ₃				
	FeO				
	MnO				
	MgO				
	CaO				
	Na ₂ O				
	K ₂ O				
	P ₂ O ₅				
	Total				
	H ₂ O ⁺				
	H ₂ O ⁻				
	Total				

¹Mainly clay and mica.

^{*}Laboratorio de Química, Instituto de Geología, Universidad Nacional Autónoma de México.

^{**}Mainly magnetite; probably some ilmenite.

^{***}Modal analysis denotes a vitric tuff.

APPENDIX C
CHEMICAL ANALYSES, NORM. AND MODES OF THE IGNEOUS ROCKS, SANTA ANA QUADRANGLE, SONORA.
(Continuation)

	Norms (C. I. P. W. Classification)			
Quartz	13.86	18.36	26.90	30.96
Orthoclase	57.82	54.24	46.14	18.90
Albite	6.81	5.20	16.24	8.38
Anorthite	1.94	2.22	6.16	14.76
Corundum	1.80		0.91	3.26
Diopside			2.70	
Hypersihene		0.10		4.84
Magnetite		1.60		
Hematite	3.52	3.52	4.00	
Ilmenite	0.61	1.06	0.60	0.76
Apatite	1.10	0.33	0.33	0.33
Rutile	0.24	0.16	1.52	
Calcite	12.70	11.60		
Olivine				
Total	100.31	98.49	104.51	82.16
Normative plagioclase	An 22	An 28	An 27	An 65
Rock classification				

¹Mainly clay and mica.
^{*}Laboratorio de Química, Instituto de Geología, Universidad Nacional Autónoma de México.
^{**}Mainly magnetite; probably some ilmenite.
^{***}Modal analysis denotes a vitric tuff.

APPENDIX C
CHEMICAL ANALYSES, NORM, AND MODES OF THE IGNEOUS ROCKS, SANTA ANA QUADRANGLE, SONORA.
(Continuation)

	Modes				
Quartz		9.5	5.1	4.4	
Microcline-perthite					
Myrmekite					
Plagioclase	28 (andesine?)	45 (andesine)	33.5 (An 38)	8.5 (An 17)	
Hornblende					
Biotite	3	5.3	7.1	trace	
Chlorite		4.1		trace	
Pyroxene			?		
Opaque minerals**	1.5	20	20	trace	
Apatite		trace?			
Zircon			trace		
Sphene					
Ilmenite					
Epidote					
Calcite	13	4.5	2	trace	
Hematite	10	3	5	1.5	
Olivine				trace	
Garnet					
Zeolite					
Serpentine?					
Igneous rock fragments					5
Volcanic glass					40
Feldspar alteration ¹	trace	10.5	2		
Groundmass, undifferentiated	31		20	40	

¹Mainly clay and mica.

*Laboratorio de Química, Instituto de Geología, Universidad Nacional Autónoma de México.

**Mainly magnetite; probably some ilmenite.

***Modal analysis denotes a vitric tuff.