Conclusion. The data obtained indicate a statistically confirmed positive effect of the developed therapeutic complex on the outcomes of unstable angina pectoris against the background of type II diabetes mellitus and with concomitant hyperventilation syndrome. This is confirmed by the duplicating criteria for the integral assessment of the outcome – the formation of the functional class of angina pectoris and the level of tolerance in physical activity – at the end of the period of inpatient treatment (while the results of assessing the outcome using the two alternative methods coincide). The results achieved in this study make it possible to recommend the developed therapeutic complex for implementation in healthcare practice.

1. Cardiology: national manual / E.V. Shlyakhto [et al.]. – Moscow, 2021. – 800 p. – ISBN 978-5-9704-6092-4.
2. Chazov E.I. The problem of treatment of patients with ischemic heart disease / E.I. Chazov // Therapeutic archive. – 2000. – № 9. – P. 59.
3. Starodubov V.I. Public health and health care: national manual / V.I. Starodubov. – Moscow, 2014. – 624 p. – ISBN: 9978-5-9704-2909-9.
4. Marchenko V.N. A comorbid patient in the practice of a therapist for IHD and COPD [electronic resource] / V.N. Marchenko – <https://petrsu.ru/files/user/3324613444dafceb8c73c8ded946eed1/Marchenko%20VN%20Comorbid%20pdf> (date of application: 05/28/2021).
5. Mitkovskaya N.P. Cardiology and comorbidity / N.P. Mitkovskaya // «I International Congress of Cardiologists and Therapists (Minsk, May 12-13)»: collection of scientific papers, 2016 – Minsk, 2016. – P. 173-177.
6. Endocrinology: national manual / I.I. Dedov [et al.]. – Moscow, 2013. – 752 p.
7. Diabetes mellitus and coronary heart disease: search for solutions / A.A. Aleksandrov [et al.] // Diabetes mellitus. – 2005. – № 1. – P. 34-38.
8. Da Costa J.M. On Irritable Heart: a Clinical Study of a Form of Functional Cardiac Disorder and its Consequences / J.M. Da Costa // The American journal of the medical sciences. – 1871. – Vol. 121, № 1. – P. 2-52.
9. Agadzhanian N.A. Hypocapnic and hypercapnic states / N.A. Aghajanyan. – Moscow, 2003. – 35 p.
10. Measurement of breathlessness in advanced disease: a systematic review / C. Bausewein [et al.] // Respiratory Medicine. – 2007. – Vol. 101. – P. 399-410.
11. Dyspnea catastrophizing and neural activations during the anticipation and perception of dyspnea / M.S. Stoeckel [et al.] // Psychophysiology. – 2017. – Vol. 55, № 2. – P. 1-10. – DOI: 10.1111/psyp.13004.
12. Abrosimov V.N. Hyperventilation syndrome in the clinic of internal diseases: dis. ... MDPhD / V.N. Abrosimov. – Ryazan, 1991. – 239 p.
13. Tokareva N.A. Hyperventilation syndrome in somatic pathology (for example, bronchial asthma, hypertension) and organ neurosis – features of the course and treatment: dis. ... PhD / N.A. Tokareva. – Moscow, 2004. – 137 p.
14. Bakhmutova Yu.V. Diagnostics and correction of hyperventilation syndrome in patients with diabetes mellitus by means of BFB-CO₂ technology / Yu.V. Bakhmutova, A.F. Pyatakovich, T.I. Yakunchenko // International Journal of Applied and Fundamental Research. – 2009. – № 6. – P. 36-39.
15. Borzenko S.Ya. Hyperventilation syndrome (vegetative-vascular dystonia syndrome) [electronic resource] / S.Ya. Borzenko. – <http://moi03.ru/node/42> (date of application: 06.06.2021).
16. Brezgunov A.V. Evaluation of the outcomes of acute coronary syndrome based on the results of the introduction of modern standards of therapy / A.V. Brezgunov. – 69th scientific-practical conference of students and young scientists with international participation «Actual problems of modern medicine and pharmacy-2015»: collection of scientific papers. – Minsk, 2015. – P. 306-310.
17. Rebrova O.Yu. Statistical analysis of medical data: application of the STATISTICA software package / O.Yu. Rebrova. – Moscow, 2002. – 312 p.
18. Fletcher R. Clinical epidemiology: evidence-based medicine / R. Fletcher, S. Fletcher, E. Wagner. – Moscow, 1998. – 352 p.

Zhirov A.I., Zhirova V.G., Galkina O.P.

Electronic microscopic research of the surface of the teeth after professional oral hygiene

Institute "Medical Academy named S.I. Georgievsky" FGAOU VO "KFU named V. I. Vernadsky" (Russia, Simferopol)

doi: 10.18411/gqws-01-2022-10

Abstract

The article presents the results of electronic microscopic and X-ray spectral studies on the study of the surface microstructure and chemical composition of enamel and dentin during

professional oral hygiene. Differences in the mineral composition of hard tissues of the tooth surface were determined for various types of ultrasonic treatment.

Keywords: electronic microscopic examination, microanalysis, types of ultrasonic treatment, mineral composition, dentin sensitivity.

The high prevalence of periodontal disease in the world as a whole (80% according to WHO) and Russia (60-100%), detected during epidemiological research in last years, indicates the need for regular professional hygiene.

The significant prevalence of periodontal diseases, the severity of the course of some of its forms determine the significance of the problem and serve as the basis for attributing the issues of early prevention and therapy to one of the topical issues of modern dentistry.

Ultrasonic, air or manual removal of subgingival dental plaque is of prime importance during the prevention and treatment of periodontal diseases. During this treatment, changes in the cement of the tooth root occur, which is a constant source of intoxication of the periodontal tissues as a result of the retention of bacterial conglomerates and their waste products. After its removal, the subsequent polishing of the tooth surface is very important, which, however, is not always used.

Purpose of the work: to study microscopic changes with a quantitative determination of microelements (Ca and P) in hard tissues, after ultrasonic removal of dental plaque and subsequent treatment with an air-abrasive device.

To achieve this goal, we examined 28 samples of human teeth removed for clinical indications. The study of the surface was carried out using a scanning electron microscope "SELMIT" REM-106, at magnifications of 12-5000 times. The teeth were divided into three groups. The first A group included 12 teeth, divided into 3 subgroups according to age.

The second group included 12 freshly removed teeth, which were divided into 3 subgroups. B1 subgroup (4 teeth) performed ultrasonic removal of hard dental plaque. B2 - removal of hard dental plaque with an ultrasonic sceller, without cooling. B3 - removal of hard dental deposits with an ultrasonic sceller followed by polishing. Group C (4 teeth) - removal of hard dental plaque with Sandman Futura treatment and polishing.

According to the results of electronic microscopic examination, in the first A group, significant differences were found in the diameter of the dentinal tubules of root dentin in terms of age. The diameter of the tubules in the group of patients aged 21-30 years, according to our observations, was $0.92 \pm 0.08 \mu\text{m}$ ($p < 0.05$).

In the teeth samples of the A2 study subgroup (age group 31-40 years old), these values corresponded to $0.69 \pm 0.04 \mu\text{m}$, the difference between them was statistically significant ($p < 0.05$). When examining samples of teeth of subgroup A3 (age group 41-50), the data for diameters were $0.52 \pm 0.05 \mu\text{m}$, the discrepancies were significant ($p < 0.05$).

Age-related changes in the number of dentinal tubules were also statistically significant. In particular, the number of dentinal tubules of the root part of the teeth of subgroup A1 was $27.9 \pm 1.6 \times 10^3 \text{ units} / \text{mm}^2$ ($p < 0.05$).

In the samples of teeth A3, which corresponded to the age category 44-59, the difference in the density of dentinal tubules was statistically significant ($p < 0.05$) and amounted to $13.6 \pm 1.1 \times 10^3 \text{ units} / \text{mm}^2$.

The data obtained showed that the number and diameter of dentinal tubules decrease with age due to their obliteration with mineral substances of the tooth. The difference in dentine mineralization in different age periods was established.

According to the results of the study, the presence of lateral branches of the processes in the predentinum was established. Branches usually branch off at right angles from the main branch stem (Picture 1). The system of lateral branches of odontoblast processes ensures the transfer of nutrients and ions, and promotes the lateral spread of microorganisms and acids, provoking the development of caries.

Picture 1. Electron diffraction pattern of a sample of group A. The process of odontoblasts with branches in the dentinal tubule

The number of dentinal tubules per mm² of surface area decreased with age in the direction from the pulp to the dentin-enamel junction.

In the study of samples of subgroup B1 after treatment with sceller, significant changes in the microrelief of the surface of hard tissues were found. As shown in picture 2, abrasive instruments leave grooves to expose the dentinal tubule openings.

Picture 2 Electronogram of a sample of group B1

We observed small, dusty fragments covering a significant area of the treated tooth surface, microcracks, chips (Picture 3).

Picture 3. Electron diffraction pattern of a sample of group B1. An area of enamel after ultrasonic treatment with a large amount of microparticles of hard tooth tissues

Picture 4. Electronogram of a sample of group B1. The area of the tooth in the area of the dentine-enamel border

To solve the set tasks under the experimental conditions, microanalysis of the chemical composition of dentin and the concentration of its mass fraction was also carried out (Picture 5; Table 1).

Picture 5. The spectrum of chemical elements enamel of group B1

Table 1

Chemical composition enamel of group B1

№	Element	(KeV)	Mass%	Error%
1	F Ka	0.677	0.39	0.46
2	P Ka	2.013	17.24	0.22
3	Ca Ka	3.690	43.29	0.40
4	Other elements		39.08	0.36
	Total		100.00	

The results of the study in the B1 observation group showed that the quantitative content of calcium and phosphorus in the root dentin was 36.94% and 13.87% (Picture 6).

The data obtained included the total quantitative content of the studied elements included in the composition of both organic and inorganic compounds (Table 2).

Picture 6. Spectrum of chemical elements dentine of group B

Table 2

Chemical composition dentin of group B.

№	Element	(KeV)	Mass%	Error%
1	F Ka	0.677	0.26	0.38
2	P Ka	2.013	13.87	0.18
3	Ca Ka	3.690	36.94	0.36
4	Other elements		48.95	0.34
	Total		100.00	

In the experimental treatment of the surface of teeth of subgroup B2 (without water cooling), a significant number of cracks were observed on the samples, leading to a change in the shape and structure of dentin (Picture 7).

Picture 7. Electronogram of a sample of group B2. Change in the shape of dentinal tubules after ultrasonic treatment without cooling

According to the results of electron microscopic examination, it was found that treatment with a sceller is accompanied by the formation of a smeared dentin layer. Moreover, a violation of the treatment regimes (without water cooling) will lead to irreversible changes in the structure of the dentin of the teeth in the form of exposure of the dentinal tubules, which initiates a long-term increase in the sensitivity of the tooth (Picture 7).

Picture 8. Electronogram of a sample of group B.2. Dentin surface after ultrasonic treatment without cooling

Violation of the integrity of enamel prisms leads to non-viability, demineralization, a decrease in the chemical composition of enamel and dentin of calcium and phosphorus by 6-7%, and fluorine by 20%, which is the cause of hyperesthesia of hard tissues of living teeth.

To establish the nature of the prepared surface after finishing treatment with an air abrasion apparatus, an electron microscopic study of samples of group C was carried out.

Picture 9. Electronogram of the dentin section of the sample of group C

In the samples of group C, the preservation of the integrity of the enamel prisms at the processing border was revealed. After treatment with the Sandman Future apparatus and polishing, the number of hard microparticles of dentinal tubules was significantly reduced. At the same time, the dentin surface was similar to unchanged dentin, and the chemical composition of the enamel was distinguished by an increased fluorine content of 20% against the background of a decrease in calcium by 9% (Picture 10, Table 3).

Picture 10. Spectrum of chemical elements of dentin in samples of group C

Table 3

Chemical composition enamel of group C

№	Element	(KeV)	Mass%	Error%
1	F Ka	0.677	0.39	0.35
2	P Ka	2.013	16.04	0.17
3	Ca Ka	3.690	40.72	0.32
4	Other elements		42.75	0.28
	Total		100.00	

The results of studies of the chemical composition of enamel in the third C group of observations showed that the quantitative content of calcium - 43.29%, fluorine - 0.49%, and phosphorus - 17.14% (Picture 9, 10, Table 3). It was found that after finishing the tooth surface with the Sandman Futura apparatus, the amount of fluoride in dentin increased by 20%, and phosphorus and calcium by 5-7%. It can be assumed that the relative stability (and in some cases the excess of certain indicators) of the quantitative composition of chemical elements of hard tooth tissues correlated with a decrease in the severity of post-preparation hypersensitivity when using the Sandman Futura air abrasion apparatus. At the same time, the concentration of fluorine trace elements in the studied samples of hard dental tissues significantly differed depending on the method of removing dental deposits.

The results of electron microscopy show that ultrasonic treatment leads to changes in the structure of the dentin of the teeth. It was noted that a smear layer is formed on the surface of the treated dentin, which contains residues of hydroxyapatites, denatured collagen fibers, preparation grooves, damaged odontoblast processes, leading to increased sensitivity of hard tissues.

The number of the above changes after processing justifies the need for the use of modern types of sealers with obligatory surface polishing, removal of the smeared dentin layer, and deep fluoridation of dental hard tissues.

Conclusions: During the electron microscopic examination of samples of unpolished cement in group B, foci of micro destruction and numerous remains of dental calculus were revealed on its surface. In group C, a clean and smooth surface with a clear relief was revealed on

samples of polished tooth cement. The study of the chemical composition of dentin in the 2nd group showed a sharp release of mineral substances after treatment with a sceller. The data obtained revealed the regularity of the predicted demineralization of the hard tissues of the teeth after various methods of mechanical action. This fact dictates the need for remineralizing therapy after removing hard and soft dental plaque.

The studies carried out give reason to recommend additional polishing of the bare surface of the cement of the tooth root, after the removal of dental plaque and deep fluoridation.

1. Borysenko A.V. German Science Herald // Electronic microscopic research of tooth enamel surface after usage of different methods of home bleaching. №1. 2019. S. 3-9.
2. Makeeva I.M. Stomatology // Electron microscopic study of wedge-shaped defects of teeth on initial stage. №4. 2009. S 39-42
3. Karammaeva M.R. Endodontia today // Influence of ultrasound on hard tissue of the tooth (Electronic-microscopic research). №2. 2014. S. 25-27
4. Zhadko S.I. Institute of dentistry // Experimental justification for the preservation of vital pulp when using fixed dental prostheses. №3. 2015. S. 84-85
5. Pavlenko A.V. Dentist. Minsk // Selection of surface protectors for preparation of hard tissues of vital teeth during orthopedic treatment (Experimental study). №2(9). 2013. S. 12-16