CONVERGENCE OF A SPECIAL SET OF TRIANGLES

Dedicated to Sherman Kopald Stein (11 August 1926-...) for his pleasant sense of humor and mathematical wisdom that the third author enjoyed as a colleague during the late 1980s at the University of California, Davis

MOSHE STUPEL¹, AVI SIGLER² and JAY JAHANGIRI³

¹Shannan Academic College of Education

Gordon Academic College of Education

Haifa

Israel

e-mail: stupel@bezeqint.net

²Shannan Academic College of Education

Haifa

Israel

e-mail: avibsigler@gmail.com

³Mathematical Sciences

Kent State University

Kent, Ohio

USA

e-mail: jjahangi@kent.edu

Abstract

An explicit expression for the convergence point of a given set of triangles is introduced leading to an open interval locus of convergence that contains the centroid of the triangles.

2010 Mathematics Subject Classification: Primary: 97G40; Secondary: 97G50.

Keywords and phrases: geometric convergence, barycentric coordinates, Ceva's theorem, Pappus configuration.

Received March 8, 2018; Revised April 16, 2018

© 2018 Scientific Advances Publishers

1. Introduction

Using coordinate systems such as rectangular, polar, spherical and cylindrical require an origin but at times we need to locate points relative to existing points or local coordinates rather than the origin [1-4].

Mobius assigned coordinates (barycentric coordinates) to the point at which the center of gravity, or barycentre, was located, reflecting the ratio of weights attached to the ends of a given rod. He then extended this idea to a system of three weighted points, forming a triangle and pointing out that the center of gravity will remain the same if the weights are increased or decreased by a common factor [5, 6].

Central to barycentric coordinates are the concept of concurrency of lines in a triangle and Ceva's theorem that "Given any triangle, the segments from each vertex to the opposite sides of the triangle are concurrent precisely when the product of the ratios of the pairs of segments formed on each side of the triangle is equal to 1" [5, 7].

Ceva's theorem is used in the proofs of many well-known theorems including the theorem of Napoleon's point, the theorem of Fermat's point, and the theorem of the Nagel point [8]. Barycentric coordinates are a fundamental tool for a wide variety of applications employing triangular meshes, underlining the definition of higher-order basis functions, the Bézier triangle in computer aided-design, and many interpolation and shading techniques in computer graphics [9, 10].

In the present paper, we use the convergence of a special set of triangles leading to the discovery of an explicit expression for the convergence point and the understanding that this point can be constructed using a compass and a straightedge ruler, even if the convergence process is infinite. These results are then generalized to Pappus triangles and generalized Pappus triangles [11-13]. Finally, we obtain an interesting open interval locus of convergence that contains the centroid of the triangle.

2. Terminology and Notation

The interior point D of the triangle $\triangle ABC$ (Figure 1)

Figure 1. Representation of the point D .

shall be represented by two methods.

(a) Using the barycentric coordinates $(\lambda_1, \lambda_2, \lambda_3)$, we obtain

$$x_D = \lambda_1 x_A + \lambda_2 x_B + \lambda_3 x_C,$$

$$y_D = \lambda_1 y_A + \lambda_2 y_B + \lambda_3 y_C, \text{ where}$$

$$\lambda_1 + \lambda_2 + \lambda_3 = 1, \text{ and}$$

$$\lambda_1 \geq 0, \lambda_2 \geq 0, \lambda_3 \geq 0.$$

(b) Using the Ceva coordinates (α, β, γ) , we obtain

AA_1 , BB_1 , and CC_1 which are three cevians that pass through D so

that $\frac{BA_1}{A_1C} = \alpha$, $\frac{CB_1}{B_1A} = \beta$, and $\frac{AC_1}{C_1B} = \gamma$, where

$$\alpha\beta\gamma = 1, \quad \alpha \geq 0, \quad \beta \geq 0, \quad \gamma \geq 0.$$

Known are the relations between the barycentric coordinates and the Ceva coordinates

$$\left. \begin{aligned} \lambda_1 &= \frac{\alpha\beta}{\alpha\beta + \alpha + 1} \\ \lambda_2 &= \frac{1}{\alpha\beta + \alpha + 1} \\ \lambda_3 &= \frac{\alpha}{\alpha\beta + \alpha + 1} \end{aligned} \right\}, \quad (1)$$

$$\left. \begin{aligned} \alpha &= \frac{\lambda_3}{\lambda_2} \\ \beta &= \frac{\lambda_1}{\lambda_3} \\ \gamma &= \frac{\lambda_2}{\lambda_1} \end{aligned} \right\}. \quad (2)$$

3. The Convergence Process

The convergence process used here are presented in two phases.

Case I: The Pappus configuration

Select the points C_1, B_1, A_1 , respectively, on the sides CA, BC , and AB so that $\frac{AA_1}{A_1B} = \frac{BB_1}{B_1C} = \frac{CC_1}{C_1A} \equiv \alpha$. Similarly, select the points C_2, B_2, A_2 , respectively, on the sides C_1A_1, B_1C_1 and A_1B_1 so that $\frac{A_1A_2}{A_2B_1} = \frac{B_1B_2}{B_2C_1} = \frac{C_1C_2}{C_2A_1} \equiv \alpha$.

Repeating this process $(N + 1)$ -times (N is a whole number), we arrive at the points $C_{N+1}, B_{N+1}, A_{N+1}$, which are, respectively, on the sides C_NA_N, B_NC_N , and A_NB_N , so that they divide the sides of the triangle $A_NB_NC_N$ at a ratio that equals α .

Note that the triangle $A_0B_0C_0$ is the original triangle ABC and the sides A_0B_0, B_0C_0 , and C_0A_0 are the sides of the original triangle ABC .

The famous Pappus' theorem claims that the triangles $\triangle ABC$ and $\triangle A_1B_1C_1$ have a common centroid [14]. Therefore, it is clear that the triangles $\triangle A_1B_1C_1$ and $\triangle A_2B_2C_2$ also have a common centroid, and so on and so forth, up to the triangles $A_NB_NC_N$ and $A_{N+1}B_{N+1}C_{N+1}$. A continuous repetition of this process leads to the convergence of the triangles $A_{N+1}B_{N+1}C_{N+1}$ to a single centroid point as N approaches infinity and therefore we have the following:

Claim 1. $\cap \triangle A_{N+1}B_{N+1}C_{N+1}$ is the centroid point of $\triangle ABC$, as N tends to infinity.

Case II: Generalization

The points C_1, B_1, A_1 lie, respectively, on the sides CA, BC , and AB of the triangle $\triangle ABC$ so that

$$\frac{BB_1}{B_1C} = \alpha, \quad \frac{CC_1}{C_1A} = \beta, \quad \frac{AA_1}{A_1B} = \gamma \text{ (Figure 2).}$$

Figure 2. Convergence of Pappus triangles.

The points C_2, B_2, A_2 lie, respectively, on the sides C_1A_1, B_1C_1 , and A_1B_1 of the triangle $\Delta A_1B_1C_1$ so that

$$\frac{B_1B_2}{B_2C_1} \equiv \alpha, \quad \frac{C_1C_2}{C_2A_1} \equiv \beta, \quad \frac{A_1A_2}{A_2B_1} \equiv \gamma.$$

Repeating this process $(N+1)$ -times, we conclude that the points $C_{N+1}, B_{N+1}, A_{N+1}$, respectively, lie on the sides C_NA_N, B_NC_N, A_NB_N so that

$$\frac{B_NB_{N+1}}{B_{N+1}C_N} \equiv \alpha, \quad \frac{C_NC_{N+1}}{C_{N+1}A_N} \equiv \beta, \quad \frac{A_NA_{N+1}}{A_{N+1}B_N} \equiv \gamma.$$

Next we state and prove.

Claim 2. $\cap \Delta A_{N+1}B_{N+1}C_{N+1} = S$; a point of ΔABC , with the barycentric coordinates

$$(\lambda_1^S, \lambda_2^S, \lambda_3^S) = \left(\frac{\alpha\beta(\gamma+1)}{\alpha\beta(\gamma+1) + \beta\gamma(\alpha+1) + \alpha\gamma(\beta+1)}, \right. \\ \left. \frac{\beta\gamma(\alpha+1)}{\alpha\beta(\gamma+1) + \beta\gamma(\alpha+1) + \alpha\gamma(\beta+1)}, \frac{\alpha\gamma(\beta+1)}{\alpha\beta(\gamma+1) + \beta\gamma(\alpha+1) + \alpha\gamma(\beta+1)} \right), \quad (3)$$

or with the Ceva coordinates

$$(\lambda_1^S, \lambda_2^S, \lambda_3^S) = \left(\frac{\alpha(\beta+1)}{\beta(\alpha+1)}, \frac{\beta(\gamma+1)}{\gamma(\beta+1)}, \frac{\gamma(\alpha+1)}{\alpha(\gamma+1)} \right). \quad (4)$$

Proof. We transform the triangle ABC by an affine transformation to a triangle in which A, B , and C have the coordinates $A(0, 1)$, $B(0, 0)$, and $C(1, 0)$.

Next, we transform the points A, C, B by an affine transformation to respective points C_1, B_1, A_1 so that $\frac{BB_1}{B_1C} = \alpha$, $\frac{CC_1}{C_1A} = \beta$, and $\frac{AA_1}{A_1B} = \gamma$.

Repeating this process $(N+1)$ -times, we note that these affine transformations preserve the simple ratio where the triangle $\Delta A_NB_NC_N$

shall be transformed into the triangle $\Delta A_{N+1}B_{N+1}C_{N+1}$ so that the division ratios of the corresponding sides are α , β , and γ , as required.

The affine transformation f that transforms A, C, B to C_1, B_1, A_1 , is denoted by $f(\Delta ABC) \equiv \Delta A_1B_1C_1$ (Figure 3). This is a continuous function from the convex domain ΔABC to the convex domain $\Delta A_1B_1C_1$ that is contained in ΔABC . Therefore f has a fixed point S . This point S shall be the only common point of the triangles $\cap \Delta A_{N+1}B_{N+1}C_{N+1}$.

Figure 3. Affine transformation $f(\Delta ABC) = \Delta A_1B_1C_1$.

To derive the coordinates of the point S , observe that the coordinates of A_1 , B_1 , and C_1 are

$$A_1 = \left(0, \frac{1}{\gamma + 1}\right), B_1 = \left(\frac{\alpha}{\alpha + 1}, 0\right), C_1 = \left(\frac{1}{\beta + 1}, \frac{\beta}{\beta + 1}\right).$$

The fixed point for f has the same barycentric coordinates in ΔABC and $\Delta A_1B_1C_1$. Therefore it is a solution of the following system of equation:

$$x_S = \underbrace{\lambda_3 \cdot 1 + \lambda_1 \cdot 0 + \lambda_2 \cdot 0}_{\Delta ABC} = \underbrace{\lambda_1 \cdot 0 + \lambda_2 \cdot \frac{\alpha}{\alpha + 1} + \lambda_3 \cdot \frac{1}{\beta + 1}}_{\Delta A_1B_1C_1}.$$

In the other words,

$$\lambda_3 = \lambda_2 \cdot \frac{\alpha}{\alpha + 1} + \frac{\lambda_3}{\beta + 1}, \quad (5)$$

$$\lambda_1 = \frac{\lambda_1}{\gamma + 1} + \frac{\lambda_3 \beta}{\beta + 1}. \quad (6)$$

From (5), it follows that

$$\alpha_S = \frac{\lambda_3}{\lambda_2} = \frac{\alpha(\beta + 1)}{\beta(\alpha + 1)}. \quad (5^*)$$

From (6), it follows that

$$\beta_S = \frac{\lambda_1}{\lambda_3} = \frac{\beta(\gamma + 1)}{\gamma(\beta + 1)}. \quad (6^*)$$

And clearly, we obtain

$$\gamma_S = \frac{\lambda_2}{\lambda_1} = \frac{\gamma(\alpha + 1)}{\alpha(\gamma + 1)}.$$

A simple calculation, using Equation (1) gives $\lambda_1^S = \frac{\alpha\beta(\gamma + 1)}{\alpha\beta(\gamma + 1) + \beta\gamma(\alpha + 1) + \alpha\gamma(\beta + 1)}$. The process for λ_2^S and λ_3^S are similar and will be omitted.

Note. For given α , β , and γ , the point S can be constructed using a compass and a straightedge ruler (though the convergence process is infinite). An accurate description of segment construction, by using compass and straightedge, based on expressions whose representation is by segments, can be found in references [1, 2].

Next we deal with three particular cases (Figure 4).

Figure 4. The case of the middle of the segments.

First particular case

If $\alpha = \beta = \gamma = 1$, then the convergence is at the centroid of $\triangle ABC$.

Second particular case (Pappus)

If $\alpha = \beta = \gamma = k$, where $k(0 < k < \infty)$ is fixed, then S is the centroid of $\triangle ABC$ because $\alpha_S = \beta_S = \gamma_S = 1$.

Third particular case (Generalized Pappus)

If $\frac{BB_1}{B_1C} = k\alpha$, $\frac{CC_1}{C_1A} = k\beta$ and $\frac{AA_1}{A_1B} = k\gamma$, where $k(0 < k < \infty)$ is fixed, then

$$\left. \begin{aligned} \alpha_S &= \frac{\alpha(k\beta + 1)}{\beta(k\alpha + 1)} \\ \beta_S &= \frac{\beta(k\gamma + 1)}{\gamma(k\beta + 1)} \\ \gamma_S &= \frac{\gamma(k\alpha + 1)}{\alpha(k\gamma + 1)} \end{aligned} \right\}. \quad (7)$$

In the other words, aside from the Pappus' case, the point S depends on k and is therefore denoted by $S_{(k)}$.

Finally, we state and prove our

Claim 3. The locus of $S_{(k)}$ where k ($0 < k < \infty$) is fixed, is the open interval straight-line segment bounded by two points M_1 and M_2 where M_1 is the centroid of the triangle and M_2 has the barycentric coordinates

$$\lambda_1^{M_2} = \frac{\alpha\beta}{\alpha\beta + \beta\gamma + \gamma\alpha}, \lambda_2^{M_2} = \frac{\beta\gamma}{\alpha\beta + \beta\gamma + \gamma\alpha}, \lambda_3^{M_2} = \frac{\alpha\gamma}{\alpha\beta + \beta\gamma + \gamma\alpha}.$$

Proof. When k tends to zero in (7), we obtain the Ceva coordinates

$$M_2 = \left(\frac{\alpha}{\beta}, \frac{\beta}{\gamma}, \frac{\gamma}{\alpha} \right), \text{ which means the barycentric coordinates}$$

$$\left(\lambda_1^{M_2}, \lambda_2^{M_2}, \lambda_3^{M_2} \right) = \left(\frac{\alpha\beta}{\alpha\beta + \beta\gamma + \gamma\alpha}, \frac{\beta\gamma}{\alpha\beta + \beta\gamma + \gamma\alpha}, \frac{\alpha\gamma}{\alpha\beta + \beta\gamma + \gamma\alpha} \right).$$

When k tends to infinity, we obtain $(1, 1, 1) = (\alpha_{M_1}, \beta_{M_1}, \gamma_{M_1})$, or

barycentric coordinates $\left(\lambda_1^{M_2}, \lambda_2^{M_2}, \lambda_3^{M_2} \right) = \left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3} \right)$. We now prove

that $S_{(k)}$ is located on the segment connecting M_1 and M_2 . It is enough to prove that

$$\begin{vmatrix} \lambda_1^{M_1} & \lambda_2^{M_1} & \lambda_3^{M_1} \\ \lambda_1^{M_2} & \lambda_2^{M_2} & \lambda_3^{M_2} \\ \lambda_1^{S(k)} & \lambda_2^{S(k)} & \lambda_3^{S(k)} \end{vmatrix} = 0,$$

or

$$\begin{vmatrix} 1 & 1 & 1 \\ \alpha\beta & \beta\gamma & \alpha\gamma \\ \alpha\beta(k\gamma + 1) & \beta\gamma(k\alpha + 1) & \alpha\gamma(k\beta + 1) \end{vmatrix} = 0,$$

or

$$\begin{vmatrix} 1 & 1 & 1 \\ \alpha\beta & \beta\gamma & \alpha\gamma \\ k\alpha\beta\gamma & k\alpha\beta\gamma & k\alpha\beta\gamma \end{vmatrix} = 0.$$

We will show that $S_{(k)}$ is located between M_1 and M_2 . We connect A to M_1 and M_2 , and obtain the respective points of intersection D_1 and D_2 , (Figure 5).

Figure 5. The location of the point $S_{(k)}$.

Without loss of generality, we assume that M_2 and D_2 are more to the left of M_1 and D_1 , in the other words $\frac{\alpha}{\beta} < 1$.

In this case $\frac{\alpha}{\beta} < \alpha_{S_{(k)}} = \frac{\alpha(\beta k + 1)}{\beta(\alpha k + 1)} < 1$ and therefore $S_{(k)}$ is located between M_1 and M_2 .

References

- [1] G. E. Martin, *Geometric Constructions*, Springer, New York (1998), 131-142.
- [2] V. Oxman, M. Stupel and A. Sigler, Geometric constructions for geometric optics using a straightedge only, *Journal for Geometry and Graphics* 18(1) (2014), 73-79.
- [3] M. Stupel, V. Oxman and A. Sigler, More on geometrical constructions of a tangent to a circle with a straightedge only, *The Electronic Journal of Mathematics and Technology* 8(1) (2014), 17-30.
- [4] A. Sutton, *Ruler and Compass: Practical Geometric Constructions*, Walker & Company, 2009.
- [5] P. Lidberg, Barycentric and Wachspress Coordinates in two Dimensions: Theory and Implementation for Shape Transformations, Uppsala Universitet, U.U.D.M. Project Report, February 2011.
- [6] S. K. Stein, Archimedes: What did he do Besides Cry Eureka, *The Mathematical Association of America*, Page 15, Washington, 1999.
- [7] D. A. Brannan, M. F. Esplen and J. J. Gray, *Geometry*, Cambridge University Press, page 75, Cambridge, 2004.
- [8] D. N. V. Krishna, The fundamental property of Nagel point: A new proof, *Journal of Mathematical Sciences & Mathematics Education* 12(2) (2017), 21-28.
<http://www.msme.us/2017-2-4.pdf>
- [9] R. M. Rustamov, Y. Lipman and T. Funkhouser, Interior distance using barycentric coordinates, *Eurographics Symposium on Geometry Processing* 28(5) (2009), 1279-1288.
 DOI: <https://doi.org/10.1111/j.1467-8659.2009.01505.x>
- [10] J. Vince, *Mathematics for Computer Graphics*, (4th Edition), Springer, London, Heidelberg, New York, Dordrecht, 2013. ISBN: 978-1-4471-6289-6.
- [11] H. Eves, *An Introduction to the History of Mathematics*, Philadelphia: Saunders College Publishing, 1992.
- [12] A. Kryftis, *A Constructive Approach to Affine and Projective Planes*, Ph.D. Thesis, University of Cambridge, 2015.
- [13] B. D. S. McConnell, A Six-Point Ceva-Menelaus Theorem, 3 March 2014.
<https://arxiv.org/pdf/1403.0478.pdf>
- [14] A. W. Goodman and G. Goodman, Generalizations of the theorems of Pappus, *The American Mathematical Monthly* 76(4) (1969), 355-366.