HOMOCLINIC ORBIT FOR A FOURTH-ORDER SYSTEM

XIAOLEI REN

School of Mathematical Science

Nanjing Normal University

Nanjing, Jiangsu 210097

P. R. China

e-mail: rwxiaolei@163.com

Abstract

In this paper, homoclinic orbit of a fourth-order system will be concerned, several assumptions are made to prove a fourth-order system has a homoclinic orbit q emanating from 0 where $q \in C^4(\mathbf{R}, \mathbf{R}^n)$. Under these assumptions, we can find the homoclinic orbit of the system by defining a function $I_k(q)$ via the Mountain Pass Theorem.

1. Introduction and Main Results

In this paper, we will deal with the existence of homoclinic orbit of the fourth-order system:

$$-q^{(4)} + Cq'' - L(t)q + V_q'(t, q) = 0, \quad (1.1)$$

where $q \in \mathbf{R}^n$, C is positive definite symmetric matrix, L and V satisfy:

2010 Mathematics Subject Classification: 37C60, 37B55, 34L30, 49J35, 49K15.

Keywords and phrases: homoclinic orbit, the Mountain Pass Theorem, critical point, periodic solutions.

Received September 17, 2019

(L) $L(t)$ is a continuous T -periodic matrix valued function and is a positive definite symmetric for all $t \in [0, T]$.

(V₁) $V \in C^1(\mathbf{R} \times \mathbf{R}^n, \mathbf{R})$ is T -periodic in t .

(V₂) There is a $\mu > 2$ such that $0 < \mu V(t, q) \leq (q, V'_q(t, q))$ for all $q \in \mathbf{R}^n \setminus \{0\}$.

(V₃) $V(t, q) = o(|q|^2)$ as $q \rightarrow 0$ uniformly for $t \in [0, T]$.

We say that a solution $q(t)$ of (1.1) is a homoclinic orbit if $q(t) \in C^4(\mathbf{R}, \mathbf{R}^n)$ such that $q(t) \rightarrow 0$ and $q'(t) \rightarrow 0$ as $|t| \rightarrow \infty$.

Theorem 1.1. *Assume (L), (V₁)–(V₃) hold. Then (1.1) has a nonzero homoclinic orbit.*

This paper is motivated by Rabinowitz [1], who has showed that under (L), (V₁)–(V₃), the system

$$q'' - L(t)q + V'_q(t, q) = 0 \quad (1.2)$$

has a nontrivial solution $q \in C^2(\mathbf{R}, \mathbf{R}^n)$ satisfying $|q(t)| + |q'(t)| \rightarrow 0$ as $|t| \rightarrow \infty$.

In Section 2, we will prove Theorem 1.1 using methods in [1].

2. Proof of Theorem 1.1

Similar to [1], for each $k \in N$, let $E_k = W_{2kT}^{2,2}(\mathbf{R}, \mathbf{R}^n)$ with the norm:

$$\|q\|_k^2 = \int_{-kT}^{kT} [\|q''(t)\|^2 + (Cq'(t), q'(t)) + (L(t)q(t), q(t))] dt,$$

and let

$$\begin{aligned} I_k(q) &= \int_{-kT}^{kT} \left[\frac{1}{2} |q''(t)|^2 + \frac{1}{2} (Cq'(t), q'(t)) + \frac{1}{2} (L(t)q(t), q(t)) - V(t, q) \right] dt \\ &= \frac{1}{2} \|q\|_k^2 - \int_{-kT}^{kT} V(t, q) dt. \end{aligned} \quad (2.1)$$

It is easy to check $I_k \in C^1(E_k, \mathbf{R})$ and

$$I_k(q')u = \int_{-kT}^{kT} [(q'(t), u''(t)) + (Cq'(t), u'(t)) + (L(t)q(t), u(t)) - (V'_q(t, q(t), u(t)))] dt.$$

Lemma 2.1. *If $I_k(q') = 0$, then $q \in C^4(\mathbf{R}, \mathbf{R}^n)$ and $q = q(t)$ is a solution of (1.1).*

Proof. If q is a critical point of I_k , then

$$\int_{-kT}^{kT} [(q''(t), h''(t)) + (Cq'(t), h'(t)) + (L(t)q(t) - V'_q(t, q(t)), h(t))] dt = 0,$$

for all $h \in E_k$. Letting $\delta(t) = L(t)q(t) - V'_q(t, q(t))$ means that $\int_{-kT}^{kT} \delta(t) dt = 0$

since $\mathbf{R}^n \subset E_k$. So there is $x \in E_k$ satisfying $x''(t) = \delta(t)$ and

$$\int_{-kT}^{kT} ((q''(t) - Cq(t) + x(t)), h(t)) dt = 0, \quad \forall h \in E_k.$$

Thus, there is a constant $\xi \in \mathbf{R}^n$ such that $q''(t) - Cq(t) + x(t) = \xi$ a.e. $x \in [-kT, kT]$. So

$$q^{(4)}(t) - Cq''(t) + L(t)q(t) - V'_q(t, q(t)) = 0,$$

since $\delta(t) = L(t)q(t) - V'_q(t, q(t))$ is continuous. The proof is complete. $\square$

In order to prove Theorem 1.1 as in [1], we will use the Mountain Pass Lemma.

Proposition 2.1. *Let E be a real Banach space and $I \in C^1(E, \mathbf{R})$ satisfy (PS). (PS) says whenever $I(u_m)$ is bounded and $I'(u_m) \rightarrow 0$ as $m \rightarrow \infty$, the sequence u_m possesses a convergent subsequence.*

If further $I(0) = 0$,

(I_1) There exist constants $\rho, \alpha > 0$ such that $I|_{\partial B_\rho(0)} \geq \alpha$;

and

(I_2) There exists $e \in E \setminus \overline{B_\rho(0)}$ such that $I(e) \leq 0$, then I possesses a critical value $c \geq \alpha$ given by

$$c = \inf_{g \in \Gamma} \max_{s \in [0,1]} I_k(g(s)), \quad (2.2)$$

where

$$\Gamma = \{g \in C([0, 1], E) | g(0) = 0 \text{ and } g(1) = e\}, \quad (2.3)$$

$B_\rho(0)$ denotes an open ball of radius ρ about 0. $\square$

We will obtain a critical point of I_k via the Mountain Pass Theorem. Moreover, the critical points of I_k in E_k are $2kT$ periodic solutions of (1.1). First we need to prove $I_k(q)$ satisfies (PS). Recall that we say I satisfied (PS) if that $I(u_m)$ is bounded and $I'(u_m) \rightarrow 0$ imply that $u_m \rightarrow u$ for some $u \in E$ along a subsequence.

Lemma 2.2. $I_k(q)$ satisfies (PS).

Proof. To verify (PS), define first

$$\|q\|_\beta = \left(\|q''\|_{L^2}^2 + C\|q'\|_{L^2}^2 + (L(t)q, q)_R + \beta\|q\|_{L^2}^2 \right)^{\frac{1}{2}},$$

where $\beta > 0$. Thus $\|\cdot\|_\beta$ is equivalent to $\|\cdot\|_{E_k}$. Suppose q_m is a sequence in E_k , $|I(q_m)| \leq K$ and $I'(q_m) \rightarrow 0$. For m large enough,

$$\begin{aligned} & |I'(q_m)q_m| \\ &= \left| \|q_m\|_\beta^2 - \beta \|q_m\|_{L^2}^2 - \int_{-kT}^{kT} (q_m, V'_q(t, q_m)) d\tau \right| \\ &\leq \|q_m\|_\beta. \end{aligned} \quad (2.4)$$

Therefore,

$$\left| \int_{-kT}^{kT} (q_m, V'_q(t, q_m))_R d\tau \right| \leq \|q_m\|_\beta + \|q_m\|_\beta^2. \quad (2.5)$$

For $y \in R$ large enough, we have

$$|y|^2 \leq \mu(y, V'_q(t, q_m)). \quad (2.6)$$

Hence, by (V_2) and (2.5)–(2.6) for a constant γ independent of m , we have

$$\begin{aligned} \int_{-kT}^{kT} V(t, q_m) d\tau &\leq \frac{1}{\mu} \int_{-kT}^{kT} (q_m, V'_q(t, q_m)) d\tau \\ &\leq \frac{1}{\mu} (\|q_m\|_\beta + \|q_m\|_\beta^2), \\ K &\geq \frac{1}{2} \|q_m\|_\beta^2 - \frac{1}{2} \beta \|q_m\|_{L^2}^2 - \int_{-kT}^{kT} V(t, q_m) d\tau \\ &\geq \frac{1}{2} \|q_m\|_\beta^2 - \left(\frac{1}{2} \beta + \frac{1}{\mu} \right) (\|q_m\|_\beta + \|q_m\|_\beta^2) - \gamma. \end{aligned} \quad (2.7)$$

Choosing $\beta \in \left(0, 1 - \frac{2}{\mu}\right)$, (2.7) shows $\|q_m\|_{E_k}$ is uniform bound, and then

there is a subsequence of q_m converges weakly in E_k and strongly in L^∞

to $q \in E_k$. The mean value of q_m converges to the mean value of $\bar{q}$. By (2.4), we have $I(q') = q - \phi(q)$, where $\phi(q)$ is compact, hence $I(q') \rightarrow 0$, then $q_m \rightarrow \phi(\bar{q})$, (PS) is verified. The proof is complete. $\square$

Lemma 2.3. *If (L) and (V₁)–(V₃) hold, then I_k defined in (2.2) possesses a critical point q_k with the corresponding critical value c_k .*

Proof. By (2.2), we have $I_k(0) = 0$. By (V₂), let $f(s) = V(t, s\xi)$; for any fixed point $\xi \in \mathbf{R}^n$, we have

$$s f'(s) \geq \mu f(s) = \mu V(t, s\xi).$$

For $s \geq 1$, $s f'(s) \geq \mu f(s)$. Integrate it in $[1, s]$, then $f(s) \geq s^\mu f(1)$.

That is $V(t, s\xi) \geq s^\mu V(t, \xi)$, for this, we can get

$$\begin{aligned} V(t, \xi) &\leq V\left(t, \frac{\xi}{|\xi|}\right) |\xi|^\mu, \text{ for } 0 < |\xi| \leq 1, \\ &\geq V\left(t, \frac{\xi}{|\xi|}\right) |\xi|^\mu, \text{ for } |\xi| \geq 1. \end{aligned} \quad (2.8)$$

Then there are constants $\alpha_k, \varepsilon_k > 0$ such that

$$\alpha_k \|q\|_{L^\mu[-kT, kT]} \leq \|q\|_{L^x[-kT, kT]} \leq \varepsilon_k \|q\|_k. \quad (2.9)$$

Moreover by (2.1) and (2.8), we have

$$I_k(\beta\varphi) = \frac{1}{2} \|\beta\varphi\|_k^2 - \int_{-kT}^{kT} V(t, \beta\varphi) dt \leq \frac{1}{2} \beta^2 \|\varphi\|_k^2 - \int_{-kT}^{kT} V\left(t, \frac{\beta\varphi}{|\beta\varphi|}\right) |\beta\varphi|^\mu dt.$$

Then there are constants $\alpha_1, \alpha_2 > 0$ such that

$$I_k(\beta\varphi) \leq \frac{1}{2} \beta^2 \|\varphi\|_k^2 - \alpha_1 |\beta|^\mu \int_{-kT}^{kT} |\varphi| dt + \alpha_2, \quad (2.10)$$

for all $\beta \in \mathbf{R}$ and $\varphi \in E_k \setminus \{0\}$. By (2.1) and (V₃), $e = e_k$ and any $e_k \in E_k \setminus \{0\}$ shows $I_k(e_k) \leq 0$. Consequently by Proposition 2.1, I_k

possesses a positive critical value c_k given by (2.2) and (2.3) with $E = E_k$ and $\Gamma = \Gamma_k$. Let q_k denote the corresponding critical point of I on E_k . The proof is complete. $\square$

Remark. Note that $q_k \neq 0$ since $c_k > 0$.

Lemma 2.4. *The c_k and q_k obtained in Lemma 2.3 are independent of k .*

Proof. As in [1], we have $\varphi \in E_1 \setminus \{0\}$ such that

- (i) $\varphi(\pm T) = \varphi'(\pm T) = 0,$
- (ii) $I_1(\varphi) \leq 0.$

By (2.10), we can find ϕ satisfies (i), any sufficiently large multiple φ of ϕ satisfies (i) and (ii), by Lemma 2.3, let

$$\begin{aligned} e_k(t) &= \varphi(t) \quad |t| \leq T \\ &= 0 \quad T \leq |t| \leq kT. \end{aligned}$$

Then by (2.8), we have $e_k \in E_k \setminus \{0\}$ and $I_k(e_k) = I_1(e_1) \leq 0$. Note also that $g_k(s) = se_k \in \Gamma_k$ for all $k \in N$ and $I_k(g_k(s)) = I_1(g_1(s))$. Therefore by (2.2),

$$c_k \leq \max_{s \in [0,1]} I_1(g_1(s)) \equiv M. \quad (2.11)$$

So c_k is independent of k and (2.11) leads to a priori bounds for q_k . Since $I'(q_k) = 0$, by (V_2) ,

$$\begin{aligned} c_k &= I_k(q_k) - \frac{1}{2} I'_k(q_k) q_k \\ &= \int_{-kT}^{kT} \left[\frac{1}{2} (q_k, V'_q(t, q_k) - V(t, q_k)) \right] dt \\ &\geq \left(\frac{\mu}{2} - 1 \right) \int_{-kT}^{kT} V(t, q_k) dt. \end{aligned} \quad (2.12)$$

Hence by (2.1) and (2.12), we have

$$\begin{aligned}
\|q_k\|_k^2 &= \int_{-kT}^{kT} \left[|q''(t)|^2 + (Cq'(t), q'(t)) + ((L(t)q(t), q(t))) \right] dt \\
&= 2I_k(q) + 2 \int_{-kT}^{kT} V(t, q) dt \\
&= 2c_k + \frac{4c_k}{\mu - 2} \\
&\leq 2M \left(1 + \frac{2}{\mu - 2} \right) \equiv M_1.
\end{aligned} \tag{2.13}$$

By (2.9), we have a k -dependent bound on $\|q_k\|_{L^\infty[-kT, kT]}$. The proof is complete. $\square$

Proof of Theorem 1.1. We can complete this theorem under the previous lemmas. By these lemmas, we obtain a critical point of I_k and we know the critical points of I_k in E_k are periodic solutions of (1.1).

As in [1], for $q_k \in E_k$, $t, \tau \in [-kT, kT]$

$$|q(t)| \leq q(\tau) + \left| \int_{\tau}^t q'(s) ds \right|.$$

We have

$$\begin{aligned}
|q(t)| &\leq \int_{t-\frac{1}{2}}^{t+\frac{1}{2}} |q(t)| d\tau + \int_{t-\frac{1}{2}}^{t+\frac{1}{2}} \left| \int_{\tau}^t q'(s) ds \right| d\tau \\
&\leq 2 \left(\int_{t-\frac{1}{2}}^{t+\frac{1}{2}} (|q'(\tau)|^2 + |q(\tau)|^2) d\tau \right)^{\frac{1}{2}}, \\
&\leq c \|q\|_k,
\end{aligned} \tag{2.14}$$

where c depends on L . Hence

$$\|q\|_{L^\infty[-kT, kT]} \leq c\|q\|_k. \quad (2.15)$$

And so, (2.13) and (2.15) imply

$$\|q_k\|_{L^\infty[-kT, kT]} \leq cM_1^{\frac{1}{2}} \equiv M_2. \quad (2.16)$$

Similarly, (2.16) will also hold if we replace q_k with q'_k . Because $q = q_k(t)$ is a solution of (1.1), q_k is bounded in $C^4[a, b]$ for any $a < b$, and hence there is a subsequence of q_k converging in $C^4_{\text{loc}}(\mathbf{R}, \mathbf{R}^n)$ to a solution q of (1.1) satisfying

$$\int_{-\infty}^{\infty} \left[|q''(t)|^2 + (Cq'(t), q'(t)) + (L(t)q(t), q(t)) \right] dt < \infty.$$

It follows that

$$|q(t)| + |q'(t)| \rightarrow 0 \text{ as } |t| \rightarrow \infty.$$

Now it is sufficient show $q(t) \neq 0$. By the inner product of (1.1) with q_k and then integration by parts shows:

$$\begin{aligned} & \int_{-kT}^{kT} \left(-q_k^{(4)} + Cq_k'' - L(t)q_k + V'_q(t, q_k), q_k \right) d\tau = 0 \\ & \int_{-kT}^{kT} -|q_k''|^2 - (Cq'_k, q'_k) - (L(t)q_k, q_k) - V(t, q) + (V_q(t, q_k), q_k) d\tau = 0 \\ & \|q_k\|_k^2 = \int_{-kT}^{kT} (q_k, V'_q(t, q_k)) dt. \end{aligned} \quad (2.17)$$

As in [1], set

$$Y(0) = 0 \text{ and for } s > 0$$

$$Y(s) = \max_{\substack{t \in [0, T] \\ |\xi| \leq s}} \frac{(\xi, V'_q(t, \xi))}{|\xi|^2}.$$

By $(V_1)-(V_2)$, $Y(s) > 0$ if $s > 0$, $Y \in C(\mathbf{R}^+, \mathbf{R}^+)$, Y is monotone non-decreasing and $Y(s) \rightarrow \infty$ as $s \rightarrow \infty$. Also by the definition of Y

$$\frac{(q_k, V'_q(t, q_k(t)))}{|q_k(t)|^2} \leq Y(\|q_k\|_{L^\infty[-kT, kT]}) \quad \text{for all } t \in [-kT, kT]. \quad (2.18)$$

Hence by (2.17) and (2.18),

$$\begin{aligned} \|q_k\|_k^2 &\leq Y(\|q_k\|_{L^\infty[-kT, kT]}) \int_{-kT}^{kT} |q_k|^2 dt \\ &\leq l_0^{-1} Y(\|q_k\|_{L^\infty[-kT, kT]}) \|q_k\|_k^2, \end{aligned}$$

where $l_0 > 0$ satisfies $L(t) \geq l_0 I_{2n}$, $\forall t$. And

$$Y(\|q_k\|_{L^\infty[-kT, kT]}) \geq l_0.$$

By the properties of Y , there is a $M_3 > 0$ such that

$$\|q_k\|_{L^\infty[-kT, kT]} \geq M_3. \quad (2.19)$$

Because L and V are T -periodic in t and $q_k(t)$ is a $2kT$ -periodic solution, $q(t + jT)$, $j \in [-k, k] \cap \mathbf{Z}$ is also a solution of (1.1), and we may assume that the maximum of $|q_k(t)|$ occurs in $[0, T]$. Therefore if $q_k(t) \rightarrow 0$ in C_{loc}^1 along our subsequence,

$$\|q_k\|_{L^\infty[-kT, kT]} = \max_{t \in [0, T]} |q_k(t)| \rightarrow 0.$$

Contrary to (2.19). The proof is complete. $\square$

References

- [1] P. H. Rabinowitz, Periodic solutions of hamiltonian systems, Communications on Pure and Applied Mathematics 31(2) (1978), 157-184.

DOI: <https://doi.org/10.1002/cpa.3160310203>

- [2] V. C. Zelati and P. H. Rabinowitz, Homoclinic orbits for second order Hamiltonian systems possessing superquadratic potentials, *Journal of the American Mathematical Society* 4(4) (1991), 693-727.
DOI: <https://doi.org/10.1090/S0894-0347-1991-1119200-3>
- [3] G.-W. Chen, Superquadratic or asymptotically quadratic Hamiltonian systems: ground state homoclinic orbits, *Annali di Matematica Pura ed Applicata* 194(3) (2015), 903-918.
DOI: <https://doi.org/10.1007/s10231-014-0403-9>
- [4] P. H. Rabinowitz, Homoclinic orbits for a class of Hamiltonian systems, *Proceedings of the Royal Society of Edinburgh Section A: Mathematics* 114(1-2) (1990), 33-38.
DOI: <https://doi.org/10.1017/S0308210500024240>
- [5] Y. Xu and M. Huang, Homoclinic orbits and Hopf bifurcations in delay differential systems with T-B singularity, *Journal of Differential Equations* 244(3) (2008), 582-598.
DOI: <https://doi.org/10.1016/j.jde.2007.09.003>
- [6] A. R. Champneys and G. J. Lord, Computation of homoclinic solutions to periodic orbits in a reduced water-wave problem, *Physica D: Nonlinear Phenomena* 102(1-2) (1997), 101-124.
DOI: [https://doi.org/10.1016/S0167-2789\(96\)00206-0](https://doi.org/10.1016/S0167-2789(96)00206-0)
- [7] A. Szulkin and W. M. Zou, Homoclinic orbits for asymptotically linear Hamiltonian systems, *Journal of Functional Analysis* 187(1) (2001) 25-41.
DOI: <https://doi.org/10.1006/jfan.2001.3798>
- [8] G. A. Leonov, N. V. Kuznetsov and T. N. Mokaev, Hidden attractor and homoclinic orbit in Lorenz-like system describing convective fluid motion in rotating cavity, *Communications in Nonlinear Science and Numerical Simulation* 28(1-3) (2015), 166-174.
DOI: <https://doi.org/10.1016/j.cnsns.2015.04.007>
- [9] G. A. Leonov, N. V. Kuznetsov and T. N. Mokaev, Homoclinic orbits, and self-excited and hidden attractors in a Lorenz-like system describing convective fluid motion, *European Physical Journal-Special Topics* 224(8) (2015), 1421-1458.
DOI: <https://doi.org/10.1140/epjst/e2015-02470-3>
- [10] M. Izydorek and J. Janczewska, Homoclinic solutions for a class of the second order Hamiltonian systems, *Journal of Differential Equations* 219(2) (2005), 375-389.
DOI: <https://doi.org/10.1016/j.jde.2005.06.029>

- [11] A. R. Champneys, Y. A. Kuznetsov and B. Sandstede, A numerical toolbox for homoclinic bifurcation analysis, *International Journal of Bifurcation and Chaos* 6(5) (1996), 867-887.

DOI: <https://doi.org/10.1142/S0218127496000485>

- [12] A. R. Champneys, Homoclinic orbits in reversible systems and their applications in mechanics, fluids and optics, *Physica D: Nonlinear Phenomena* 112(1-2) (1998), 158-186.

DOI: [https://doi.org/10.1016/S0167-2789\(97\)00209-1](https://doi.org/10.1016/S0167-2789(97)00209-1)