**EXISTENCE AND CONTINUOUS DEPENDENCE
OF THE LOCAL SOLUTION OF NON-HOMOGENEOUS
KdV-K-S EQUATION IN PERIODIC
SOBOLEV SPACES**

**YOLANDA SILVIA SANTIAGO AYALA
and SANTIAGO CESAR ROJAS ROMERO**

Facultad de Ciencias Matemáticas
Universidad Nacional Mayor de San Marcos
Av. Venezuela Cdra. 34, Lima 01
Lima
Perú
e-mail: ysantiago@unmsm.edu.pe
srojasr@unmsm.edu.pe

2020 Mathematics Subject Classification: 35G10, 35Q53, 47D06, 35B40.

Keywords and phrases: uniqueness solution, KdV-Kuramoto-Sivashinsky equation, non-homogeneous equation, periodic Sobolev spaces, Fourier theory, calculus in Banach spaces.

Received November 25, 2020

© 2021 Scientific Advances Publishers

This work is licensed under the Creative Commons Attribution International License (CC BY 3.0).

http://creativecommons.org/licenses/by/3.0/deed.en_US

Open Access

Abstract

In this article, we prove that initial value problem associated to the non-homogeneous KdV-Kuramoto-Sivashinsky (KdV-K-S) equation in periodic Sobolev spaces has a local solution in $[0, T]$ with $T > 0$, and the solution has continuous dependence with respect to the initial data and the non-homogeneous part of the problem. We do this in an intuitive way using Fourier theory and introducing a C_0 -semigroup inspired by the work of Iorio [2] and Ayala and Romero [8].

Also, we prove the uniqueness solution of the homogeneous and non-homogeneous KdV-K-S equation, using its dissipative property, inspired by the work of Iorio [2] and Ayala and Romero [9].

1. Introduction

First, we want to comment that from Theorem 3.1 in [7], we have that the KdV-K-S homogeneous problem is globally well posed and, in addition to the inequality (3.2) in [7], we have the continuous dependence of the solution of homogeneous problem. In this work, in Theorem 3.2, we will prove the existence and uniqueness of the local solution for the non-homogeneous problem, and from inequality (3.8), we will get the continuous dependence of the solution with respect to the initial data and respect to the non-homogeneous part.

Thus, in both homogeneous and non-homogeneous cases, the estimatives are made from the explicit form of the solution, that is, by applying the Fourier transform to the respective equation.

Another result in this work is about the dissipative property of the homogeneous problem and some estimates of it, using differential calculus in H_{per}^s . This is included in Theorem 3.3 which we will develop in Subsection 3.2. So, using Theorem 3.3, we deduce the results of continuous dependence and uniqueness of solution for both homogeneous and non-homogeneous problems, respectively.

Finally, we obtain the continuous dependence and uniqueness of solution for the non-homogeneous case, applying the corresponding properties from [7].

We cite some works about KdV-K-S equation [1], [3] and for dissipative properties of systems [4].

Our article is organized as follows. In Section 2, we indicate the methodology used and cite the references used. In Section 3, we proved the main results.

Finally, in Section 4, we give the conclusions of our study.

2. Methodology

As a theoretical framework in this article we use the existence and regularity results of [7]. Also, we use the references [2], [6], [7], [8], [9], [10] and [5] for the Fourier theory in periodic Sobolev spaces, and differential and integral calculus in Banach spaces.

3. Main Results

First, using the Fourier transform, we will prove that the non-homogeneous problem has a unique solution and it continuously depends respect to the initial data and the non-homogeneity.

Second, we will study the uniqueness of the solution for homogeneous case using another technique that involves the dissipative property of the problem.

Finally, we will remark two additional ways leading to the continuous dependence of the solution for the non-homogeneous problem.

3.1. The non-homogeneous problem (P_c^F) is locally well posed

Theorem 3.1. *Let $\phi \in H_{\text{per}}^s$, $s \in \mathbb{R}$, $\mu > 0$, $F \in C([0, T], H_{\text{per}}^s)$, where $T > 0$, $\{S(t)\}_{t \geq 0}$ the semigroup of class C_o of homogeneous case ($F = 0$), introduced in the Theorem 3.2 from [3], and*

$$u_p(t) := \int_0^t S(t - \tau)F(\tau)d\tau,$$

then $u_p \in C([0, T], H_{\text{per}}^s) \cap C^1([0, T], H_{\text{per}}^{s-4})$ and satisfies

$$\begin{cases} \partial_t u_p + \partial_x^3 u_p + \mu(\partial_x^4 u_p + \partial_x^2 u_p) = F(t) \in H_{\text{per}}^{s-4}, \\ u_p(0) = 0, \end{cases} \quad (3.1)$$

with the derivative given by

$$\lim_{h \rightarrow 0} \left\| \frac{u_p(t+h) - u_p(t)}{h} + \partial_x^3 u_p + \mu(\partial_x^4 u_p + \partial_x^2 u_p) - F(t) \right\|_{s-4} = 0. \quad (3.2)$$

Proof. We remark that $S(t-\tau)F(\tau) \in H_{\text{per}}^s, \forall \tau \in (0, t)$ and $\tau \mapsto S(t-\tau)F(\tau)$

is continuous in $[0, t]$ then $\underbrace{\exists \int_0^t S(t-\tau)F(\tau)d\tau}_{u_p(t):=} \in H_{\text{per}}^s$.

Now, we will prove $u_p \in C([0, T], H_{\text{per}}^s)$, that is, $\|u_p(t+h) - u_p(t)\|_s \rightarrow 0$ when $h \rightarrow 0$.

Let $h > 0$

$$\begin{aligned} u_p(t+h) - u_p(t) &= \int_0^{t+h} S(t+h-\tau)F(\tau)d\tau - \int_0^t S(t-\tau)F(\tau)d\tau \\ &= \int_0^t \{S(t+h-\tau) - S(t-\tau)\}F(\tau)d\tau \\ &\quad + \int_t^{t+h} S(t+h-\tau)F(\tau)d\tau, \end{aligned}$$

taking the norm $\|\cdot\|_s$ we obtain

$$\begin{aligned} \|u_p(t+h) - u_p(t)\|_s &\leq \underbrace{\left\| \int_0^t \{S(t+h-\tau) - S(t-\tau)\}F(\tau)d\tau \right\|_s}_{I_1:=} \\ &\quad + \underbrace{\left\| \int_t^{t+h} S(t+h-\tau)F(\tau)d\tau \right\|_s}_{I_2:=}. \end{aligned}$$

Using the M -Test of Weierstrass, we get

$$\begin{aligned} I_1 &\leq \int_0^t \|S(t+h-\tau)F(\tau) - S(t-\tau)F(\tau)\|_s d\tau \\ &\leq \frac{\epsilon}{2T} \int_0^t d\tau = \frac{\epsilon t}{2T} \leq \frac{\epsilon T}{2T} = \frac{\epsilon}{2}, \end{aligned}$$

since $\exists \delta > 0$ such that:

$$\text{if } |h| < \delta, \text{ then } \|S(t+h-\tau)F(\tau) - S(t-\tau)F(\tau)\|_s < \frac{\epsilon}{2T}, \forall \tau \in (0, t).$$

Using the mean value Theorem in H_{per}^s , we obtain

$$\frac{1}{h} \int_t^{t+h} S(t+h-\tau)F(\tau) d\tau \rightarrow S(0)F(t) = F(t),$$

when $h \rightarrow 0$. Then

$$\int_t^{t+h} S(t+h-\tau)F(\tau) d\tau = \underbrace{\underbrace{h}_{\rightarrow 0} \cdot \frac{1}{h} \int_t^{t+h} S(t+h-\tau)F(\tau) d\tau}_{\rightarrow F(t)} \rightarrow 0,$$

when $h \rightarrow 0$. So, we have

$$\left\| \int_t^{t+h} S(t+h-\tau)F(\tau) d\tau \right\|_s \rightarrow 0,$$

when $h \rightarrow 0$. That is, $I_2 < \frac{\epsilon}{2}$ whenever $|h| < \delta^*$.

Therefore, $\|u_p(t+h) - u_p(t)\|_s \leq I_1 + I_2 < \epsilon$ whenever $|h| < \min\{\delta, \delta^*\}$.

From definition of $u_p(t)$ we have $u_p(0) = 0$.

Now, we will prove that $\exists \partial_t u_p(t)$ in H_{per}^s . In effect,

$$\begin{aligned}
& \frac{u_p(t+h) - u_p(t)}{h} \\
&= \frac{1}{h} \left\{ \int_0^{t+h} S(t+h-\tau)F(\tau)d\tau - \int_0^t S(t-\tau)F(\tau)d\tau \right\} \\
&= \frac{1}{h} \left\{ \int_0^t \{S(t+h-\tau)F(\tau) - S(t-\tau)F(\tau)\}d\tau + \int_t^{t+h} S(t+h-\tau)F(\tau)d\tau \right\} \\
&= \frac{1}{h} \int_0^t \{S(t+h-\tau)F(\tau) - S(t-\tau)F(\tau)\}d\tau + \frac{1}{h} \int_t^{t+h} S(t+h-\tau)F(\tau)d\tau.
\end{aligned}$$

Using the mean value Theorem in H_{per}^{s-4} , we obtain

$$\frac{1}{h} \int_t^{t+h} S(t+h-\tau)F(\tau)d\tau \rightarrow S(0)F(t) = F(t), \quad (3.3)$$

when $h \rightarrow 0$.

Since

$$\frac{S(t+h-\tau)F(\tau) - S(t-\tau)F(\tau)}{h}$$

converges uniformly to $\partial_t \{S(t-\tau)F(\tau)\}$ in H_{per}^{s-4} , $\forall \tau \in [0, t]$, we obtain that

$$\int_0^t \frac{S(t+h-\tau)F(\tau) - S(t-\tau)F(\tau)}{h} d\tau$$

converges to $\int_0^t \partial_t \{S(t-\tau)F(\tau)\} d\tau$ when $h \rightarrow 0$.

Now, we remark that $\partial_t\{S(t-\tau)F(\tau)\} = (-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))S(t-\tau)F(\tau)$ in H_{per}^{s-4} . Since $(-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))$ is a closed operator, then

$$\int_0^t (-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))S(t-\tau)F(\tau)d\tau = (-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))\int_0^t S(t-\tau)F(\tau)d\tau.$$

Therefore,

$$\int_0^t \partial_t\{S(t-\tau)F(\tau)\}d\tau = (-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))\underbrace{\int_0^t S(t-\tau)F(\tau)d\tau}_{u_p(t):=}$$

That is,

$$\int_0^t \frac{S(t+h-\tau)F(\tau) - S(t-\tau)F(\tau)}{h} d\tau \rightarrow (-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))u_p(t), \quad (3.4)$$

when $h \rightarrow 0$.

Finally, in H_{per}^{s-4} , using (3.3) and (3.4), we get

$$\underbrace{\exists \lim_{h \rightarrow 0} \frac{u_p(t+h) - u_p(t)}{h}}_{\partial_t u_p(t)=} = (-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))u_p(t) + F(t).$$

Using that $\|\partial_x^m f\|_{s-m} \leq \|f\|_s$, $\forall m \in \mathbb{Z}^+$, $\forall f \in H_{\text{per}}^s$, we obtain

$$\begin{aligned} & \|\partial_t u_p(t+h) - \partial_t u_p(t)\|_{s-4} \\ &= \|(-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))u_p(t+h) + F(t+h) \\ &\quad - \{(-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))u_p(t) + F(t)\}\|_{s-4} \\ &\leq \|F(t+h) - F(t)\|_{s-4} + \|(-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2))\{u_p(t+h) - u_p(t)\}\|_{s-4} \\ &\leq \|F(t+h) - F(t)\|_{s-4} + \|-\partial_x^3\{u_p(t+h) - u_p(t)\}\|_{s-4} \\ &\quad + \|\mu\partial_x^4\{u_p(t+h) - u_p(t)\}\|_{s-4} + \|\mu\partial_x^2\{u_p(t+h) - u_p(t)\}\|_{s-4} \end{aligned}$$

$$\begin{aligned} &\leq \|F(t+h) - F(t)\|_{s-4} + \|u_p(t+h) - u_p(t)\|_{s-1} \\ &\quad + \mu \|u_p(t+h) - u_p(t)\|_s + \mu \|u_p(t+h) - u_p(t)\|_{s-2}. \end{aligned}$$

Since $u_p(t) \in H_{\text{per}}^s \subset H_{\text{per}}^{s-1} \subset H_{\text{per}}^{s-2}$, then

$$\begin{aligned} &\|\partial_t u_p(t+h) - \partial_t u_p(t)\|_{s-4} \\ &\leq \|F(t+h) - F(t)\|_{s-4} + (1+2\mu)\|u_p(t+h) - u_p(t)\|_s \rightarrow 0, \end{aligned}$$

when $h \rightarrow 0$.

Now, since F is continuous in H_{per}^{s-4} and u_p is continuous in H_{per}^s , then $\partial_t u_p$ is continuous in H_{per}^{s-4} for $t \in [0, T]$, that is, $\partial_t u_p \in C([0, T], H_{\text{per}}^{s-4})$.

Therefore, $u_p \in C([0, T], H_{\text{per}}^s) \cap C^1([0, T], H_{\text{per}}^{s-4})$. $\square$

Theorem 3.2. Let $\phi \in H_{\text{per}}^s$, $s \in \mathbb{R}$, $\mu > 0$, $F \in C([0, T], H_{\text{per}}^s)$, where $T > 0$, and $\{S(t)\}_{t \geq 0}$ the semigroup of class C_o of contraction in H_{per}^s as in Theorem 3.1, then

(1) The function

$$u^F(t) := S(t)\phi + \underbrace{\int_0^t S(t-\tau)F(\tau)d\tau}_{u_p(t)=}, \quad t \in [0, T] \quad (3.5)$$

belongs to $C([0, T], H_{\text{per}}^s) \cap C^1([0, T], H_{\text{per}}^{s-4})$ and

(2) $u^F(t)$ is the unique solution of

$$(P_c^F) \quad \left\{ \begin{array}{l} u_t + u_{xxx} + \mu(u_{xxxx} + u_{xx}) = F(t) \in H_{\text{per}}^{s-4}, \\ u(0) = \phi, \end{array} \right. \quad (3.6)$$

with the derivative given by

$$\lim_{h \rightarrow 0} \left\| \frac{u(t+h) - u(t)}{h} + u_{xxx} + \mu(u_{xxxx} + u_{xx}) - F(t) \right\|_{s-4} = 0. \quad (3.7)$$

(3) Let $\psi_j \in H_{\text{per}}^s$, $F_j \in C([0, T], H_{\text{per}}^s)$, $j = 1, 2$. The map $\psi \rightarrow u$ is continuous in the following sense. Let u_1 and u_2 the corresponding solutions to initial data ψ_1 and ψ_2 , and with non-homogeneity F_1 and F_2 , respectively. Then

$$\|u_1(t) - u_2(t)\|_s \leq \|\psi_1 - \psi_2\|_s + T\|F_1 - F_2\|_{\infty, s-4}, \quad t \in [0, T], \quad (3.8)$$

$$\underbrace{\sup_{t \in [0, T]} \|u_1(t) - u_2(t)\|_s}_{\|u_1 - u_2\|_{\infty, s} :=} \leq \|\psi_1 - \psi_2\|_s + T\|F_1 - F_2\|_{\infty, s-4}, \quad (3.9)$$

$$\begin{aligned} \|\partial_t u_1(t) - \partial_t u_2(t)\|_{s-4} &\leq (1 + 2\mu)\|u_1(t) - u_2(t)\|_s + \|F_1 - F_2\|_{\infty, s-4}, \quad t \in [0, T], \\ &\leq (1 + 2\mu)\|u_1 - u_2\|_{\infty, s} + \|F_1 - F_2\|_{\infty, s-4} \\ &\leq (1 + 2\mu)\|\psi_1 - \psi_2\|_s + [(1 + 2\mu)T + 1]\|F_1 - F_2\|_{\infty, s-4}, \end{aligned} \quad (3.10)$$

where we have used the notation

$$\|h\|_{\infty, r} = \sup_{t \in [0, T]} \|h(t)\|_r, \quad h \in C([0, T], H_{\text{per}}^r). \quad (3.11)$$

Proof. We work the following steps:

(1) Let $u(t) := u^F(t) = S(t)\phi + u_p(t)$, we will prove that $u \in C([0, T], H_{\text{per}}^s) \cap C^1([0, T], H_{\text{per}}^{s-4})$. In effect, as $S(\cdot)\phi \in C([0, T], H_{\text{per}}^s)$ and $u_p(\cdot)\phi \in C([0, T], H_{\text{per}}^s)$, then $u(\cdot) = S(\cdot)\phi + u_p(\cdot) \in C([0, T], H_{\text{per}}^s)$.

Moreover, as $S(\cdot)\phi \in C^1([0, +\infty), H_{\text{per}}^{s-4})$ and $u_p(\cdot) \in C^1([0, T], H_{\text{per}}^{s-4})$, then $u(\cdot) = S(\cdot)\phi + u_p(\cdot) \in C^1([0, T], H_{\text{per}}^{s-4})$.

(2) We will prove that u is the solution of (P_c^F) . In effect, we know that $\exists \partial_t S(t)\phi$ and $\exists \partial_t u_p(t)$ in H_{per}^{s-4} , then

$$\begin{aligned} \partial_t u(t) &= \partial_t \underbrace{S(t)\phi}_{u_h(t)} + \partial_t u_p(t) \\ &= -\partial_x^3 u_h(t) - \mu(\partial_x^4 u_h(t) + \partial_x^2 u_h(t)) \\ &\quad - \partial_x^3 u_p(t) - \mu(\partial_x^4 u_p(t) + \partial_x^2 u_p(t)) + F(t) \\ &= -\partial_x^3 \{u_h(t) + u_p(t)\} - \mu(\partial_x^4 \{u_h(t) + u_p(t)\} \\ &\quad + \partial_x^2 \{u_h(t) + u_p(t)\}) + F(t) \\ &= -\partial_x^3 u(t) - \mu(\partial_x^4 u(t) + \partial_x^2 u(t)) + F(t) \end{aligned}$$

in H_{per}^{s-4} , where $u_h(\cdot)$ is solution of the homogeneous equation (P_c) .

(3) Also, $u(0) = u_h(0) + u_p(0) = \phi + 0 = \phi$.

(4) Let $\psi_j \in H_{\text{per}}^s$ and $F_j \in C([0, T], H_{\text{per}}^s)$ for $j = 1, 2$, then

$$u_j(t) = S(t)\psi_j + \int_0^t S(t-\tau)F_j(\tau)d\tau$$

is solution of $(P_c^{F_j})$ with initial data $u_j(0) = S(0)\psi_j = \psi_j$, for $j = 1, 2$.

Then

$$u_1(t) - u_2(t) = S(t)\{\psi_1 - \psi_2\} + \int_0^t S(t-\tau)\{F_1(\tau) - F_2(\tau)\}d\tau.$$

From where we obtain, for $t < T$:

$$\begin{aligned}
\|u_1(t) - u_2(t)\|_s &\leq \|S(t)\{\psi_1 - \psi_2\}\|_2 + \int_0^t \|S(t-\tau)\{F_1(\tau) - F_2(\tau)\}\|_s d\tau \\
&\leq \|\psi_1 - \psi_2\|_s + \int_0^t \|F_1(\tau) - F_2(\tau)\|_s d\tau \\
&\leq \|\psi_1 - \psi_2\|_s + \sup_{\tau \in [0, T]} \|F_1(\tau) - F_2(\tau)\|_s \int_0^t d\tau \\
&\leq \|\psi_1 - \psi_2\|_s + T \cdot \sup_{\tau \in [0, T]} \|F_1(\tau) - F_2(\tau)\|_s.
\end{aligned}$$

Therefore,

$$\sup_{t \in [0, T]} \|u_1(t) - u_2(t)\|_s \leq \|\psi_1 - \psi_2\|_s + T \cdot \sup_{\tau \in [0, T]} \|F_1(\tau) - F_2(\tau)\|_s.$$

(5) On the other hand, in H_{per}^{s-4} , we have

$$\begin{aligned}
\partial_t u_j(t) &= \partial_t u_{h,j}(t) + \partial_t u_{p,j} \\
&= [-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2)]u_{h,j} + [-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2)]u_{p,j} + F_j(t) \\
&= [-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2)]u_j + F_j(t),
\end{aligned}$$

for $j = 1, 2$. So,

$$\partial_t u_1(t) - \partial_t u_2(t) = [-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2)]\{u_1 - u_2\} + \{F_1(t) - F_2(t)\}.$$

Taking norm, we obtain

$$\begin{aligned}
\|\partial_t u_1(t) - \partial_t u_2(t)\|_{s-4} &= \|[-\partial_x^3 - \mu(\partial_x^4 + \partial_x^2)]\{u_1 - u_2\} + \{F_1(t) - F_2(t)\}\|_{s-4} \\
&\leq \|\partial_x^3\{u_1 - u_2\}\|_{s-4} + \mu\|\partial_x^4\{u_1 - u_2\}\|_{s-4} \\
&\quad + \mu\|\partial_x^2\{u_1 - u_2\}\|_{s-4} + \|F_1(t) - F_2(t)\|_{s-4}.
\end{aligned}$$

Using $\|\partial_x^m \psi\|_{s-m} \leq \|\psi\|_s$, for $m = 1, 2, 3$ and $H_{\text{per}}^s \subset H_{\text{per}}^{s-1} \subset H_{\text{per}}^{s-2}$, we get

$$\begin{aligned}
\|\partial_t u_1(t) - \partial_t u_2(t)\|_{s-4} &\leq \|u_1 - u_2\|_{s-1} + \mu \|u_1 - u_2\|_s \\
&\quad + \mu \|u_1 - u_2\|_{s-2} + \|F_1(t) - F_2(t)\|_{s-4} \\
&\leq (1 + 2\mu) \|u_1 - u_2\|_s + \|F_1(t) - F_2(t)\|_{s-4} \\
&\leq (1 + 2\mu) \|u_1(t) - u_2(t)\|_s + \sup_{t \in [0, T]} \|F_1(t) - F_2(t)\|_{s-4} \\
&\leq (1 + 2\mu) \|u_1 - u_2\|_{\infty, s} + \|F_1 - F_2\|_{\infty, s-4} \\
&\leq (1 + 2\mu) \|\psi_1 - \psi_2\|_s + [(1 + 2\mu)T + 1] \|F_1 - F_2\|_{\infty, s-4}.
\end{aligned}$$

□

Remark 3.1. Inequality (3.8) says that the solution of the non-homogeneous problem (P_c^F) continuously depends on the initial data and the non-homogeneity F .

Corollary 3.1. (P_c^F) has a unique solution.

Proof. This follows by applying inequality (3.8) with $\psi_1 = \psi_2$ and $F_1 = F_2$. □

Corollary 3.2. The unique solution of (P_c^F) is

$$\begin{aligned}
u^F(t, x) &= \sum_{k=-\infty}^{+\infty} e^{-\{-ik^3 + \mu(k^2-1)k^2\}t} \hat{\phi}(k) e^{ikx} \\
&\quad + \int_0^t \sum_{k=-\infty}^{+\infty} e^{-\{-ik^3 + \mu(k^2-1)k^2\}(t-\tau)} \hat{F}(k, \tau) e^{ikx} d\tau.
\end{aligned}$$

3.2. Dissipative property of the homogeneous problem

Let $\mu > 0$, $s \in \mathbb{R}$ and the homogeneous problem

$$(P_c) \left\{ \begin{array}{l} w \in C([0, \infty), H_{\text{per}}^s) \cap C^1([0, \infty), H_{\text{per}}^{s-4}) \\ \partial_t w + \partial_x^3 w + \mu(\partial_x^4 w + \partial_x^2 w) = 0 \in H_{\text{per}}^{s-4} \\ w(0) = \phi \in H_{\text{per}}^s. \end{array} \right.$$

Theorem 3.3. *Let w the solution of (P_c) with initial data $\phi \in H_{\text{per}}^s$, then we obtain the following results:*

$$(1) \partial_t \|w(t)\|_{s-4}^2 = -2\mu \langle \partial_x^2 w(t) + \partial_x^4 w(t), w(t) \rangle_{s-4} \leq 0.$$

$$(2) \|w(t)\|_{s-4} \leq \|\phi\|_{s-4} \leq \|\phi\|_s, \quad t \geq 0.$$

Proof. As $H_{\text{per}}^s \subset H_{\text{per}}^{s-4}$ then the following expressions are well defined: $\langle \partial_t w, w \rangle_{s-4}$ and $\langle w, \partial_t w \rangle_{s-4}$.

So, we have

$$\begin{aligned} \partial_t \|w(t)\|_{s-4}^2 &= \partial_t \langle w(t), w(t) \rangle_{s-4} \\ &= \langle \partial_t w(t), w(t) \rangle_{s-4} + \langle w(t), \partial_t w(t) \rangle_{s-4} \\ &= 2 \operatorname{Re} \langle \partial_t w(t), w(t) \rangle_{s-4}. \end{aligned} \quad (3.12)$$

For the other hand,

$$\begin{aligned} \langle \partial_x^2 w + \partial_x^4 w, w \rangle_{s-4} &= 2\pi \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} \{ \widehat{\partial_x^2 w + \partial_x^4 w}(k) \} \overline{\widehat{w}(k)} \\ &= 2\pi \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} \underbrace{k^2 (k^2 - 1)}_{M(k):=} \widehat{w}(k) \overline{\widehat{w}(k)} \\ &= 2\pi \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} M(k) |\widehat{w}(k)|^2. \end{aligned} \quad (3.13)$$

We remark that the series in (3.13) is convergent since $M(k) \leq k^4 \leq (k^2)^4 \leq (1+k^2)^4$, $\forall k \in Z$ and $w(t) \in H_{\text{per}}^s$.

Moreover, as

$$M(k) = \begin{cases} 0, & \text{if } k \in \{-1, 0, 1\}, \\ k^2(k^2 - 1) > 0, & \text{if } k \in Z - \{-1, 0, 1\}, \end{cases}$$

and $M(k) \geq 12$, $\forall k \in Z - \{-1, 0, 1\}$, then the convergent series (3.13) is positive.

That is,

$$\langle \partial_x^2 w + \partial_x^4 w, w \rangle_{s-4} \geq 0.$$

So, we have

$$- \langle \partial_x^2 w + \partial_x^4 w, w \rangle_{s-4} \leq 0. \quad (3.14)$$

Also, we obtain

$$\begin{aligned} \langle \partial_x^3 w, w \rangle_{s-4} &= 2\pi \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} \widehat{\partial_x^3 w}(k) \cdot \overline{\widehat{w}(k)} \\ &= 2\pi \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} (ik)^3 \widehat{w}(k) \cdot \overline{\widehat{w}(k)} \\ &= -i \underbrace{2\pi \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} k^3}_{b:=} |\widehat{w}(k)|^2. \end{aligned} \quad (3.15)$$

Now, we will prove that the series of the equality (3.15) is convergent. In effect, using the inequality: $|k|^3 \leq |k|^8 = (|k|^2)^4 \leq (1+k^2)^4$ and $w(t) \in H_{\text{per}}^s$, we have

$$\begin{aligned}
\left| \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} k^3 |\widehat{w}(k)|^2 \right| &\leq \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} |k|^3 |\widehat{w}(k)|^2 \\
&\leq \sum_{k=-\infty}^{+\infty} (1+k^2)^{s-4} (1+k^2)^4 |\widehat{w}(k)|^2 \\
&= \sum_{k=-\infty}^{+\infty} (1+k^2)^s |\widehat{w}(k)|^2 = \frac{1}{2\pi} \|w(t)\|_s^2 < \infty.
\end{aligned}$$

Then the series is convergent, that is,

$$\langle \partial_x^3 w, w \rangle_{s-4} = -ib, \text{ with } b \in \mathbb{R}. \quad (3.16)$$

From (3.12), using $\partial_t w = -\mu(\partial_x^2 w + \partial_x^4 w) - \partial_x^3 w$, $\mu > 0$, the inequality (3.14) and the equality (3.16), we get

$$\begin{aligned}
\partial_t \|w(t)\|_{s-4}^2 &= 2 \operatorname{Re} \langle \partial_t w(t), w(t) \rangle_{s-4} \\
&= 2 \operatorname{Re} \langle -\mu(\partial_x^2 w(t) + \partial_x^4 w(t)) - \partial_x^3 w(t), w(t) \rangle_{s-4} \\
&= 2\mu \operatorname{Re} \{ -\langle \partial_x^2 w(t) + \partial_x^4 w(t), w(t) \rangle_{s-4} \} \\
&\quad - 2 \operatorname{Re} \langle \partial_x^3 w(t), w(t) \rangle_{s-4} \\
&= -2\mu \operatorname{Re} \langle \partial_x^2 w(t) + \partial_x^4 w(t), w(t) \rangle_{s-4} - 0 \\
&= -2\mu \operatorname{Re} \langle \partial_x^2 w(t) + \partial_x^4 w(t), w(t) \rangle_{s-4} \leq 0.
\end{aligned}$$

Therefore, $\|w(t)\|_{s-4}^2$ is not increasing. Then, $\|w(t)\|_{s-4}^2 \leq \|w(0)\|_{s-4}^2$, $\forall t \geq 0$.

As $(\|w(t)\|_{s-4} - \|w(0)\|_{s-4})(\|w(t)\|_{s-4} + \|w(0)\|_{s-4}) \leq 0$, we have

$$\|w(t)\|_{s-4} \leq \|w(0)\|_{s-4} \leq \|w(0)\|_s, \quad \forall t \geq 0.$$

That is, $\|w(t)\|_{s-4} \leq \|\phi\|_{s-4} \leq \|\phi\|_s$, $\forall t \geq 0$. □

Corollary 3.3 (Continuous dependence of the solution of (P_c)). *Let u and v solutions of (P_c) with initial data ϕ_1 and ϕ_2 in H_{per}^s , respectively. Then the following assertions hold:*

$$\partial_t \|u(t) - v(t)\|_{s-4}^2 = -2\mu < \partial_x^2 \{u(t) - v(t)\} + \partial_x^4 \{u(t) - v(t)\}, \quad u(t) - v(t) >_{s-4} \leq 0,$$

and

$$\|u(t) - v(t)\|_{s-4} \leq \|\phi_1 - \phi_2\|_{s-4} \leq \|\phi_1 - \phi_2\|_s, \quad t \geq 0. \quad (3.17)$$

Proof. Define $w := u - v$ then w satisfies:

$$\begin{cases} \partial_t w + \partial_x^3 w + \mu \{ \partial_x^2 w + \partial_x^4 w \} = 0, \\ w(0) = \phi_1 - \phi_2. \end{cases}$$

We conclude using the Theorem 3.3. $\square$

Corollary 3.4 (Uniqueness of solution of (P_c)). *(P_c) has a unique solution.*

Proof. In effect, let u and v solutions of (P_c) with the same initial data, that is, $\phi_1 = \phi_2 = \phi$.

From (3.17), we obtain $\|u(t) - v(t)\|_{s-4} \leq \|0\|_s = 0$. Then, $\|u(t) - v(t)\|_{s-4} = 0$.

So, $u(t) = v(t)$, $\forall t \geq 0$, that is, $u = v$. $\square$

3.3. Continuous dependence and uniqueness solution of (P_c^F) by Theorem 3.3

The Theorem 3.3 will allow us to prove the following results.

Corollary 3.5 (Continuous dependence of the solution of (P_c^F)). *Let u and v solutions of (P_c^F) with initial data ϕ_1 and ϕ_2 in H_{per}^s , respectively. Then $u - v$ is solution of (P_c) with initial data $\phi_1 - \phi_2$ and satisfies:*

$$\partial_t \|u(t) - v(t)\|_{s-4}^2 = -2\mu < \partial_x^2 \{u(t) - v(t)\} + \partial_x^4 \{u(t) - v(t)\}, u(t) - v(t) >_{s-4} \leq 0,$$

and

$$\|u(t) - v(t)\|_{s-4} \leq \|\phi_1 - \phi_2\|_{s-4} \leq \|\phi_1 - \phi_2\|_s. \quad (3.18)$$

Proof. It is an immediate consequence from Theorem 3.3. $\square$

Corollary 3.6 (Uniqueness solution of (P_c^F)). (P_c^F) has a unique solution.

Proof. In effect, let u and v solutions of (P_c^F) with the same initial data, that is $\phi_1 = \phi_2 = \phi \in H_{\text{per}}^s$, then using the Corollary 3.5, we have $u - v$ is solution of (P_c) with initial data 0 and $\|u(t) - v(t)\|_{s-4} \leq \|0\|_s = 0$.

Therefore, $\|u(t) - v(t)\|_{s-4} = 0$. Thus, $u(t) = v(t)$, $\forall t \geq 0$, that is, $u = v$.

$\square$

3.4. Continuous dependency and uniqueness solution of (P_c^F) through results from [7]

In this subsection, we will use results from [7] to prove the following corollary:

Corollary 3.7 (Continuous dependency of the solution of (P_c^F)). Let u and v solutions of (P_c^F) with initial data ϕ_1 and ϕ_2 in H_{per}^s , respectively. Then $u - v$ is solution of (P_c) with initial data $\phi_1 - \phi_2$ and satisfies:

$$\|u(t) - v(t)\|_s \leq \|\phi_1 - \phi_2\|_s, \quad t \in [0, T], \quad (3.19)$$

$$\|u(t) - v(t)\|_r \leq C\|\phi_1 - \phi_2\|_s, \quad \forall r \in \mathbb{R}, \quad t \in (0, T]. \quad (3.20)$$

Proof. From [7]: inequality (3.2) and inequality of Theorem 3.1, we conclude. $\square$

Remark 3.2. As a consequence of the Corollary 3.7, taking $\phi_1 = \phi_2$, we obtain the uniqueness of solution of the problem (P_c^F) .

4. Conclusion

From our study of the KdV-K-S equation in periodic Sobolev spaces, for the homogeneous (P_c) and non-homogeneous (P_c^F) case, we have obtained the following results:

(1) In Theorem 3.2, using the Fourier transform, we proved that the non-homogeneous problem (P_c^F) is locally well posed in compacts, obtaining continuous dependence with respect to the initial data and the non-homogeneity.

(2) We showed the dissipative property of the homogeneous problem, where an estimate for the norm of global solution was obtained, which allowed us to deduce the continuous dependence (with respect to the initial data) and uniqueness solution of (P_c) .

(3) We prove the continuous dependence respect to the initial data, and uniqueness solution of (P_c^F) by using Theorem 3.3.

(4) We prove the continuous dependence respect to the initial data, and uniqueness solution of (P_c^F) by using results from [7].

(5) In this paper, we have obtained results that can be applied to the KdV-K-S models with two parameters, and these promote the analysis of the corresponding convergence.

References

- [1] H. A. Biagioni, J. L. Bona, R. Iorio and M. Scialom, On the Korteweg-de Vries-Kuramoto-Sivashinsky equation, *Advances in Differential Equations* 1(1) (1996), 1-20.
- [2] R. J. Iorio, *Fourier Analysis and Partial Differential Equations*, Cambridge University, 2002.
- [3] P. Guzman, S. Marx and E. Cerpa, Stabilization of the linear Kuramoto-Sivashinsky equation with a delayed boundary control, *IFAC Papers Online* 52(02) (2019), 70-75.
- [4] Z. Liu and S. Zheng, *Semigroups Associated with Dissipative System*, Chapman Hall/CRC, 1999.
- [5] H. K. Pathak, *An Introduction to Nonlinear Analysis and Fixed Point Theory*, Springer, Singapur, 2018.
- [6] Yolanda Santiago Ayala, Santiago Rojas Romero and Teófanés Quispe Mendez, Espacios de sobolev periódico y unproblema de Cauchy asociado a un modelo de ondas en un fluido viscoso, *Theorema, Segunda Epoca* 3(04) (2016), 7-23.
- [7] Yolanda Santiago Ayala and Santiago Rojas Romero, Regularity and well-posedness of a problem to one parameter and its behavior at the limit, *Bulletin of the Allahabad Mathematical Society* 32(02) (2017), 207-230.
- [8] Yolanda Santiago Ayala and Santiago Rojas Romero, Existencia y regularidad de solución de la ecuación del calor en espacios de Sobolev periódico, *Selecciones Matemáticas* 06(1) (2019), 49-65.
DOI: <http://dx.doi.org/10.17268/sel.mat.2019.01.08>
- [9] Yolanda Santiago Ayala and Santiago Rojas Romero, Unicidad de solución de la ecuación del calor en espacios de Sobolev periódico, *Selecciones Matemáticas* 07(1) (2020), 172-175.
DOI: <http://dx.doi.org/10.17268/sel.mat.2020.01.16>
- [10] D. Stuart, *Partial Differential Equations Example Sheet 2*, 2014.
<http://www.damtp.cam.ac.uk/user/examples/D15b.pdf>