Research Article

Physicochemical variations in the honey produced by *Apis dorsata* from Punjab, Pakistan

Rizwana Kousar¹ and Samina Qamer^{2*}

1. Department of Zoology, Government College University, Faisalabad-Pakistan

2. Honey Bee Research Institute, National Agricultural Research Centre, Islamabad-Pakistan

*Corresponding author's email: saminabee@yahoo.com

Citation

Rizwana Kousar and Samina Qamer. Physicochemical variations in the honey produced by *Apis dorsata* from Punjab, Pakistan. Pure and Applied Biology. Vol. 6, Issue 2, pp733-739.

<http://dx.doi.org/10.19045/bspab.2017.60078>

Received: 27/10/2016

Revised: 09/03/2017

Accepted: 15/03/2017

Online First: 19/03/2017

Abstract

A total of 24 samples was collected from three different localities (n= 3, Lahore, n= 10, Jhang and n=11, Kalar Kahar). These honey samples were analyzed for moisture content, pH, electrical conductivity, free acidity, lactone content, total acidity, proline and HMF content for quality evaluation of *Apis dorsata* honey. The average moisture content was 22.86%, 22.97% and 21.13% from Lahore, Jhang and Kalar Kahar, respectively. The pH values fell within the prescribed range of 3.42-6.1. The mean free acidity of all honey samples was well with international limits. Lactone content was 33.44meq/kg from Lahore, 15.53meq/kg from Jhang and 12.77meq/kg from Kalar Kahar. Total acidity calculated for honey samples of Lahore was 60.21meq/kg, 55.46meq/kg for Jhang and 50.36meq/kg for Kalar Kahar region. Proline content was estimated at an average of 5146.62mg/kg of honey samples from Lahore, 4328.44mg/kg from Jhang and 3554.68mg/kg from Kalar Kahar honey samples which were higher than minimum International Honey Standards limit. HMF content was measured as 36.41mg/kg from Lahore, 30.44mg/kg from Jhang and 89.93mg/kg from Kalar Kahar. The HMF Content was within the maximum International limit.

Keywords: *A.dorsata* honey; Proline and HMF content; EC value

Introduction

Apis dorsata, the giant honeybee, is found in the tropical regions of Asia [1]. It is also one of the native honeybee species of Pakistan commonly known as “dommna” and wild in nature. It's yellow in color with black stripes on each adnominal segment. It usually builds a single comb which is 1-1.6 m wide and 0.8-1.5m long under the stout branches of tall trees or buildings or water tower or cliff to protect their nests from top predators [2]. The comb is formed by several layers of

protective curtains of bees. These protective curtains keep a constant temperature for brood nest between 0-33 °C. *Apis dorsata* seasonally migrates back and forth between low (10 m) and uplands (1100 m) to escape from harsh environmental conditions. When it migrates from low agriculture lands, they always aggregate in masses from 25-120 colonies (depend on nesting space availability). At this time of the year different types of honey crops are bloom. A single fully-grown nest may weigh over 20

kg including brood, honey, pollen and adult bee. However, in wet season they never aggregate in masses probably due to scarcity of food, strong wind, heavy rain fall, storms, and the low number of individuals in the colony [3]. Although, physical properties and chemical composition of honey has been investigated by many workers [4, 5]. However, it may vary with the bee species responsible for its production. In Pakistan commercial beekeeping is done by keeping *A.mellifera* (European honeybee species) hives and quality evaluation of honey in the country is mostly carried out by either buying or collecting honey available in local market or from beekeepers [6, 7]. The composition of honey also depends on the type of plant source utilized by the bees as well as regional and climatic conditions [4, 8, 9]. Honey is generally evaluated by a physicochemical analysis of constituents. Several of these constituents are of great importance to the honey industry as they influence the storage quality, granulation, texture, flavor and the nutritional and medicinal quality of the honey. The present study was aimed firstly to find *Apis dorsata* colonies on different floral sources and secondly to assess the physicochemical properties of *Apis dorsata* honey from Pakistan in order to increase the scientific knowledge about the honey produced by this honey bee species. A total of 24 samples were collected from three different localities

Nankan Sahib, Faisalabad and Kalar Kahar. The predominant vegetation in these areas were *Guava*, *Jaman Eucalyptus* in Nankana Sahib, *Citrus*, *Phulai*, *Mulberry*, *Neem*, *Guava*, *Eucalyptus*, *Peepal*, *Acacia* in Faisalabad and *Jojoba*, *Phulai*, *Neem*, *Mulberry*, *Citrus*, *Mango*, *Acacia*, *Sheesham* in Kalar Kahar.

Materials and methods

Fresh Pakistani *Apis dorsata* honey samples were collected from different areas of Punjab, Pakistan: Lahore (Figure 1), Jhang (Figure 2) and Kalar Kahar (Figure 3) (Table 1). All these samples, after collecting, were stored in tightly capped jars under ordinary room temperature (25°C - 29°C) in the laboratory until completely analysis. There was no sign of granulation in honey samples. No preservative or any heating was applied at any stage. Honey samples were diluted in distilled water at room temperature according to the requirements of the test. All tests were performed in triplicate at the Research laboratory of Zoology department of GC University, Faisalabad. The physicochemical methods used for the characterization of honey are those as recommended by International Honey Commission [10]. The data was subjected to statistical analysis in which different physicochemical parameters were compared with international reference limits using student's "t" test [11].

Figure 1. Colony of *Apis dorsata* on guava tree in Lahore

Figure 2. Colony of *Apis dorsata* on citrus tree in Jhang

Figure 3. Colony of *Apis dorsata* on Acacia tree in Kalar Kahar

Table 1. Samples collection sites of *Apis dorsata* honey

S.No	Nectar source (predominant)	Locality	No. of samples
1	Guava, Jaman Eucalyptus	Lahore	3
2	Citrus, Phulai, Mulberry, Neem, Guava, Eucalyptus, Peepal, Acacia	Jhang	10
3	Jojoba, Phulai, Neem, Mulberry, Citrus, Mango, Acacia, Sheesham	Kalar Kahar	11

Results and discussion

Moisture Content

The moisture content in all samples of *Apis dorsata* are exceeding (Table 2) the limits of International Honey Standards and of

Directive, 2001/110/EC from council of European Union, according to which water content of honey should not be more than 21%. This higher moisture content in *Apis dorsata* honey has also been recoded

previously by [12] from four (23.5%, 23.99%, 23.91% and 22.2%) different forests of Nepal, 20.9% (18.9-24.2) by [13], 21% by [14], 23.1% and 27.8% by [15] and [16] from Philippines. This high water content in *Apis dorsata* honey could be because of the possible dilution of honey by

rain water as this honey bee species forms comb high in open air [17]. Therefore, uniformly high moisture content seems to be the peculiarity of *Apis dorsata* honey and could be used to evaluate the honey on the basis on honey bee species.

Table 2. Comparison of physico-chemical characteristics of *Apis dorsata* honey from Lahore, Jhang and Kalar Kahar

S. No	Nesting site →	Jhang		Lahore		Kalar Kahar		Codex draft 1999	EU draft 1999	Directive 2001/EC **
	Parameters ↓	Range	Mean	Range	Mean	Range	Mean			
1	pH	3.57-4.62	4.11 ±0.14	3.44-3.73	3.48 ±0.06	3.52-4.74	4.62 ±0.26			
2	Moisture content%	22.20-25.00	22.97 ±0.26	21.80-25.00	22.86 ±0.26	21.67-25.00	21.13 ±0.87	<21%	<21%	<21%
3	EC (μS/cm)	277.67-1347.87	911.534 ±7.23	233.00-272.67	249 ±1.12	258.67-524.33	459.97 ±3.84	0.80*mS/cm		
4	Free acidity (meq/kg)	21.67-58.00	42.64 ±3.85	19.00-33.33	25.77 ±1.48	26.83-58.17	37.11 ±2.44	≤50meq/kg	≤40meq/kg	≤50meq/kg
5	Lactone content (meq/kg)	4.00-34.16	15.53 ±1.62	30.00-31.83	33.44 ±2.46	6.00-21.27	12.77 ±1.69			
6	Total acidity (meq/kg)	51.20-82.47	55.46 ±5.21	51.20-69.67	60.21 ±7.55	31.67-74.83	50.36 ±3.93			
7	Proline content (mg/kg)	3415.73-6344.61	4328.44 ±342.67	4465.13-8760.51	5146.62 ±2.46	2808.38-7939.09	3554.68 ±359.81	≥180*mg/kg		
8	HMF content (mg/kg)	14.82-43.76	30.44 ±5.29	28.54-43.33	36.41 ±4.46	48.54-149.76	89.93 ±5.40	≤60mg/kg		

“t”test. Probability values at 5% for different physicochemical parameters of Faisalabad, Nankana Shib and Kalar Kahar area. For moisture content $p>0.05$, $p>0.05$ for pH, $p>0.05$ for EC, $p>0.05$ for free acidity, Lactone and total acidity, $p>0.05$ for proline content and HMF content.

* The suggested values for proline content and electrical conductivity for new honey standards (Bogdanov *et al.*, 1999)

** Council of the European Union. Council Directive 2001/110/EC of Dec. 20, 2001, relating to honey. *Off. J. Eur. Commun.*, 2002, Jan 12, L10/47-L10/52

pH

The average pH from Lahore, Jhang and Kalar Kahar *Apis dorsata* honey samples were found to be 3.48 (3.44-3.73), 4.11 (3.57-4.62), 4.02 (3.52-4.74), respectively. These values showed no significant difference and fell within the prescribed range of 3.42 - 6.1 [18]. The pH values obtained in present study showed close resemblance with the values (3.66 and 4.06) obtained by [17] in the honey samples of *Apis dorsata* harvested from two different

nesting sites (trees) in Chitwan district, central Nepal. These pH values, however, lower as compares to the ones obtained by [12] in *Apis dorsata* honey samples harvested from four 4.68 (4.3-5.1), 4.58 (4.3-4.7), 4.39 (3.7-4.6) and 3.8 (3.7-4.3) Terai forests, Nepal.

Electrical conductivity

Average values of electrical conductivity (EC) of *Apis dorsata* honey were 249μS/cm (233.0-272.67μS/cm), 911.53μS/cm (277.67-1347μS/cm) and 459.97μS/cm

(258.67-524.33 $\mu\text{S/cm}$), from Lahore, Jhang and Kalar Kahar. All these values of electrical conductivity value of were higher than least limits of international honey standards and of directive 2001/110/EC from council of European Union for blossom honey ($\sim 0.8 \text{ mS/cm}$). [17] recorded EC values of *Apis dorsata* honey as 0.9 mS/cm. The conductivity values found by [12] were 0.060 (0.44-0.70), 0.61 (0.51-0.74), 0.48 (0.19-0.54) and 0.22mS/cm (0.18-0.42). EC value 0.558 mS/cm was reported by [4]. All these EC values were lower than the value found in *Apis dorsata* honey samples from Punjab.

Free acidity

The Free acidity in *Apis dorsata* honey from Lahore, Jhang and Kalar Kahar were 25.77meq/kg (19.00-33.33 meq/kg), 42.65meq/kg (21.67-58.00 meq/kg) and 37.11meq/kg (26.83-58.17meq/kg), respectively. Although average free acidity was found less than 50meq/kg, a maximum limit for acidity prescribed by international honey standards and by the directive 2001/110/EC from the Council of European Union. However, few (samples no 11, 13 and 16) from Jhang and samples no 7 and 8 from Kalar Kahar had free acidity value above the permissible limit. [19] from Pakistan, [20] from Calcutta, India reported formic acid value in *Apis dorsata* honey. [17] did not report free or total acidity in *Apis dorsata* honey samples collected from Chitwan district, Central Nepal. [12] reported free acidity values 44.459meq/kg (26.5-51.59meq/kg), 43.169meq/kg (35.00-47.519meq/kg), 43.149meq/kg (33.5-60.009meq/kg) and 48.9meq/kg (39.5-61.00) of Nepal's *Apis dorsata* honey. The free acidity values obtained from present study are closer to the values of *Apis dorsata* honey reported by [12].

Lactone content

Lactone content was measured at an average of 33.44meq/kg (30-32.83meq/kg)

of honey sample collected from Lahore, 15.53meq/kg (4.00-34.16meq/kg) from Jhang and 12.77meq/kg (6.00-21.67meq/kg) from Kalar Kahar. All these values of lactone content of *Apis dorsata* honey were found L.A vales closer to the values were 13.2 meq/kg (0.00-20.5 meq/kg), 18.79 meq/kg (11.5-18.00 meq/kg), 15.14 meq/kg (12.5-18.5 meq/kg) and 3.5meq/kg (7.00-21.00 meq/kg) determined by [12]. [4] noticed the lactone value in Spanish honey at an average of 5.08meq/kg.

Total acidity

The total acidity calculated for honey samples of Lahore 60.21meq/kg (51.20-69.67 meq/kg), Jhang 55.46 meq/kg (51.20-82.47meq/kg), and Kalar Kahar was 50.36meq/kg (31.67-74.83 meq/kg). [12] reported total acidity value of 61.74meq/kg (30.00-71.50meq/kg), 67.98meq/kg (46.5-64.00meq/kg), 60.02meq/kg (46.00-77.00meq/kg) and 56.39meq/kg (53.00-79.50meq/kg) in *Apis dorsata* honey collected from four Teari forest, Nepal. [16] from Philippines reported total acidity value of 40.2meq/kg as an average in *Apis dorsata* honey. [4] reported 33.23meq/kg mean total acidity in honey samples from Central Spain. High acidity has been considered as an indicator of purity of honey by [21]. Possible reason of elevated acidity values in *Apis dorsata* honey is unlike that of *A. mellifera* honey is because of its multi-floral foraging behavior. Different types of nectars with various concentrations of acids may ultimately result into high level of acidity. These high acidity values of *Apis dorsata* honey could be due to limited or short termed shelf life.

Proline content

Proline content were estimated at an average of 5146.62 mg/kg (4465.13-8760.51mg/kg) from Lahore, 4328.44mg/kg (3415.73-6344.61mg/kg) Jhang and 3554.68mg/kg (2808.38-7939.09mg/kg) of *Apis dorsata* honey samples from Kalar Kahar. [12]

reported 98.38mg/kg (23.38-153.6mg/kg), 100.8mg/kg (20.65-330.00mg/kg), 119.98mg/kg (42.15-193.18mg/kg) and 160mg/kg (101.97-198.81mg/kg) of proline content of *A. dorsata* honey samples from Nepal. [17] reported 875.82ppm of proline content in *A. dorsata* honey samples from Nepal. [22] reported 76.0mg/100g and 81.1mg/100g proline in honey samples from two different geographical regions of Spain. The proline content of *A. dorsata* honey samples from Punjab (Lahore, Jhang and Kalar Kahar) was higher from all of them. These results show no resemblance with previous determination of proline content in *Apis dorsata* honey samples. However, proline content of Jhang and Kalar Kahar were close. [23] determined proline content varied from 437.82 to 2169mg/kg in *A. mellifera* honey. Some authors have reported that high values for proline are typical for honeydew honeys [24]. The proline content has been shown to vary considerably between different honeys [10]. [25] Louveaux (1985) believe that the majority of the proline comes from bee salivary secretion.

HMF Content

HMF is one of the important quality criterions to determine the freshness and overheating of honey. In order to check the freshness of Pakistani *Apis dorsata* honey average HMF Content were measured as 36.41mg/kg (28.54-43.13mg/kg) from Lahore, 30.74mg/kg (16.32-53.76mg/kg) Jhang and 89.93mg/kg (48.45-149.76mg/kg) from Kalar Kahar. HMF Content reported by [12] was 33.36mg/kg (7.68-61.28mg/kg), 30.36mg/kg (3.84-57.60mg/kg), 36.48mg/kg (7.68-59.52mg/kg) and 56.10mg/kg (21.12-76.6mg/kg) from four Terai forests, Nepal. The present study showed that HMF content in Lahore and Jhang honey samples were within the limit of International honey Standards i-e 40mg/kg, but the HMF content of *Apis dorsata* honey samples collected

from Kalar Kahar exceeds the maximum limit described for HMF Content by Codex Alimentarius. The reason of high HMF content might be that, all the three areas from where these honey samples were collected, have sub-tropical climate and according to [26] the honey produced in subtropical climate have high HMF value which may exceeds 40mg/kg.

Conclusions

The present investigations conclude that honey produce by *Apis dorsata* showed various quality parameters close to the International honey standards. Moisture content is beyond the Standard limits and seems to be specific in the honey produced by this honeybee species.

Authors' contributions

Conceived, designed the experiments: S Qamer, Performed the experiments: R Kousar, Analyzed the data: R Kousar, Wrote the paper: S Qamer.

References

1. Ruttner F (1988). Biogeography and taxonomy of honey bees, *Springer-Verlag, Berlin Heidelberg* 284.
2. Wongsiri S, Thapa R, Oldroyd B & Burgett MD (1996). A magic bee tree: Home of *Apis dorsata*. *Am Bee J* 136: 196-199.
3. Thapa R, Wongsiri S & Prawn S (1999b). Colony migration of *Apis dorsata* in the northern parts of Thailand. *Asian bees and bee keeping in: proceedings of Res and Dev* 39-43.
4. Iglesias MT, De lorenzo C, Polo MDC, Martin-alvarez PJ & Pueyo E (2004). Usefulness of amino acid composition to discriminate between honeydew and floral honeys. Application to honeys from a small geographic area. *J Agric Fd Chem* 52: 84-8.
5. Qamer S, Muzaffar N, Ali SS & Shakoori AR (2005). Physicochemical analysis of Pakistani unifloral honey from different nectar sources. *Proc Pakistan Congr Zoo* 25: 123-130.
6. Khan KA, Al-Ghamdi AA & Ansari MJ (2016). The characterization of blossom honeys from two provinces of Pakistan.

- Italian Journal of Food Science* 28: 625-637.
7. Iftakhar F, Mahmood R, Islam N, Sarwar G, Masood MA & Shafiq H (2014). Physicochemical Analysis of Honey Samples Collected from Local Markets of Rawalpindi and Islamabad, Pakistan. *American Journal of Biochemistry* 4 (2): 35-40.
 8. Abu-tarboush HM, Al-kahtani HA & El-sarrage MS (1993). Floral type identification and quality evaluation of some honey types. *Fd Chem* 46:13-17.
 9. Bonod L, Sandoz JC, Loublier Y & Pham-delegue MH (2003). Learning and discrimination of honey odours by the honey bees. *Apidologie* 34: 147-159.
 10. Bogdanov S, Lullmann C, Martin P, Ohe Von Der W, Russmann H, Vorwohl G, Marcazzan GL, Piro R, Flamini C, Morlot M, Lehritier J, Borneck R, Marioleas P, Tsigouri A, Kerkuliet J, Ortiz A, Ivanov T, Darcy B, Mossel B & Vit P (1999). Honey quality and international regulatory standards: review by the International Honey Commission. *Bee Wld* 80: 61-69.
 11. Steel RGD & Torrie JH (1981). Analysis of variance factorial experiments. In: Principal and procedure of statistics. 2nd Ed McGraw Hill International Book Company Tokyo, pp. 336-375.
 12. Qamer S, Ahmad F, Latif F, Ali SS & Shakoori AR (2008). Physicochemical Analysis of *Apis dorsata* honey from Terai forests, Nepal. *Pakistan Journal of Zoology* 40: 53-58.
 13. Phadke RP (1968). Studies on Indian honey. Proximate composition and physicochemical characterization of honey from wild honey bee *Apis dorsata*, *Apis florea* and *Trigona*. *Indian Bee J* 30: 3-8.
 14. Malkar A (1997). Comparative pollen and chemical analysis of West Bengal honey. Ph.D. thesis, University Calcutta, India.
 15. Laude VT, Naegel L & Horn H (1991). The physicochemical properties of some Philippine honey. *Apidologie* 22: 371-257.
 16. Minah HV, Mendoza BG & Laigo FM (1971). The chemical composition of honey produced by *Apis dorsata*. *J Apic Res* 10: 91-97.
 17. Joshi SR, Pechhaker H, Willam A & Vonderohe W (2000). Physicochemical characteristic of *Apis dorsata*, *Apis cerana* and *Apis mellifera* From Chitwan district, Nepal. *Apidologie* 31: 367-375.
 18. White JRJW, Riethof ML, Subers MH & Kushnir I (1962). Composition of American honey. *Tech Bull U. S. Dept Agric* 126: 144.
 19. Latif A, Qayyum A & Mazoor-ul-haq (1956). Research on the composition of Pakistan honey. *Pakistan. J scient Res* 8: 163-380.
 20. Mitra SN & Mathew TV (1968). Studies on the characterizations of honey. *J Proc Inst Chem India* 40: 26-30.
 21. Suarez-luque S, Mato I & Huidobro JF (2002). Rapid determination of minority organic acids in honey by high performance liquid chromatography. *J Chromatogr* 955: 207-214.
 22. Sanchez MDP, Huidobro JF, Matp I, MuniateguI S & Sancho MT (2001). Correlation between Proline content of honey and botanical origin. *Deutsche. Lebensum. Rundschau* 5: 171-175.
 23. Meda ACE, Lamien JMM, Romito OG & Nacoulma (2005). Physicochemical analysis of Burkana Fasan honey. *Acta Vet Brno* 74: 147-152.
 24. Diez MJ, Andres C & Terrab A (2004). Physicochemical parameters and pollen analysis of Moroccan honeydew honeys. *International Journal of Food Science and Technology* 39: 167-176.
 25. Louveaux J (1985). Le miel Cah. *Nutr Diet* 20: 57-70.
 26. Lagrange V & Sanders SW (1988). Honey in cereal-based new food products. *Cereal Fd Wld* 33:833-838.