Research Article

Antimicrobial properties of hydrogen peroxide and potash alum alone and in combination against clinical bacterial isolates

Huma Khurshid¹, Muhammad Rafiq², Fahad Nazir³, Imran Ali⁴, Muneer Ahmed⁵, Bilal Akbar⁶, Mushtaq Ahmed¹ and Anser Ali^{1*}

1. Department of Zoology, Mirpur University of Science and Technology (MUST), Mirpur-10250 (AJK)-Pakistan

2. Department of Physiology & Biochemistry, Cholistan University of Veterinary & Animal Sciences, Bahawalpur-Pakistan

3. Department of Entomology, Faculty of Crop Protection, Sindh Agriculture University Tandojam-Pakistan

4. Department of Biotechnology, Mirpur University of Science and Technology (MUST), Mirpur-10250 (AJK)-Pakistan

5. School of Biological Sciences, University of Punjab, Quaid-e-Azam Campus, Lahore-Pakistan

6. Department of Mechanical Engineering, Mirpur University of Science and Technology (MUST), Mirpur-10250 (AJK)-Pakistan

*Corresponding author's email: anser.zoology@must.edu.pk

Citation

Huma Khurshid, Muhammad Rafiq, Fahad Nazir, Imran Ali, Muneer Ahmed, Bilal Akbar, Mushtaq Ahmed and Anser Ali. Antimicrobial properties of hydrogen peroxide and potash alum alone and in combination against clinical bacterial isolates. Pure and Applied Biology. Vol. 8, Issue 4, pp2238-2247.

<http://dx.doi.org/10.19045/bspab.2019.80169>

Received: 20/05/2019

Revised: 16/07/2019

Accepted: 24/07/2019

Online First: 03/08/2019

Abstract

Hydrogen peroxide (H₂O₂) and potash alum (ALM) are considered important for wide range of biological activities including microbial disinfection. However, comparison of H₂O₂ and ALM antibacterial effect alone and in combination on different clinical bacterial isolates with respect to post treatment time is still not well studied. Thus, in present study we tested susceptibility of five bacterial isolates; *Enterococcus faecalis* (*E. faecalis*), *Enterococcus faecium* (*E. faecium*), *Escherichia coli* (*E. coli*), *Staphylococcus aureus* (*S. aureus*) and *Klebsiella pneumoniae* (*K. pneumoniae*) against H₂O₂ and ALM alone and in combination which showed concentration dependent and incubation dependent effect. The H₂O₂ bacterial susceptibility trend was *E. coli* > *K. pneumoniae* > *S. aureus* > *E. faecium* > *E. faecalis* and ALM susceptibility trend was as *K. pneumoniae* > *S. aureus* > *E. faecalis* > *E. coli* > *E. faecium* at highest tested concentration (35mg/ml) and 24hr of incubation period. Comparatively, antimicrobial activity was higher with H₂O₂ than ALM however in the order of 24hr > 48hr > 72hr. Both H₂O₂ and ALM showed more zones of inhibition at highest tested concentration than positive control azithromycin (AZI) against *E. faecalis*, *E. faecium* and *S. aureus* at all incubation periods. Moreover, H₂O₂ and ALM in combination (1:1) showed increased zone of inhibition than ALM alone (against all bacteria), H₂O₂ alone (against *E. faecalis* and *E. faecium*) and, AZI (against *E. faecalis*, *E. faecium* and *S. aureus*) without increasing final concentration. Thus, combination treatment might be more effective disinfection and antisepsis strategy which may help us in minimizing dose dependent side effects without compromising efficacy.

Keywords: Combination; Hydrogen peroxide; Potash alum; Zones of inhibition

Introduction

Antimicrobial agents are extensively utilized in health care, industry and the environment to control and treat microbial infection [1, 2]. Combining antimicrobial agent may improve their activities by synergism and may provide an effective alternative to microbial resistance against single chemical [1, 3]. Additionally, a combination strategy is cost effective and time-saving which does not require long laboratory work to design, synthesize or discover new drugs more frequently. Thus, encouraging the use of drugs in combination for the management of bacterial infections.

Hydrogen peroxide (H_2O_2), our study compound, retains bactericidal efficacy which is being observed in H_2O_2 producing bacteria and phagocytic cells that inhibit other bacterial species and can kill invading microorganism, respectively [4]. The ability of hydrogen peroxide to produce hydroxyl radical contributes in the oxidation process of biomolecules. Hydrogen peroxide conversion to cytotoxic compounds is supported by peroxidases and by reducing agents [4]. Hydrogen peroxide along with produced free hydroxyl radicals may contribute in microbial management [1]. H_2O_2 is common disinfectant used to control gingival plaques [5], treating biofilms [6] and Fournier's gangrene [7]. It is also used for the sterilization of other surfaces [8] including industrial fish egg [9, 10] and fruits treatments [11].

Alum (ALM), our second study compound is a salt known as aluminium potassium sulfate ($KAl(SO_4)_2$) [12] and is suggested as a class 1 dynamic compound in mouthwashes [13], and used for the treatment of pediatric cough, hemorrhagic cystitis [14] ulcers, oral cavities and burns with anticarcinogenic effect [15], cosmetics [16], and domestic and industrial water treatments [17]. It has potential inhibitory effects on microbes [18].

In infection treatment, side effects associated with an overload of drugs in the human body and reduced microbial susceptibility to a single drug could be controlled by using them in combination even at lower concentrations. A synergistic or additive effect in mixtures is a key point in decreasing concentrations of drugs without compromising outcomes. Similarly, the synergy of hydrogen peroxide was found with many compounds including rifampicin [19], chlorhexidine [20], neucoproine [21], hypothiocyanite [22], sodium bicarbonate [23], iodine [24-26], different organic acids [27, 28], as well as with UV-irradiation [29]. Likewise, alum in combination with leaf extract of guava (*Psidium guajava*) has been utilized topically as an antiperspirant, an antibacterial and for astringent purposes [30]. However, alum and H_2O_2 incubation wise and concentration wise in combination against various microorganisms were not studied. Thus, in this study, we compared incubation-dependent and concentration-dependent inhibitory effects of alum and hydrogen peroxide alone and in mixtures against five clinical bacterial isolates.

Materials and methods

Culture media and chemicals

Nutrient agar (Neogen Company, UK), nutrient broth (Merck, Germany), and dimethyl sulfoxide (BIOCHEM) were used in experiments. Test compounds hydrogen peroxide (Sigma) and potash alum (purchased from the local market of Mirpur, AJK) were diluted with saline to acquire desired concentrations.

Strains and culture conditions

The clinical bacterial isolates *Enterococcus faecalis* (*E. faecalis*), *Enterococcus faecium* (*E. faecium*), *Escherichia coli* (*E. coli*), *Staphylococcus aureus* (*S. aureus*) and *Klebsiella pneumoniae* (*K. pneumoniae*) were taken from the Combined Military Hospital (CMH) Rawalpindi (Pakistan) for this study. Bacteria were sub-cultured on

nutrient agar plates and incubated at 37°C for 24hr in this experiment.

Anti-bacterial activity of alum and hydrogen peroxide separately

Fresh bacterial cultures were prepared by spreading 70µl bacterial suspension (0.060 OD at 600 wavelengths) on agar plate homogeneously. After short air dry (10min), 4 wells (6mm diameter) were made on each plate and loaded with 50µl of test compounds having 4.4mg/ml, 8.7mg/ml, 17.5mg/ml and 35mg/ml concentrations, respectively. Later, the plates were incubated at 37°C for 24hr, 48hr and 72hr. Finally, plates were photographed and zones of inhibition were measured in cm.

Anti-bacterial activity of alum and hydrogen peroxide in combination

In combination test, alum and hydrogen peroxide were used in 1:1 ratio with two-fold increasing concentration from well-1 to well-4, respectively. Briefly, well-1, well-2, well-3 and well-4 were loaded with 50µl of alum and hydrogen peroxide in 1:1 ratio from 4.4mg/ml, 8.7mg/ml, 17.5mg/ml and 35mg/ml concentrations for testing their synergistic effects. Later, the plates were incubated, photographed and zones of inhibition were measured the same as in the screening of alum and hydrogen peroxide separately.

Statistical Analysis

All experiments were performed thrice in triplicates. Student's t-test was applied to check significant differences using the lowest zone of inhibition (except zero) VS each tested zone of inhibition (^ap<0.05; ^bp<0.005, ^cp<0.0001).

Results and Discussion

The present study was carried out to evaluate the antibacterial activity of

hydrogen peroxide (H₂O₂) and potash alum (ALM) alone and in combination. For this, well diffusion method was used against five pathogenic bacterial isolates namely *E. coli*, *E. faecium*, *E. faecalis*, *S. aureus* and *K. pneumoniae*. Later, zones of inhibition were measured as a representative of bacterial susceptibility at different concentrations and different incubation periods at 24hr, 48hr and 72hr.

Individual antibacterial effect of alum and hydrogen peroxide

The hydrogen peroxide tested concentrations, 35mg/ml, 17.5mg/ml, 8.7mg/ml and 4.4mg/ml showed 0.783cm, 0.693cm, 0.581cm, 0.461cm zones of inhibition against *E. faecalis*, 0.872cm, 0.725cm, 0.612cm, 0.522cm zones of inhibition against *E. faecium*, 0.891cm, 0.712cm, 0.583cm, 0.421cm zones of the inhibition against *S. aureus*, 1.129cm, 0.976cm, 0.806cm, 0.647cm zones of inhibition against *E. coli* and 0.998cm, 0.821cm, 0.705cm, 0.579cm zones of inhibition against *K. pneumoniae*, respectively at 24 hr incubation (Table 1). Hydrogen peroxide is known as an effective disinfectant and it retains the ability to quickly kill microorganisms [31]. In addition to antimicrobial activity of hydrogen peroxide because of its oxidative properties it generates more powerful oxidant, called hydroxyl radical [32]. Explaining hydrogen peroxide's mode of action, it is reported that hydrogen peroxide exposure to logarithmically growing *E. coli* to kill by damaging DNA at low concentration whereas higher concentration (> 10 mM) may kill by damaging all macromolecules [33-36].

Table 1. Antibacterial susceptibility of selected bacteria against H₂O₂, ALM and control AZI at indicated concentrations and incubation periods. Student's t-test was applied to check significant where ^ap<0.05; ^bp<0.005, ^cp<0.0001

Tested compound	Concentration (mg/ml)				Concentration (mg/ml)				Concentration (mg/ml)				Test bacteria
	4.4	8.7	17.5	35	4.4	8.7	17.5	35	4.4	8.7	17.5	35	
	Zone of inhibition (cm) at 24 hr incubation				Zone of inhibition (cm) at 48 hr incubation				Zone of inhibition (cm) at 72 hr incubation				
AZI	0.00±0.00	0.09±0.02	0.28±0.01 _c	0.33±0.01 _c	0.00±0.00	0.09±0.02 _a	0.23±0.00 _b	0.29±0.03 ^c	0.00±0.00	0.13±0.02 _a	0.22±0.00 _b	0.29±0.01 _c	<i>E. faecalis</i>
H ₂ O ₂	0.46±0.01	0.58±0.01 _c	0.69±0.01 ^c	0.78±0.01 _c	0.46±0.01	0.58±0.01 _c	0.69±0.01 _c	0.78±0.01 ^c	0.45±0.01	0.53±0.02 _a	0.63±0.02 _c	0.71±0.02 _c	
ALU	0.01±0.00	0.09±0.02 _a	0.26±0.02 ^c	0.44±0.02 _c	0.01±0.01	0.10±0.02 _a	0.23±0.02 _c	0.40±0.02 ^c	0.01±0.00	0.09±0.03 _a	0.25±0.02 _c	0.43±0.02 _c	
AZI	0.00±0.00	0.11±0.02	0.20±0.02	0.33±0.03 _a	0.00±0.00	0.10±0.02	0.19±0.02	0.29±0.01 ^a	0.00±0.00	0.10±0.01	0.19±0.03	0.28±0.03 _a	<i>E. faecium</i>
H ₂ O ₂	0.52±0.01	0.61±0.01 _a	0.73±0.01 ^c	0.87±0.01 _c	0.52±0.01	0.61±0.01 _a	0.71±0.01 _c	0.86±0.00 ^c	0.51±0.01	0.60±0.01 _a	0.70±0.01 _c	0.82±0.01 _c	
ALU	0.00±0.00	0.02±0.02	0.18±0.02 ^c	0.42±0.02 _c	0.00±0.00	0.03±0.01	0.17±0.02	0.42±0.02 ^c	0.01±0.00	0.03±0.01	0.16±0.02 _c	0.39±0.02 _c	
AZI	0.00±0.00	0.10±0.02 _c	0.26±0.00 ^b	0.34±0.03 _c	0.00±0.00	0.09±0.01 _a	0.24±0.01 _a	0.31±0.03 ^c	0.00±0.00	0.09±0.01 _b	0.3±0.01 ^a	0.31±0.03 _b	<i>S. aureus</i>
H ₂ O ₂	0.42±0.01	0.58±0.01 _c	0.71±0.02 ^c	0.89±0.02 _c	0.42±0.01	0.58±0.01 _c	0.70±0.01 _c	0.87±0.01 _c	0.41±0.01	0.54±0.01 _a	0.66±0.02 _c	0.80±0.02 _c	
ALU	0.01±0.00	0.12±0.02 _a	0.23±0.02 ^c	0.46±0.02 _c	0.01±0.00	0.11±0.03 _a	0.23±0.02 _c	0.43±0.02 _c	0.01±0.01	0.09±0.02 _a	0.21±0.01 _c	0.42±0.02 _c	
AZI	1.09±0.03	1.29±0.02 _b	1.42±0.01 ^c	1.56±0.01 _c	1.02±0.02	1.24±0.02 _a	1.36±0.02 _b	1.52±0.01 _c	1.04±0.00	1.24±0.02 _a	1.38±0.03 _c	1.50±0.04 _c	<i>E. coli</i>
H ₂ O ₂	0.65±0.01	0.81±0.02 _b	0.98±0.02 ^c	1.13±0.02 _c	0.64±0.01	0.80±0.01 _b	0.96±0.01 _c	1.11±0.01 _c	0.64±0.01	0.79±0.01 _c	0.96±0.02 _c	1.11±0.02 _c	
ALU	0.01±0.01	0.13±0.03 _a	0.26±0.02 ^c	0.43±0.03 _c	0.01±0.00	0.11±0.02 _a	0.25±0.02 _c	0.42±0.02 _c	0.01±0.01	0.09±0.02 _a	0.21±0.02 _c	0.40±0.01 _c	
AZI	1.10±0.03	1.23±0.02 _a	1.38±0.02 ^c	1.49±0.02 _c	1.04±0.01	1.19±0.03 _a	1.33±0.01 _b	1.46±0.03 ^c	0.10±0.03	1.18±0.04 _a	1.30±0.02 _b	1.43±0.02 _b	<i>K. pneumoniae</i>
H ₂ O ₂	0.58±0.02	0.71±0.02 _c	0.82±0.01 ^c	1.00±0.02 _c	0.57±0.02	0.69±0.00 _c	0.81±0.01 _c	0.99±0.02 _c	0.42±0.01	0.55±0.02 _a	0.66±0.01 _c	0.81±0.03 _c	
ALU	0.03±0.00	0.11±0.02 _a	0.26±0.02 ^c	0.49±0.02 _c	0.02±0.00	0.1±0.03	0.29±0.02 _b	0.47±0.02 _c	0.01±0.00	0.07±0.02	0.21±0.03 _b	0.45±0.03 _c	

Similarly, alum concentrations, 35mg/ml, 17.5mg/ml, 8.7mg/ml and 4.4mg/ml showed 0.438cm, 0.263cm, 0.089cm, 0.005cm zones of inhibition against *E. faecalis*, 0.421cm, 0.184cm, 0.019cm, 0cm zones of inhibition against *E. faecium*, 0.455cm, 0.234cm, 0.122cm, 0.011cm zones of inhibition against *S. aureus*, 0.430cm, 0.258cm, 0.126cm, 0.013cm zones of inhibition against *E. coli*, and 0.485cm, 0.264cm, 0.106cm, 0.025cm zones of inhibition against *K. pneumoniae*, respectively at 24hr incubation (Table 1). These results concur with the previous findings which reported protective effect of the 100 PPM alum solution as less than 1000 PPM alum or more [37]. In 2014, Bnyan et al [38], additionally observed a paramount bactericidal effect of alum however, the mechanism of bactericidal effect of alum is not prominent [39]. Some postulations attribute the antibacterial effect of alum to reduction in acidity or deleterious effects on bacterial cell wall [40].

Thus, H₂O₂ bacterial susceptibility trend was *E. coli*>*K. pneumoniae*>*S. aureus*>*E. faecium*>*E. faecalis* and ALM susceptibility trend were as *K. pneumoniae*>*S. aureus*>*E. faecalis*>*E. coli*>*E. faecium* at highest tested concentration and 24hr of incubation period (Table 1). However, H₂O₂ showed more zone of inhibition than alum against all tested bacteria at all tested concentrations and incubation. Interestingly, zones were higher at lower incubation time which reduced later in the order of 24hr > 48hr > 72hr incubation times even at same H₂O₂ and ALM concentration (Table 1).

Similarly, Saranraj *et al.*, 2012, [31] reported hydrogen peroxide as an effective disinfectant but this effect was a short term which means hydrogen peroxide has no long-term or preserving effect. Similar, incubation effect was observed for azithromycin (AZI), used as a positive control as shown in (Table 1). However, both H₂O₂ and ALM showed more zones of inhibition at the highest tested concentration (35mg/ml) than AZI against *E. faecalis*, *E. faecium* and *S. aureus* at 24hr, 48hr and 72hr of incubation.

Combined antibacterial effect of alum and hydrogen peroxide

To check combination effect, alum and hydrogen peroxide were used in 1:1 ratio with two-fold increase in concentration from well-1 to well-4 in well diffusion method, respectively, as explained in method section. The combination of alum and hydrogen peroxide in 1:1 ratio forming 4.4mg/ml, 8.7mg/ml, 17.5mg/ml and 35mg/ml total concentrations showed 0.508cm, 0.635cm, 0.779cm and 1.052cm zones of inhibition against *E. faecalis*, 0.497cm, 0.656cm, 0.802cm and 0.979cm zones of inhibition against *E. faecium*, 0.406cm, 0.552cm, 0.672cm and 0.843cm zones of inhibition against *S. aureus*, 0.414cm, 0.547cm, 0.735cm and 0.870cm zones of inhibition against *E. coli* and, 0.397cm, 0.575cm, 0.770cm and 0.968cm zones of inhibition against *K. pneumoniae*, respectively at 24hr of inhibition (Table 2). In combination test, zones were higher at lower incubation time which reduced later in order of 24hr > 48hr > 72hr incubation times same as H₂O₂ and ALM in individual test.

Table 2. Antibacterial susceptibility of selected bacteria against H₂O₂ and ALM in combination (1:1) at indicated concentrations and incubation periods. Student's t-test was applied to check significance where ^ap<0.05; ^bp<0.005, ^cp<0.0001

Combination	Concentration (mg/ml)				Concentration (mg/ml)				Concentration (mg/ml)				Test Bacteria
	4.4	8.7	17.5	35	4.4	8.7	17.5	35	4.4	8.7	17.5	35	
	Zone of inhibition (cm) at 24 hr incubation				Zone of inhibition (cm) at 48 hr incubation				Zone of inhibition (cm) at 72 hr incubation				
H ₂ O ₂ + ALM	0.51±0.02	0.64±0.01 ^a	0.78±0.01 ^c	1.05±0.02 _c	0.46±0.02	0.57±0.02 _a	0.68±0.02 _c	0.92±0.00 _c	0.45±0.05	0.53±0.02	0.64±0.02 _a	0.88±0.02 _c	<i>E. faecalis</i>
H ₂ O ₂ + ALM	0.50±0.05	0.66±0.06 ^a	0.81±0.03 ^b	0.98±0.04 _c	0.48±0.03	0.63±0.06 _a	0.75±0.04 _c	0.91±0.02 _c	0.48±0.04	0.62±0.04 _a	0.73±0.03 _c	0.88±0.03 _c	<i>E. faecium</i>
H ₂ O ₂ + ALM	0.41±0.02	0.55±0.03 ^a	0.67±0.02 ^b	0.84±0.02 _c	0.38±0.03	0.51±0.03	0.61±0.01 _b	0.78±0.01 _c	0.39±0.02	0.51±0.02	0.60±0.02 _a	0.76±0.01 _c	<i>S. aureus</i>
H ₂ O ₂ + ALM	0.41±0.01	0.55±0.01	0.74±0.01 ^c	0.87±0.03 _c	0.37±0.01	0.49±0.01	0.66±0.01 _{2c}	0.79±0.02 _c	0.39±0.01	0.53±0.01 _a	0.68±0.01 _c	0.78±0.03 _c	<i>E. coli</i>
H ₂ O ₂ + ALM	0.40±0.03	0.58±0.04 _c	0.77±0.02 ^c	0.97±0.02 _c	0.37±0.04	0.50±0.04 _a	0.69±0.02 _c	0.86±0.01 _{1c}	0.36±0.02	0.50±0.03 _a	0.69±0.03 _c	0.83±0.01 _c	<i>K. pneumonia</i>

Interestingly, H₂O₂ and ALM in combination showed more zones of inhibition at the highest tested concentration (35mg/ml) than AZI against *E. faecalis*, *E. faecium* and *S. aureus* at 24hr, 48hr and 72hr of incubation (Table 1 & 2). Moreover, H₂O₂ and ALM in the combination showed more zone of inhibition than H₂O₂ alone against *E. faecalis* and *E. faecium* and, also more effect than ALM alone against all tested bacteria i.e *E. faecalis*, *E. faecium*, *S. aureus*, *E. coli* and *K. pneumoniae* at all incubations (Table 1 & 2), summarized in (Table 3). Amadi *et al.* [30] found similarly increased antibacterial impacts of alum with guava leaf extract in combination. Another study reported that the combination of alum with antibiotics (tetracycline and cefotaxime) was more effective than antibiotic combination and

the combination effect was more efficient than an individual effect [40]. Similarly, hydrogen peroxide in combination with lactoperoxidase and thiocyanate is reported much more effective than hydrogen peroxide alone as an inhibitor of bacterial metabolism and growth [41]. Thus, the combination strategy provides us with an opportunity to improve their efficacy at low concentration. Therefore, side effects associated with a higher load of drugs can be reduced without compromising their efficacy [42]. Due to these advantages, combination therapies have become a standard in many areas including cancer treatment [43], hypertension [44], asthma [45], and AIDS [46, 47] which open a new horizon for research and new hope for disease control including bacterial infections in future.

Table 3. Comparison of antibacterial effect at highest tested concentration (35 mg/ml) of test compounds and their combinations against selected bacteria

Bacteria	Susceptibility trend against tested compounds
<i>E. faecalis</i>	H ₂ O ₂ +ALU > H ₂ O ₂ > ALU > AZI
<i>E. faecium</i>	H ₂ O ₂ +ALU > H ₂ O ₂ > ALU > AZI
<i>S. aureus</i>	H ₂ O ₂ > H ₂ O ₂ +ALU > ALU > AZI
<i>E. coli</i>	AZI > H ₂ O ₂ > H ₂ O ₂ +ALU > ALU
<i>K. pneumoniae</i>	AZI > H ₂ O ₂ > H ₂ O ₂ +ALU > ALU

Conclusion

This study found that the increasing concentration of H₂O₂ and ALM have increasing antibacterial effects on tested bacteria which reduced with time. Moreover, H₂O₂ and ALM in combination showed increased zone inhibition than ALM alone (against all bacteria) and H₂O₂ alone (against *E. faecalis* and *E. faecium*) without increasing final concentration. Thus, a combination strategy might help us in minimizing dose-dependent side effects of drugs without compromising effectiveness.

Authors' contributions

Conceived and designed the experiments: A Ali & H Khurshid, Performed the experiments: H Khurshid & A Ali, Analyzed the data: A Ali & B Akbar,

Contributed materials/ analysis/ tools: M Rafiq, F Nazir & M Ahmed, Wrote the paper: I Ali & M Ahmed.

Acknowledgement

This work was financially supported by Mirpur University of Science & Technology (MUST), AJK, Pakistan. The authors are thankful to Combined Military Hospital (CMH) Rawalpindi (Pakistan) for providing us bacterial isolates for this study. The authors declare no financial or any other conflict of interest.

References

- McDonnell G & Russell AD (1999). Antiseptics and disinfectants: activity, action, and resistance. *Clin Microbiol Rev* 12(1): 147–179.
- Bowler PG, Duerden BI & Armstrong DG (2001). Wound microbiology and

- associated approaches to wound management. *Clin Microbiol Rev* 14(2): 244–269.
3. Foucquier J & Guedj M (2015). Analysis of drug combinations: current methodological landscape. *Pharma Res Per* 3(3): e00149.
4. Juven JB & Pierson MD (1996). Antibacterial effects of hydrogen peroxide and methods for its detection and quantitation. *J of Food Prot* 59(11): 1233–1241.
5. Prabhu R, Kohale B, Agrawal AA, Wagle SV, Bhartiya G & Chaudhari D (2017). A comparative clinical study to evaluate the effect of 1.5% hydrogen peroxide mouthwash as an adjunct to 0.2% chlorhexidine mouthwash to reduce dental stains and plaque formation. *IJCMR* 4(10): 2181–2184.
6. Presterl E, Suchomel M, Eder M, Reichmann S, Lassnigg A, Graninger W & Rotter M (2007). Effects of alcohols, povidone-iodine and hydrogen peroxide on biofilms of *Staphylococcus epidermidis*. *J Antimicrob Chemother* 60(2): 417–420.
7. Atakan IH, Kaplan M, Kaya E, Aktoz T & Inci O (2002). A life-threatening infection: Fournier's gangrene. *Int Urol Nephrol* 34(3): 387–392.
8. Malik DJ, Shaw CM, Rielly CD & Shama G (2013). The inactivation of *Bacillus subtilis* spores at low concentrations of hydrogen peroxide vapour. *J Food Eng* 114(3): 391–396.
9. Arndt RE, Wagner EJ & Routledge MD (2001). Reducing or withholding hydrogen peroxide treatment during a critical stage of rainbow trout development: effects on eyed eggs, hatch, deformities, and fungal control. *North Am J Aquaculture* 63(2): 161–166.
10. Wagner EJ, Oplinger RW & Bartley RWM (2012). Laboratory and production scale disinfection of salmonid eggs with hydrogen peroxid. *North Am J Aquaculture* 74(1): 92–99.
11. Ukuku DO, Bari ML, Kawamoto S & Isshiki K (2005). Use of hydrogen peroxide in combination with nisin, sodium lactate and citric acid for reducing transfer of bacterial pathogens from whole melon surfaces to fresh-cut pieces. *Int J Food Microbiol* 104(2): 225–233.
12. Chang R (2005). *Chemistry*. 8th edition. McGraw-Hill Companies. Inc New York USA.
13. Olmez A, Can H, Ayhan H & Olur H (1998). Effect of an alum-containing mouthrinse in children for plaque and salivary levels of selected oral microflora. *J Clin Pediatr Dent* 22(4): 335–440.
14. Bestoon MF (2012). Evidence of feasibility of aluminum potassium sulphate (alum) solution as a root canal irrigant. *J Bagh College Dentistry* 24(1): 1–5.
15. Mourugan K & Suryakanth MP (2004). Evaluation of an alum-containing mouthrinse for inhibition of salivary *Streptococcus mutans* levels in children- A controlled clinical trial. *J Indian Soc Ped Prev Dent* 22(3): 100–105.
16. Alzormor AK, Moharram AS & AlAbsi NM (2014). Formulation and evaluation of potash alum as deodorant lotion and after shaving astringent as cream and gel. *Int Curr Pharm J* 3(2): 228–233.
17. Potter NN & Hotchkiss JH (2007). *Food Science*. 5th edition. CBS Publishers and Distributors, Daryaganji, New Delhi India.
18. Ahmed KT (2011). Inhibition of swarming in *Proteus mirabilis* by alum (hydrated aluminum potassium sulphate). *J University of Anbar for Pure Science* 5(2): 20–24.
19. Humphrey TJ (1990). The synergistic inhibition of *Campylobacter jejuni* by rifampicin and hydrogen peroxide. *Lett Appl Microbiol* 10(2): 97–100.

20. Steinberg D, Heling I, Daniel I & Ginsburg I (1999). Antibacterial synergistic effect of chlorhexidine and hydrogen peroxide against *Streptococcus sobrinus* *Streptococcus faecalis* and *Staphylococcus aureus*. *J Oral Rehabil* 26(2): 151–156.
21. Almeida CEB, Felicio DL, Galhardo RS, Cabral-Neto JB & Leita AC (1999). Synergistic lethal effect between hydrogen peroxide and neocuproine (2,9-dimethyl 1,10-phenanthroline) in *Escherichia coli*. *Mutat Res DNA Repair* 433(1): 59–66.
22. Carlsson J, Edlund MB & Hanstrom L (1984). Bactericidal and cytotoxic effects of hypothiocyante-hydrogen peroxide mixtures. *Infect Immun* 44(3): 581–586.
23. Miyasaki KT, Genco RJ & Wilson ME (1986). Antimicrobial properties of hydrogen peroxide and sodium bicarbonate individually and in combination against selected oral, gram-negative, facultative bacteria. *J Dent Res* 65(9): 1142–1148.
24. Voboril R & Weberova J (2004). Successful treatment of infected vascular prosthetic grafts in the groin using conservative therapy with povidoneiodine solution. *Ann Vasc Surg* 18(3): 372–375.
25. Ulivieri S, Toninelli S, Petrini C, Giorgio A & Oliveri G (2011). Prevention of postoperative infections in spine surgery by wound irrigation with a solution of povidone-iodine and hydrogen peroxide. *Arch Orthop Trauma Surg* 131(9): 1203–1206.
26. SM Ng & Wieckowski S (1989). Stable hydrogen peroxide dental gel, US Patent-4839156 Colgate-Palmolive Company. 1199.
27. Li Y & Wu C (2013). Enhanced inactivation of *Salmonella Typhimurium* from blueberries by combinations of sodium dodecyl sulfate with organic acids or hydrogen peroxide. *Food Res Int* 54(2): 1553–1559.
28. Martin H & Maris P (2012). Synergism between hydrogen peroxide and seventeen acids against six bacterial strains. *J Appl Microbiol* 113(3): 578–590.
29. Shama G (1992). Inactivation of *Escherichia coli* by ultraviolet light and hydrogen peroxide in a thin film contactor. *Lett Appl Microbiol* 15(6): 259–260.
30. Amadi LO, Wanabia D & Amadi V (2016). Combination effects of alum and guava (*Psidium guajava*) leaf extracts on some pathogens from clinical samples. *Int J Curr Res* 8(5): 31354–31358.
31. Cadnum JL, Mana TS, Jencson A, Thota P, Kundrapu S, Donskey CJ (2015). Effectiveness of a hydrogen peroxide spray for decontamination of soft surfaces in hospitals. *Am J Infect Control* 43(12):1357–1359.
32. Juven BJ & Pierson MD (1996). Antibacterial effects of hydrogen peroxide and methods for its detection and quantitation. *J Food Prot* 59(11): 1233–1241.
33. Uhl L, Gerstel A, Chabalier M & Dukan S (2015). Hydrogen peroxide induced cell death: One or two modes of action? *Heliyon* 1(4): e00049.
34. Imlay JA (2003). Pathways of oxidative damage. *Annu Rev Microbiol* 57: 395–418.
35. Imlay JA & Linn S (1986). Bimodal pattern of killing of DNA-repair-defective or anoxically grown *Escherichia coli* by hydrogen peroxide. *J Bacteriol* 166(2): 519–527.
36. Imlay JA & Linn S (1988). DNA damage and oxygen radical toxicity. *Science* 240(4857): 1302–1309.
37. Putt MS & Kleber CJ (1995). Effect of aluminum concentration on dental caries formation in the rat. *Oral Dis* 1(2): 80–5.

38. Bnyan IA, Alta'ee AH & Kadhum NH (2014). Antibacterial activity of aluminum potassium sulfate and syzygium aromaticum extract against pathogenic microorganisms. *J Nat Sci Res* 4(15): 11-4.
39. Arul PF, Babu GD, Lavanya M, Vidhya KS & Devasena T (2011). Toxicity studies of aluminium oxide nanoparticles in cell lines. *Int J Nanotechnol Appl* 5(2): 99-107.
40. Ali ZM (2018). Synergistic antibacterial interaction between an alum and antibiotics on some microorganism. *Sci J Med Res* 2(5): 47-51.
41. Thomas EL, Milligan TW, Joyner RE & Jefferson MM (1994). Antibacterial activity of hydrogen peroxide and the lactoperoxidase hydrogen peroxide-thiocyanate system against oral streptococci. *Infect Immun* 62(2): 529-535.
42. Danesh A, Ljungh Å, Mattiasson B & Mamo G (2016). Synergistic effect of haloduracin and chloramphenicol against clinically important gram-positive bacteria. *Biotechnol Rep* 13: 37-41.
43. Humphrey RW, Brockway LM, Bonk DT, Dohoney KM, Doroshow JH & Meech SJ (2011). Opportunities and challenges in the development of experimental drug combinations for cancer. *J Natl Cancer Inst* 103(16): 1222-1226.
44. Glass G (2004). Cardiovascular combinations. *Nat Rev Drug Discov* 3(9): 731-732.
45. Nelson HS (2001). Combination treatment with fluticasone propionate/salmeterol in the treatment of asthma, *J Allergy Clin Immunol* 107(2): 398-416.
46. Larder BA, Kemp SD & Harrigan PR (1995). Potential mechanism for sustained antiretroviral efficacy of AZT-3TC combination therapy. *Science* 269(5224): 696-699.
47. Oversteegen L, Shah M & Rovini H (2007). HIV combination products. *Nat Rev Drug Discov* 6(12): 951-959.