Review Article

Elemental analysis of medicinal plants and their impacts on human health

Abdul Sattar^{1,2}, Abdul Baqi^{1,3}, Muhammad Suleman⁴, Muhammad Tahir Shah^{1,4}, Samiullah^{1*}, Naqeebullah Khan¹ and Attiq-Ur-Rehman¹

1. Department of Chemistry, University of Balochistan, Quetta 87300-Pakistan

2. National Vocational & Technical Training Commission (NAVTTTC), Regional Office Balochistan, Quetta 87300-Pakistan

3. Colleges Higher and Technical Education Balochistan, Quetta 87300-Pakistan

4. Balochistan Residential College Loralai, Balochistan-Pakistan

*Corresponding author's email: sami435889@gmail.com

Citation

Abdul Sattar, Abdul Baqi, Muhammad Suleman, Muhammad Tahir Shah, Samiullah, Naqeebullah Khan and Attiq-Ur-Rehman. Elemental analysis of medicinal plants and their impacts on human health. Pure and Applied Biology. Vol. 9, Issue 1, pp249-255. <http://dx.doi.org/10.19045/bspab.2020.90029>

Received: 17/07/2019

Revised: 01/10/2019

Accepted: 05/10/2019

Online First: 23/10/2019

Abstract

The medicinal plants are the origin of biologically active elements, which are essential for both plants and animals. Trace elements help various biochemical mechanisms in the body of living organisms. The major reason for numerous medical issues in the human being is either a deficiency or toxicity of essential elements. Therefore, investigating the distinct biodynamic elements of medicinal plants and their evaluation is very much important. The therapeutic plants have been utilized in traditional drugs for the treatment of ailments, such as hypertension, minimizing cholesterol level, malignancy, fever, leprosy and eye problems. The most abundant necessary elements found in such plants are Magnesium, Manganese, Copper, Selenium, Iron, Calcium, Sodium and Zinc which are known for their capability to treat different diseases. This review generally discusses the information related to the elemental composition of medicinal plants and their impacts on the living body. The outcomes increase the values of medicinal plants for the preparation of conventional drugs and pharmaceutical factories. This review explores the requirement of further research on therapeutic plants to prepare more effective medicines as different plants have variable values of elements.

Keywords: Essential elements; Human health; Medicinal plants; Prescription; Treatment

Introduction

Background

Therapeutic plants are the most primitive health care items utilized for the treatment of different diseases. They are the wealthy reservoir of antioxidants and other necessary elements [1]. Remedial herbs have reliably been esteemed as a method for the treatment of several diseases in the old era and have imparted a vital role for the preparation of recently advanced drugs with the addition of some chemical constituents. The effectiveness of

medicinal plants for the intention of cure is usually assumed regarding their natural parts like fundamental oils, nutrients, glycosides, vitamins and different components of the living body. It is a known fact that larger quantity or continued intake of restorative plants may prompt a higher collection of various minerals that result in different fitness issues [2]. Necessary metals can likewise deliver poisonous impacts when ingested in larger amounts although non-essential trace metals are lethal even in reduced amount

for human health [3-5]. The quantity of trace minerals in therapeutic herbs on the far side admissible extent is larger anxiety for the people everywhere throughout the world [6]. The shortage of essential elements also results in various disorders [7].

Herbs may assimilate heavy metals from earth, atmosphere or water. The magnitude of necessary minerals in medicinal plants differs from species to species and composition of soil [8]. However, conventional homegrown prescription has been broadly utilized in the progressing as well as in the developed world as an alternate source. Information about the essential elements in curative plants is very important along with organic constituents. Many necessary minerals have important roles in the production of dynamic particles that play a vital function in the treatment of various diseases [9, 10]. Elements play an important role in the medicinal values of plants and treatment and in health and ailment. Likewise, nutrients and biochemical's such as carbohydrates, fats, proteins also play a vital part in satisfying human needs for energy and life processes [11]. Herbs are the vital sources of numerous medications like emetics, hostile to malignancy, antispasmodics, anti-microbes and so on. Therapeutic plants have anti-toxin characteristics and extensively used by the people living in far-off areas around the world [12].

Various scientists exposed the values of trace elements that enhanced the importance of medicinal plants [13]. Rare minerals are the components that are present in fewer amounts but are necessary for the body. It is proved from the latest knowledge that a sufficient measure of trace elements is needed in the maintenance of metabolism and activities of biocatalysts [14, 15]. The familiarity with essential elements is necessary to explain the importance of medicinal herbs to cure various disorders through their medicinal activities and to spread knowledge about the treatment of therapeutic herbs [16].

Flora of Pakistan is rich in medicinal plants, which are therapeutically used. More than 1,000 species possess medicinal properties, which are used by local communities to cure different diseases [17].

The medicinal plants are ordinarily applied because of their easy approach and fewer reactions. According to World Health Organization (WHO 2002), 80% of the people primarily use medicinal plants for their treatment and it is necessary to examine such herbs for their effectiveness, security, and efficiency [18]. The conventional drugs bring improvement in the immune system. Furthermore, they raise chemotherapy that demolishes a wide range of malignant growth [19]. So, various trace elements have been considered as the essential components of the food that repair insulin metabolic process in forbearing of diabetes [20].

Several analysts have likewise reported that necessary minerals usefully influence the complexities of diabetes mellitus [21].

Impacts of elements on human health

Importance of some of the elements found in medicinal plants are given as:

Copper and human health

Copper is found in several parts of human body parts and various enzymes and is thought to be a significant necessary trace element. However, too much ingestion of copper results in hepatic destruction and Wilson's disease. Wilson's disease is caused due to the addition of free copper in the liver, kidney, and brain [22]. The shortage of Cu has bad fitness issues, including aneurysms, harm the circulatory system, hernias, hemorrhages and may affect the passage of nutrients through cell walls [23].

Copper is necessary to the whole living creatures as a trace nutritional constituent as it is a main component of the respirational enzyme systems. Cu is a part of the bloodstain hemocyanin that is substituted by hemoglobin complexed with iron. However, copper is an important trace element required in a small amount for the suitable working of tissues and

organizations but the higher quantity may be poisonous and fatal [24]. The site of storage of copper in the human body is muscles, bones and liver. Cu mixes are utilized as a bacteriostatic agent, fungi killer and preservation of wood [25]. The shortage of copper in humans occurs very rarely as it is commonly found in ordinary food items such as legumes, nuts, grains, etc. Copper is an obligatory trace element in some vegetations and animals but not for all [26]. Scarcity of copper results in anomalies of bones, reduced development, defects in the metabolism of sugar and fats. Copper assistances the body for the utilization of iron. It plays a vital role in the nervous system, development of bones, maintenance of glucose and protects the plasma membrane from the effects of free radicals. A lot of cardiac problems and blood irregularities may be caused due to the insufficiency of copper [27, 28].

Iron and human health

Iron is an essential element, which is the main constituent of hemoglobin and is present in numerous human enzymes. Its concentration in red blood cells and muscle tissue is very high. Larger intake of iron results in hepatotoxicity. Severe hepatic failure has been observed due to overdose of ferrous sulphate in an earlier age of adulthood [29]. Anemia, a nutritional disease that is caused due to a shortage of iron [30].

Iron plays a significant role in oxygen and electron transference in the human body. Its deficit reasons in gastric issues, bleeding from the nose and cardiovascular diseases [31]. Mixes of iron and protein have a dynamic function in the metabolic process of all living things [32].

Manganese and human health

Manganese is an essential element that is the main constituent of metalloenzymes, which plays an important role in the oxidation of fats and cholesterol [33]. Shortage of manganese may result in bleeding issues but high quantity causes talking problems, limb spasms, and encephalitis [34].

Manganese is necessary for the normal working of reproduction and the central nervous system. Its insufficiency results in heart problems and bone issues in youngsters and young [35].

It has been investigated that Manganese is tangled in normal immune activities, maintenance of blood glucose and cellular vitality and the resistance against free radicals [36]. It stimulates specific enzymes, which show vital roles in the breakdown of starches, protein, and cholesterol. It is also necessary for regular insulin production, its discharge and a variation in its metabolic rate has been involved in diabetes growth [37]. Shortage of manganese can cause weakened glucose forbearance, transformed starch and fat metabolism, decreased insulin emission and skeletal irregularities [38]. Mn is a very significant element while maintaining the glucose of body fluid. It has strong estrogenic characteristics and it is very important for the treatment of menopausal indications, bone issues and post-delivery depression. Insufficient Manganese consumption has been linked with a parenteral diet, causing dermatitis, alters hair coloration and decelerated hair growing. Steady displacement of joints (mainly knee junctions) is also related to a deficiency of Manganese. Similarly, an excess amount of Mn causes rupture of tendon/ligament and pneumonia. Moreover, damage to sex drive and sperm destruction has been investigated in males. Excessive manganese affects the assimilation of dietetic iron. It has outstanding antioxidant characters which are important during oxidative pressure. For example, Manganese superoxide dismutase is the prime antioxidant enzyme in mitochondria that takes almost totally the oxygen used by cells. The investigation has exposed that numerous Mn-activated enzymes play significant functions in the breakdown of carbohydrates, proteins and fats [39]. Mn is the best co-factor of enzymes (glycosyl transferases) needed for the production of proteoglycans necessary

for the growth of solid bones and cartilage. It is also accountable for the excretion of prolinase which is responsible for amino acid, the main constituent of collagen manufacture and repairing of injuries that use closely all the oxygen consumed by cells [40].

Zinc and human health

Zinc is a necessary element, and cellular Zn stimulates the homeostatic mechanism to evade the storage of extra zinc. Excess zinc causes a shortage of Cu and cell apoptosis. Moreover, zinc deficit has been connected to a repressed immune system [41]. Zn and Cu both are considered significant for breakdown sugar and depressing cholesterol [42].

The part of Zn in the creation of insulin and catalysis of several enzymes catalytic reactions has been investigated [43]. The low level of Zn in plasma unfavorably distresses the capability of islet cells to secrete and produce insulin [44]. Furthermore, it may exaggerate the insulin confrontation in type 2 diabetes and so reasons further problems [45].

Zinc is a nutritional constituent for humans and animals. The human body has about two gm to three gm. It has an important function in DNA duplication, prenatal period, adolescence and menopause. The greater amount of Zinc protects people from cadmium harming although the side effects of Zn over dosage include vomiting, dizziness, sicknesses and motions [46].

Calcium and human health

Inadequacy can prompt low bone mass (osteopenia), bone breaks, deadness and shivering in the fingers and unusual heart rhythms. Some of the plants show the satisfactory dimension of Ca collection [47, 48].

Chromium and human health

It maintains insulin and blood sugar levels by activating insulin signing and metabolic path and consequently it might improve insulin affectability. The balance of lipid digestion by Cr in marginal tissues may signify an extra novel system of activity [49]. Insufficiency of Cr or its biodynamic

type has been entangled in the pathogenesis of insulin obstruction and diabetes [50].

Magnesium and human health

Magnesium is a cofactor of numerous enzymes in the starch oxidative breakdown and has a vital function in glucose carrying process of the plasma membrane. It has also a role in insulin production, combination and working [51]. It takes part in the phosphorylation reactions of sugar and its breakdown and it may affect the secretion and function of the hormones that aid to regulate blood sugar stages [52]. It was also investigated that a lack of Magnesium might reduce insulin facilitated glucose take in and have been linked with the increase of insulin conflict [53].

Mg is broadly dispersed in animals and plants, diets and drinks. Green leafy herbs, for example, spinach, beans, nuts, seeds and whole grains are good sources. Magnesium, a plentiful element in the body, is part of various foods items, added to further nutrition, available as a dietetic component and present in drugs [54, 55]. A sufficient dose of Mg from food does not cause a health problem in fit people since the kidneys remove extra amounts in the urine. Though, high dosages of Mg from diet or medicines usually cause diarrhea, nausea and stomach hampering [56].

Analytical Techniques

The most extensively and generally used methods of elemental investigation, providing standard ranks of accuracy and precision, comprise flame atomic absorption spectrometry (FAAS), UV/Vis spectrophotometer, flame photometry, graphite furnace atomic absorption spectrometry (GFAAS), inductively coupled plasma atomic emission spectrometry (ICP-AES), and inductively coupled plasma mass spectrometry (ICP-MS) [57].

Conclusion

It is concluded from the above review/study to examine the specific quantity of those elements from their permissible limits that were set up by the world health organization (WHO). If the amount of these elements in medicinal plants is high than the standard level, it may cause severe

health-related issues, particularly in human beings. The majority of the general population believes that herbal prescriptions are safe and nontoxic. Even current chemotherapeutics are unaware of metals available in therapeutic plants. Moreover, the accumulation of toxic metals in the soil causes different types of contaminations that transfer through roots and arial parts of plants by the food chain.

Therefore, it is necessary to investigate the medicinal plants properly before the manufacturing of the medicines and further, such kind of investigation on the plant will help in choosing different dynamic constituents and in dealing with the helping of a specific formulation.

Authors' contributions

Conceived and designed the experiments: A Sattar, Attiq-Ur-Rehman & A Baqi Performed the experiments: A Sattar, Samiullah & A Baqi, Analyzed the data: Samiullah, N Khan & M Suleman, Contributed materials/ analysis/ tools: Samiullah, N Khan & Muhammad Tahir Shah, Wrote the paper: A Sattar, A Baqi & Samiullah.

References

- Smith-Hall C, Larsen HO & Pouliot M (2012). People, plants and health: a conceptual framework for assessing changes in medicinal plant consumption. *J of Ethno Biol and Ethno Medi* 8(1): 43.
- Sharma RK, Agrawal M & Marshall FM (2009). Heavy metals in vegetables collected from production and market sites of a tropical urban area of India. *Food and Chem Toxicol* 47(3): 583-591.
- Sheded MG, Pulford ID & Hamed AI (2006). Presence of major and trace elements in seven medicinal plants growing in the South-Eastern Desert, Egypt. *J of Arid Environ* 66(2): 210-217.
- Koc H & Sari H (2009). Trace metal contents of some medicinal, aromatic plants and soil samples in the Mediterranean region, Turkey. *J of Appl Chem Res* 8(4) 52-57.
- Basgel S & Erdemoglu SB (2006). Determination of mineral and trace elements in some medicinal herbs and their infusions consumed in Turkey. *Sci of the Total Environ* 359(1-3): 82-89.
- Van Wyk BE & Wink M (2018). Medicinal plants of the world. CABI.
- Hashmi DR, Ismail S & Shaikh GH (2007). Assessment of the level of trace metals in commonly edible vegetables locally available in the markets of Karachi city. *Pak J of Bot* 39(3): 747-751.
- Khan SA, Khan L, Hussain I, Marwat KB & Akhtar N (2008). Profile of heavy metals in selected medicinal plants. *Pak J of Weed Sci Res* 14(1-2): 101-110.
- Yamashita CI, Saiki M, Vasconcellos MBA, & Sertie JAA (2005). Characterization of trace elements in Casearia medicinal plant by neutron activation analysis. *Appl Radiation and Isotopes* 63(5-6): 841-846.
- Okasha TA (2019). Major trends of Psychosomatic Medicine in North Africa and Middle East. In Global Psychosomatic Medicine and Consultation-Liaison Psychiatry. *Springer, Cham* pp 147-161.
- Adnan M, Hussain J, Tahir Shah M, Shinwari Z K, Ullah F, Bahader A & Watanabe T (2010). Proximate and nutrient composition of medicinal plants of humid and sub-humid regions in North-west Pakistan. *J of Medi Plants Res* 4(4): 339-345.
- Prashant T, Bimlesh K, Mandeep K, Gurpreet K & Harleen K (2011). A review on phytochemical screening and extraction in medicinal plants. *Inter Pharmac Scie I*(1): 98-106.
- Milani RF, Silvestre LK, Morgano MA & Cadore S (2019). Investigation of twelve trace elements in herbal tea commercialized in Brazil. *J of Trace Elements in Medi and Biol* 52: 111-117.
- Abdul Sattar S, Seetharami Reddy B, Koteswara Rao V, Ramnarayana K, Naga Raju GJ, Bhuloka Reddy S & Madhusudhana Rao PV (2012). Estimation of trace elements in some medicinal plants used in anti-cancer drugs by PIXE. *X-Ray Spectrom* 41(2): 111-116.
- Singh NK, Devi CB, Singh TS & Singh NR (2010). Trace elements of some selected medicinal plants of Manipur. *Indian J of Natural Products and Reso* 1(2): 227-231.
- Anal JMH & Chase P (2016). Trace elements analysis in some medicinal plants using graphite furnace-atomic absorption spectroscopy. *Environ Engi Res* 21(3): 247-255.

17. Hussain J, Khan FU, Ullah R, Muhammad Z, Rehman N, Shinwari ZK & Hussain SM (2011). Nutrient evaluation and elemental analysis of four selected medicinal plants of Khyber Pakhtoonkhwa, Pakistan. *Pak J of Bot* 43(1): 427-434.
18. Saeed M, Muhammad N & Khan H (2011). Assessment of heavy metal content of branded Pakistani herbal products. *Trop J of Pharma Res* 10(4): 499-506.
19. Muhammed A (2015). A personal discussion held in May 2015 with a horticulturist and traditional herbalist, staying in Kano City.
20. Clark TA, Deniset JF, Heyliger CE & Pierce GN (2014). Alternative therapies for diabetes and its cardiac complications: role of vanadium. *Heart Failure Rev* 19(1): 123-132.
21. Dey P & Khaled KL (2015). Quantitative Estimation of Some Essential Minerals of *Gymnema Sylvestre* as a Potential Herb in Counteracting Complications of Diabetes. *Inter J of Res Stud in Biosci* 3(1): 71-74.
22. NIH National Institutes of Health. *Trace Elements and Metals*. June 2017. <https://livertox.nih.gov/TraceElementsAndMetals.htm>.
23. Araya M, Pizarro F, Olivares M, Arredondo M, Gonzalez M, & Méndez M (2006). Understanding copper homeostasis in humans and copper effects on health. *Biol Res* 39(1): 183-187.
24. Aweto AO & Enaruvbe GO (2010). Catenary variation of soil properties under oil palm plantation in south western Nigeria. *Ethiopian J of Environ stud and Manag* 3(1).
25. Fry RS, Ashwell MS, Lloyd KE, O'Nan AT, Flowers WL, Stewart KR & Spears JW (2012). Amount and source of dietary copper affect small intestine morphology, duodenal lipid peroxidation, hepatic oxidative stress, and mRNA expression of hepatic copper regulatory proteins in weanling pigs. *J of Ani Sci* 90(9): 3112-3119.
26. Zhang P & Gao H (1999). Determination of copper in waste water by beta-correction new spectrophotometry. *Guang pu xue yu guang pu fen xi= Guang pu* 19(3): 471-473.
27. Liang Q & Zhou B (2007). Copper and manganese induce yeast apoptosis via different pathways. *Mol Biol of the Cell* 18(12): 4741-4749.
28. Lawrence AL, Fox J & Castille JrFL (1981). Decreased toxicity of copper and manganese ions to shrimp nauplii (*Penaeus stylirostris* Stimpson) in the presence of EDTA. *J of the World Mariculture Soc* 12(1): 271-280.
29. Daram SR & Hayashi PH (2005). Acute liver failure due to iron overdose in an adult. *The Southern Med J* 98(2): 241-245.
30. Fu ZH, Xie MY, Zhang ZM & Guo L (2004). Determination of inorganic elements in *Plantago* by ICP-AES. *Guang pu xue yu guang pu fen xi= Guang pu* 24(6): 737-740.
31. Ullah R, Khader JA, Hussain I, Talha NMA & Khan N (2012). Investigation of macro and micro-nutrients in selected medicinal plants. *African J of Phar and Pharmacol* 6(25): 1829-1832.
32. Swaminathan S, Fonseca VA, Alam MG & Shah SV (2007). The role of iron in diabetes and its complications. *Diabetes Care* 30(7): 1926-1933.
33. Rehnberg GL, Hem JF, Carter SD, Linko RS & Laskey JW (1982). Chronic ingestion of Mn3O4 by rats: tissue accumulation and distribution of manganese in two generations. *J of Toxicol and Environ Heal, Part A Curr Iss* 9(2): 175-188.
34. EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) (2013). Scientific opinion on dietary reference values for manganese. *EFSA J* 11(11): 3419.
35. Khan SA, Khan L, Hussain I, Marwat KB & Akhtar N (2008). Profile of heavy metals in selected medicinal plants. *Pak J of Weed Sci Res* 14(1-2): 101-110.
36. Koh ES, Kim SJ, Yoon HE, Chung JH, Chung S, Park CW & Shin SJ (2014). Association of blood manganese level with diabetes and renal dysfunction: a cross-sectional study of the Korean general population. *BMC Endocrine Disord* 14(1): 24.
37. Kazi TG, Afridi HI, Kazi N, Jamali MK, Arain MB, Jalbani N & Kandhro GA (2008). Copper, chromium, manganese, iron, nickel, and zinc levels in biological

- samples of diabetes mellitus patients. *Biol Trace Element Res* 122(1): 1-18.
38. Sium M, Kareru P, Keriko J, Girmay B, Medhanie G & Debretsion S (2016). Profile of trace elements in selected medicinal plants used for the treatment of diabetes in Eritrea. *The Scientific World Journal*, 2016.
39. Marschner H (2011). *Marschner's mineral nutrition of higher plants*. Academic Press.
40. Mose DG, Mushrush GW & Chrosniak C (1990). The radioactive hazard of potable water in Virginia and Maryland. *Bull of Environ Contamin and Toxicol* 44(4): 508-513.
41. Plum LM, Rink L & Haase H (2010). The essential toxin: impact of zinc on human health. *International journal of environmental research and public health*, 7(4), 1342-1365.
42. Krupanidhi S, Sreekumar A & Sanjeevi CB (2008). Copper & biological health. *Indian J of Medi Res* 128(4): 448.
43. Maret W (2013). Zinc biochemistry: from a single zinc enzyme to a key element of life. *Adv in Nutri* 4(1): 82-91.
44. Brender JR, Hartman K, Nanga RPR, Popovych N, de la Salud Bea R, Vivekanandan S & Ramamoorthy A (2010). Role of zinc in human islet amyloid polypeptide aggregation. *J of the American Chem Soc* 132(26): 8973-8983.
45. Hashemipour M, Kelishadi R, Shapouri J, Sarrafzadegan N, Amini M, Tavakoli N & Poursafa P (2009). Effect of zinc supplementation on insulin resistance and components of the metabolic syndrome in prepubertal obese children. *Hormones*, 8(4): 279-285.
46. Alloway BJ (2003). *Heavy metals in soils*, John Wiley and Sons. Bud rot on oil palm plantations: Link to soil physical properties and nutrient status. *Better Crops Inter* 2: 22-5.
47. Ghani A, Ali Z, Ishtiaq M, Maqbool M & Parveen S (2012). Estimation of macro and micronutrients in some important medicinal plants of Soon Valley, District Khushab, Pakistan. *African J of Biotechnol* 11(78): 14386-14391.
48. Shirin K, Imad S, Shafiq S & Fatima K (2010). Determination of major and trace elements in the indigenous medicinal plant *Withania somnifera* and their possible correlation with therapeutic activity. *J of Saudi Chem Soc* 14(1): 97-100.
49. Qiao W, Peng Z, Wang Z, Wei J & Zhou A (2009). Chromium improves glucose uptake and metabolism through upregulating the mRNA levels of IR, GLUT4, GS, and UCP3 in skeletal muscle cells. *Biol Trace Elem Res* 131(2): 133-142.
50. Niamat R, Khan MA, Khan KY, Ahmad M, Ali B, Mazari P & Ahmad H (2012). Element content of some ethnomedicinal *Ziziphus Linn.* species using atomic absorption spectroscopy technique. *J of Appl Pharma Sci* 2(3): 96.
51. Velayutharaj A, Saraswathi R, Shivakumar R, Saha S, Niranjan G, Ramesh R & Sreenivasan AR (2016). Association of serum magnesium with glycemic control and insulin resistance in patients with type 2 diabetes mellitus. *Inter J of Curr Res and Rev* 8(13): 17.
52. Chaudhary DP, Sharma R & Bansal DD (2010). Implications of magnesium deficiency in type 2 diabetes: a review. *Biol Trace Elem Res* 134(2): 119-129.
53. Viktorinova A, Toserova E, Krizko M & Durackova Z (2009). Altered metabolism of copper, zinc, and magnesium is associated with increased levels of glycated hemoglobin in patients with diabetes mellitus. *Metabol* 58(10): 1477-1482.
54. Olafisoye OB, Oguntibeju OO & Osibote OA (2017). Trace elements and radionuclides in palm oil, soil, water, and leaves from oil palm plantations: A review. *Critical Rev in Food Sci and Nutr* 57(7): 1295-1315.
55. Vinceti M, Wei ET, Malagoli C, Bergomi M & Vivoli G (2001). Adverse health effects of selenium in humans. *Rev on Environ Heal* 16(4): 233-252.
56. Goldhaber SB (2003). Trace element risk assessment: essentiality vs. toxicity. *Regul Toxicol and Pharmacol* 38(2): 232-242.
57. Bolann BJ, Rahil-Khazen R, Henriksen H, Isrenn R & Ulvik RJ (2007). Evaluation of methods for trace-element determination with emphasis on their usability in the clinical routine laboratory. *Scandinavian J of Clin and Lab Invest* 67(4): 353-366.