ARNALDO BORDONI (*)

NEW DATA ON THE PALAEARCTIC XANTHOLININI.
 14. SPECIES IN THE ZOOLOGICAL MUSEUM OF THE MOSCOW LOMONOSOV
 STATE UNIVERSITY (COLEOPTERA STAPHYLINIDAE)
 277th Contribution to the knowledge of the Staphylinidae

(*) Museo di Storia Naturale dell'Università di Firenze, sezione di Zoologia "La Specola", via Romana 17, 50125 Firenze, Italy; e-mail: arnaldo.bordoni@fastwebnet.it

Bordoni A. – New data on the palaearctic Xantholinini. 14. Species of the Zoological Museum of the Moscow Lomonosov State University (Coleoptera Staphylinidae). 277th Contribution to the knowledge of the Staphylinidae.

A large collection of Russian Xantholinini preserved in the Zoological Museum of the Moscow Lomonosov State University was studied and a checklist of the species and all records are listed. New species are described and figured (*Hypnogyra orientalis* sp. n. from East Russia and *Xantholinus samarensis* sp. n. from South Russia). The following synonymies are proposed: *Xantholinus laevigatus* Jacobsen, 1849 = *Xantholinus hochhuthi* Bordoni, 1975, syn. n. and *Hypnogyra hoffmanni* (Bernhauer, 1928) = *Xantholinus nudobiiiformis* Tikhomirova, 1973. Some species are new record for the listed regions: *Stenistoderus versicolor* (Solsky, 1871) (Astrakhan Oblast), *Leptacinus sulcifrons* (Stephens, 1833) (East Siberia), *L. othioides* Baudi, 1870 (Siberia), *Allolinus insolens* Smetana, 1967 (Russia), *Gyrohypnus wutaiahnensis* Bordoni, 2000 (Russia), *G. sichuanensis* Zheng, 1995 (Russia), *G. ochripennis* (Eppelsheim, 1892) (Siberia), *Nudobius lentus* (Gravenhorst, 1806) (Kirghisistan), *Milichilinus decorus* (Erichson, 1839) (Russia), *Hypnogyra hoffmanni* (Bernhauer, 1928) (Russia), *Xantholinus dvoraki* Coiffait, 1956 (Tajikistan), *X. audrasi* Coiffait, 1956 (Azerbaijan), *X. tricolor* (Fabricius, 1787) (Moldavia), *X. haematodes* Kolenati, 1846 (Georgia), *X. procerus* (Erichson, 1839) (Ukraine), *X. caucasicus* Bordoni, 1975 (North Ossetia), *X. vinicolor* Ushakov, 1989 (North Ossetia), and *Neoypnus mandschuricus* (Bernhauer, 1923) (North Korea).

KEY WORDS: Coleoptera, Staphylinidae, Xantholinini, Moscow Museum, checklist, records, new species, new synonymies, Russia.

INTRODUCTION

My interest in Russian Xantholinini began several decades ago, when it led me to contact local colleagues, in particular Anna Tikhomirova, which enabled me to study the types of various species described from the region.

The results of those early studies were published in BORDONI, 1975. Some records, however remained unpublished. My interest was recently rekindled when I received a large number of specimens from Dr. Aleksey Gusakov, curator of the collections of the Zoological Museum of Moscow Lomonosov State University.

The present contribution is the result of the study of this material, to which I have added specimens from the private collections of Alexey Shavrin (Institute of Systematic Biology, Daugavpils), Alexey Solodovnikov (Zoologisk Museum, Kobenhavn), Ivan Ushakov (Moscow), and very few specimens of other colleagues: Yasuhiko Hayashi (Kawanishi) and Andreas Putz (Eisenhüttenstadt) and a few from my own collection, including specimens received as a gift from Anna Tikhomirova, and few others received for study from the Museums of Budapest, Milan and St. Petersburg in the early 1970s.

This paper provides a fairly comprehensive picture of the species occurring in Russia, although it does not include a few of the less common species. It is hoped that the list, which only includes the Xantholinini kept in the Moscow Museum, will stimulate further research. The study suggests that the most common species in Russia are *Nudobius*

lentus (Gravenhorst, 1806) and *Xantholinus tricolor* (Fabricius, 1787). Not a few species are associated with decaying plants, since the observed area is rich in forests. The only ones of these Moscow Museum specimens mentioned in SMETANA, 1982 are those belonging to the genus *Zeteotomus* Jacquelin du Val, 1857 in the Kuril Islands. Some species of Xantholinini were cited from Russia by other authors but I do not know of any similar list of taxa related to this tribe for this region.

I consider it unnecessary to repeat the complete distributions of the various species, so I refer only to some contributions on the Russian Xantholinini and to the catalogues of HERMAN (2001) and SMETANA (2004). I have adopted the subdivisions of Russia suggested by LÖBL & SMETANA, 2004: ST: South European Territory; CT: Central European Territory; NT: North European Territory; WS: West Siberia; ES: East Siberia; FE: Far East, and cite potential new records.

Acronyms of depositories: cB: coll. Bordoni, Firenze, Italy; cH: coll. Hayashi, Kawanishi, Japan; cP: coll. Putz, Eisenhüttenstadt, Germany; cS: coll. Schülke, Berlin, Germany; cSha: coll. Shavrin, Daugavpils, Latvia; cSol: coll. Solodovnikov, Kobenhavn, Denmark; cU: coll. Ushakov, Moscow, Russia; ISBD: Institute of Systematic Biology, Daugavpils, Latvia; MBM: Magyar Nemzeti Múzeum, Budapest, Hungary; MCSNM: Museo civico di Storia naturale, Milano, Italy; MCSNT: Museo civico di Storia naturale, Trieste, Italy; ZIN: Zoological Institute, St. Petersburg, Russia; ZMNU: Zoological Museum of Moscow University, Russia.

TAXONOMY

Zeteotomus scripticollis (Hochhuth, 1849)

MATERIAL EXAMINED – Caucasus, Adler, V. Solotikhin 20.IX-2.X.1974, 1 ex. (cSol); S Russia, Krasnodarskyi, 1.5 km N Betta, Solodovnikov 25.IV.1995, 2 exx. (cB); S Russia, Krasnodar, Ubinskaya, V. Kovalev VI.1971, 1 ex. (cB). Azerbaijan, Mts Talish, Aurora, Danilevsky V.1979, 1 ex. (cB).

DISTRIBUTION – Hungary, Azerbaijan, Georgia (HERMAN, 2001; SMETANA, 2004). ST.

NOTE – Many specimens was collected under the bark of fir. I give a figure of the genitalia (Fig. I, 1-3).

Zeteotomus dilatipennis (Kirschenblat, 1949)

MATERIAL EXAMINED – E Russia, reg. Khabarovsky, Khehzhir, Potozkaya 25-30.IV.1976, 10 exx. (ZMNU), 5 exx. (cB); same data, 6-13.V.1976, 3 exx. (ZMNU), 2 exx. (cB); Kuril Is., Kunashir, Mendleev, Potozkaya, 10-29.IX.1972, 2 exx. (ZMNU), 1 ex. (cB); same data, 29.IX.1972, 1 ex. (ZMNU); same data, 28.IV-4.VI.1977, 4 exx. (ZMNU), 4 exx. (cB); same data, Ilmazajev, 17.IX.1972, 1 ex. (ZMNU); same data, Sernovodsk, Potozkaya 10-14.IX.1972, 5 exx. (ZMNU), 3 exx. (cB); same data, Potozkaya 14.IX.1972, 3 exx. (ZMNU), 2 exx. (cB).

DISTRIBUTION – Russian Far East, Shaanxi, Heilongjiang (HERMAN, 2001; SMETANA, 1982, 2004; BORDONI, 2006; 2009). FE, Kuril Isl.

Fig. I – 1. Tergite of the male genital segment; 2. Sternite of the same; 3. Aedeagus of *Zeteotomus scripticollis* (Hochhuth); 4. Tergite of the male genital segment; 5. Sternite of the same; 6. Aedeagus of *Zeteotomus dilatipennis* (Kirschenblat) (scale bar: 0.1 mm).

NOTE – The specimens were collected under bark of fir, alder and cherry. I give a figure of the genitalia (Fig. I, 4-6) which was already illustrated by SMETANA (l. c.).

Zeteotomus maximus (Bernhauer, 1907)

MATERIAL EXAMINED – Kuril Is., Kunashir, Mendleevo, Potozkaya, 23.IV.1977, 1 ex. (ZMNU), 1 ex. (cB); same data 23.IV-14.IX.1972, 14 exx. (ZMNU), 6 exx. (cB); same data, 10.IX.1972, 1 ex. (ZMNU), 1 ex. (cB); same data, Tretjakovo, Potozkaya, 6.V.1977, 1 ex. (ZMNU).

DISTRIBUTION – Japan (BORDONI, 1997); Russian Far East, China (HERMAN, 2001; SMETANA, 1982, 2004). FE, Kuril Isl.

Stenistoderus versicolor (Solsky, 1871)

MATERIAL EXAMINED – S Russia, Astrakan, Dosan', K. Grebennikov 16.V.1997, 1 ex. (ISBD), Caucasus, Krasnodarsky, Dolgaja, Solodovnikov, 1 ex. (ZMNU); Caucasus, Bakhejisaraysky, Schastlivae, Solodovnikov 2.VI.1996, 1 ex. (cB).

DISTRIBUTION – Georgia, Tadjikistan (USHAKOV, 1989; HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). ST. Probably a new record for Astrakhan Oblast.

NOTE – The specimens were collected under bark of fir and alder.

Phacophallus parumpunctatus (Gyllenhal, 1827)

MATERIAL EXAMINED – Caucasus, Krasnodarsky, Bagovskaya, 600 m, Solodovnikov 15.V.1996, 1 ex. (cSol).

DISTRIBUTION – Widespread species in the Palaearctic Region and introduced in some other regions (HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). CT, ST.

Leptacinus batychrus (Gyllenhal, 1827)

MATERIAL EXAMINED – Russia, Mosca, Filippov 10.VII.1918, 1 ex.; same data, Mjavkovo, leg. ? 25.VI.1906, 1 ex.; Russia, Volynsk, Tushino 25.VII.1906, 1 ex. (ZMNU); Central Siberia, Irkutsk, Zalarinska, Tungui, Shavrin 20-24.VI.1997, 2 exx. (cSha). S Turkmenistan, Diumak, V. Savizky 7-12.X.1993, 6 exx. (cSol), 1 ex. (cB).

DISTRIBUTION – Widespread species in the Palaearctic Region (HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). ST.

Leptacinus sulcifrons (Stephens, 1833)

MATERIAL EXAMINED – Russia, Studenez, Mosca, Kozhevnikov leg., 1 ex. (cB); E Siberia, Cità, Krasno-Cikoisky, Cikoï river, Anishenko 21.VIII.1998, 1 ex.; Central Siberia, Irkutsk, Zalarinska, Bazir, Shavrin 20-24.VI.1997, 1 ex.; same data, Alarsky, Alar, Shavrin 26-28.VI.1997, 1 ex.; Central S Siberia, Buriatia, Goryachinsk, Termal, Shavrin 12.VIII.1997, 1 ex.; SE Siberia, Shelenkhov, Shavrib 2.IX.1992-1.V.1994, 1 ex. (cSha); SE

Russia, Orenburgskaya, Buzluskyy park, 2 km N Koltubaysky, I. Goreslavev 4.VIII.1998, 1 ex. (cSol); Central Russia, Tatarstan, Stolbishensky, Pescanye Kovali, Utrobina 9.V.1957, 4 exx.; same data, Knibishev, Utrobina 4.IX.1957, 4 exx.; same data, Balinsky, Sherganovo, Utrobina 2.VIII.1959, 1 ex.; SW Russia, South Don, Dshanibsky steppe, Dernina 7.X.1964, 1 ex.; Perm'skaya, I. Shavrov VIII.1921, 1 ex. (ZMNU); Caucasus, Krasnodarsky, Solodovnikov 27.VII.1995, 4 exx.; Krasnodarsky, Bagovskaya, 600 m, Solodovnikov 15.V.1996, 1 ex. (cSol), 1 ex. (cB); Kanaskaya, Solodovnikov 27.VII.1995, 1 ex. (cSol); same data, Kuban' river, 50 km NWW Krasnodar, Solodovnikov 16.VI.1994, 1 ex. (cB); same data, Novopokhladnoe, 1000 m, Solodovnikov 2.VI.1994, 2 exx.; same data, Saratovskaya, Solodovnikov 24.IV.1996, 1 ex. (cSol). Turkmenistan, Ashgabat, A. Yagdeyev 2.VI.1967, 1 ex. (ZMNU).

DISTRIBUTION – Widespread species in the Palaearctic Region (HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). CT. New record for East Siberia.

Leptacinus pusillus (Stephens, 1833)

MATERIAL EXAMINED – Russia, Central S Siberia, Buriatia, Il'inka, Selenga river, Shavrin 1.VIII.1997, 1 ex. (cSha).

DISTRIBUTION – Widespread species in the Palaearctic Region (HERMAN, 2001; SMETANA, 2004). CT, NT, ST, WS.

Leptacinus formicetorum Märkel, 1841

MATERIAL EXAMINED – Russia, Moscovskaya, Zvenigorod, Nicolina gora, Uijasniov 21.X-25.XI.1951, 2 exx. (ZMNU); Central Siberia, Irkutsk, Ust-Ordinsky, Zagi, Shavrin 23.VIII.1996, 1 ex.; S Siberia, S Buryatia, Chikoj, 6 km S Zingan-Debe, Shavrin 1-3.VI.2002, 2 exx.; SE Siberia, Shelekhov, Podkamsoe, Shilenkov 1.VI.1995, 1 ex. (cSha). Turkmenistan, Akibai, Tamarin (?), Potozkaya 20.III-9.IV.1971, 3 exx. (ZMNU), 1 ex. (cB); same data, 14.IV.1971, 1 ex. (cB); same data, Ipai-Kalà, Potozkaya 23.V.1971, 2 exx. Georgia, Batumi, D. Romashov 5.IV.1928, 1 ex. (ZMNU).

DISTRIBUTION – Widespread species in the Palaearctic Region (HERMAN, 2001; SMETANA, 2004). CT, NT, WS.

Leptacinus othioides Baudi, 1870

MATERIAL EXAMINED – Caucasus, Krasnodarsky, Novopokhladnoe, 1000 m, Solodovnikov 2.VI.1994, 1 ex. (cSol); Russia, Central Siberia, Irkutsk, c/o Baikal lake, Shavrin 17.IV.1995, 2 exx. (cSha); SE Siberia, Shelekhov, Shavrin 2.IX.1992-1.V.1994, 5 exx. (cSha); Central E Russia, Samara, Chernigovsky, Goreslavev 10.VIII.1991, 1 ex. (cSol), 1 ex. (cB); same data, Samara river, Goreslavev 2.VII.1995, 5 exx. (cSol), 1 ex. (cB); same data, Bakhilova, Goreslavev 3.V.1993, 1 ex. (cSol), 1 ex. (cB); same data, Poliana, Solodovnikov 27.VII.1995, 4 exx. (cSol).

DISTRIBUTION – Europe and Mediterranean (HERMAN, 2001; SMETANA, 2004). ST. New record for Siberia.

Allolinus peliopterus (Solsky, 1875)

MATERIAL EXAMINED – Russia, E Siberia, Krasno-Cikoisky, Khilok river, Ceremushka, Shavrin 26.VIII.1998, 1 ex. (cSol); same data, Irkutsk, Alazsky, Alar', Shavrin 26-28.VI.1997, 1 ex. (cB); same data, Buryatia, Dzhidinskiy, Dzhida river, Shavrin XII.1974, 1 ex. (cSol), 1 ex. (cB); same data, Buryatia, Ust-Kiryan Mt, leg. ? 17.VII.1931, 1 ex. (cSol); same data, Belozyorsk, Shavrin 4-9.VII.1997, 1 ex. (cB); same data, Kuyutunskij, Ust'-Kada, Shavrin 24.VI.2009, 1 ex. (cSol).

DISTRIBUTION – Russian Far East, Mongolia (HERMAN, 2001; SMETANA, 2004). ES.

Allolinus insolens Smetana, 1967

MATERIAL EXAMINED – S Russia, Krasnodar, Semenskhko Mt., 1000 m, Solodovnikov 11.VI.1992, 1 ex.; SE Siberia, Chitinskaya, Krasno-Chihojskij, Khilok, Shavrin 26.VIII.1998, 1 ex. (cSol); E Siberia, Irkutsk, S. Rodionoff 30.VII.1917, 1 ex.; E Russia, Primorskij Kraj, Nat. Res. Kedrowaja Padj, 40 km WSW Wladiwostok, F. Hieke 6-8.VIII.1991, 3 exx. (cB).

DISTRIBUTION – Mongolia (HERMAN, 2001; SMETANA, 2004). New record for Russia: South Russia, Siberia and Far East.

Gauropterus fulgidus (Fabricius, 1787)

MATERIAL EXAMINED – Mosca, Bogorodskoe, leg. ? 18.IV.1938, 1 ex.; E Siberia, Sakha, K. Satunin 7.VIII.1988, 1 ex. (ZMNU).

DISTRIBUTION – Palaearctic Region (HERMAN, 2001; SMETANA, 2004). CT, NT, ST.

Gauropterus sanguinipennis (Kolenati, 1846)

MATERIAL EXAMINED – Central East Russia, Samara, Samara river, Goreslavetz 24.IV.1994, 1 ex.; Caucasus, Krasnodarsky, 10 km S Ubinskaya, 200 m, Solodovnikov 22.VIII.1995, 1 ex.; same data, B. & M. Savizkie 28.V.1992, 1 ex. (cSol).

DISTRIBUTION – Eastern Europe and Western Asia (HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). ST.

Gauropterus sanguinipes Reitter, 1889

MATERIAL EXAMINED – Azerbaijan, Shemacha, 1 ex.; Caucasus, Krasnodar, 1 ex. (ZMNU).

DISTRIBUTION – East Palaearctic Region (HERMAN, 2001; SMETANA, 2004).

Gyrophypnus wutaishanensis Bordoni, 2000

MATERIAL EXAMINED – E Russia, Primorskij Kraj, Lazovskij distr., 5 km NE Lazo, J. Sundukov 1-5.VI.1995, 1 ex. (cP).

DISTRIBUTION – Shaanxi, Shanxi (HERMAN, 2001; SMETANA, 2004). New record for Russia.

Gyrophypnus sichuanensis Zheng, 1995

MATERIAL EXAMINED – E Russia, S Primorye, Lazovskij Res., Pereakalnaya river, Y. Sundukov 8-11.VII.2007, 1 ex. (cB).

DISTRIBUTION – Sichuan, Korea (HERMAN, 2001; SMETANA, 2004; BORDONI, 2000). First record for Russia.

Gyrophypnus angustatus Stephens, 1833

MATERIAL EXAMINED – Russia, Mosca, Studener, 1 ex.; Mosca, leg ? 3.V.1922, 2 exx. (ZMNU), 1 ex. (cB); Mosca, Bykovo, O. Kryzanosky 2.V.1934, 1 ex.; Mosca, Vorobyevy gory, B. Stark 11.IV.1948, 2 exx.; Moskovskaya, Mytishinskiy, Kliazma, Stark 24.IV-25.VIII.1947, 3 exx. (ZMNU), 2 exx. (cB); same data, Zvenigorod, Nicolina gora, Stark 17.VIII.1944-13.V.1945, 1 ex. (ZMNU); Central Siberia, Irkutsk, Anishenko VIII.1995, 1 ex.; same data, B. Kon. (?), Shavrin 17.VII.1993, 2 exx.; E Siberia, Krasno-Cikoisky, Gonacikhi river, Anishenko 22.VIII.1998, 1 ex. (cSha); Russia Central E, Samara, Bolshoy Cernigousky, I. Greslavev 8.VIII.1998, 1 ex. (cSol); Siberia, Novosibirsk, Obskoe, Dvdko 27.IV.1997, 1 ex. (cB); Caucasus, Krasnodarsky, Novopokhladoe, 700 m, V. Savizky 3.VI.1995, 1 ex.; same data, Guzerepl, Solodovnikov 9-10.VI.1992, 3 exx. (cSol).

DISTRIBUTION – Widespread species in the Palaearctic Region, introduced in other regions (USHAKOV, 1988; GUSAROV, 1989; SOLODOVNIKOV, 1998; HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). CT, NT, ST, WS.

Gyrophypnus fracticornis (Müller, 1776)

MATERIAL EXAMINED – Russia, Moskovskaya, Elino, Gusakov 19.VIII-4.X.1995, 2 exx. (ZMNU); Central Siberia, 17 km Irkutsk, Olkha river, Shavrin 17.VIII.1993, 1 ex. (cSha); S Russia, Krasnodar, Semashkho Mt, 1000 m, Solodovnikov 11.VI.1990, 1 ex.; NW Caucasus, Maly Trach Mt, 1600-2000 m, Novopokhladoe, Solodovnikov 3.VI.1994, 1 ex. (cSol). Turkmenistan, Kouchka Broussailles (?), leg. ? 16.V.1963, 1 ex. (cSha).

DISTRIBUTION – Widespread species in the Palaearctic Region, introduced in other regions (HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). CT, NT, ST, WS.

Gyrophypnus punctulatus (Paykull, 1789)

MATERIAL EXAMINED – Russia, Leningrad, Filippov 20.IV.1918, 1 ex. (ZMNU); same data, Divenskaya, Filippov 12.VII.1916, 1 ex. (cB); Pushkin (St. Petersburg), Filippov 2.V.1918, 4 exx.; W Russia, Yaroslavl', Filippov 28.VI. 1918, 4 exx.; Skhodnja, Mosca, leg. ? 8.IX.1906-20.VIII.1908. 5 exx. (ZMNU), 2 exx. (cB); Mosca, Zarizino park, leg. ? 22.V.1930, 1 ex.; Mosca, Marjina rosha, Dazkevic 11.IV. ?, 1 ex.; Mosca, Iljinskaya, D. Romashov 14.IV.1921, 1 ex. (ZMNU), 1 ex. (cB); Caucasus, Maikop, Tolardwsky, Shaposhnikov 15.V.1932, 1 ex. (ZMNU).

DISTRIBUTION – Widespread species in the Palearctic Region (HERMAN, 2001; SMETANA, 2004). CT, NT, ST.

***Gyrophypnus ochripennis* (Eppelsheim, 1892)**

MATERIAL EXAMINED – Russia, S Siberia, Altai, Tscholesmantal, S. Wache & G. Biedermann 3.X.1907, 1 ex.; E Siberia, Irkutsk, Angarsk, I. Enustschenko 30.V.2008, 2 exx. (cB); same data, Khamar-Daban Mts, Snezhnaya valley, Shavrin 5-8.VI.2007, 1 ex. (cSha). Uzbekistan, Andizhan, Filippov 27.IV.1924, 1 ex. (ZMNU). S Tagikistan, Muminabad Sharan, Tikhomirova 10.V.1969, 9 exx. (cB).

DISTRIBUTION – Kavkaz, Western Asia (HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). ST. First record for Siberia.

***Gyrophypnus atratus* (Heer, 1839)**

MATERIAL EXAMINED – Ostankino, Mosca, leg. ? 10.X.1906, 1 ex. (ZMNU), 1 ex. (cB); Vorobjevygori, Mosca, 2 exx. (ZMNU).

DISTRIBUTION – Europe (HERMAN, 2001; SMETANA, 2004). NT.

***Nudobius lentus* (Gravenhorst, 1806)**

MATERIAL EXAMINED – NW Russia, Petropol (St. Petersburg), 2 exx. (cB); Russia, Leningradskaya, Boksitogorsky, 7 km Mosilevo, Raguva river, R. Filomov 13.III.1994, 1 ex. (cSol); Moskovskaya, Zvenigorod, Nicolina gora, Stark 17.VIII.1944-13.V.1945, 2 exx.; same data, Zelenograd, A. Gusakov 4.X.1995-23.IV.1998, 1 ex.; same data, Mamontovka, Filippov 26.IV.1926, 1 ex. (ZMNU); Moscow reg., Chekhovskii distr., Luch vill., O. Mosolov 10.IV.2004, 4 exx. (cB); Central E Russia, Samara Obl., Bakhilova Polyana vill., Zhiguli Nat. Res., I. Goreslavetz 29.VIII.1997, 1 ex. (cSol); SE Siberia, Shelekhov, Shavrin 2-IX.1992-1.V.1994, 7 ex. (cSha), 1 ex. (cB); E Siberia, Citá, Krasno-Cikoisky, Zagi, 42 km W Iamarovka (Mogron), Shavrin 24.VIII.1998, 1 ex. (cB); Central Siberia, Irkutsk, Shavrin VIII.1992, 1 ex. (cSha); Central South Siberia, Buriatia, Tazny, Potozkaya 16-30.VI.1976-2.VII.1976, 6 exx. (ZMNU), 4 exx. (cB); S Siberia, Baikal lake, Hama-Daban, Potozkaya 9.VI.1976, 1 ex.; same data, Babuskin, Potozkaya 5.VI.1976, 1 ex.; SE Russia, Suputinka, Potozkaya 9.IX.1968-27.V.1969, 9 exx. (ZMNU), 3 exx. (cB); SE Russia, Khabarovsk, Mehzi, Potozkaya 10-13.V.1976, 14 exx. (ZMNU), 1 ex. (cB); Primorsky, Lasovsky Res., Potozkaya 15.VII.1979, 2 exx. (ZMNU), 1 ex. (cB); Primorsk Krai, Arsevev env., Strab VI.1991, 1 ex. (cB); E Russia, S Primorye, Lazovskij Res., Perekatnaya river, Y. Sundukov 15-16.VIII.2007, 2 exx. (cSha); Kuril Is., Kunashir, Mendeleevo, Potozkaya 28.IX.1972-1.X.1972, 9 exx. (ZMNU), 2 exx. (cB); same data, 11.IX.1972, 1 ex.; same data, Sernovodsk, Potozkaya 20.IX.1972, 1 ex. (ZMNU); Kundur, Potozkaya 22-27.VI.1975, 1 ex. (ZMNU), 1 ex. (cB). Kirghizistan, Sujan, Filippov 5.V.1924, 1 ex. (ZMNU); same data, Teplokljucenka, leg. ? 29.VII.1976, 1 ex. (cB).

DISTRIBUTION – Palearctic Region (Jacobson, 1909; Kryzhanovskii *et al.*, 1973; HERMAN, 2001; SMETANA, 1982,

2004). CT, NT, ST, WS. First record for Kyrgyzstan.

NOTE – The specimens were collected under bark of spruce, pine, poplar and larch.

This species is very variable in size, coloration (some specimens pale, with yellowish elytra in Buriatia), shape of head (some specimens with longer and wider head, and denser punctation on head), and in the series of punctures on pronotum.

***Nudobius fasciatus* (Hochhuth, 1849)**

MATERIAL EXAMINED – Azerbaijan, Mts Talish, Aurora, Danilevsky 13.IV.1980, 1 ex. (cB).

DISTRIBUTION – Azerbaijan, Iran (BORDONI, 1975; HERMAN, 2001; SMETANA, 2004; BORDONI, 2010).

NOTE – The species was included in the genus *Calontholinus* Reitter, 1908 before being assigned to *Nudobius* Thomson, 1860 (BORDONI, 2010).

***Nudobius umbratus* (Motschulsky, 1860)**

MATERIAL EXAMINED – Republic Altai, Onggudayskiy distr., 10 km SE Semlinsky pass, A. Kirejchuk 15.VI.1999, 3 exx. (cSol), 2 exx. (cB); NW Caucasus, Novoprohladnoe, 700 m, V. Savitsky 25.VI.1995, 1 ex.; Krasnodar Terr., Aigba Mt, Krasnaya Poljana, 800-900 m, A. Solodovnikov 13.VIII.1994, 3 exx. (cSol); same data, Semashkho Mt., Solodovnikov 8.IV.1990, 1 ex.; same data, Ubinskaya Mt., Solodovnikov 8.IV.1990, 1 ex. (cB).

DISTRIBUTION – Western Asia (HERMAN, 2001; SMETANA, 2004). ST.

***Hypnogyra angularis* (Ganglbauer, 1895)**

MATERIAL EXAMINED – Russia, Mosca, Ostankino, Filippov 1918, 1 ex.; Mosca, Nicolina gora, S. Nikulin 6.VIII.1938, 1 ex.; Moskovskaya, Zvenigorod, Abramxevo, Stark 13.VIII.1949-3.VIII.1958, 1 ex. (ZMNU). Ukraina, Poltava, 1 ex. (ZIN).

DISTRIBUTION – Europe, North Africa (HERMAN, 200; SMETANA, 2004; BORDONI, 2011). CT.

***Hypnogyra hoffmanni* (Bernhauer, 1928)**

The type of *Xantholinus nudobiiiformis* Tikhomirova, 1973 from Khasan (Khasansky Distr., Far East Russia) on the border of North Korea and China was not found in the Zoological Institute of St. Petersburg, but the external characters and the small figure in the description (in particular the peculiar shape of parameres and the structure of the inner sac of the aedeagus) correspond to those of *Hypnogyra hoffmanni*, so I propose the following synonymy: *Hypnogyra hoffmanni* (Bernhauer, 1928) = *Xantholinus nudobiiiformis* Tikhomirova, 1973, syn. n.

DISTRIBUTION – This species is known from Northeastern China (Heilongjiang and Hebei) (BORDONI, 2000) and North Korea (BORDONI, in press). First record for Russia.

NOTE – In recent catalogues (HERMAN, 2001 and SMETANA, 2004) *nudobiiiformis* is referred to the genus *Megalinus* Mulsant & Rey, 1877.

Hypnogyra orientalis sp. n.

MATERIAL EXAMINED – Holotype ♂: NE Russia, Primorsky, Lasuvoky Res., A. Tikomirova 9.VI.1967 (cB); paratype ♀: SE Russia, Khabarovsk prov., Slevianka, 20 km N Troistkoye, Amur bank, W. Shawaller 14-19.VI.1990 (cB).

DESCRIPTION – Length of body 7 mm; from anterior margin of head to posterior margin of elytra: 3.3 mm. Body (Fig. III, 1) pale reddish-brown with dark head and yellowish posterior margins of elytra; antennae brown with the apex of scape and 2nd-3rd segments light brown. Very similar to *H. tenebrosa* Bordoni, 1997 from Japan but more slender, narrower and paler; in particular the head is narrower, with less rounded posterior angles; the eyes a little larger, and with finer punctation and shorter and narrower pronotum and shorter and narrower elytra. Posterior margin of the 6th visible abdominal tergite usually dentate, the protrusions thick and rather short.

Tergite and sternite of the male genital segment as in Fig. II, 1-2. Aedeagus (Fig. II, 3; Fig. III, 2) 0.7 mm long, shorter than in *H. tenebrosa*, with different parameres and inner sac, which is narrow and long, slightly dilated in the proximal portion, covered with fine and very dense spinulae.

Fig. II – 1. Tergite of the male genital segment; 2. Sternite of the same; 3. Aedeagus of *Hypnogyra orientalis* sp. n.; 4. Tergite of the male genital segment; 5. Sternite of the same of *Xantholinus samarensis* sp. n.; 6. Tergite of the male genital segment; 7. Sternite of the same of *Xantholinus vinicolor* Ushakov (scale bar: 0.1 mm).

Fig. III – 1. Habitus (total length: 7 mm); 2. Aedeagus (total length: 0.7 mm) of *Hypnogyra orientalis* sp. n.; 3. Habitus; 4. Aedeagus of *Xantholinus samarensis* sp. n. (total length respectively: 9.7 mm and 1.1 mm); 5. Habitus; 6. Aedeagus of *Xantholinus vinicolor* Ushakov (total length respectively: 7.5 mm and 1.25 mm).

ETYMOLOGY – The specific epithet refers to the Eastern part of Russia.

DISTRIBUTION – The species is known only from the Far East Russia.

NOTE – The new species differs from the congeners especially by the structure of the inner sac of the aedeagus.

Milichilinus decorus (Erichson, 1839)

MATERIAL EXAMINED – Russia, Donhovka, Mosca, leg. ? 7.IX.1980, 1 ex.; NW Russia, 18 km Kalinin, Putilevo, Putilosky prison, Ushakov 19-27.VII.1982, 3 exx. (ZMNU), 3 exx. (cB); Urali, Visimsky Nat. Res., leg. ? I.1996, 1 ex. (ZMNU). Ukraina, Zakarpatskaya, Vinogradovsky, Tysa river, A. Koval 19.VII.1981, 1 ex.; same data, Kinodievo, Mucaciovo, Kurbatov 31.VIII.1983, 1 ex. (ZMNU)-

DISTRIBUTION – Europe (HERMAN, 2001; SMETANA, 2004). New record for Russia.

Xantholinus crassicornis Hochhuth, 1851

MATERIAL EXAMINED – Caucasus, Leder-Reitter, 1 ex. (MCSNT); Caucasus, 1 ex. (ZIN). Georgia, Abastuman, Reitter, 1 ex. (MCSNT), 1 ex. (cB); same data, Swanetien, 1 ex. (MCSNT); Caucasus, (Georgia), Abastuman, Leder, 1 ex. (MBM), 1 ex. (cB); Georgia, Banis-heri, Gori, Kirschenblat 16.V.1929, 1 ex. (ZIN).

DISTRIBUTION – Georgia, Turkey, South Russia (BORDONI, 1975 (sub *X. lederi* Coiffait, 1966); HERMAN, 2001; SMETANA, 2004).

Xantholinus linearis (Olivier, 1795)

MATERIAL EXAMINED – Russia, Mosca, K. Arnoldi 28.VI.1958, 1 ex.; Moskovskaya, Mytishinskiy, Klizma, Stark 24.IV-25.VIII.1946, 1 ex.; same data, Zvenigorod, Nicolina gora, Stark 29.IV-28.V.1945, 10 exx. (ZMNU); Russia Central E, Samara, Siniy Syrt, 13.5 km SW Costino, Rostash' river, Smeliansky 25.VIII.1994, 1 ex.; Central Russia, Kirovskaya, Svevca, Inferev 11.VI.1999, 1 ex. (cSha); same data, Kirov, Svecha, G. Juferov 27.V.1999, 2 exx.; S Urali, Curganskaya, Tikhonovka, Gorikoe lake, R. Filimonov 8.VIII.1995, 1 ex. (cSol); Central South Siberia, Buriatia, Selenga river, Tataurovo, Shavrin 2.VIII.1997, 1 ex.; same data, Tunkinskay valley, 2 km SW Turan, Shavrin 21.VIII.2006, 2 exx.; Central Siberia, Irkutsk, Olkha river, Shavrin 17.VIII.1993, 1 ex.; SE Siberia, Shelekhov, Olkhinskaya Mt., Olkha river, Shavrin 22.VIII.1993, 1 ex.; S Siberia, Baikal lake, Svjaty Nos peninsula, Mt Markov, 1800 m, Shavrin 11.VIII.1997, 1 ex. (cSha); E Siberia, Buryatia, Tunkinskij, Arshan, Kyngarga river, Shavrin 12.VIII.2009, 1 ex. (cSol); same data, Tunkinskaya valley, Moigoti, Shavrin 21.VIII.2006, 5 exx.; same data, Tunkinskaya valley, 2 km SW Turan, Shavrin 21.VIII.2006, 2 exx. (cSha), 2 exx. (cB); same data, Irkutsk, Usolskiy, Kitoi river, Shavrin 30.VII.1998, 1 ex. (cSol); Irkutsk prov., Shelenkhov, Irkai, Shavrin 2.VIII.2005, 4 exx.; same data, Erbogachyon, Nizhnyaya, Tunguska river, Shavrin & Enustschenko 18-27.VIII.2006, 2 exx. (cSha), 1 ex. (cB); Irkutsk, Cheremkhovskij, Sredmj Balaj, Shavrin 1-2.VIII.2006, 2 exx. (cB); same data, Katangskij, 15 km

Podvoloshino-Kirensk, Lena river, Shavrin & Enustschenko 1.VIII.2008, 1 ex. (cSol).

DISTRIBUTION – Palaearctic Region, introduced in other regions (BORDONI, 1975; HERMAN, 2001; SMETANA, 2004; SEMIONENKOV *et al.*, 2015). CT, NT, ST, WS.

Xantholinus longiventris Heer, 1839

MATERIAL EXAMINED – S Russia, Rostovskaya, Kosharsky, Podtelkovo, Minoransky 5.VI.1979, 3 exx. (cU), 1 ex. (cB); S Russia, Kamensk Ros., Glubokaya, Arnoldi 3.VIII.1949, 1 ex. (cU); W Russia, Belgorod, Borisovka, Lesna Vorskla Res., Solodovnikov 25.VI.1995, 1 ex.; Crimea, Simferopol, Salghir river, A. Benkovskiy 16.IV.1996, 1 ex. (cSol); Tauria (Crimea), Sebastopol, V. Pliginski 3.II.1911, 1 ex.; Crimea, Inkerman, V. Pliginski 3.XII.1911, 1 ex. (cB).

DISTRIBUTION – Palaearctic Region, introduced in other regions (HERMAN, 2001; SMETANA, 2004; SEMIONENKOV *et al.*, 2015). CT, NT, ST, WS.

Xantholinus dvoraki Coiffait, 1956

MATERIAL EXAMINED – SW Russia, Samara, 1.5 km Liplen stat., Goreslavetz 21.V.1994, 1 ex. (cSol); Caucasus, Leder, 1 ex. (MCSNT); Rostov reg., Sholokhovskii distr., Kalininskii vill., E. Khachikov 12.VI.2000, 1 ex.; same data, 18-23.VII.2005, 1 ex.; same data, Elanskaya vill. E. Khachikov 17-20.IX.2001, 1 exx. (cB). Taghikistan, Gissarskiy Mt., 1300-1800 m, Solodovnikov 18.V.1988, 2 exx. (cSol). Ukraine, Mukacevo, leg. ? I.1936, 2 exx. (MCSNT).

DISTRIBUTION – Europe (HERMAN, 2001; SMETANA, 2004; BORDONI, 2011; SEMIONENKOV *et al.*, 2015). CT, ST, WS. New record for Tajikistan.

Xantholinus audrasi Coiffait, 1956

MATERIAL EXAMINED – Russia, Mosca, Uzkoe, Arnoldi 19.IV.1961, 1 ex. (cB); Moscovskaya, Ceshnikovo, Tikhomirova 11.VIII.1963, 1 ex. (ZMNU); Central East Russia, Samara, Abdul-Zavod, I. Goreslavev 19.VIII.1995, 1 ex. (cSol); S Russia, Rostov, Oblivsky, 1 ex.; same data, Nadciciovka, Kisarizkaya, 5.VIII.1978, 1 ex. (cU); S Russia, Krasnodar, Mt Semashko, 1000 m, Solodovnikov 11.VI.1992, 1 ex.; same data, Guzeripl, Solodovnikov 10.VI.1992, 1 ex.; same data, Novopokhladnoe, 1400 m, V. Savinsky 19.VI.1995, 2 exx. (cB); Crimea, Simferopol, Salghir valley, leg. ? 5.IV.1957, 1 ex. (ZMNU); N Ossezia, Zeiskoe gorge, Mt Kalpersky, 1800 m, S. Alekseev 6.V.1994, 1 ex. (cB); same data, 9.VII.1982-16.IV.1984, 18 exx. (cU), 8 exx. (cB); N Ossezia, Dsvigis, Fiagdon river, Alekseev 9.X.1983, 1 ex.; W Caucasus, Krasnaja Poljana, Rivkin 17.V.1985, 1 ex. (cU). Ukraine, Mukacevo, leg. ? I.1936, 1 ex. (MCSNT). Armenia, Nts Mehri, Tashtungsky pass, Rivkin 27.IV.1963, 1 ex. (cB); same data, Mts Mehri, Kuris, Rivkin 26.IV.1963, 1 ex. (cU). Turkmenistan, Kopetdag Est, Ai-dere, 800 m, Eremin 25.X.1980, 1 ex. (cU). Azerbaijan, Talish, Lerik, Belov 20-25.VIII.1985, 4 exx. (cU), 1 ex. (cB); same data, Talish, Gosmalian 18.IV.1980, 1 ex. (cU).

DISTRIBUTION – Europe, Turkey (BORDONI, 1975 (sub. *X. armeniacus* Coiffait, 1956); HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). New record for Azerbaijan.

Xantholinus tricolor (Fabricius, 1787)

MATERIAL EXAMINED – Estonia, Saaremaa, Tagomoisa 24.VII.1829, 1 ex. (ZMNU). NW Russia, 18 km Kalinin, Putilovo, Volga river, Putilovsky prison, Ushakov 26.VI.1982, 5 exx. (cU); Russia, Pushkin (St. Petersburg), Filippov 3.V-30.VII.1918, 3 exx.; W Russia, Yaroslavl', Filippov 15.VI.1918, 2 exx. (cSha); Leningradskaya, Mininskaya, 30 km N Luga, Solodovnikov 24.IV.1994, 2 exx. (cSol); Leningradskaya, Lebiazy Res., Ushakov 19.V.1985, 1 ex.; same data, 1.V.1977, 2 exx.; Vologodskaya, Kadnilovsky Res., Ushakov 3.VII.1962, 3 exx. (cU), 1 ex. (cB); Russia, Mosca, Yanskyi Res., V. Savinsky 3.VI.1969, 1 ex. (cSol); same data, Staraja Kashira, Kozhevnikov 26.V.1902, 1 ex.; Mosca, Ostankino, Filippov 6.VIII.1918, 2 exx. (cSha); Moskovskaya, Bakovka, Dimovich VI.1985, 1 ex.; same data, N. Beresina, Chernogolovka 3.V.1984, 1 ex.; same data, Solnechogorodskiy, Peshki, Suvorov 1984, 2 exx.; Kaludskaya, Zukovskiy, Ushakov 15.V.1987, 3 exx.; Okskiy Res., Peredelsky, Eskov 19.X.1980, 1 ex.; same data, Bitzy, Ushakov 2.V.1985, 1 ex. (cU), 1 ex. (cB); same data, S Voskessenskaya, Peski, Ushakov 1.VI.1985, 1 ex.; same data, 33 km Kazanskaya, Ushakov 21.VII.1985, 1 ex.; Mliupino, Smirnova 7.VIII.1983, 1 ex.; same data, Krasnopahrinsky, Malinskoe Res., Belov 4.V.1980, 1 ex.; same data, Puskinsky, Gorenki, S. Kurbatov 18.IV.1976, 1 ex. (cU), 1 ex. (cB); same data, Serpuhovo, 2 km W Prioksko-Terrasny Res., Ushakov 29-30.VII.1983, 1 ex.; same data, Serebrioniy bor., Gruntal 5.IX.1984, 3 exx.; same data, Malino, Kriukovskiy, U. Meshkov 21.VIII.1984, 2 exx.; same data, Shodnia, Meshkov 13.VII.1984, 1 ex.; same data, 12 km Zvenigord, Mikailov 25.VI-22.VII.1981, 11 exx. (cU), 2 exx. (cB); same data, Donovka, Ushakov 7.IX.1980, 1 ex. (cU); Moscovskaya, Pushino, Solodovnikov 21.VI.1996, 1 ex.; same data, Zvenigorod, Abramzevo, Stark 13.VIII.1949-3.VIII.1958, 5 exx.; same data, Zvenigorod, Nicolina gora, Stark 17.VIII.1944-13.V.1945, 4 exx. (cSha); Mosca, Vorobyey gory, 2 exx.; Mosca, Zelenograd, Gusakov 31.V.1997, 1 ex.; Luban Station, Mosca, Shavrov, 1 ex.; Bykovo Station, leg. ? 24.VI.1908, 2 exx.; Mosca, Petrovsko-Razumovskoe, E. Smirnov 23.IV.1934, 1 ex.; Mosca, Losiny ostrov, leg. ? 23.VII.1919, 2 exx. (ZMNU), 2 exx. (cB); Central Russia, Voronezhskiy, Usmahka, K. Arnold 16.VI-29.VII.1964 and 23-31.VIII.1962, 5 exx.; Permskaya, Lysvensky, Kamenka, L. Voronova 26.VII.1967, 2 exx.; same data, Tatarka, L. Veronova 28.VII.1966, 1 ex. (cU), 1 ex. (cB); Verbilky, Dmitrievsky nerd, V. Lindgolm 28.VIII.1916, 1 ex. (ZMNU); Urali, Visimsky Res., 1 ex. (cU); W Russia, Belgorod, Borisovka, V. Savizkin 23.VII.1992, 1 ex. (cSol); same data, 23.XI.1992, 1 ex. (ZMNU); S Russia, Krasnodarskiy, 1.5 km N Betta, Solodovnikov 25.IV.1995, 1 ex. (cSol); same data, Eskov 10.VI.1980, 1 ex. (cU); W Siberia, 8 km S Tjumen', Solodovnikov 1998, 1 ex. (cB); same data, Armizon, Poriadina, 1 ex. (cU); S Siberia, 17 km Irkutsk, c/o Baikal lake, Anishenko 30.VI.1994-15.VI.1995, 1 ex.; same data, Alarsky, Alar, Shavrin 26-28.VI.1997, 1 ex.; same data, 17 km Irkutsk, A. Anishenko 1.VIII.1991, 1 ex. (cSha); Irkutsk, Ust'Kutsiky, 50 km SE Shumilovo, Shavrin & Enutschenko 24-25.VI. 2007, 2 exx.; same data, Shelekhov, Olkha, Shavrin 2.VIII.2005, 1 ex.; same data, 17

km Baikal lake, Shavrin 7.VII.1994, 2 exx. (cB); same data, Telezkoe lake, Altaysky Res., Erohov 17.VI.1979, 1 ex. (cU); same data, Tavshetskij, Solonezkoe lake, Shelaevo, 630 m, Shavrin 21-22.VI.2009, 1 ex. (cSol); SE Siberia, Taishetskij, Birjusa river, Patrikha, Shavrin 24.VI.1998, 1 ex. (cSha); Altai Rep., Turochakskij distr., Yailu vill., R. Dudko 29.VII.1994, 3 exx.; E Siberia, Khaman, Talsinka trib. of Snozhnyya river, A. Shavrin 5.VIII.2007, 3 exx. (cSol) same data, Khaman-Daban Mts, Snezhnaya R, valley, Talzinka river, Shavrin 5-8.VI.2007, 10 exx. (cSha), 2 exx. (cB); Central S Siberia, Barguzinskaya valley, 20 km E Ust' Barguzin, Gusikha, Shavrin 9.VIII.1997, 2 exx.; E Russia, Yakutia, Kamaz, Duban, 500 m, Osinovka river, Shavrin 25-31.VII.1995, 1 ex.; same data, Tankhoi, Osinovka river, Shavrin 24.VII.1995, 2 exx.; Crimea, Simferopol, Ghir valley, leg. ? 5.VIII.1957, 1 ex. (cSha). Ukraina, Uzgorod, Mukacevo, Dubrovin 14.VI.1973, 1 ex.; same data, Zakarpatsky, Tiacev, Koval 4-25.VII.1981, 2 exx.; same data, Svaliasky, Kerezki, 800 m, Koval 8.IV-24.VIII.1981, 6 exx. (cU), 1 ex. (cB); same data, Borgiava river, Koval 24-29.VII.1981, 1 ex.; same data, Beregovsky, Velikaya Bakta, Koval 8.X-4.XI.1981, 3 exx.; same data, Mukacevo, Ushakov 9.VII.21985, 5 exx. (cU), 1 ex. (cB); same data, Carpazi, Volovez, Pedetloka, Kurbatov 12.VIII.1986, 1 ex. (cU). Bielorusia, 3 km SE Vitebsk, Solodovnikov 9.VI-2.VII.1990, 1 ex.; same data, Vitebsk, Solodovnikov 10.IV.1994, 1 ex. (cSol); same data, 2-8.V.1989, 1 ex. (cSol), 1 ex. (cB). Moldavia, Ris. Codry, Ushakov 26.VI.1985, 1 ex. (cU), 1 ex. (cB); Kadry Lozovo, Janushev 3.VII.1968, 1 ex. (cU).

DISTRIBUTION – Europe, Tadjikistan (BORDONI, 1975; HERMAN, 2001; SMETANA, 2004; SEMIONENKOV *et al.*, 2015). CT, NT, ST, WS, ES. First record for Moldavia.

Xantholinus reitteri Coiffait, 1966

MATERIAL EXAMINED – Caucasus, (Georgia), Batumi, Ianushev 11.X.1974, 1 ex. (cB); Abkhazia, N Sukhumi, Ahalsheni, Ushakov 22.IX.1985, 1 ex.; same data, Gumistinsky Res., c/o Nizniy Zumur, Ushakov 25.IX.1985, 3 exx. (ZMNU), 2 exx. (cB); same data, Codor ridge, Muskvaza mouth, Ushakov 29.VI.1986, 1 ex. (cB). Azerbaijan, Mts Talish, 1 ex. (ZIN), 1 ex. (cB); same data, Aurora, leg. ? 7.V.1980, 1 ex (ZMNU).

DISTRIBUTION – Azerbaijan, Georgia (BORDONI, 1975; HERMAN, 2001; SMETANA, 2004).

Xantholinus laevigatus Jacobsen, 1849

MATERIAL EXAMINED – Russia, Petropoli (St. Petersburg), 1 ex. (cB); Mosca, Ryazan, 3 exx. (cB); Mosca, Tichomirova 2.VIII.1963-7.VII.1967, 13 exx.; Moskovskaya, 12 km Zvenigord, K. Khailov 14.VI-25.VIII.1991, 6 exx. (ZMNU), 2 exx. (cB); same data, Finashin 25.VIII.1973, 1 ex.; same data, Skhodja, Meshkov 13.VII.1984, 1 ex. (ZMNU), 1 ex. (cB); Central East Russia, Samara, Togliatti, I. Panteleev VI.1994, 1 ex. (cB); W Caucasus, Krasnaja Poljana, Roubal VII.1910, 1 ex. (MCSNM). Ukraina, Kiev, 4 km Krugli, Ushakov 25.VI.1988, 1 ex. (ZMNU), 1 ex. (cB); Zakarpatskaya, Svaliavsky, Kerezky, Borgoriava river, A. Koval 24.VI-3.X.1981, 2 exx.; same data, Sinevir lake, Sloboda river, 900 m, A. Koval 14.VIII.1981, 1 ex. (ZMNU).

DISTRIBUTION – Europe, Turkey (BORDONI, 1975; HERMAN, 2001; SMETANA, 2004; SEMIONENKOV *et al.*, 2015). CT, NT, ST.

NOTE – The structure of the inner sac of the aedeagus shows that the specimens from Russia identified in the past as *X. hochhuthi* Bordini, 1975 belong to the range of variability of *X. laevigatus*, so I propose the following synonymy: *Xantholinus laevigatus* Jacobsen, 1849 = *Xantholinus hochhuthi* Bordini, 1975, syn. n.

***Xantholinus haematodes* Kolenati, 1846**

EMATERIAL EXAMINED – Georgia, Gudauri, leg. ? 20.VIII.1929, 1 ex. (ZIN); Abkhazia, Sukhumi, Tichomirova ? 1.VII.1974, 1 ex. (cB). Caucasus, Teberda, Roubal, 1 ex. (MCSNT).

DISTRIBUTION – Azerbaijan, South Russia (BORDONI, 1975; HERMAN, 2001; SMETANA, 2004). ST. New record for Georgia.

***Xantholinus procerus* (Erichson, 1839)**

MATERIAL EXAMINED – SW Ukraina, Zakarpatskaya, Vinogradovsky, Tysa river, K. Koval 19.VII.1981, 10 exx. (ZMNU), 4 exx. (cB). Georgia, Abkhazia, Gagra, Belov 2-3.IX.1977, 2 exx. (ZMNU).

DISTRIBUTION – Romania, Azerbaijan, Georgia, South Russia (HERMAN, 2001; SMETANA, 2004). ST. New record for Ukraina.

***Xantholinus fortispunctatus* Motschulsky, 1860**

MATERIAL EXAMINED – S Russia, Rostov, Krasnosulinskij, Domleskhoz, E. Khachikov 2-4.V.1999, 1 ex. (cB); Crimea, Sebastopoli, 1 ex. (ZIN); Caucasus, Krasnodar, Kuban river, A. Solodovnikov 4.VII.1992, 1 ex. (cB); same data, Kutschevskij Distr., Krasnoe, D. Forminykh 18.VII.2007, 1 ex. (cSol). Azerbaijan, Baku, 1 ex. (ZMNU). Turkmenistan, Kopet Dag, Kara-Kola, 1 ex. (cB).

DISTRIBUTION – Eastern Europe, Western Asia (BORDONI, 1975 (sub *X. semirufus* Reitter, 1901); HERMAN, 2001; SMETANA, 2004; BORDONI, 2011).

***Xantholinus caucasicus* Bordini, 1975**

MATERIAL EXAMINED – Caucasus, Zaghedak, Bolshaya Saba river, 1700 m, V. Savinsky 17.VIII.1995, 1 ex.; same data, Brerpia river, 8-10 km NW Krasnaja Poljana, 800-900 m, V. Savinsky 7.VIII.1994, 1 ex. (cSol), 1 ex. (cB); same data, Acipse river, 7 km NW Krasnaja Poljana, 700 m, Solodov-nokov 27.VII.1994, 1 ex. (cSol); same data, Mzymta river, Krasnaja Poljana, 550-650 m, V. Savinsky 5.IX.1995, 1 ex. (cB); Teberda, Gonokhzoa and Klukhora gorge, D. Romashov 1.IX.1939, 1 ex. (ZMNU); Teberda, 1 ex.; N Caucasus, Karachevo-Cherkessia, Teberdinskij Res., Dvujazychnyi glacier, Y. Arzanov 20, VI.1997, 2 exx. (cB). North Ossetia, Ardon Tsei, 2000 m, A. Xlimenko 13.IX.1990, 2 exx. (cB).

DISTRIBUTION – Georgia, South Russia, Turkey (HERMAN, 2001; SMETANA, 2004; BORDONI, 2011). ST. First record for North Ossetia.

***Xantholinus (Helicophallus) samarensis* sp. n.**

MATERIAL EXAMINED – Holotype ♂: SE Russia, Samara, Sinyi Syrt, 13.5 km SW Kostino, I. Smeljansky VII-VIII. 1993 (cSol); paratypes: same data, 1 ♂ (cSOL), 2 ♂ (cB); same data, Pokhovistnevsky, Abdul-Zavod, Goreslavev 19.VIII.1995, 1 ♂ (cSOL).

DESCRIPTION – Length of body 9.7 mm; from anterior margin of head to posterior margin of elytra: 5 mm. Body (Fig. III, 3) entirely pale yellowish-brown. Head sub-ovoid, with rounded sides and broadly rounded posterior angles. Eyes very small and flat. Surface of head with traces of transverse micro-striation on the sides and fine, widely-spaced punctation, apart from on a median strip. Pronotum as long and wide as head, narrow posteriad, with oblique anterior margins, rounded anterior angles and sides not sinuate. Surface shiny, with dorsal series of 10-11 punctures and lateral series of 8-9 irregular punctures; other punctures near the anterior angles; all the punctures similar to those of head. Elytra a little shorter and much wider than pronotum, dilated posteriad, with slightly rounded humeral angles. Surface with fine, dense, shallow punctation, arranged in numerous series. Abdomen with very fine and dense, transverse micro-striation and fine, sparse punctation.

Tergite and sternite of the male genital segment as in Fig. II, 4-6. Aedeagus (Fig. III, 4) 1.1 mm long, with the long inner sac of usual shape; the spines of the distal series and the scales and spines of the rolled portion are large and dense.

ETYMOLOGY – The specific epithet refers to the type locality.

DISTRIBUTION – The species is known only from the cited localities.

NOTE – *X. samarensis* is the most north-eastern species of the western-central Russia belonging to the subgenus *Helicophallus* Coiffait, 1956. It differs from *X. araxis* Reitter, 1898 from Azerbaijan and Iran by its coloration, punctation, and structure of the larger and dense scales and spines of the inner sac of the aedeagus.

***Xantholinus (Helicophallus) vinicolor* Ushakov, 1989**

MATERIAL EXAMINED – N Ossetia, Dunta, A. Zamotaylov 2.VIII.1993, 2 ♂ (cSOL), 2 ♂ (cB).

REDESCRIPTION – Length of body 7.5 mm; from anterior margin of head to posterior margin of elytra: 3.5 mm. Body (Fig. III, 5) dark reddish-brown with infuscate head and paler elytra; antennae dark brown, legs brown. Head oblong, narrow anteriad, with slightly rounded sides and broadly rounded posterior angles. Eyes small and flat. Surface of head shiny, with traces of transverse micro-striation on the sides and deep and sparse punctation, apart from on a very wide median portion. Pronotum shiny, as long and wide as head, dilated posteriad, with narrowly rounded anterior angles and sides not sinuate. Surface with dorsal series of 8-9 fine, spaced punctures and lateral series of 6-7 irregular, more large punctures; some punctures near the widely- anterior angles. Elytra sub-rectangular, shorter and narrower than pronotum, a little dilated posteriad, with marked humeral angles. Surface wrinkled, with polygonal micro-reticulation and wide, superficial punctation, not

particularly dense (the distance between the punctures is equal to their diameter), arranged in numerous series. Abdomen with very fine and dense, transverse microstriation and fine, sparse punctation.

Tergite and sternite of the male genital segment as in Fig. II, 6-7. Aedeagus (Fig. III, 6) 1.25 mm long, with inner sac usually curled; the first portion of it covered with long, dense spines, after the distal two series of spines.

DISTRIBUTION – The species is known from Chechnya-Ingushetia. First record for North Ossetia. This is the first record since the description of the species.

NOTE – This species differs from *X. variabilis* Hochhuth, 1851 from Armenia and Georgia especially in the structure of the inner sac that in *X. variabilis* is covered with fine and sparse scales.

SEMIONENKOV *et al.*, 2015 also cites *Xantholinus (Helicophallus) distans* Mulsant & Rey, 1853 from Russia. The presence of this species in the south-western part of Russia is possible, considering its general distribution (cfr SMETANA, 2004).

Neoypnus mandschuricus (Bernhauer, 1923)

MATERIAL EXAMINED – Russian Far East, Janka Lake, Barabash-Levada, A. Gorodinski 1.VII.2004, 1 ex. (cH); Ussurijski, Ornoe Primorje, Kamenushka 1.VII.1980, 1 ex. (ZMNU).

DISTRIBUTION – Far East Russia, Liaoning, South Korea (KRYZHANOVSKII *et al.*, 1973 (sub. *X. ussuricus* Tikhomirova, 1973); HERMAN, 2001; SMETANA, 2004; BORDONI, 2003). FE.

NOTE – I have also seen this species from Korea, Mt Onjisan, Hwanghae Bukdo, leg. ? 21-27.1994, 2 exx. (cH), 1 ex. (cB). First record for North Korea.

ACKNOWLEDGEMENTS

I wish to thank the following colleagues for providing informations and the material to this study: I. Bucciarelli, in memoriam (Museo civico di Storia naturale, Milan), A. Colla (Museo civico di Storia naturale, Trieste), Y. Hayashi (Kawanishi), A. Putz (Eisenhüttenstadt), A. Ryvkin (Moscow), M. Schülke (Berlin), A. Shavrin (Institute of Systematic Biology, Daugavpils), A. Solodovnikov (Zoologisk Museum, København), G. Szél (Magyar Nemzeti Múzeum, Budapest), A. Tikhomirova, in memoriam (Zoological Institute, St. Petersburg), I. Ushakov (Moscow). A particular thank to A. Gusakov (Zoological Museum of Moscow University, Moscow) for the very copious material, and Anna Chernyaeva (Florence) for the translation of the very numerous localities labels in Cyrillic, and to S. Cuoco (Livorno) for the nice photos.

REFERENCES

BORDONI A., 1975 – *Studi sulla sistematica e la geonemia degli Xantholinus. VIII. Le specie eurocentroasiatiche e caucasiche in particolare. Revisione dei tipi e descrizione di nuove entità (Col. Staphylinidae)* - Memorie della Società entomologica italiana, 53 (1974): 56-96.

BORDONI A., 2000 – *Contribution to the knowledge of the Xantholinini from China. I (Coleoptera, Staphylinidae)* - Mitteilungen aus dem Museum für Naturkunde in Berlin (Zool. Reihe), 76: 121-133.

BORDONI A., 2003 – *Note su alcuni Xantholinini euroasiatici e descrizione di un nuovo genere della Manciuuria (Coleoptera Staphylinidae)* - Bollettino della Società entomologica italiana, 134, 3: 219-228.

BORDONI A., 2006 – *Contributo alla conoscenza degli Xantholinini della Cina. VI. Zeteotomus dilatipennis (Kirshenblatt, 1948) nello Shaanxi e descrizione di Erymus dalianus sp. n. dello Yunnan (Coleoptera: Staphylinidae)* - Animma x, 12: 16-23.

BORDONI A., 2009 – *Contribution to the knowledge of the Xantholinini of China. XIV. Nudobius linanensis n. sp. from Zhejiang, notes and new records on some interesting species (Insecta Coleoptera Staphylinidae)* - Quaderno di Studi e Notizie di Storia Naturale della Romagna, 28: 105-109.

BORDONI A., 2010 – *Male genitalia of palaearctic species of the genus Nudobius (Coleoptera, Staphylinidae)* - Bollettino della Società entomologica italiana, 142, 1: 19-26.

BORDONI A., 2011 – *Notes on Palaearctic Xantholinini. VI. New species and new records from Caucasus (Coleoptera, Staphylinidae)* - Fragmenta entomologica, 43, 1: 41-56.

GUSAROV V., 1989 – *Fauna i ekologiya stafilinid (Coleoptera, Staphylinidae) Kryma. Podsemeistva Metopsiinae, Proteinae, Omaliinae, Piestinae, Tachyporinae, Oxyporinae, Steninae, Paederinae, Xantholininae, Staphylininae (in Russian)* - Vestnik Leningradskogo Universiteta. Biologii, 3: 3-17.

HERMAN L.H., 2001 – *Catalogue of the Staphylinidae (Insecta: Coleoptera). 1758 to the end of the second millennium. Part VI. Staphylininae Group (Part 3). Staphylininae: Staphylinini (Quediina, Staphylinina, Tanygnathinina, Xanthopygina), Xantholinini. Staphylinidae Icetae sedis, Fossils, Protactinae†* - Bulletin of the American Museum of Natural History, 265: 3021-3840.

JACOBSON G.G., 1908-1909 – *Il semeistvo Staphylinidae, stafiliny* - Devrien. St-Petersburg, 6-7: 443-569.

KRYZHANOVSKII O.L., TIKHOMIROVA A.L., FILATOVA L.D., 1973 – *Staphylinidy (Coleoptera, Staphylinidae) Yuzhnogo Primor'ia* - In Ekologia pochvennykh bezepoozvononochnykh, Moskva: 144-173.

SEMIONENKOV O.J., SEMENOV V.B., GILDENKOV M.Y., 2015 – *Rove beetles (Coleoptera: Staphylinidae) of the West of the European part of Russia (excepting subfamilies Pselaphinae, Scydmeninae and Scaphidiinae)* - Smolensk. Universum, 392 pp.

SOLODOVNIKOV A.Y., 1998 – *Fauna of Staphylinidae (Coleoptera, Staphylinidae) of the North Western Caucasus. Subfamilies Staphylininae, Xantholininae, Paederinae, Steninae and Oxytelinae (in Russian)* - Entomologicheskoe Obozrenie, 77 (2): 331-354.

SMETANA A., 1982 – *Remarks on some Xantholinini from the Kuril Islands (Coleoptera: Staphylinidae)* - The Coleopterist Bulletin, 36, 2: 309-313.

SMETANA A., 2004 – *Staphylinidae*, pp. 237-698 - In I. Löbl & A. Smetana (editors): Catalogue of Palaearctic Coleoptera. vol. 2 - Stenstrup: Apollo Books, 942 pp.

USHAKOV I.A., 1989 – *K poznaniyu Xantholininae (Coleoptera, Staphylinidae) fauny SSSR* - Zoologicheskii Zhurnal, 68: 127-130.

64 - Blank Page