Sex-related differences in predicting choledocholithiasis using current American Society of Gastrointestinal Endoscopy risk criteria

Ankit Chhoda^a, Deepanshu Jain^b, Shashideep Singhal^c

Yale-Waterbury Internal Medicine Program, Yale School of Medicine, Waterbury Connecticut; Albert Einstein Medical Center, Philadelphia; University of Texas Health Science Center at Houston, Houston, Texas, USA

Abstract

Background American Society of Gastrointestinal Endoscopy (ASGE) criteria is widely used to predict probability of choledocholithiasis among patients with symptomatic gallstone disease but does not take sex into consideration.

Methods The cohort study included patients who underwent either endoscopic retrograde cholangiopancreatography or intraoperative cholangiography for suspected choledocholithiasis at our medical center. Clinical, laboratory and radiological data were collected for each patient and used to stratify them based on the ASGE risk criteria for choledocholithiasis.

Results A total of 646 patients composed the final study cohort. The composite incidence rate of choledocholithiasis for male and female groups was 47.5% and 46.2% respectively. Total bilirubin (>4 mg/dL), cholangitis, abnormal liver function tests (transaminitis), dilation of the common bile duct (>6 mm) on ultrasound examination, and biliary pancreatitis were individually associated with choledocholithiasis in 73.5%, 68.4%, 61.1%, 60.0%, and 51.7% of males, respectively. Total bilirubin >4 mg/dL and 1.8-4 mg/dL, cholangitis, and transaminitis were individually associated with choledocholithiasis in 82.6%, 64.0%, 58.2%, and 50.0% of females, respectively. The distribution in probability group/incidence of choledocholithiasis was as follows: for males, low probability 3.3%/0%, intermediate probability 56.7%/33.8%, high probability 40%/77.8%; and for females, low probability 5.3%/14.3%, intermediate probability 70.2%/39.6%, high probability 24.5%/72.1%.

Conclusions The composite incidence for choledocholithiasis was similar across male and female patients. A significantly higher proportion of females compared to males were in the intermediate probability group. Better sex stratification can help improve the positive and negative predictive values of ASGE risk stratification criteria. Improve patient outcomes and reduce associated healthcare cost.

Keywords Gallstones, choledocholithiasis, sex distribution

Ann Gastroenterol 2018; 31 (1): 1-6SS

^aDepartment of Internal Medicine, Yale-Waterbury Internal Medicine Program, Yale School of Medicine, Waterbury Connecticut (Ankit Chhoda);

^bDivision of Gastroenterology and Hepatology, Department of Internal Medicine, Albert Einstein Medical Center, Philadelphia, PA (Deepanshu Jain); ^cDivision of Gastroenterology, Hepatology and Nutrition, University of Texas Health Science Center at Houston, Houston, Texas (Shashideep Singhal), USA

Conflict of Interest: None

Correspondence to: Shashideep Singhal, MD, Division of Gastroenterology, Hepatology and Nutrition, University of Texas Health Science Center at Houston, 6431 Fannin, MSB 4.234, Houston, Texas, USA 77030, e-mail: sdsinghal@gmail.com

Received 31 July 2017; accepted 4 September 2017; published online 10 October 2017

DOI: <https://doi.org/10.20524/aog.2017.0198>

Introduction

Gallstone disease is a widespread disorder affecting 20 million Americans. It is diagnosed by demonstration of a stone in the gallbladder using transabdominal ultrasound (US) or, based on a history of cholecystectomy [1]. Female sex has been identified as one of the major risk factor for gallstone disease [2,3]. The increased incidence, however, decreases by the fifth decade and the incidence in both sexes becomes comparable [4].

Along with cholecystitis, cholangitis, biliary pancreatitis and choledocholithiasis are complications of gallstone disease, caused either by migration of the stone from the gallbladder, or by intraductal precipitation due to bile stasis, biliary diverticulum or a large bile duct. It has been estimated that the frequency of stones in the common bile duct (CBD) during cholecystectomy ranges from 5-20%; however, the exact prevalence remains unknown [5-8].

The criteria of the American Society of Gastrointestinal Endoscopy (ASGE) are widely used for the prediction of choledocholithiasis and management based on risk stratification (Table 1) [9]. The aim of our study was to determine sex-related differences in the prediction of choledocholithiasis in patients with symptomatic gallstones.

Patients and methods

Study location

University hospital and a community hospital under the University of Texas Health Science Center, Houston.

Table 1 ASGE recommended criteria for predicting choledocholithiasis

Predictors of choledocholithiasis	
Very strong	
<ul style="list-style-type: none"> • CBD stone on transabdominal US • Clinical ascending cholangitis • Bilirubin >4 mg/dL 	
Strong	
<ul style="list-style-type: none"> • Dilated CBD on US (6 mm with gallbladder <i>in situ</i>) • Bilirubin level 1.8-4 mg/dL 	
Moderate	
<ul style="list-style-type: none"> • Abnormal liver biochemical test other than bilirubin • Age >55 years • Clinical gallstone pancreatitis 	
Assigning a likelihood of choledocholithiasis based on clinical predictors	
• Presence of any very strong predictor	High
• Presence of both strong predictors	High
• No predictors present	Low
• All other patients	Intermediate

ASGE, American Society of Gastrointestinal Endoscopy; CBD, common bile duct; US, ultrasound examination

Study duration

The study covered the 3 years from November 1, 2012 to August 1, 2015.

Inclusion criteria

1. Patients undergoing endoscopic retrograde cholangiopancreatography (ERCP) for suspected choledocholithiasis: high probability based on ASGE criteria, or patients with intermediate probability where immediate cholecystectomy was not feasible due to their medical condition.
2. Patients who underwent intraoperative cholangiography (IOC) at the time of cholecystectomy: low/intermediate probability for choledocholithiasis. All patients with a positive IOC subsequently underwent ERCP.

IOC during cholecystectomy is a reliable modality recommended in ASGE guidelines for the evaluation of choledocholithiasis [9]. At our institution, most patients with intermediate or low probability for choledocholithiasis underwent IOC routinely during cholecystectomy. In our experience, this approach can save time and avoid additional investigations, such as endoscopic US or magnetic resonance cholangiopancreatography. Only a small proportion of patients with intermediate probability who were at high risk for cholecystectomy underwent ERCP as a temporary measure, while awaiting medical optimization for cholecystectomy. Fig. 1 illustrates the numbers of patients in each category.

Exclusion criteria

Patients who were being followed up for incomplete stone clearance from previous ERCP and stent replacement were excluded.

Figure 1 Study flow diagram
CBD, common bile duct

Data collection

Clinical data at presentation (age, sex, symptomatology, laboratory and imaging parameters) and procedural (IOC and ERCP) details with complications were collected. The pre-ERCP imaging, which included abdominal US, was reviewed to collect data regarding CBD stone visualization with acoustic shadowing and CBD diameter.

Methodology

Our study was retrospective and observational in nature. After the relevant data had been collected, the study cohort was divided into two groups based on sex. Each group was further stratified into low-, intermediate- or high-risk category for choledocholithiasis based on the ASGE risk factors. The true incidence rate for choledocholithiasis for each category in our study was then compared with the predicted risk to ascertain any clinically significant differences.

Statistical analysis

Statistical analysis was performed using SPSS version 23.0. Quantitative data were represented as mean $\pm$ standard deviation and qualitative data as numbers and percentages. Statistical differences were investigated using the *t*-test for continuous variables and the chi-square test for categorical variables. Any difference or association between two variables with a *P*-value <0.05 was considered to be statistically significant. Beta error was set at 20% and hence power at 80%.

Variable definition

Ascending cholangitis was defined as fever (temperature $>37.8^{\circ}\text{C}$) with abdominal pain and jaundice. Biliary pancreatitis was defined as elevated lipase levels (three times

the institutional cutoff) in the absence of other risk factors, such as alcohol ingestion or medications associated with pancreatitis.

ERCP procedure

The ERCP was performed using Olympus TJF 160 VR or TJF 145 side-viewing endoscopes. All patients provided written consent to undergo ERCP and were informed of the risks and benefits of the procedure. Patients underwent propofol sedation assisted by an anesthesiologist or conscious sedation from midazolam or ketamine administered by the endoscopist. An initial cholangiogram was performed to determine the maximum diameter of the CBD and the number and size of the stones. The patients underwent endoscopic sphincterotomy over a guidewire. The stones were retrieved via a balloon catheter or a Dormia basket. The bile duct clearance was documented with a balloon catheter occlusion cholangiogram at the end of the procedure. A biliary stent was placed for residual CBD stone and follow-up ERCP was scheduled. The patients were followed up for post-procedural complications on the following day.

Results

A total of 646 patients satisfied the inclusion and exclusion criteria. The patients in our study were predominantly female ($n=526$) rather than male ($n=120$). The median age was significantly less for males (36.3 years) than for females (49 years) ($P<0.0001$). ASGE risk factors among male and female patients were also compared for the incidence of CBD stones (Table 2). None of the risk factors showed a significant sex-related difference in the incidence of CBD. Logistic univariate analysis was performed to determine the association of the risk factors with choledocholithiasis for both sexes (Supplemental Tables 1 & 2). Apart from abnormal liver function tests (LFTs) among males and dilated CBD

Table 2 Distribution of ASGE risk factors across male and female patients

ASGE risk factors	Male patients (n=120)		Female patients (n=526)		P-value risk factors	P-value CBD stone incidence
	Total patients (incidence)	CBD stones (incidence)	Total patients (incidence)	CBD stones (incidence)		
Cholangitis	19 (15.83%)	13 (68.4%)	55 (10.45%)	32 (58.2%)	0.179	0.59
Total bilirubin >4 mg/dL	34 (28.3%)	25 (73.5%)	46 (8.74%)	38 (82.6%)	<0.0001	0.41
Total bilirubin 1.8-4 mg/dL	31 (25.83%)	15 (48.4%)	100 (19.01%)	64 (64.0%)	0.09	0.14
Biliary pancreatitis	29 (24.16%)	15 (51.7%)	132 (25.09)	64 (48.5%)	0.907	0.84
Total bilirubin <1.8 mg/dL	55 (45.83%)	17 (30.9%)	380 (72.24%)	139 (36.6%)	<0.001	0.45
CBD size on US >6 mm	30 (25%)	18 (60.0%)	280 (53.23%)	113 (40.4%)	<0.001	0.051
Abnormal LFT	72 (60%)	44 (61.1%)	414 (78.7%)	207 (50%)	0.00018	0.09
Age >55 years	38 (31.67)	18 (47.37%)	445 (84.6%)	198 (44.5%)	<0.0001	0.74

ASGE, American Society of Gastrointestinal Endoscopy; CBD, common bile duct; LFT, liver function test; US, ultrasound examination

among females, all risk factors lacked a statistically significant association. Male patients with abnormal LFTs had significantly lower odds (odds ratio [OR] 0.33, 95% confidence interval [CI]=0.131- 0.827, $P=0.02$) for choledocholithiasis, whereas female patients with a CBD size >6 mm had significantly higher odds (OR 1.9, 95% CI=1.135-3.048, $P=0.01$) for choledocholithiasis. These parameters were further subjected to receiver operator curve analysis to evaluate their accuracy in determining choledocholithiasis in the two sexes. Among all the parameters, CBD size on US was found to have a significant area under the curve (area under the curve 0.786; 95%CI 0.644-0.927; $P=0.006$). A cutoff of 4.5 mm for CBD size on US would have 65% sensitivity and 78% specificity, compared to 35% sensitivity and nearly 100% specificity for a 6 mm cutoff.

The patients included in the study were divided into high, intermediate and low likelihood for choledocholithiasis based on the ASGE criteria (Table 3). Of composite study population, 40.0% males and 24.5% females were categorized as high risk. In the high-likelihood category, 70.8% of males and 72.1% of females had CBD stones, reflecting the high accuracy of the ASGE criteria in predicting CBD stones for this category ($P=0.85$). The majority of males (56.7%) and females (70.2%) were categorized as having intermediate risk ($P=0.0025$). In the intermediate-likelihood category, 33.8% of male and 39.6% of female patients had CBD stones ($P=0.42$). A very small fraction of the study group was categorized as low likelihood for both males (3.3%) and females (5.3%); 0% of males and 14.3% of females in this category were found to have CBD stones ($P=0.99$).

Discussion

Choledocholithiasis is a common problem in the USA and other western cultures. The burden of disease secondary to stones in the CBD is significant. Significant morbidity and mortality is associated with complications of CBD stones, including conditions such as cholangitis and biliary pancreatitis. Several studies have questioned the real world utility of the ASGE criteria in predicting choledocholithiasis [10,11]. Female sex is a known risk factor for gallstone disease, but the ASGE risk criteria for predicting the likelihood of choledocholithiasis do not take sex into consideration. Our cohort study aimed to determine any sex-related differences in the prediction of choledocholithiasis using the ASGE risk criteria.

The composite incidence of choledocholithiasis for males (47.5%) and females (46.2%) differed only minimally. The incidence of choledocholithiasis was higher for female patients than for males across the low (14.3% vs. 0%), intermediate (39.6% vs. 33.8%) and high (72.1% vs. 70.8%) risk ASGE categories ($P>0.05$). Five out of eight ASGE risk factors among males, in contrast to four of eight among females, were individually associated with a $>50\%$ incidence of choledocholithiasis. Total bilirubin >4 mg/dL, cholangitis, abnormal LFTs (transaminitis), CBD dilation >6 mm on US, and biliary pancreatitis were individually associated with choledocholithiasis in 73.5%, 68.4%, 61.1%, 60.0%, and 51.7% of males, respectively. The association was not in the same order for the female cohort. Total bilirubin >4 and 1.8-4 mg/dL, cholangitis, and transaminitis were individually associated with choledocholithiasis in 82.6%, 64.0%, 58.2%, and 50.0% of females, respectively. No statistically significant difference between male and female patients was observed in the incidence of choledocholithiasis across individual ASGE-defined risk factors. Despite the lack of statistical significance, these trends suggest there is a need to re-evaluate the strength of association of individual risk factor for choledocholithiasis separately for males and females. A study by Suarez *et al* showed that a second set of laboratory tests failed to improve the diagnostic accuracy of ASGE criteria in predicting choledocholithiasis [10]. In addition, the authors also concluded that even a significant decline in LFTs did not reliably predict spontaneous stone passage [10].

Further stratification based on the ASGE risk factors for choledocholithiasis revealed an interesting distribution pattern across the low-, intermediate- and high-likelihood categories for each sex. The majority of male (56.7%) and female (70.2%) patients with suspected CBD stones were categorized by the ASGE criteria as having intermediate likelihood. More than one third of these patients (33.8% of males, 39.8% of females) were eventually found to have CBD stones. This casts doubt upon the true utility of these criteria in helping physicians decide the best next step in the clinical workup of these patients, potentially delaying therapy and increasing the risk of complications in this, the largest subgroup. More than 70% of subjects (either sex) in the high-likelihood group were found to have CBD stones, clearly demonstrating the high specificity of the ASGE criteria for this subgroup. Interestingly, these criteria showed a high negative predictive value for CBD stones among male patients classified as low likelihood, but failed to do the same for the respective female cohort. A small but significant

Table 3 Incidence of CBD stones across ASGE risk categories (for choledocholithiasis) in male and female patients

ASGE risk strata	Male patients		Female patients		P-value for CBD stone incidence
	Total patients (n=120)	CBD stone incidence (47.5%)	Total patients (n=526)	CBD stone incidence (46.2%)	
Low likelihood	4 (3.3%)	0	28 (5.3%)	4 (14.3%)	0.99
Intermediate likelihood	68 (56.7%)	23 (33.8%)	369 (70.2%)	146 (39.6%)	0.42
High likelihood	48 (40.0%)	34 (70.8%)	129 (24.5%)	93 (72.1%)	0.85

ASGE, American Society of Gastrointestinal Endoscopy; CBD, common bile duct

Summary Box

What is already known:

- Gallstone disease is a widespread disorder
- Female sex is one of the major risk factors for gallstone disease
- American Society of Gastrointestinal Endoscopy (ASGE) criteria are widely used to risk stratify patients with symptomatic cholelithiasis for concomitant choledocholithiasis
- Current ASGE criteria do not take sex into account in predicting the risk of choledocholithiasis

What the new findings are:

- The incidence of choledocholithiasis was higher for female patients than for male patients across the low-, intermediate- and high-likelihood ASGE categories
- The ASGE criteria stratify the majority of patients as intermediate risk, with more than one third found to have choledocholithiasis
- The ASGE criteria showed a high negative predictive value for choledocholithiasis among the low-risk category for males, but not for females

percentage (14.8%) of females in the low-likelihood group were found to have CBD stones, which again puts these criteria in question.

Our study had some limitations. Its retrospective nature limited our ability to extract information regarding length of hospital stay, or additional data from modalities such as endoscopic US or magnetic resonance cholangiopancreatography. Despite the large sample size, the trends we observed failed to reach statistical significance. In our study, the observed incidence of CBD stones was similar for both sexes. However, a sex-related difference in the association of clinical, radiological and laboratory risk factors with the risk

of choledocholithiasis was noted. The current ASGE criteria, which do not take sex into consideration, predominantly stratifies both male and female patients as having intermediate risk, which in the majority of cases fails to help the physician manage these patients in the real world. Larger prospective studies are needed to improve the current ASGE criteria so as to achieve better risk stratification of patients with suspected choledocholithiasis, improve patient outcomes, and reduce the associated healthcare costs.

References

1. Everhart JE, Khare M, Hill M, Maurer KR. Prevalence and ethnic differences in gallbladder disease in the United States. *Gastroenterology* 1999;**117**:632-639.
2. Völzke H, Baumeister SE, Alte D, et al. Independent risk factors for gallstone formation in a region with high cholelithiasis prevalence. *Digestion* 2005;**71**:97-105.
3. Diehl AK. Epidemiology and natural history of gallstone disease. *Gastroenterol Clin North Am* 1991;**20**:1-19.
4. Jørgensen T. Gallstones and plasma lipids in a Danish population. *Scand J Gastroenterol* 1989;**24**:916-922.
5. Collins C, Maguire D, Ireland A, Fitzgerald E, O'Sullivan GC. A prospective study of common bile duct calculi in patients undergoing laparoscopic cholecystectomy: natural history of choledocholithiasis revisited. *Ann Surg* 2004;**239**:28-33.
6. Prat F, Meduri B, Ducot B, Chiche R, Salimbeni-Bartolini R, Pelletier G. Prediction of common bile duct stones by noninvasive tests. *Ann Surg* 1999;**229**:362-368.
7. O'Neill CJ, Gillies DM, Gani JS. Choledocholithiasis: overdiagnosed endoscopically and undertreated laparoscopically. *ANZ J Surg* 2008;**78**:487-491.
8. Petelin JB. Laparoscopic common bile duct exploration. *Surg Endosc* 2003;**17**:1705-1715.
9. Maple JT, Ben-Menachem T, Anderson MA, et al; ASGE Standards of Practice Committee. The role of endoscopy in the evaluation of suspected choledocholithiasis. *Gastrointest Endosc* 2010;**71**:1-9.
10. Nárvaez Rivera RM, González González JA, Monreal Robles R, et al. Accuracy of ASGE criteria for the prediction of choledocholithiasis. *Rev Esp Enferm Dig* 2016;**108**:309-314.
11. Suarez AL, LaBarre NT, Cotton PB, Payne KM, Coté GA, Elmunzer BJ. An assessment of existing risk stratification guidelines for the evaluation of patients with suspected choledocholithiasis. *Surg Endosc* 2016;**30**:4613-4618.

Supplement Tables

Supplemental Table 1 Logistic univariate analysis of ASGE risk factors for choledocholithiasis among male patients

Parameters	Coefficient	Standard error	P-value	Odds ratio	95% confidence interval
Age >55 years	−0.485	0.763	0.525	0.616	0.138-2.749
Total bilirubin	0.208	0.185	0.261	1.231	0.857-1.767
Abnormal LFT	1.113	0.471	0.018	0.329	0.131-0.827
Biliary pancreatitis	0.098	0.505	0.846	1.103	0.410-2.970
CBD size on US >6 mm	0.089	0.068	0.192	1.093	0.956-1.250
Cholangitis	−0.778	0.508	0.126	0.459	0.170-1.244

ASGE, American Society of Gastrointestinal Endoscopy; CBD, common bile duct; LFT, liver function test

Supplemental Table 2 Logistic univariate analysis of ASGE risk factors for choledocholithiasis among female patients

Parameters	Coefficient	Standard error	P-value	Odds ratio	95% confidence interval
Age >55 years	0.968	0.726	0.183	2.632	0.634-10.927
Total bilirubin	0.019	0.137	0.891	1.019	0.779-1.332
Abnormal LFT	−0.926	0.910	0.309	0.396	0.067-2.359
Biliary pancreatitis	−3.93	0.768	0.618	0.682	0.151-3.070
CBD size on US >6 mm	0.621	0.252	0.014	1.860	1.135-3.048
Cholangitis	0.912	1.103	0.409	2.4889	0.286-21.66

ASGE, American Society of Gastrointestinal Endoscopy; CBD, common bile duct; LFT, liver function test