New Banach Sequence Spaces That Is Defined By The Aid Of Lucas Numbers

Murat KARAKAŞ¹, Ayşe Metin KARAKAŞ¹

ABSTRACT: In this work, we establish a new matrix by using Lucas numbers and define a new sequence space. Besides, we give some inclusion relations and investigate the geometrical properties such as Banach-Saks type, weak fixed point property for this space.

Keywords: Banach-Saks property, difference matrix, difference sequence spaces, lucas numbers, weak fixed point property

Lucas Sayıları Yardımıyla Tanımlanan Yeni Banach Dizi Uzayları

ÖZET: Bu makalede, Lucas sayılarını kullanarak yeni bir matris oluşturuyoruz ve yeni bir dizi uzayı tanımlıyoruz. Ayrıca bu uzay için bazı kapsama bağıntıları veriyoruz ve uzayın p tipi Banach-Saks, zayıf sabit nokta gibi geometrik özelliklerini araştırıyoruz.

Anahtar Kelimeler: Banach-Saks özelliği, fark matrisi, fark dizi uzayları, lucas sayıları, zayıf sabit nokta özelliği

¹ Murat KARAKAŞ (0000-0003-1968-2040), Ayşe Metin KARAKAŞ (0000-0003-3552-0105), Bitlis Eren Üniversitesi, Fen Edebiyat Fakültesi, İstatistik, Bitlis, Türkiye
Sorumlu yazar/Corresponding Author: Murat KARAKAŞ, mkarakas@beu.edu.tr

INTRODUCTION

Recently, (Kara, 2013) has defined and studied Fibonacci difference sequence spaces. According to his study, we have examined the Lucas difference sequence space and have shown that this space is a Banach -space and also is linear isomorphic to ℓ_p for $1 \leq p \leq \infty$.

All real and complex valued sequences are represented by ω . A sequence space is linear subspace of ω . Throughout the paper $\ell_\infty, \ell_p (1 \leq p < \infty), c, c_0$,

characterise the spaces of all bounded, p -absolutely summable, convergent and null sequences.

Let $B = (b_{nk})$ be an infinite matrix of real numbers $b_{nk} (n, k = 1, 2, \dots)$ and X, Y be two sequence spaces. It is said that the matrix B describes a matrix transformation from X into Y , and we symbolize it $B: X \rightarrow Y$, if the sequence $Bx = (B_n(x))$ is in Y for every $x = (x_k) \in X$ where

$$B_n(x) = \sum_k b_{nk} x_k \quad (1)$$

The sequence Bx is said to be the B -transform of x by the matrix B . The notation (X, Y) shows the class of matrices B such that $B: X \rightarrow Y$. Therefore $B \in (X, Y)$ iff the series on the equality (1) converges for each $n \in \mathbb{N}$ and every $x \in X$, and $Bx \in Y$ for all $x \in X$.

The matrix domain of B for a sequence space X is given by

$$X_B = \{x \in \omega: Bx \in X\} \quad (2)$$

which is a sequence space.

For the sequence whose n^{th} term is e_n and others are 0 for each $n \in \mathbb{N}$, we'll write $e^{(n)}$ and also use $e = (1, 1, \dots)$.

Recently, several authors have made use of the view of constituting sequence space by the help of matrix domain of an infinite triangle matrix, e.g., (Mursaleen and Noman, 2010; Mursaleen and Noman, 2010; Mursaleen and Noman, 2011), (Mursaleen et al., 2006), (Altay et al., 2006), (Başar and Altay, 2003), (Altay and Başar, 2005), (Savaş et al., 2009), (Karakas, 2015), (Kara and Başarır, 2012).

The matrix domain λ_Δ is called the difference sequence space if λ is a normed or paranormed sequence space where Δ symbolizes the backward difference matrix $\Delta = (\Delta_{nk})$ and $\Delta' = (\Delta'_{nk})$ symbolizes the transpose of the matrix Δ , the forward difference matrix, which are identified by

$$\Delta_{nk} = \begin{cases} (-1)^{n-k}, & n \geq k \geq n-1 \\ 0, & k > n \text{ or } 0 \leq k < n-1 \end{cases}$$

and

$$\Delta'_{nk} = \begin{cases} (-1)^{n-k}, & n \leq k \leq n+1 \\ 0, & k > n+1 \text{ or } 0 \leq k < n \end{cases}$$

for all $k, n \in \mathbb{N}$; respectively. (Kızmaz, 1981) first presented the concept of difference sequence spaces and determined the following sequence spaces:

$$X(\Delta) = \{x \in w: (x_k - x_{k+1}) \in X\}, \quad X = \ell_\infty, c, c_0.$$

(Altay and Başar, 2007) examined the difference sequence space bv_p which involves sequences (x_k) such that $(x_k - x_{k-1})$ in the case $0 < p < 1$. For $1 \leq p \leq \infty$, (Çolak et al., 2004) examined this difference space. In addition, authors analyzed certain difference sequence spaces, see (Et, 1993), (Mursaleen, 1996), (Gaur and Mursaleen, 1998), (Et and Esi, 2000), (Et and Çolak, 1995), (Et and Başarır, 1997), (Karakaş et al., 2013).

The primary aim of this note is to establish the Lucas difference matrix $\hat{E}$ and introduce new sequence spaces $\ell_p(\hat{E})$ and $\ell_\infty(\hat{E})$ connected with matrix domain of $\hat{E}$ for sequence spaces ℓ_p and ℓ_∞ where $1 \leq p < \infty$.

In the second part of this paper, we give some formulas and basic notions about Lucas sequence and a BK -space. In the third part, by using Lucas numbers, we introduce the sequence spaces $\ell_p(\hat{E})$ and $\ell_\infty(\hat{E})$, and establish some inclusion relations. Also, we construct the basis of $\ell_p(\hat{E})$ for $1 \leq p < \infty$. Finally, we analyze some geometric properties for the space $\ell_p(\hat{E})$, $1 \leq p < \infty$.

$$\sum_{i=0}^n L_i = L_{n+2} - 1,$$

$$\sum_{i=1}^n L_i^2 = L_n L_{n+1} - 2,$$

MATERIAL AND METHODS

A K -space is a sequence space with a linear topology provided each maps $\tau_k: X \rightarrow \mathbb{C}$, $\tau_k(x) = x_k$ is continuous for all $k \in \mathbb{N}$. If a K -space X is a complete linear metric space, then X is called a FK -space. A Banach K -space is a normed FK -space. The sequence spaces ℓ_∞, c, c_0 are Banach -spaces with the norm $\|x\|_\infty = \sup_k |x_k|$. Also ℓ_p , is a Banach -space with $\|x\|_{\ell_p} = (\sum_k |x_k|^p)^{1/p}$, $1 \leq p < \infty$.

For a normed linear space X , the sequence $(b^{(n)})_{n=0}^\infty$ is called Schauder basis if there is a unique sequence $(\gamma_n)_{n=0}^\infty$ of scalars such that $x = \sum_{n=0}^\infty \gamma_n b^{(n)}$ for every $x \in X$, that is,

$$\lim_{m \rightarrow \infty} \|x - \sum_{n=0}^m \gamma_n b^{(n)}\|_X = 0$$

The Lucas sequence (L_n) is defined by using Fibonacci recurrence relation and different initial conditions as:

$$L_0 = 2, L_1 = 1 \text{ and } L_n = L_{n-1} + L_{n-2}, n \geq 2.$$

Lucas numbers are closely related to Fibonacci numbers. In spite of this relation, they exhibit distinct properties. Some fundamental characteristics of Lucas sequences are given as follows (Koshy, 2001), (Vajda, 2008):

$$\lim_{n \rightarrow \infty} \frac{L_n}{L_{n-1}} = \varphi \text{ (golden section),}$$

$$L_{n-1}L_{n+1} - L_n^2 = -5 \cdot (-1)^n.$$

It can be easily derived by placing in the last equality that $L_{n-1}^2 + L_n L_{n-1} - L_n^2 = -5(-1)^n$.

RESULTS AND DISCUSSION

In the present section, we'll firstly give the following new double band matrix $\hat{E} = (\hat{L}_{nk})$ by means of the sequence (L_n) of Lucas numbers for all $n, k \in \mathbb{N} - \{0\}$:

$$\hat{L}_{nk} = \begin{cases} -\frac{L_n}{L_{n-1}}, & k = n-1 \\ \frac{L_{n-1}}{L_n}, & k = n \\ 0, & \text{other} \end{cases}$$

The inverse of the above Lucas matrix can be easily calculated and it is given by

$$\hat{L}^{-1} = \begin{cases} \frac{L_n^2}{L_{k-1}L_k}, & n \geq k > 0 \\ 0, & n < k \end{cases}.$$

Now, we define the $\hat{E}$ -transform of a sequence $x = (x_n)$ in the form:

$$y_n = \hat{E}_n(x) = \frac{L_{n-1}}{L_n} x_n - \frac{L_n}{L_{n-1}} x_{n-1}, \quad n \geq 1. \quad (3)$$

And now, let's introduce the Lucas difference sequence spaces $\ell_p(\hat{E})$ and $\ell_\infty(\hat{E})$ in the form of

$$\ell_p(\hat{E}) = \left\{ x \in w: \sum_n \left| \frac{L_{n-1}}{L_n} x_n - \frac{L_n}{L_{n-1}} x_{n-1} \right|^p < \infty \right\}, \quad 1 \leq p < \infty$$

$$\ell_\infty(\hat{E}) = \left\{ x \in w: \sup_n \left| \frac{L_{n-1}}{L_n} x_n - \frac{L_n}{L_{n-1}} x_{n-1} \right| < \infty \right\}.$$

These spaces can be redefined with the help of equality (2) by

$$\ell_p(\hat{E}) = (\ell_p)_{\hat{E}}, \quad 1 \leq p < \infty \text{ and } \ell_\infty(\hat{E}) = (\ell_\infty)_{\hat{E}}. \quad (4)$$

Theorem 3.1. The sets $\ell_p(\hat{E})$ and $\ell_\infty(\hat{E})$ are Banach -spaces with

$$\|x\|_{\ell_p(\hat{E})} = \left(\sum_n |\hat{E}_n(x)|^p \right)^{1/p}, 1 \leq p < \infty \text{ and } \|x\|_{\ell_\infty(\hat{E})} = \sup_n |\hat{E}_n(x)|.$$

Proof. By the Theorem 4.3.12 of (Wilansky, 1984), we obtain that $\ell_p(\hat{E})$ and $\ell_\infty(\hat{E})$ are Banach K -spaces with the above norms since the matrix is triangle and equality (4) holds.

$\ell_p(\hat{E})$ and $\ell_\infty(\hat{E})$ are the sequence spaces of non-absolute type. So indeed,

$$\|x\|_{\ell_p(\hat{E})} \neq \| |x| \|_{\ell_p(\hat{E})} \text{ and } \|x\|_{\ell_\infty(\hat{E})} \neq \| |x| \|_{\ell_\infty(\hat{E})}.$$

This means that the absolute property is not valid on the spaces $\ell_p(\hat{E})$ and $\ell_\infty(\hat{E})$ for fewest one sequence where $|x| = (|x_k|)$.

Theorem 3.2. The Lucas difference sequence space $\ell_p(\hat{E})$ of non-absolute type is linear isomorphic to ℓ_p for $1 \leq p \leq \infty$.

Proof. Let $1 \leq p \leq \infty$ and consider the transformation Z from $\ell_p(\hat{E})$ to ℓ_p defined by $x \rightarrow y = Zx$ with the notation of equality (3). Then, we have $Zx = y = \hat{E}x \in \ell_p$ for every $x \in \ell_p(\hat{E})$. It is trivial that Z is linear. Additionally, it is easy to see that if $x = 0$ and so $Zx = 0$ is injective.

Besides, let $y = (y_k) \in \ell_p$ and consider the sequence $x = (x_k)$ by

$$x_k = \sum_{j=1}^k \frac{L_k^2}{L_{j-1}L_j} y_j. \quad (5)$$

Next, we obtain that

$$\begin{aligned} \|x\|_{\ell_p(\hat{E})} &= \left(\sum_k \left| \frac{L_{k-1}}{L_k} x_k - \frac{L_k}{L_{k-1}} x_{k-1} \right|^p \right)^{\frac{1}{p}} \\ &= \left(\sum_k \left| \frac{L_{k-1}}{L_k} \sum_{j=1}^k \frac{L_k^2}{L_{j-1}L_j} y_j - \frac{L_k}{L_{k-1}} \sum_{j=1}^{k-1} \frac{L_{k-1}^2}{L_{j-1}L_j} y_j \right|^p \right)^{\frac{1}{p}} \\ &= (\sum_k |y_k|^p)^{\frac{1}{p}} = \|y\|_{\ell_p} < \infty \text{ and} \end{aligned}$$

$$\|x\|_{\ell_\infty(\hat{E})} = \sup_k \left| \frac{L_{k-1}}{L_k} x_k - \frac{L_k}{L_{k-1}} x_{k-1} \right| = \sup_k |y_k| = \|y\|_{\ell_\infty} < \infty.$$

This means that $x \in \ell_p(\hat{E}), 1 \leq p \leq \infty$. $\ell_p(\hat{E})$ and ℓ_p are linear isomorphic. Hence, we see that Z is surjective and preserves the norm. Therefore, Z is a linear bijection and so $\ell_p(\hat{E}) \cong \ell_p$ holds.

Theorem 3.3. If $1 < p \leq \infty$, then the inclusion $\ell_p \subset \ell_p(\hat{E})$ strictly

Proof. Let $x \in \ell_p, 1 < p \leq \infty$. It can be easily seen that the inequalities $\frac{L_{k-1}}{L_k} \leq 2$ and

$\frac{L_k}{L_{k-1}} \leq 3$ hold. By means of these inequalities and equality (3), we have

$$\sum_k |\hat{E}_k(x)|^p \leq \sum_k 6^{p-1} (|2x_k|^p + |3x_{k-1}|^p) \leq 6^{2p-1} (\sum_k |x_k|^p + \sum_k |x_{k-1}|^p),$$

$$\sup_k |\hat{E}_k(x)| \leq 5 \sup_k |x_k|.$$

Hence, for $1 < p \leq \infty$, we obtain

$$\|x\|_{\ell_p(\hat{E})} \leq 36 \|x\|_p \text{ and } \|x\|_{\ell_\infty(\hat{E})} \leq 5 \|x\|_\infty. \quad (6)$$

Additionally, the sequence $x = (x_k) = (L_k^2) = (1, 3^2, 4^2, 7^2, \dots)$ is in $\ell_p(\hat{E}) - \ell_p$. This gives that the inclusion $\ell_p \subset \ell_p(\hat{E})$ is strict in the case $1 < p \leq \infty$. Also, the inequality (6) holds for $p = 1$.

Theorem 3.4. $\ell_p(\hat{E}) \subset \ell_s(\hat{E})$ for $1 \leq p < s$.

Proof. Let y be the sequence given by equality (3). If we take $x \in \ell_p(\hat{E})$, then we have $y \in \ell_p$. Since the inclusion $\ell_p \subset \ell_s$ holds, we obtain that $y \in \ell_s$. This yields us $x \in \ell_s(\hat{E})$ and so, the inclusion $\ell_p(\hat{E}) \subset \ell_s(\hat{E})$ holds.

Theorem 3.5. The sequence $(b^{(k)})_{k=1}^\infty$ which is established by

$$(b^{(k)})_n = \begin{cases} \frac{L_n^2}{L_{k-1}L_k}, & n \geq k \\ 0, & n < k \end{cases} \quad (7)$$

forms a basis for $\ell_p(\hat{E})$ in the case $1 \leq p < \infty$. Also, every $x \in \ell_p(\hat{E})$ has a unique representation of the form

$$x = \sum_k \hat{E}_k(x) b^{(k)}. \quad (8)$$

Proof. From equality (7), $\hat{E}(b^{(k)}) = e^{(k)} \in \ell_p$ and this means that $b^{(k)} \in \ell_p(\hat{E})$. Now, let's take $x \in \ell_p(\hat{E})$ and put $x^{(m)} = \sum_{k=1}^m \hat{E}_k(x) b^{(k)}$ for every non-negative integer m .

Thus, it is obtained that

$$\hat{E}(x^{(m)}) = \sum_{k=1}^m \hat{E}_k(x) \hat{E}(b^{(k)}) = \sum_{k=1}^m \hat{E}_k(x) e^{(k)}$$

and also

$$\hat{E}_n(x - x^{(m)}) = \begin{cases} \hat{E}_n(x), & m < n \\ 0, & m \geq n \geq 0 \end{cases}.$$

So, there is a non-negative integer m_0 such that $\sum_{n=m_0+1}^{\infty} |\hat{E}_n(x)|^p \leq \left(\frac{\varepsilon}{2}\right)^p$ for any $\varepsilon > 0$. Hence, we have that

$$\|x - x^{(m)}\|_{\ell_p(\hat{E})} = \left(\sum_{n=m+1}^{\infty} |\hat{E}_n(x)|^p \right)^{1/p} \leq \left(\sum_{n=m_0+1}^{\infty} |\hat{E}_n(x)|^p \right)^{1/p} \leq \frac{\varepsilon}{2} < \varepsilon$$

for every $m \geq m_0$, that is,

$$\lim_{m \rightarrow \infty} \|x - x^{(m)}\|_{\ell_p(\hat{E})} = 0.$$

Last, let's assume that $x = \sum_k \beta_k(x) b^{(k)}$ to demonstrate the uniqueness of equality (8) for $x \in \ell_p(\hat{E})$. By using the continuity of the linear transformation Z , we have

$$\hat{E}_n(x) = \sum_k \beta_k(x) \hat{E}_n(b^{(k)}) = \sum_k \beta_k(x) \delta_{nk} = \beta_n(x).$$

Therefore, the proof is completed.

$$\hat{E}_n(x - x^{(m)}) = \begin{cases} \hat{E}_n(x), & m < n \\ 0, & m \geq n \geq 0 \end{cases}.$$

So, there is a non-negative integer m_0 such that $\sum_{n=m_0+1}^{\infty} |\hat{E}_n(x)|^p \leq \left(\frac{\varepsilon}{2}\right)^p$ for any $\varepsilon > 0$. Hence, we have that

$$\|x - x^{(m)}\|_{\ell_p(\hat{E})} = \left(\sum_{n=m+1}^{\infty} |\hat{E}_n(x)|^p \right)^{1/p} \leq \left(\sum_{n=m_0+1}^{\infty} |\hat{E}_n(x)|^p \right)^{1/p} \leq \frac{\varepsilon}{2} < \varepsilon$$

for every $m \geq m_0$, that is,

$$\lim_{m \rightarrow \infty} \|x - x^{(m)}\|_{\ell_p(\hat{E})} = 0.$$

Last, let's assume that $x = \sum_k \beta_k(x) b^{(k)}$ to demonstrate the uniqueness of equality (8)

for $x \in \ell_p(\hat{E})$. By using the continuity of the linear transformation Z , we have

$$\hat{E}_n(x) = \sum_k \beta_k(x) \hat{E}_n(b^{(k)}) = \sum_k \beta_k(x) \delta_{nk} = \beta_n(x).$$

Therefore, the proof is completed.

Theorem 3.6. The spaces $\ell_p(\hat{E})$ and bv_p do not include each other for $1 \leq p < \infty$.

Proof. If we take $x = (x_k) = (L_k^2) = (1, 3^2, 4^2, 7^2, \dots)$ and $e = (1, 1, 1, \dots)$, then we result that $x \in \ell_p(\hat{E})$ and $x \notin bv_p$ by reason of $\hat{E}x = (2, 0, 0, \dots) \in \ell_p$ and $\Delta x = (1, L_0 L_3, L_1 L_4, \dots, L_{k-2} L_{k+1}, \dots) \notin \ell_p$. Now, we take the equation

$$\left| \frac{L_{k-1}}{L_k} - \frac{L_k}{L_{k-1}} \right| = \frac{|L_{k-1}^2 - L_k^2|}{L_{k-1} L_k} = \frac{|(-5)(-1)^k - L_{k-1} L_k|}{L_{k-1} L_k}$$

into consideration. If k is even, then $L_{k-1} L_k < |(-5)(-1)^k - L_{k-1} L_k|$. Hence, the

series $\sum_k \left| \frac{L_{k-1}}{L_k} - \frac{L_k}{L_{k-1}} \right|^p$ is not convergent for $1 \leq p < \infty$. Thus, $\hat{E}e = \left(\frac{L_{k-1}}{L_k} - \frac{L_k}{L_{k-1}} \right) \notin$

ℓ_p in the case $1 \leq p < \infty$ and it is clear that $e \in \ell_p$. In conclusion, the spaces $\ell_p(\hat{E})$ and bv_p coincide but they do not contain one another.

A Banach space Y possess Banach-Saks property if any bounded sequence in Y approves a subsequence

$$R(X) = \sup \left(\lim_{n \rightarrow \infty} \inf \|x_n + x\| \right)$$

He also proved that a Banach space X with $R(X) < 2$ has the weak fixed point property.

Of late years, some studies about geometrical properties of a sequence space can be seen in (Et and Karakaya, 2014), (Karakaya and Altun, 2014), (Mursaleen et al., 2007).

Theorem 3.7. $\ell_p(\hat{E})$ holds the Banach-Saks property of type p .

whose arithmetic mean converges in norm. Similarly, a Banach space Y holds weak Banach-Saks property if any weakly null sequence in X admits a subsequence whose arithmetic mean strongly converges in norm.

Let X be a Banach space. (Garcia-Falset, 1994) defined the coefficient $R(X)$ as:

Proof. It can be proved by standard technic which is available in (Karaş et al., 2013).

Remark 3.1. As $\ell_p(\hat{E})$ is linearly isomorphic to space ℓ_p , we take in consideration .

$$R(\ell_p(\hat{E})) = R(\ell_p) = 2^{1/p}$$

Since $R(\ell_p(\hat{E})) < 2$, it can be given the below theorem:

Theorem 3.9. The space $\ell_p(\hat{E})$ possess the weak fixed point property for $1 < p < \infty$.

REFERENCES

- Kara EE, 2013. Some topological and geometrical properties of new Banach sequence spaces. *J. Inequal. Appl.*, 38: 15 pp.
- Mursaleen M, Noman AK, 2011. On some new sequence spaces of non-absolute type related to the spaces and I. *Filomat*, 25: 33-51.
- Mursaleen M, Noman AK, 2010. On the spaces of -convergent and bounded sequences. *Thai J. Math.*, 8: 311-329.
- Mursaleen M, Noman AK, 2010. On some new difference sequence spaces of non-absolute type. *Math. Comput. Modelling*, 52: 603-617.
- Mursaleen M, Başar F, Altay B, 2006. On the Euler sequence spaces which include the spaces and II. *Nonlinear Analysis*, 65: 707-717.
- Altay B, Başar F, Mursaleen M, 2006. On the Euler sequence spaces which include the spaces and I. *Inform. Sci.*, 176: 1450-1462.
- Başar F, Altay B, 2003. On the space of sequences of -bounded variation and related matrix mappings. *Ukrainian Math. J.*, 55: 136-147.
- Altay B, Başar F, 2005. Some Euler sequence spaces of non-absolute type. *Ukrainian Math. J.*, 57: 1-17.
- Savaş E, Karakaya V, Şimşek N, 2009. Some -type new sequence spaces and their geometric properties. *Abstr. Appl. Anal.*, Art. ID 696971: 12 pp.
- Karaş M, 2015. A new regular matrix defined by Fibonacci numbers and its applications. *BEU Journal of Science*, 4: 205-210.
- Kara EE, Başarır M, 2012. An application of Fibonacci numbers into infinite Toeplitz matrices. *Caspian Journal of Mathematics Sciences*, 1: 1-6.
- Kızmaz H, 1981. On certain sequence spaces. *Canad. Math. Bull.*, 24: 169-176.
- Altay B, Başar F, 2007. The fine spectrum and the matrix domain of the difference operator Δ on the sequence space . *Commun. Math. Anal.*, 2: 1-11.
- Çolak R, Et M, Malkowsky E, 2004. Some topics of sequence spaces, *Lecture notes in Mathematics*. *Firat Univ. Press*, 1-63.
- Et M, 1993. On some difference sequence spaces. *Doğa Mat.*, 17: 18-24.
- Mursaleen M, 1996. Generalized spaces of difference sequences. *J. Math. Anal. Appl.*, 203: 738-745.
- Gaur AK, Mursaleen M, 1998. Difference sequence spaces. *Internat. J. Math. Math. Sci.*, 21: 701-706.
- Et M, Esi A, 2000. On Köthe-Toeplitz duals of generalized difference sequence spaces. *Bull. Malays. Math. Sci. Soc. (2)*, 23: 25-32.
- Et M, Çolak R, 1995. On some generalized difference sequence spaces. *Soochow J. Math.*, 21: 377-386.
- Et M, Başarır M, 1997. On some new generalized difference sequence spaces. *Period. Math. Hungar.*, 35: 169-175.

Karakaş M, Et M, Karakaya V, 2013. Some geometric properties of a new difference sequence space involving lacunary sequences. *Acta Math. Sci. Ser. B Engl. Ed.*, **33**: 1711-1720.

Koshy T, 2001. Fibonacci and Lucas Numbers with Applications. John Wiley&Sons, Canada. 672 p.

Vajda S, 2008. Fibonacci and Lucas Numbers, and the Golden Section Theory and Applications. Dover Edition, NewYork, USA. 189 p.

Wilansky A, 1984. Summability Through Functional Analysis. Elsevier Science Publishers BV, Amsterdam, The Netherlands. 317 p.

García-Falset J, 1994. Stability and fixed points for nonexpansive mappings. *Houston J. Math.*, **20**: 495-506.

Et M, Karakaya V, 2014. A new difference sequence set of order α and its geometrical properties. *Abstr. Appl. Anal.*, Art. ID 278907: 4 p.

Karakaya V, Altun F, 2014. On some geometric properties of a new paranormed sequence space. *J. Funct. Spaces*, Art. ID 685382: 8 p.

Mursaleen M, Çolak R, Et M, 2007. Some geometric inequalities in a new Banach sequence space. *J. Inequal. Appl.*, Art. ID 86757: 6 p.

Karakaş M, Çınar M, Et M, 2013. Some geometric properties of a new sequence space. *J. Comput. Anal. Appl.*, **15**: 23-31.