iJRASET

International Journal For Research in
Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6   Issue: I   Month of publication: January 2018

DOI: <http://doi.org/10.22214/ijraset.2018.1142>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Conductivity of Mixed Ligand Complexes of Alkali Metal Salts

Shahnawaz Mahmood¹

¹Associate Professor and Head of the Department of Applied Sciences & Humanities Mewat Engineering College MP alla, Nuh, HARYANA - 122107

Abstract: Our present work shows that hydrogen bonding is not necessary for formation of adducts between a chelating ligand and the alkali metal salt of a chelate ion.

Keywords: hydroxyacetophenone phenylhydrazone,

I. INTRODUCTION

Only two bands are observed : one at 738 cm^{-1} (assigned to out – of – plane motion of the H – atoms on the heterocyclic rings) and the other at 854 cm^{-1} (assigned to out of plane motion of hydrogen atoms on the central ring). On complication, the multiple splitting of these two bands arise from out – of – plane motions other than those in phase and also probably from overtones of low lying fundamentals in resonance. The splitting of these two bands appears to be metal sensitive.

These facts suggest the coordination of the alkali metals with 1,10 – phenanthroline through the nitrogen atoms of its pyridine fragments. Three corners of the triangles shown in the above table represent the alkali metals Li, Na and K. Symbols indicate that the corresponding complexes were obtained. Only two bands are observed : one at 738 cm^{-1} (assigned to out – of – plane motion of the H – atoms on the heterocyclic rings) and the other at 854 cm^{-1} (assigned to out of plane motion of hydrogen atoms on the central ring). On complication, the multiple splitting of these two bands arise from out – of – plane motions other than those in phase and also probably from overtones of low lying fundamentals in resonance. The splitting of these two bands appears to be metal sensitive. These facts suggest the coordination of the alkali metals with 1,10 – phenanthroline through the nitrogen atoms of its pyridine fragments. Our present work shows that hydrogen bonding is not necessary for formation of adducts between a chelating ligand and the alkali metal salt of a chelate ion.

II. CONDUCTIVITIES

Molar conductivities of all the complexes were measured in N – methyl – 2 – pyrrolidone at 250°C at a concentration of 10^{-3} M . The solvent has been calibrated by Banerjee et.al¹⁶ and they have observed that the values of molar conductivities in the range $35 - 40\text{ cm}^{-1}$ correspond to 1:1 electrolyte. Detail studies^{125,126} of this solvent have shown that significantly different values as obtained with different anions, even for tetra – alkylammonium cations, and that 1:1 electrolytes may give values as low as $25\text{ ohm}^{-1}\text{ cm}^{-1}\text{ mole}^{-1}$. From the results (Table – 7.2), it is observed that none of the values approach either ideal or 1:1 electrolyte. Low values of molar conductivities of these complexes would seem to indicate neutral complexes. However, low values are also obtained for the simple salts, which may be due to ion-pair formation and the consequent preliminary chelation. The trend $\text{Li} > \text{Na} > \text{K}$ would seem to indicate that the lithium complexes is the strongest in solution. Significantly high values of molar conductivities of these complexes in N – methyl – 2 – pyrrolidone suggest that they have undergone dissociation

III. STRUCTURE AND BONDING

Analytical data lead to the general molecular formula M1.L' for these complexes, where M = alkali metal salt of the Q-Hydroxyacetophenone hydrazone, Q-Hydroxyacetophenone phenyl hydrazone or Q-Hydroxyacetophenone – 2, 4 - dinitrophenyl hydrazone and L' = 1,10 – phenanthroline. Their IR spectra suggest the coordination of the ligand L' with alkali metals through nitrogen atom of the pyridine fragments. Accordingly, these complexes should have the following structure :

Fig 7.5

IV. EXPERIMENTAL

Ligand 1,10 - phenanthroline

The ligand 1,10 – phenanthroline of E. Merck quality was used as such.

A. Adducts of 1,10 – phenanthroline

1) *Adducts with alkali metal salts of O-Hydroxyacetophenone hydrazine:* To the hot suspension of an alkali metal salt of O-Hydroxyacetophenone hydrazone (OHAHz) in acetone, excess of solid 1,10 – phenanthroline was added. On stirring, the contents went into solution and almost instantaneously, pale yellow coloured adduct came apart. It was redissolved by warming the contents on a steam bath for 5 minutes. On cooling, the pale yellow adduct reappeared. It was filtered, washed with cold benzene dried in a desiccator over anhydrous CaCl_2 .

a) *Li OHAHz.phen*

Found : C, 71.38 ; H, 5.00; N, 16.62%

$\text{C}_{20}\text{H}_{17}\text{N}_4\text{OLi}$ requires

: C, 71.42 ; H, 5.06 ; N, 16.67%

b) *NaOHAHz.phen*

Found : C, 68.13 ; H, 4.87; N, 15.85 ; Na, 6.50%

$\text{C}_{20}\text{H}_{17}\text{N}_4\text{ONa}$ requires

: C, 68.18 ; H, 4.83; N, 15.91 ; Na, 6.53%

c) *KOHAHz.phen*

Found : C, 65.17 ; H, 4.58; N, 15.20 ; K, 10.52%

$\text{C}_{20}\text{H}_{17}\text{N}_4\text{OK}$ requires

: C, 65.22 ; H, 4.62; N, 15.22 ; K, 10.59%

2) *Adducts with alkali metal salts of O-Hydroxyacetophenone phenylhydrazone :* These adducts were obtained by the method strictly analogous to that described for the preparation of adducts of 1,10 – phenanthroline with alkali metal salts of O-Hydroxyacetophenone phenylhydrazone . Ease of formation of the adducts was found to be greater in comparison to the formation of the adducts in case of OHAHz.

a) *Li OHAPz.phen*

Found : C, 75.69 ; H, 5.01; N, 13.51%

$\text{C}_{26}\text{H}_{21}\text{N}_4\text{OLi}$ requires

: C, 75.73 ; H, 5.09; N, 13.59%

b) *Na OHAPz.phen*

Found : C, 72.82 ; H, 4.89; N, 13.02 ; Na, 5.32%

$C_{26}H_{21}N_4O$ Na requires

:C, 72.89 ; H, 4.91; N, 13.08 ; Na, 5.37%

c) *KOHADNPz.phen*

Found : C, 70.22 ; H, 4.69; N, 12.66; K, 8.74%

$C_{26}H_{21}N_4O$ K requires

: C, 70.27 ; H, 4.73; N, 12.61; K, 8.78%

3) *Adducts with alkali metal salts of O-Hydroxyacetophenone - 2,4 – dinitrophenylhydrazine*

These adducts too could be obtained by the procedure as described above for the preparation of adducts of OHAPz.

a) *LiOHADNPz.phen*

Found : C, 62.23 ; H, 3.62; N, 16.72%

$C_{20}H_{18}N_6O_5Li$ requires

: C, 62.28 ; H, 3.59 ; N, 16.77%

b) *NaOHADNPz.phen*

Found : C, 60.30 ; H, 3.52; N, 16.20 ; Na, 4.50%

$C_{20}H_{18}N_6O_5Na$ requires

: C, 60.35 ; H, 3.48; N, 16.25 ; Na, 4.45%

c) *KOHADNPz.phen*

Found : C, 58.50 ; H, 3.42; N, 15.72; K, 7.36%

$C_{20}H_{18}N_6O_5K$ requires

: C, 58.54 ; H, 3.48; N, 15.76; K, 7.32%

REFERENCES

- [1] D.P. Craig and R.S. Nyholm : Chelating Agents and Metal Chelates', F.P. Dwyer and D.P. Mellor, eds., Academic Press, New York, 1964
- [2] Born : Z. Physick : 1820, 1, 45
- [3] T.C. Shields : Chem. Comm : 1968, 832
- [4] R.G. Pearson : J. Chem. Educ. : 1968, 45, 581, 643
- [5] N.V. Sidgwick and S.G.P. Plant : J.Chem. Soc. : 1925, 127, 209
- [6] R.N. Tichane and W.E. Benett : J. Amer. Chem. Soc. ; 1957, 79, 1293
- [7] D.E. Fenton, C. Nave and M.R. Truter : J. Chem, Soc. (Dalton), 1973, 2188
- [8] D.L. Hughes A.K. Banerjee, D. Prakash, P. Kejariwal and S.K. Roy : J. Chem, Soc. (Dalton), 1973, 2347
- [9] D.E. Fenton, C. Nave and M.R. Truter : J. Chem, Soc. (Dalton), 1973, 2188
- [10] M.A. Bush, H. Luth and M.R. Truter : J. Chem. Soc. (A), 1971, 740
- [11] F.M. Brewer : J. Chem. Soc., 1931, 361

10.22214/IJRASET

45.98

IMPACT FACTOR:
7.129

IMPACT FACTOR:
7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)