iJRASET

International Journal For Research in
Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: I Month of publication: January 2018

DOI: <http://doi.org/10.22214/ijraset.2018.1209>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Synthesis and Biological Screening of N-{4'-[5''-(ARYL)-1''-H-PYRAZOL-3''-yl] PHENYL}-DIBENZO [b,f][1,4] THIAZEPIN-11-AMINE.

M.V.Gondaliya¹, D. M .Purohit²

¹Shri JSB & KMB Arts, Shri ANS Science and Shri NFS Commerce College – Kholwad (Surat)

²ShriM. & N. Virani Science College – Kalawad Road, Rajkot-360005 (GUJ) INDIA

Abstract: Pyrazoline derivatives shows broad spectrum of pharmacological activity with a view of getting to synthesized and evaluated biological screening of N-[4'-(5''-aryl-1''-H-pyrazol-3''-yl)phenyl]dibenzo[b,f][1,4]thiazepin-11-amine(5a-5j).The compound(5a-5j) have been synthesized by the condensation of 1''-[4'-diabenzo[b,f][1,4] thiazepin-11-yl)aminophenyl]-3''-aryl-2-ene-1''-ones with hydrazine hydrate. The constitutions of the synthesized products were supported by IR, ¹H NMR and mass spectral data. The products were screened for their antimicrobial activity at 50 µg concentration by cup-plate method.

Keywords: yrazolines., Anti-microbial activity

I. INTRODUCTION

Nitrogen containing five membered heterocycles, pyrazolines have proved to be the most useful frame work for therapeutic activities, pyrazolines have attracted attention of medical chemistry for both with regard to heterocyclic chemistry and the pharamacological activities and therapeutic activity, which have been studied extensively for their industrial applications. Pyrazolines gave antidiabetic¹, tranquilizer², analgesic³⁻⁴, bactericidal⁵⁻⁶, cardiovascular⁷, antitumor⁸, antineoplastic⁹.

II. MATERIALS AND METHODS

All the melting points were determined in open glass capillary tubes and are uncorrected. IR spectra recorded on SHIMADZU-FT-IR-8400 Spectrophotometer; 4000-400 cm⁻¹ (KBr disc). ¹HNMR spectra were recorded on a BRUKER spectrometer (400MHz) instrument using TMS as an internal standard. Mass spectra were recorded on waters QDA spectrophotometer. All the compounds gave satisfactory elemental analysis.

III. EXPERIMENTAL

A. Step-I: Synthesis of 4'-Acetylphenyl[dibenzo[b,f][1,4]thiazepines-11-yl]amine

A mixture of 11-chlorodibenzo[b,f][1,4] thiazepine (2.45g, 0.01M) 4-amino acetophenone (1.35g, 0.01M) in methanol (25ml) and pyridine (2 ml) was refluxed on an oil bath at 130°C at 8 hrs. The content was cooled and poured into crushed ice-filtered and washed with water. The isolated product was crystallized from ethanol yield 87.32%, M.P:195⁰C (Found: C:73.20; H:4.60; N:8.11; C₂₁H₁₆N₂O₃ required C: 73.25; H: 4.65; N:8.13%)

B. Step-II: Synthesis of 1''-[4'-(dibenzo[b,f][1,4]thiazepines-11-yl)aminophenyl]-3''-(3''',4'''-dimethoxy phenyl)-2-ene-1''-one

A Mixture of (4-acetyl phenyl[b,f][1,4] thiazepine-11-yl)amine (3.44g, 0.01M) 3,4-dimethoxy benzaldehyde (1.66g, 0.01M), ethanol(15 ml), 20% NaOH till the solution vigorously stirring at basic medium at 24 hrs. The contents were poured into crushed ice, acidified filtered and crystallized from ethanol yield 72.32% M.P.208⁰C, (Found:C:73.00; H:4.71; N:5.60; C₃₀H₂₄N₂O₃S required: C:73.17; H:4.91; N:5.69%)

IR (KBr) cm⁻¹:2966 (C-H Str.(asym) ; 2851 (C-H Str. (sym) ; 1456 (C-H def bending) ; 3047 (C-H str)1485 (C=C Ring skeleton) ; 1257 (C-H i.p.def) ; 763 (C-H o.o.p) ; 1338 (C-N Str.) ; 3321 (N-H Str.) ; 1658 (C=O Str.) ; 1232 (C-O-C Str.). ¹H NMR(ppm) : 3.82 (6H (s), Ar-OCH₃); 6.97-7.59(15H (m) Ar-H); 7.73-7.75 (1H (d)=CH); 8.23-8.73 (1H (d) =CH); 9.62 (1H (s) -NH); Mass (m/z): 492,440,369,205,77

C. Step-III: Synthesis of N-{4'-[5''-(3''',4'''-dimethoxyphenyl)-1''-H-pyrazole-3''-yl]phenyl}-dibenzo[b,f][1,4]thiazepin-11-amine (5e)

A mixture of 1''-[4'-(dibenzo[b,f][1,4] thiazepine-11-yl)amino phenyl]-3''-(3''',4'''- dimethoxy phenyl)-2-ene-1''-one. (4.92gm 0.01M); hydrazine hydrate (1.0ml) in ethanol (15ml) was refluxed for 12 hrs. The product was poured into crushed ice filtered, washed with water and crystallised from dioxane. yield: 73.11% M.P. 257⁰ C (Found: C: 70.85; H: 4.99; N: 10.95; C₃₀H₂₆N₄O₂S, Required: C: 71.14; H: 5.17; N: 11.06%) IR (KBr) cm⁻¹:2993 (C-H Str.(asym) ; 2833 (C-H Str. (sym) ; 1456 (C-H def bending) ; 3072 (C-H str.Ar)1535 (C=C Ring skeleton) ; 1257 (C-H i.p.def) ;761 (C-H oop def) ; 1338 (C-N Str.) ; 3321 (N-H Str.) ; 1232 (C-O-C Str.) ; 2594(C-S-C Str.) ;¹H NMR (ppm): 2.08(2H (d) -CH₂); 3.73(6H (s) Ar-O-CH₃) ; 4.80 (1H (t) -CH) ; 5.41 (1H (s) -NH) ; 7.01-8.27(15H (m)Ar-H) ;10.01 (1H (s)-NH) ; Mass (m/e): 507, 473, 305, 221. 77.Similarly other compounds (5a-5j) have been synthesized.

D. Reaction Scheme

Table:I Physical Constants of the compounds (5a-5j)

Sr. No.	Ar	Molecular Formula	M.P °C	Yield	% of Nitrogen	
					Calcd.	Found
5a	C ₆ H ₅ -	C ₂₈ H ₂₂ N ₄ S	157	87.63	12.55	12.35
5b	4-CH ₃ -C ₆ H ₄ -	C ₂₉ H ₂₄ N ₄ S	169	80.33	12.17	12.03
5c	2-OCH ₃ -C ₆ H ₄ -	C ₂₉ H ₂₄ N ₄ OS	186	82.73	11.76	11.57
5d	4-OCH ₃ -C ₆ H ₄ -	C ₂₉ H ₂₄ N ₄ OS	242	71.55	11.76	11.53
5e	3,4-(OCH ₃) ₂ -C ₆ H ₃ -	C ₃₀ H ₂₆ N ₄ O ₂ S	257	73.11	11.06	10.95
5f	3,4,5-(OCH ₃) ₃ -C ₆ H ₂ -	C ₃₁ H ₂₈ N ₄ O ₃ S	229	68.75	10.44	10.21
5g	2-NO ₂ -C ₆ H ₄ -	C ₂₈ H ₂₁ N ₅ O ₂ S	210	73.50	14.25	13.03
5h	3-NO ₂ -C ₆ H ₄ -	C ₂₈ H ₂₁ N ₅ O ₂ S	173	71.35	14.25	14.04
5i	4-NO ₂ -C ₆ H ₄ -	C ₂₈ H ₂₁ N ₅ O ₂ S	183	69.53	14.25	14.09
5j	4-NH ₂ -C ₆ H ₄ -	C ₂₈ H ₂₃ N ₅ S	200	83.15	15.18	15.07

E. Antibacterial Activity: ¹⁰

The products were screened for their antibacterial activity. The nutrient agar broth prepared by the usual method, was inoculated specially with 0.5 ml for 24 hours, old subculture of *B. megaterium*, *S. aureus*, *Escherichia coli*, *Staphylococcus aureus*, in separate conical flasks at 40-50°C and mixed well by gentle shaking. About 25ml of the contents of the flask were poured and evenly spread in a petridish (13 cm in diameter) and allowed to set for 2 hrs. The cups (10 mm in diameter) were formed by the help of borer in an agar medium and filled with 0.1 ml (1.0mg/ml) solution of sample in dimethyl formamide. The plates were incubated at 32°C for 24 hrs and the control was also maintained with 0.1 ml of DMF in similar manner and the zone of inhibition of the bacterial growth are measured in mm diameter and are recorded in Table No. II

F. Antifungal Activity: ¹¹

Aspergillus niger was employed for testing fungicidal activity using cup-plate method the cultures were maintained on Sabouraud's agar slants. Purified compounds were used for testing the fungicidal activity, sterilized Sabouraud's agar medium was inoculated with 72 hours old 0.5 ml suspension of fungal spores in a sterilized separate flask. About 25 ml of the inoculated medium was evenly spreader in a Petridish and allowed to set for 2 hrs. The cups (10 mm in a diameter) were punched in Petridish and loaded with 0.1 ml (1.0 mg/ml) of solution of a sample in dimethyl formamide. The plate was incubated at room temperature 37°C for 48hrs. After the completion of incubation period. The zone of inhibition or growth in the form of diameter in mm was measured. Along the test solution in each Petridish one cup was filled up with solvent acts as control. The zones of inhibition are recorded in Table No. 2

Table : II

Anti-microbial studies of N-{4'-[5''-(aryl)-1''-H-pyrazole-3''-yl] phenyl}-dibenzo[b,f][1,4]thiazepin-11-amine (5a-5j)

Sr.No.	Ar	Antibacterial activity Zone of inhibition in m.m.				Antifungal activity Zone of inhibition in m.m.
		B. megaterium	S. aureus	S. taphimarium	E. coli	A. niger
5a	C ₆ H ₅ -	16	19	14	16	20
5b	4-CH ₃ -C ₆ H ₄ -	18	18	18	13	21
5c	2-OCH ₃ -C ₆ H ₄ -	18	13	19	15	23
5d	4-OCH ₃ -C ₆ H ₄ -	27	21	17	17	24
5e	3,4-(OCH ₃) ₂ -C ₆ H ₃ -	25	23	18	19	20
5f	3,4,5-(OCH ₃) ₃ -C ₆ H ₂ -	17	17	19	17	22
5g	2-NO ₂ -C ₆ H ₄ -	15	15	20	13	18
5h	3-NO ₂ -C ₆ H ₄ -	19	23	18	12	24
5i	4-NO ₂ -C ₆ H ₄ -	23	17	20	14	20
5j	4-NH ₂ -C ₆ H ₄ -	21	20	23	15	24

IV. RESULTS AND DISCUSSION

Compounds (5a-5j) were screened for their in vitro antibacterial activity using cup-plate agar diffusion method at a concentration of 50 µg/ml using Gram positive bacterial strains such as *B. megaterium* and *S. aureus* and Gram negative bacterial strain such as *Escherichia coli* and *S. taphimarium*. Known antibiotics like ampicillin, Chloramphenicol, norfloxacin, and Griseofulvin were used for comparison purpose. Comparable antimicrobial activity represented in Table –III

Table ; III
Comparable antimicrobial activity with known standard drugs

	Antibacterial activity Zone of inhibition in m.m.				Antifungal activity Zone of inhibition in m.m.
	B. megaterium Ar	S. aureus Ar	S.taphimarium Ar	E.coli Ar	A.niger Ar
	4-OCH ₃ -C ₆ H ₄ -27	4-OCH ₃ -C ₆ H ₄ - 21	2-NO ₂ -C ₆ H ₄ – 20	4-OCH ₃ -C ₆ H ₄ -17	2-OCH ₃ -C ₆ H ₄ - 23
	3,4-(OCH ₃) ₂ -C ₆ H ₃ - 25	3,4-(OCH ₃) ₂ -C ₆ H ₃ - 23	4-NO ₂ -C ₆ H ₄ - 20	3,4-(OCH ₃) ₂ -C ₆ H ₃ - 19	4-OCH ₃ -C ₆ H ₄ - 24
	4-NO ₂ -C ₆ H ₄ -23	3-NO ₂ -C ₆ H ₄ - 23	4-NH ₂ -C ₆ H ₄ - 23	3,4,5-(OCH ₃) ₂ -C ₆ H ₃ - 17	3-NO ₂ -C ₆ H ₄ -24
	4-NH ₂ -C ₆ H ₄ - 21				4-NH ₂ -C ₆ H ₄ -24
Comparable activity with known standard drugs					
Ampicilin50µg	29	28	31	33	-
Chloramphenicol50µg	28	31	28	29	-
Norfloxacin 50µg	32	30	26	29	-
Griseofulvin 50µg	-	-	-	-	26

REFERENCES

- [1] H. G. Garg and P. P. Singh;Chem. Soc., 2, 1141 (1936).
- [2] B. Hans, R. Rolf and R. Rudolf;US.Pat., 3, 822, 283 (1974); Chem. Abstr., 81, 105494r (1974).
- [3] Delay Francois(Felfnenich S. A.) Patent Schrift (Switz) Appl 3, 21, 765(993)
- [4] Aysel G. Seret D., Gultaze C., Keyser E., Kamil V.;Ear J.Med. Chem., 35, 359-64 (2000).
- [5] P. Desaea; A. Nunrich; M. Capdemy and G. Devaux; J. Med. Chem., 25, 285 (1990).
- [6] KallurayaBlakrishna, R. Chimabalkar, G. Rai, R. Gururaja, S. Shenoy; J. Indian Coun. Chemi.,18(2), 39-42 (2001); Chem. Abstr, 138, 238061(2003).
- [7] Y. Hiroyuti, O. Mocoto et al.; Eur Pat. Appl. Ep 295695 (C1 C07D 401/6) (1988); Chem. Abstr, 111,23510 (1989).
- [8] W. I. Ronald, A. Adriano; Chem. Abstr.,126, 181346f (1997).
- [9] H. M. Mokhtar, H. M. Faidailah;Pharmazie, 42, 482 (1987).
- [10] A. L. Barry; "The Antimicrobial Suceptibility test, principle and practices (ELBS-4th Edition) 180-193 (1976)
- [11] F. Simoncini R. R.Rengone and C. Calann.Farmaco D. Part., 23 (10), 559-70 19680; ChelllAbstr. **59**, 109851d (1968).

10.22214/IJRASET

45.98

IMPACT FACTOR:
7.129

IMPACT FACTOR:
7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)