iJRASET

International Journal For Research in
Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: IV Month of publication: April 2018

DOI: <http://doi.org/10.22214/ijraset.2018.4642>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Strong form of Fuzzy Closed Sets in Fuzzy Topological Space

D. Ananthi¹, K. Ygapriya², G. Bhuvana³, N. Sathishkumar⁴, R. Ragupathiraja⁵

^{1, 2, 3, 4, 5}Asst. Prof Edayathangudy G. S., Pillay Arts and Science College Nagapattinam

Abstract: In this paper we have introduced a new class of fuzzy sets called fuzzy strongly (gsp)*-closed sets, properties of this set are investigated and we introduce new fuzzy spaces namely, fuzzy $T_s(gsp)^*$ -space, fuzzy $gT_s(gsp)^*$ -space, fuzzy $g^*T_s(gsp)^*$ -space and fuzzy $g^{**}T_s(gsp)^*$ -space. Further strongly (gsp)*-continuous mappings are also introduced and investigated.

Keywords: fuzzy Strongly (gsp)*-closed sets, fuzzy Strongly (gsp)*-continuous maps, fuzzy $T_s(gsp)^*$ -space, fuzzy $gT_s(gsp)^*$ -space, fuzzy $g^*T_s(gsp)^*$ -space and fuzzy $g^{**}T_s(gsp)^*$ -space.

I. PRELIMINARIES

Throughout this paper (X, τ) and (Y, σ) represent non-empty fuzzy topological spaces on which no separation axioms are assumed unless otherwise mentioned. For a fuzzy subset A of a space (X, τ) , $cl(A)$ and $int(A)$ denote the fuzzy closure and the fuzzy interior of A respectively.

A. Definition 1.1

A Subset A of fuzzy topological space (X, τ) is called;

- 1) Fuzzy semi open set if $A \subseteq cl(int(A))$ and a fuzzy semi-closed set if $int(cl(A)) \subseteq A$.
- 2) Fuzzy semi pre-open set if $A \subseteq cl(int(cl(A)))$ and a fuzzy semi-pre closed set if $int(cl(int(A))) \subseteq A$
- 3) Fuzzy regular -open set if $int(cl(A)) = A$ and a fuzzy regular -closed.

B. Definition 1.2

A Subset A of fuzzy topological space (X, τ) is called;

- 1) Fuzzy generalized closed set (briefly fuzzy g-closed) if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy open in (X, τ)
- 2) Fuzzy g^* -closed set if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy g open in (X, τ)
- 3) Fuzzy g^{**} -closed set if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy g^* open in (X, τ)
- 4) Fuzzy wg - closed set if $cl(int(A)) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy open in (X, τ)
- 5) Fuzzy regular generalized closed set (briefly fuzzy rg-closed) if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy regular open in (X, τ)
- 6) Fuzzy sg^{**} - closed set if $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy g^{**} open in (X, τ)
- 7) Fuzzy sg^* - closed set if $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy g^* open in (X, τ)
- 8) Fuzzy generalized semi-closed set (briefly fuzzy gs-closed) if $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy open in (X, τ)
- 9) Fuzzy gsp - closed set if $spcl(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy open in (X, τ)
- 10) Fuzzy (gsp)*- closed set if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U fuzzy gsp is open in (X, τ)

C. Definition 1.3

A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called ;

- 1) fuzzy g - continuous if $f^{-1}(V)$ is a fuzzy g -closed set of (X, τ) for every fuzzy closed set V of (Y, σ)
- 2) fuzzy g^* - continuous if $f^{-1}(V)$ is a fuzzy g^* -closed set of (X, τ) for every fuzzy closed set V of (Y, σ)
- 3) fuzzy g^{**} - continuous if $f^{-1}(V)$ is a fuzzy g^{**} -closed set of (X, τ) for every fuzzy closed set V of (Y, σ)
- 4) fuzzy rg - continuous if $f^{-1}(V)$ is a fuzzy rg -closed set of (X, τ) for every fuzzy closed set V of (Y, σ)
- 5) fuzzy wg - continuous if $f^{-1}(V)$ is a fuzzy wg -closed set of (X, τ) for every fuzzy closed set V of (Y, σ)
- 6) fuzzy (gsp)* - continuous if $f^{-1}(V)$ is a fuzzy (gsp)* -closed set of (X, τ) for every fuzzy closed set V of (Y, σ)

D. Definition 1.4

A fuzzy topological space (X, τ) is said to be;

- 1) Fuzzy $T_{1/2}^*$ space if every fuzzy g^* -closed set in it is fuzzy closed.
- 2) Fuzzy T_d space if every fuzzy g_s -closed set in it is fuzzy g -closed.

II. FUZZY STRONGLY (GSP)* - CLOSED SETS IN FUZZY TOPOLOGICAL SPACE

We introduce the following definition

A. Definition 2.1

A subset A of a fuzzy Topological space (X, τ) is said to be a fuzzy strongly (gsp)*-closed set if $\text{cl}(\text{int}(A)) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy gsp-open.

- 1) *Lemma 2.1:* Every fuzzy closed set is fuzzy strongly (gsp)*-closed.
- 2) *Proof:* Let A be a fuzzy closed. Then $\text{cl}(A) = A$. Let us prove that A is fuzzy strongly (gsp)* - closed. Let $A \subseteq U$ and U be fuzzy gsp-open. Then $\text{cl}(A) \subseteq U$. Since A is fuzzy closed, $\text{cl}(\text{int}(A)) \subseteq \text{cl}(A) \subseteq U$. Then $\text{cl}(\text{int}(A)) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy gsp - open. so A is fuzzy strongly (gsp)* - closed. The converse of the above proposition need not be true in general as seen in the following example.

B. Lemma 2.2

Every fuzzy g -closed set is fuzzy strongly (gsp)*-closed.

- 1) *Proof:* Let A be fuzzy g -closed. Then $\text{cl}(A) \subseteq U$ Whenever $A \subseteq U$ and U is fuzzy open in (X, τ) . To prove A is fuzzy strongly (gsp)* - closed. Then $A \subseteq U$ and U be fuzzy (gsp) open. We have $\text{cl}(A) \subseteq U$ Whenever $A \subseteq U$ and U is fuzzy open in (X, τ) . Since every fuzzy open set is (fuzzy gsp) -open. We have $\text{cl}(A) \subseteq U$ Whenever $A \subseteq U$ and U is fuzzy (gsp)-open in (X, τ) . But $\text{cl}(\text{int}(A)) \subseteq \text{cl}(A) \subseteq U$ whenever $A \subseteq U$ and U is (fuzzy gsp)- open in (X, τ) then $\text{cl}(\text{int}(A)) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy (gsp) - open in (X, τ) so A is fuzzy strongly (gsp)*-closed. The converse of the above proposition need not be true in general as seen in the following example.

C. Lemma 2.3

Every fuzzy g^* -closed set is fuzzy strongly (gsp)* - closed.

The converse of the above proposition need not be true in general as seen in the following example.

D. Lemma 2.4

Every fuzzy rg - closed set is fuzzy strongly (gsp)* - closed.

- 1) *Proof:* Let A be fuzzy rg -closed set. Then $\text{cl}(A) \subseteq U$ Whenever $A \subseteq U$ and U is fuzzy regular-open in (X, τ) . To prove A is fuzzy strongly (gsp)* - closed. Let $A \subseteq U$ and U be fuzzy (gsp) open. Since every fuzzy regular-open set is fuzzy (gsp)-open. We have $\text{cl}(A) \subseteq U$ Whenever $A \subseteq U$ and U is fuzzy (gsp)-open in (X, τ) . But $\text{cl}(\text{int}(A)) \subseteq \text{cl}(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy (gsp)-open in (X, τ) . so $\text{cl}(\text{int}(A)) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy (gsp)-open in (X, τ) . So A is fuzzy strongly (gsp)*-closed.
- 2) *Remark 2.1:* fuzzy Strongly (gsp)* - closedness is independent of fuzzy semi-closedness

E. Lemma 2.5

Every fuzzy (gsp)* - closed set is fuzzy strongly (gsp)* - closed set

- 1) *Proof:* Let A be fuzzy (gsp)* -closed set. Then $\text{cl}(A) \subseteq U$ Whenever $A \subseteq U$ and U is fuzzy gsp-open in (X, τ) . To prove A is fuzzy strongly (gsp)* - closed. Let $A \subseteq U$ and U be fuzzy (gsp) open. Since every fuzzy (gsp)* - open set is fuzzy (gsp)-open. We have $\text{cl}(A) \subseteq U$ Whenever $A \subseteq U$ and U is fuzzy (gsp)-open in (X, τ) . But $\text{cl}(\text{int}(A)) \subseteq \text{cl}(A) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy (gsp)-open in (X, τ) so $\text{cl}(\text{int}(A)) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy (gsp)-open in (X, τ) so A is fuzzy strongly (gsp)* -closed. The converse of the above proposition need not be true in general as seen in the following example.

- 2) *Theorem 2.1:* Every fuzzy g^{**} -closed set is fuzzy strongly $(gsp)^*$ -closed. The converse of the above proposition need not be true in general.
- 3) *Remark 2.2:* fuzzy Strongly $(gsp)^*$ -closed is independent of fuzzy sg^{**} -closed.
- 4) *Remark 2.3:* Strongly fuzzy $(gsp)^*$ -closed is independent of fuzzy sg^* -closed.

F. Lemma 2.6

Every fuzzy wg -closed set is fuzzy strongly $(gsp)^*$ -closed.

- 1) *Proof:* Let A be fuzzy wg -closed. Then $cl(int(A)) \subseteq U$ Whenever $A \subseteq U$ and U is fuzzy open in (X, τ) . To prove A is fuzzy strongly $(gsp)^*$ -closed. Let $A \subseteq U$ and U be fuzzy (gsp) open. Since every fuzzy wg -open set is fuzzy (gsp) -open. We have $cl(int(A)) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy (gsp) -open in (X, τ) then $cl(int(A)) \subseteq U$ whenever $A \subseteq U$ and U is fuzzy (gsp) -open in (X, τ) . A is fuzzy strongly $(gsp)^*$ -closed.

III. FUZZY STRONGLY $(GSP)^*$ -CONTINUOUS MAPS

We introduce the following definitions

A. Definition 3.1

A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called fuzzy Strongly $(gsp)^*$ -continuous if $f^{-1}(V)$ is a fuzzy strongly $(gsp)^*$ -closed set in (X, τ) for every fuzzy closed set V of (Y, σ) .

fuzzy continuous map is fuzzy strongly $(gsp)^*$ -continuous

Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a fuzzy continuous map. Let us prove that f is fuzzy strongly $(gsp)^*$ -continuous. Let F be a fuzzy closed set in (Y, σ) . Since f is fuzzy continuous $f^{-1}(F)$ is closed in (X, τ) then $f^{-1}(F)$ is fuzzy strongly $(gsp)^*$ -closed so f is fuzzy strongly $(gsp)^*$ -continuous the converse of the above Theorem is not true

B. Theorem 3.2

Every fuzzy g -continuous map is fuzzy strongly $(gsp)^*$ -continuous

- 1) *Proof:* Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a fuzzy g -continuous. Let F be a fuzzy closed set in (Y, σ) . Since f is fuzzy g -continuous $f^{-1}(F)$ is fuzzy g -closed in (X, τ) . then $f^{-1}(F)$ is fuzzy strongly $(gsp)^*$ -closed so f is fuzzy strongly $(gsp)^*$ -continuous fuzzy closed g^{**} -closed fuzzy g -closed fuzzy wg -closed fuzzy sg^{**} -closed fuzzy g^* -closed fuzzy strongly $(gsp)^*$ -closed fuzzy $(gsp)^*$ -closed fuzzy Semi-closed sg^* -closed sg -closed. The converse of the above Theorem is not true

C. Theorem 3.3

Every g^* -continuous map is strongly $(gsp)^*$ -continuous.

- 1) *Proof:* Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a g^* -continuous. Let F be a closed set in (Y, σ) . Since f is g^* -continuous $f^{-1}(F)$ is g^* -closed in (X, τ) . By Theorem (3.6), $f^{-1}(F)$ is strongly $(gsp)^*$ -closed so f is strongly $(gsp)^*$ -continuous The converse of the above Theorem is not true

D. Theorem 3.4

Every fuzzy g^{**} -continuous map is fuzzy strongly $(gsp)^*$ -continuous.

- 1) *Proof:* Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a fuzzy g^{**} -continuous. Let F be a fuzzy closed set in (Y, σ) . Since f is fuzzy g^{**} -continuous $f^{-1}(F)$ is g^{**} -closed in (X, τ) . then $f^{-1}(F)$ is fuzzy strongly $(gsp)^*$ -closed so f is fuzzy strongly $(gsp)^*$ -continuous The converse of the above Theorem is not true.

IV. APPLICATIONS OF FUZZY STRONGLY $(GSP)^*$ -CLOSED SETS

As application of fuzzy strongly $(gsp)^*$ -closed sets, new spaces, namely fuzzy $Ts(gsp)^*$ space, $gTs(gsp)^*$ space, fuzzy $g^*T(gsp)^*$ space, fuzzy $g^{**}Ts(gsp)^*$ space are introduced. We introduced the following definitions.

A. Definition 4.1

A fuzzy space (X, τ) is called a fuzzy $Ts(gsp)^*$ -space if every fuzzy strongly $(gsp)^*$ -closed set is closed.

B. Definition 4.2

A fuzzy space (X, τ) is called a fuzzy $gTs(gsp)^*$ - space if every fuzzy strongly $(gsp)^*$ - closed set is fuzzy g closed.

C. Definition 4.3

A fuzzy space (X, τ) is called a fuzzy $g^*Ts(gsp)^*$ -space if every fuzzy strongly $(gsp)^*$ - closed set is g^* -closed

D. Definition 4.4

A fuzzy space (X, τ) is called a fuzzy $g^{**}Ts(gsp)^*$ -space if every fuzzy strongly $(gsp)^*$ - closed set is fuzzy g^{**} -closed.

2) **Theorem 4.1:** Every fuzzy $Ts(gsp)^*$ -space is fuzzy $T_{1/2}^*$ -space.

3) **Proof:** Let (X, τ) be a fuzzy $Ts(gsp)^*$ - space. Let us prove that (X, τ) is a fuzzy $T_{1/2}^*$ - space. Let A be a fuzzy g^* - closed set. Since every fuzzy g^* - closed set is fuzzy strongly $(gsp)^*$ -closed, A is fuzzy strongly $(gsp)^*$ - closed. Since (X, τ) is a fuzzy $Ts(gsp)^*$ - space, A is fuzzy closed. (X, τ) is a fuzzy $T_{1/2}^*$ - space.

4) **Theorem 4.2:** Every fuzzy $Ts(gsp)^*$ -space is fuzzy $gTs(gsp)^*$ -space.

5) **Proof:** Let A be a fuzzy strongly $(gsp)^*$ -closed set. Then A is fuzzy closed. Since the fuzzy space is $Ts(gsp)^*$ - space. And every closed set is fuzzy g -closed. Hence A is fuzzy g - closed. (X, τ) is a fuzzy $gTs(gsp)^*$ - space. The converse is not true.

6) **Theorem 4.3:** Every fuzzy $g^*Ts(gsp)^*$ -space is fuzzy $gTs(gsp)^*$ -space.

7) **Proof:** Let A be a fuzzy strongly $(gsp)^*$ -closed. Then A is fuzzy g^* -closed, since the fuzzy space is a fuzzy $g^*Ts(gsp)^*$ -space since. Every fuzzy g^* -closed set is fuzzy g -closed. Hence A is fuzzy g -closed then (X, τ) is a fuzzy $gTs(gsp)^*$ -space. The converse is not true.

8) **Theorem 4.4:** Every $g^*Ts(gsp)^*$ -space is $g^{**}Ts(gsp)^*$ -space.

9) **Proof:** Let A be a strongly $(gsp)^*$ -closed set. Then A is g^* -closed, since the space is a $g^*Ts(gsp)^*$.Since Every g^{**} -closed set is g^* -closed. Hence A is g^{**} -closed. (X, τ) is a $g^{**}Ts(gsp)^*$ -space.

REFERENCE

- [1]. B. Ghosh, Semi-continuous and semi-closed mappings and semi-connectedness in fuzzy setting, fuzzy Sets and Systems 35(3) (1990), 345–355.
- [2]. C. L. Chang, Fuzzy topological spaces, J. Math. Anal. Appl. 24 (1968), 182–190.
- [3]. C.W. Baker on preserving g -super closed sets Kyungpook Math. J. 36(1996), 195-199.
- [4]. G. Balasubramanian and P. Sundaram, On some generalizations of fuzzy continuous functions, Fuzzy Sets and Systems 86(1) (1997), 93–100.
- [5]. G. Balasubramanian and V. Chandrasekar, Totally fuzzy semi continuous functions, Bull. Calcutta Math. Soc. 92(4) (2000), 305–312.
- [6]. G. Balasubramanian, On fuzzy pre-separation axioms, Bull. Calcutta Math. Soc. 90(6) (1998), 427–434.
- [7]. K. K. Azad, On fuzzy semi continuity, fuzzy almost continuity and fuzzy weakly continuity, J. Math. Anal. Appl. 82(1) (1981), 14–32.
- [8]. K. M. Abd El-Hakeim, Generalized semi-continuous mappings in fuzzy topological spaces, J. Fuzzy Math. 7(3) (1999), 577–589.
- [9]. L. A. Zadeh, Fuzzy sets, Information and Control 8 (1965), 338–353.
- [10]. M.K. Mishra et al on “Fuzzy super continuity” International Review in Fuzzy Mathematics July –December 2012.
- [11]. M.K. Mishra M. Shukla M. Fuzzy Regular Generalized Super Closed Set” International Journal of Scientific and Research December issue July December 2012.
- [12]. M.K. Mishra, et al on “ Fuzzy super closed set” International Journal International Journal of Mathematics and applied Statistics.
- [13]. P. M. Pu and Y. M. Liu Fuzzy topology I Neighborhood structure of a fuzzy point and More-Smith Convergence. J. Math. Anal. Appl. 76(1980) ,571-594.
- [14]. P. M. Pu and Y. M. Liu Fuzzy topology II Product and quotient spaces J.Math. Anal. Appl. 77(1980) 20-37.
- [15]. P. M. Pu, and Y. M. Liu, Fuzzy topology. I. Neighborhood structure of a fuzzy point and Moore-Smith convergence, J. Math. Anal. Appl. 76(2) (1980), 571–599.
- [16]. R. K. Saraf and M. Khanna, On gs -closed sets in fuzzy topology, J. Indian Acad. Math. 25(1),(2003), 133–143.
- [17]. R. K. Saraf, and M. Khanna, Fuzzy generalized semi pre closed sets, J. Tripura Math. Soc.3(2001) 59–68.
- [18]. R. K. Saraf, and S. Mishra, Fg_- -closed sets, J. Tripura Math. Soc. 2 (2000) 27–32.
- [19]. R. K. Saraf, M. Caldas and S. Mishra, Results via Fg_- -closed sets and Fg -closed sets, Preprint.
- [20]. T. H. Yalva, c, Semi-interior and semi closure of a fuzzy set, J. Math. Anal. Appl. 132(2) (1988), 356–364.

10.22214/IJRASET

45.98

IMPACT FACTOR:
7.129

IMPACT FACTOR:
7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)