iJRASET

International Journal For Research in
Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: IV Month of publication: April 2018

DOI: <http://doi.org/10.22214/ijraset.2018.4759>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Synthesis and Biological Evaluation of N-(4-(5-Aryl-4, 5-Dihydro-1H-Pyrazol-3-yl) Phenyl) Cyclopropane Carboxamide

P. M. Akbari¹, K. D. Ladva² and V. R. Shah³

^{1, 2, 3} Department of Chemistry, Kamani Science College, Amreli-365601 Gujarat, India.

Abstract: Different pyrazoline derivatives were synthesized by cyclization of substituted chalcones with Hydrazine hydrate. Some new N-(4-(3-Aryl-acryloyl)phenyl)cyclopropane carboxamide (1a-l) and N-(4-(5-Aryl-4,5-dihydro-1H-pyrazol-3-yl)phenyl)cyclopropane carboxamide (2a-l) were prepared. All the prepared compounds were characterized by their spectral (I.R., N.M.R., Mass) data and screened for their antimicrobial activities.

Keywords: Chalcones, Pyrazoline, Antimicrobial activities.

I. INTRODUCTION

The chemistry of chalcones containing an active keto-ethylenic linkage has been assumed important because of their versatility in the synthesis of many heterocyclic compounds. Pyrazoline derivatives have been found to contain wide range of therapeutic activity such as antidiabetic⁷, antiimplantation⁸, antiallergic⁹, anticonvulsant¹⁰⁻¹¹, antineoplastic¹², antiinflammatory¹³, antitumor¹⁴, analgesic¹⁵⁻¹⁶, antimicrobial¹⁷, bactericidal¹⁸⁻²⁰ etc. Pyrazolines proved to be the most useful framework for biological activities, both have attracted attention of medicinal chemists for both with regard to heterocyclic chemistry and the pharmacological activities associated with them. This inspired us to synthesize N-(4-(5-Aryl-4,5-dihydro-1H-pyrazol-3-yl)phenyl)cyclopropane carboxamide (2a-l). The structure of synthesized compounds were assigned based on Elemental analysis, I.R. ¹H-NMR and Mass Spectral data. The antimicrobial activity was assayed by using the cup-plate agar diffusion method²⁰ by measuring the zone of inhibition in mm. All the synthesized compounds have been evaluated for their antibacterial activity towards Gram positive bacterial strains such as B.subtilis and S.aureus whereas E.coli and P.aeruginosa were Gram negative bacterial strains and antifungal activity towards A.niger at a concentration of 40 µg and synthesized compounds has been compared with standard drugs. Standard drugs like Ampicillin, Chloramphenicol, Norfloxacin, and Griseofulvin were used for comparison purpose. (Table-1)

II. EXPERIMENTAL SECTION

Melting points were taken in open capillary tubes are uncorrected. IR spectra (cm⁻¹) were recorded on Shimadzu-435-IR Spectrophotometer and ¹H-NMR Spectra on Bruker Spectrometer (400MHz) using TMS as an internal standard, chemical shift in δ ppm.

A. General Procedure for the Preparation of n-(4-(3-(4-Methoxyphenyl)Acryloyl)phenyl) Cyclopropane Carboxamide (1a-l)

A mixture of N-(4-acetylphenyl)cyclopropane carboxamide 0.5 gm(0.01 mol) with 4-methoxy benzaldehydes 0.33 gm/0.29 ml (0.01 mol) using Claisen-Schmidt condensation method in presence of 40% NaOH using methanol as a solvent at room temperature under

stirring for 8 hours. Reaction was monitored by TLC. Reaction mass was poured into chilled water. Product was filtered and dried. It was recrystallized from ethanol. Yield 81.25%, M.P.162-164°C, Elemental Analysis Calculated for $C_{20}H_{19}NO_3$ Requires: C-74.75%; H-5.96%, N-4.36%; O-14.94%, Found: C-74.70%; H-5.93; N-4.31%; O-14.91%,

B. N-(4-(3-(4-Methoxyphenyl)Acryloyl)phenyl)Cyclopropane Carboxamide (1a-l)

Yield 81.25%, M.P.162-164°C; IR(KBr) : ν Alkane C-H str. (asym.) 2938, C-H def.(asym.) 1417, , C-H o.o.p.(def) 1352, Aromatic C-H str. 3040 ,C=C str.1598,1511, Amine C-N str. 1294 ,N-H str. 3241, Ether C-O-C str. 1256, Ketone C=O str. 1658, Vinyl CH=CH str. 3040, cm^{-1} ; 1H -NMR ($CDCl_3$) : δ 0.80-1.51, (m,5H, Cyclopropane) ,3.748 (s, 3H,-OCH₃), 7.19 & 7.37 (d-d, 2H, CH=CH) , 6.85-7.86 (m,8H, Ar-H), 10.48 (s, 1H, 2°Amide), Mass m/z 322.5 (M^+); .M.F.: $C_{20}H_{19}NO_3$

C. General Procedure for the Preparation of N-(4-(5-(4-Methoxyphenyl)-4,5-Dihydro-1H-Pyrazol-3-yl)Phenyl)Cyclopropane Carboxamide (2a-l)

A mixture of N-(4-(3-(4-methoxyphenyl)acryloyl)phenyl)cyclopropane carboxamide 0.5 gm(0.01 mol) with hydrazine hydrate 0.23 gm/0.22 ml (0.03 mol) in methanol was refluxed about 8 hours until complete the reaction which was monitored by TLC. The reaction mass was poured into chilled water and product was filtered and dried. It was recrystallized from ethanol. Yield 81.73%, M.P.210-212°C, Elemental Analysis Calculated for $C_{20}H_{21}N_3O_2$ Requires:C-71.62%;H-6.31%;N-12.53%; O-9.54%, Found: C-71.58%; H-6.28%; N-12.50%; O-9.49%,

D. N-(4-(5-(4-Methoxyphenyl)-4,5-Dihydro-1H-Pyrazol-3-yl)phenyl)Cyclopropane Carboxamide (2a-l)

Yield 81.73%, M.P.210-212°C IR(KBr) : Alkane C-H str. (asym.) 2942, Aromatic C=C str. 1595, 1511, C-H o.o.p.(def) 821, Ether C-O-C str. 1246, Pyrazoline C=N str. 1595, N-H str.3328 cm^{-1} .

1H -NMR (DMSO) : δ 0.80-1.79, (m,5H, Cyclopropane), 3.73 (s, 3H,-OCH₃), 4.75 (t, 1H, Pyrazoline ring), 7.4, (s, 1H,Pyrazoline – NH), 2.77, 3.39 (d, 2H,Pyrazoline ring), 10.28 (s, 1H, 2°Amide), 6.88-7.62 (m,8 H, Ar-H). .Mass m/z 336.5 (M^+) . M.F. $C_{20}H_{21}N_3O_2$

III. REACTION SCHEME

IV. ANTIBACTERIAL ACTIVITY

It has been observed from the microbiological data that all compounds (1a-l) and (2a-l) were found to be mild to moderately active against Gram positive and Gram negative bacterial strains.

V. ANTIFUNGAL ACTIVITY

The antifungal data revealed that compounds were least toxic to the fungal strain. However mild activity was shown by the compounds against A.niger. The antibacterial activity was compared with standard drug viz. Ampicillin, Chloramphenicol, Norfloxacin and antifungal activity was compared with standard drug viz. Griseofulvin.

TABLE-1

Characterization data of the compounds (2a-l) :						
Compound No.	R	Molecular Formula	Molecular Weight	M.P. (°C)	Nitrogen %	
					Found	Calcd
2a	-C ₆ H ₅	C ₁₉ H ₁₉ N ₃ O ₃	305.37	166-170	13.71	13.76
2b	-4-OCH ₃ -C ₆ H ₄	C ₂₀ H ₂₁ N ₃ O ₂	335.40	210-212	12.50	12.53
2c	-4-N(CH ₃) ₂ -C ₆ H ₄	C ₂₁ H ₂₄ N ₄ O	348.44	212-215	16.03	16.08
2d	-C ₄ H ₉ O	C ₁₇ H ₁₇ N ₃ O ₂	295.34	208-210	14.20	14.23
2e	-2-Cl-C ₆ H ₄	C ₁₉ H ₁₈ ClN ₃ O	339.82	208-210	12.32	12.37
2f	-4-F-C ₆ H ₄	C ₁₉ H ₁₈ FN ₃ O	323.36	90-92	12.95	12.99
2g	-4-OH-C ₆ H ₄	C ₁₉ H ₁₉ N ₃ O ₂	321.37	178-181	13.06	13.08
2h	-4-OH-3-OCH ₃ -C ₆ H ₃	C ₂₀ H ₂₁ N ₃ O ₃	351.40	192-194	11.95	11.96
2i	-2-OH-C ₆ H ₄	C ₁₉ H ₁₉ N ₃ O ₂	321.37	175-177	13.05	13.08
2j	-2-NO ₂ -C ₆ H ₄	C ₁₉ H ₁₈ N ₄ O ₃	350.37	165-168	15.94	15.99
2k	-4-Cl-C ₆ H ₄ -	C ₁₉ H ₁₈ ClN ₃ O	339.82	203-205	12.33	12.37

TABLE-2

Antimicrobial Activity: (Zone of inhibition in mm) :					
Compound No.	E. coli	B.subtilis	P.aeruginosa	S. aureus	A . Niger
2a	16	10	12	13	08
2b	10	15	10	12	07
2c	11	16	11	10	10
2d	12	11	12	11	11
2e	10	9	10	12	09
2f	8	11	11	10	12
2g	11	12	10	11	13
2h	12	13	9	12	11
2i	15	11	11	15	10
2j	14	16	12	16	9
2k	11	18	10	13	8
Ampicillin	18	19	13	10	0
Chloramphenicol	13	15	15	12	0
Norfloxacin	15	14	12	13	0
Griseofulvin	0	0	0	0	14

VI. RESULTS AND DISCUSSION

The synthesis of N-(4-(3-Aryl-acryloyl)phenyl)cyclopropane carboxamide (1a-l) and N-(4-(5-Aryl-4,5-dihydro-1H-pyrazol-3-yl)phenyl)cyclopropane carboxamide (2a-l) was carried out in two steps, first by the condensation of N-(4-acetylphenyl)cyclopropane carboxamide) with different aromatic aldehydes by Claisen-Schmidt condensation in presence base catalyst to give chalcone derivatives (1a-l), which in next step were refluxed with hydrazine hydrate to yield pyrazoline derivatives (2a-l) (Reaction Scheme).

The formulas of the selected compounds were confirmed by the Elemental analysis and their structures were determined by IR, ^1H -NMR, and Mass Spectral data.

VII. CONCLUSION

The recent study leads to a convenient synthetic method for the synthesis of new compounds which show significant antibacterial and antifungal activities. Further investigation with appropriate structural modification of the above compounds may result in therapeutically useful products.

VIII. ACKNOWLEDGMENT

The authors are very much thankful to authorities of Kamani Science College, Amreli for providing research facilities & Principal and Management of Shree M.& N. Virani Science College, Rajkot for providing Spectral data of IR and MASS and also thankful to Department of Chemistry Saurashtra University Rajkot for I.R., N.M.R., Mass Spectral & Elemental analysis. The authors are thankful to Pramukh Swami Science College, Kadi for providing Antimicrobial activities data.

REFERENCES

- [1] M. Fahmy, M. Hussan, A. A. Khalt; R. A. Ahmedi, Rev. Roum-Chim., 33(7), 755-61 (1988); Chem. Abstr., 111, 77898 (1989).
- [2] M. Hassaneen Hamdi, A. E. Hamad; Salter Lett., 8(5), 27582 (1989); Chem. Abstr., 111, 57611 (1989)
- [3] M. A. El. Hashah, M., El-Kady, M. A. Saiyed, A. A. Elsayy Egypt. J. Chem., 27(6), 715-21 (1985); Chem. Abstr., 105, 20868u (1986).
- [4] V. Gohanmukkala, A. Subbaraju, R. Naykula and D. Parmeswara, Indian J. Heterocyclic Chem., 4, 87-92 (1994)
- [5] V.R. Dangar, V.R. Shah ; The International Journal Of Science & Technoledge (ISSN 2321 – 919X) , Vol 2 Issue, 1 January, 2014.
- [6] V. R. Dangar, K. N. Borkhataria and V. R. Shah; Sci. Revs. Chem. Commun.: 4(1) 2014,31-37 ISSN 2277-2669.
- [7] H. G. Garg and P. P. Singh; J. Chem. Soc., 2, 1141 (1936)
- [8] D. B. Reddy, T. Senshuna and M. V. Ramma Reddy; Indian J. Chem., 30B, 46 (1991)
- [9] B. Roman; Pharmazie, 45, 214 (1990)
- [10] Z. Brozowski, E. Pormarnacka; Acta. Pol. Pharm., 37(4), 1378, 80 (1980); Chem. Abstr., 25, 80807 (1981).
- [11] Archana Shrivastava V. K.; Chandra Ramesh, Kumar Ashok; Indian J. Chem., 41B, 2371-75 (2002); Chem. Abstr., 138, 271582 (2003)
- [12] H. M. Mokhtar, H. M. Faidallah; Pharmazie, 42, 482 (1987)
- [13] Ashok Kumar, R. S. Verma and B. P. Jagu; J. Ind. Chem. Soc., 67, 120 (1990)
- [14] W. I. Ronald, A. Adriano; Chem. Abstr., 126, 181346f (1997)
- [15] Delay Francois (Fermenich S. A.) Patent Schrift (Switz); Chem. Abstr., 117, 90276f (1992)
- [16] Ayses G., Seref D., Gultaze C., Kevser E., Kamil V.; Eur. J. Med. Chem. 35, 359-64 (2002)

- [17] Panda J. Srinivas S. V., Rao M. E.; J. Indian Chem. Soc., 79(9), 770-1 (2002) Chem. Abstr., 138, 153499n (2003)
- [18] P. Desaea; A. Nunrich; M. Carderny and G. Devaux; Eur. J. Med. Chem., 25, 285 (1990)
- [19] Kalluraya Balakrishna, Chimabalkar R., Rai G., Gururaja R., Shenoy S.; J. Indian Coun. Chemi. 18(2), 39-43 (2001) Chem. Abstr., 138, 238061.
- [20] A. L. Barry; The antimicrobial susceptibility test: Principle and practices, edited by Illuslea & Febiger, (Philadelphia), USA, 180; Biol. Abstr., 1977, 64, 25183
- [21] N. N. Kansagara, V. R. Dangar, and V. R. Shah; Inter. J. of Pharma. Sc. & Research (IJPSR), Vol 6 No 01, (Jan 2015).
- [22] J. V. Dodia, V. R. Dangar and V. R. Shah; World Journal. of Pharma. Research, Issue 9, Vol 6, Issue 9, 809-814, (2017); ISSN 2277– 7105.

10.22214/IJRASET

45.98

IMPACT FACTOR:
7.129

IMPACT FACTOR:
7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)