iJRASET

International Journal For Research in
Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: XI Month of publication: November 2019

DOI: <http://doi.org/10.22214/ijraset.2019.11024>

www.ijraset.com

Call: 08813907089

E-mail ID: ijraset@gmail.com

Diseases caused by Trichophyton Tonsurans and their Treatment: A Review

Arpit Anand¹, Suresh K Jatawa², Gyaneshwar Patel³, Alok Tripathi⁴, Archana Tiwari⁵

¹Student, School of Biotechnology, Rajiv Gandhi Proudlyogiki Vishwavidyalaya, Bhopal, Madhya Pradesh, India

²Assistant Professor, School of Biotechnology, Rajiv Gandhi Proudlyogiki Vishwavidyalaya, Bhopal, Madhya Pradesh, India

³Research Assistant, Chaperon Biotech. Pvt. Ltd., Kanpur, Uttar Pradesh, India.

⁴Lab. In charge, Chaperon Biotech Pvt. Ltd., Kanpur, Uttar Pradesh, India

⁵Head of the Department, School of Biotechnology, Rajiv Gandhi Proudlyogiki Vishwavidyalaya, Bhopal, Madhya Pradesh, India

Abstract: *Trichophyton tonsurans* is an anthropophilic dermatophyte, responsible for infections of the scalp and sometimes of the glabrous skin or nails. *Tinea capitis*, the most common dermatophytosis in children, is an infection of the scalp and hair shafts. It can be transmitted by poor hygiene and overcrowding, and can occur through contaminated hats, brushes, pillowcases, and other inanimate objects. *Tinea corporis* typically appears as single or multiple, annular, scaly lesions with central clearing, a slightly elevated, reddened edge, and sharp margination (abrupt transition from abnormal to normal skin) on the trunk, extremities, or face. *Tinea unguium*, a dermatophyte infection of the nail, is a subset of onychomycosis, which also may be caused by yeast and non-dermatophyte molds. Risk factors for this infection include aging, diabetes and poorly fitting shoes. It is characterized by thickened, brittle and discolored nail. Dermatophyte infections can be readily diagnosed based on the history, physical examination, and potassium hydroxide (KOH) microscopy. Diagnosis occasionally requires Wood's lamp examination and fungal culture or histologic examination. Topical therapy is used for most dermatophyte infections. Topical Terbinafine, among all different alternatives, for about a month has all the attribute of being the treatment of decision for *tinea corporis*, *cruris* and *pedis*.

I. INTRODUCTION

Trichophyton tonsurans is an anthropophilic dermatophyte, responsible for infections of the scalp and sometimes of the glabrous skin or nails. Unlike dermatophytosis caused by other dermatophytes, the clinical features of infection due to *T. tonsurans* are not very apparent initially. *Tinea corporis* due to *T. tonsurans* appears as small, erythematous, scaly plaques, often measuring only 1-2 cm in diameter, have no central clearing, and are similar in appearance to eczema. *Tinea capitis* due to *T. tonsurans* can be subdivided into three types: the seborrheic subtype, which is primarily characterized by the presence of dandruff and crusts; the kerion celsi subtype; and the black dot subtype (1-4).

A. *Tinea Capitis*

Trichophyton tonsurans, an anthropophilic dermatophyte, is the most normal reason for *tinea capitis* in North America. Other causes: *Microsporum canis* (more often than not from pets), *Microsporum audouinii*, *Trichophyton mentagrophytes*, *Trichophyton violaceum*. *Tinea capitis* is prevalent in youngsters 3 to 10 years old; family individuals in danger; can happen in postmenopausal youngster caretakers. *T. tonsurans* is transmitted from human to human. *M. canis* is transmitted from pets (hounds, felines).

The differential diagnosis for *tinea capitis* includes alopecia areata, atopic dermatitis, folliculitis, impetigo, lupus erythematosus, psoriasis, seborrheic dermatitis, traction alopecia, and trichotillomania. [5] Standard treatment recently comprised of griseofulvin, 20 mg/kg every day (limit of 1 g) in micronized structure (accessible in suspension), or 10 to 15 mg/kg every day (limit of 750 mg) in ultramicronized structure. Since it is oftentimes the main specialist secured by protection plans, griseofulvin stays a well known, generally utilized treatment. The term of treatment for the most part is 6 to about two months. As of late, numerous suppliers have discovered that higher dosages of micronized griseofulvin (up to 25 mg/kg every day) and longer spans of treatment are here and there required for cure. [6] *M. canis* sickness frequently is more hard to treat than *T. tonsurans*, what's more, higher portion and longer span of treatment regularly are required to accomplish total resolution. [7,8] Administration of griseofulvin with a greasy supper (e.g., milk, frozen yogurt, nutty spread) improves medicate assimilation; research center tests to screen for antagonistic impacts are not required except if treatment surpasses two months. A forthcoming preliminary utilizing micronized griseofulvin (20 mg/kg/day) and topical treatment with twice-week after week selenium sulfide cleanser for about a month and a half demonstrated mycologic, clinical, and complete fix paces of 89%, 66%, and 49%, respectively. [9] Common side impacts of griseofulvin incorporate sickness, migraine, and photosensitivity responses.

B. *Tinea Corporis*

Tinea corporis is a shallow skin disease that happens worldwide and influences all age gatherings. There is a preference for guys, and about half of cases in the US happen in kids <15 long stretches of age.[10] Any dermatophyte can cause sickness, yet the prevalent pathogen differs topographically. *T. rubrum*, *T. tonsurans*, *T. mentagrophytes*, *M. canis*, and *E. floccosum* are the most widely recognized reasons for *tinea corporis* in the US. Securing happens through direct contact with tainted people or creatures (basically hounds also, felines) and, less ordinarily, through presentation to debased fomites. The infectivity and provocative potential change by species. Flare-ups among wrestlers are normal and have been named *tinea gladiatorum* (*tinea capitis* additionally can be available in influenced individuals).[11] *T. tonsurans* is the most widely recognized reason for *tinea gladiatorum* in the US.[12] Three clinical types of ailment happen: papulosquamous, incendiary/ vesicular, and granulomatous. Papulosquamous ailment is the most normal and shows as an erythematous papule or plaque that develops into an annular ring injury with well-characterized edges—the work of art "ringworm." Infection spreads to include encompassing solid skin, frequently with a focal territory of clearing. Incendiary or vesicular *tinea corporis* regularly happens with *M. canis* contamination and regularly has fine vesicular sores at the plaque's propelling edge. By and large, staying discrete sores may combine. Granulomatous *tinea corporis*, regularly alluded to as Majocchi granuloma, is the least normal type of ailment and regularly happens in kids whose underlying disease was misdiagnosed and treated with topical corticosteroid specialists, in immunocompromised hosts, or in females who have reinoculated the growth while shaving their legs.[13] The granulomas structure when dermatophytes inside hair follicles break into the dermis and cause a fiery reaction. The disease is subacute or ceaseless and shows as firm, nontender skin knobs with an overlying covering or plaque; injuries related with leg shaving are follicular or on the other hand perifollicular and regularly have a circumferential scale on an erythematous base. Most types of *tinea corporis* can be treated with on more than one occasion day by day topical antifungal treatment for 14 to 21 days. Ideal span of treatment what's more, near adequacy of topical operators have not been assessed widely. Patients with across the board cutaneous sores or a granulomatous response once in a while are dealt with fundamentally with terbinafine, griseofulvin, or elective antifungal specialists. Most suggest treatment with oral specialists, and in an examination utilizing fluconazole 200 mg week by week, 100% of influenced people were culture antagonistic by the third week.[14] Several studies have archived a slight predominance and less reactions utilizing more up to date specialists, for example, terbinafine and fluconazole, contrasted and griseofulvin.[15,16] *Tinea gladiatorum* raises specific worry of spread to different competitors. It is hazy to what extent contaminated people ought to maintain a strategic distance from contact with others, yet numerous specialists prescribe 10 to 15 days after beginning of treatment.

C. *Tinea Unguium (Onychomycosis)*

Tinea unguium is a dermatophytic disease of the nails that is regularly connected with *tinea pedis*. Commonness increments with age and is exceptional in prepubertal kids. Overall predominance ranges from 0.1% to 1%.[17] Adults have a higher predominance at 2% to 14%, and pervasiveness in old individuals approaches 40%.[17] moreover, kids with onychomycosis regularly have more seasoned influenced family members.[18] *T. rubrum*, *T. mentagrophytes*, and *E. floccosum* are the essential pathogens of *tinea unguium*. [19,20] Although clinical introduction can change, most usually the organism legitimately attacks keratin along the distal and horizontal parts of the nail and gradually spreads proximally along the nail.[21] The nail turns out to be "delicate" and deformed with subungual flotsam and jetsam and nail staining. A progressively shallow disease makes central, sporadic hazy injuries on the nail. Treatment of onychomycosis incorporates oral and topical treatments. Griseofulvin is regularly utilized in youngsters, however it has over and again been recorded in grown-ups that terbinafine prompts higher fix rates in onychomycosis than does griseofulvin. Most specialists use terbinafine if monetary also, therapeutic contemplations don't contraindicate use. Choices treatments incorporate itraconazole and fluconazole. The span of griseofulvin treatment is drawn out (6–12 months), and repeat is normal. Terbinafine and the triazoles commonly are regulated for shorter spans (3 months) or as heartbeat regimens; in any case, they are not affirmed by the FDA for the treatment of pediatric onychomycosis.[22,23] furthermore, mechanical debridement with 40% urea balm applied every day for 10 days is now and again utilized in cases hard-headed to fundamental treatment or when fundamental treatment is contraindicated. Traditionally, topical antifungal treatment has included ciclopirox and amorolfine. More up to date specialists that have been demonstrated to be sheltered and successful incorporate efinaconazole and tavaborole.[24,25–26] when all is said in done, topical specialists moreover require a drawn out course of treatment (6–12 months) yet might be less successful than oral drugs on account of failure to enter to the most reduced segments of the nail bed. In this manner topical treatment by and large is utilized just if there is no nail network (lunular) contribution. Little youngsters have slender, quickly developing nails that may take into account upgraded sedate infiltration, and preliminaries are in progress to assess the new topical operators' adequacy in

youth disease.[27] Carbon dioxide laser treatment may increment in filtration and viability in mix with topical antifungal treatment however, has not demonstrated to be exceptionally efficacious.[28]

II. CONCLUSION

Treatment of cutaneous dermatophytosis has progressively become troublesome, and dermatologists been compelled to think past standard way of thinking to counter this hazard. Despite the fact that there is adequate proof to exhibit the viability of topical antifungal in restricted infection yet, there is rare information on the recurrence of backslide once topical monotherapy stopped. Among different alternatives, topical terbinafine for about a month has all the earmarks of being the treatment of decision for restricted malady (tinea corporis/cruris/pedis). For broad sickness, the decision is less clear. Both terbinafine (250–500 mg/day for 2 a month and a half) and itraconazole (100– 200 mg/day for 2 a month) seem, by all accounts, to be successful. Be that as it may, a fitting portion and term of organization that can create mycologic fix and avoid repeat stays subtle. This audit too features the gigantic research holes in the administration of cutaneous dermatophytosis that need not stopped to give better and viable consideration to the patients. RCTs that are increasingly stringent are the need of great importance looking at the different oral antifungal treatments to give an unmistakable thought with respect to proper portion and term of treatment.

REFERENCES

- [1] Suh SB, Kim SW, Oh SH, Choi SK, Bang YJ. A case of block dot ringworm caused by trichophyton tonsurans. Korean J Dermatol 1998; 36: 918-23.
- [2] Kim HY, Kim SL, Lee WJ, Lee SJ, Kim DW, Jun JB. Intra-familial infection due to Trichophyton tonsurans in Korean general population. Korean J Med Mycol 2014; 19: 105-9.
- [3] Hiruma J, Ogawa Y, Hiruma M. Trichophyton tonsurans infection in Japan: epidemiology, clinical features, diagnosis and infection control. J Dermatol 2015; 42: 245-9.
- [4] Alshawa K, Lacroix C, Benderdouche M, Mingui A, Derouin F, Feuillade de Chauvin M. Increasing incidence of Trichophyton tonsurans in Paris, France: a 15-year retrospective study. Br J Dermatol 2012; 166:1149-50.
- [5] Slowinska M, Rucnicka L, Schwartz RA, et al. Comma hairs: a dermatoscopic marker for tinea capitis—a rapid diagnostic method. J Am Acad Dermatol 2008;59(suppl):877–879.
- [6] Kakourou T, Uksal U, European Society for Pediatric Dermatology. Guidelines for the management of tinea capitis in children. Pediatr Dermatol 2010;27:226–229.
- [7] Skerlev M, Miklic P. The changing face of Microsporum spp infections. Clin Dermatol 2010;28:146–150.
- [8] Lipozencic J, Skerlev M, Orofino-Costa R, et al. A randomized, double-blind, parallel group, duration finding study of oral terbinafine and open-label, high-dose griseofulvin in children with tinea capitis due to Microsporum species. Br J Dermatol 2002;146:816–823.
- [9] Lorch Dauk KC, Comrov E, Blumer JL, et al. Tinea capitis: predictive value of symptoms and time to cure with griseofulvin treatment. Clin Pediatr (Phila) 2010;49:280–286.
- [10] Panackal AA, Halpern EF, Watson AJ. Cutaneous fungal infections in the United States: Analysis of the National Ambulatory Medical Care Survey (NAMCS) and National Hospital Ambulatory Medical Care Survey (NHAMCS), 1995–2004. Int J Dermatol 2009;48:704–712.
- [11] Kohl TD, Martin DC, Nemeth R, et al. Fluconazole for the prevention and treatment of tinea gladiatorum. Pediatr Infect Dis J 2000;19:717–722.
- [12] Adams BB. Tinea corporis gladiatorum. Am Acad Dermatol 2002;47:286–290.
- [13] Wilson JW, Plunkett OA, Gregersen A. Nodular granulomatous perifollicle
- [14] Kohl TD, Martin DC, Nemeth R, et al. Fluconazole for the prevention and treatment of tinea gladiatorum. Pediatr Infect Dis J 2000;19:717–722.
- [15] Bourlond A, Lachapelle JM, Aussems J, et al. Double-blind comparison of itraconazole with griseofulvin in the treatment of tinea corporis and tinea cruris. Int J Dermatol 1989;28:410–412.
- [16] Cole GW, Stricklin G. A comparison of a new oral antifungal, terbinafine, with griseofulvin as therapy for tinea corporis. Arch Dermatol 1989;125:1537–1539.
- [17] Feldstein S, Totri C, Friedlander SF. Antifungal therapy for onychomycosis in children. Clin Dermatol 2015;33:333–339.
- [18] Gupta AK, Skinner AR. Onychomycosis in children: a brief overview with treatment strategies. Pediatr Dermatol 2004;21:74–79.
- [19] Hay RJ. Onychomycosis. Dermatol Clin 1993;11:161–169.
- [20] Gupta AK, Chang P, Del Rosso JQ, et al. Onychomycosis in children: prevalence and management. Pediatr Dermatol 1998;15:464–471.
- [21] Kearsle HL, Miller OF 3rd. Tinea pedis in prepubertal children: does it occur? J Am Acad Dermatol 1988;19:619–622.
- [22] Huang PH, Paller AS. Itraconazole pulse therapy for dermatophyte onychomycosis in children. Arch Pediatr Adolesc Med 2000;154:614–618.
- [23] Rosen T, Friedlander SF, Kircik L, et al. Onychomycosis: epidemiology, diagnosis, and treatment in a changing landscape. J Drugs Dermatol 2015;14:223–228.
- [24] Elewski BE, Aly R, Baldwin SL, et al. Efficacy and safety of tavaborole topical solution, 5%, a novel boron-based antifungal agent, for the treatment of toenail onychomycosis: results from 2 randomized phase-III studies. J Am Acad Dermatol 2015;73:62–69.
- [25] Bhatt V, Pillai R. Efinacozole topical solution, 10%: formulation development program of a new topical treatment of toenail onychomycosis. J Pharm Sci 2015;104:2177–2182.
- [26] Coronado D, Merchant T, Chanda S, Zane LT. In vitro nail penetration and antifungal activity of tavaborole, a boron-based pharmaceutical. J Drugs Dermatol 2015;14:609–614.
- [27] Singal A, Khanna D. Onychomycosis: diagnosis and management. Ind J Dermatol Venerol Leprol 2011;77:659–672.
- [28] Lin E, Kim H, Park Y, et al. Toenail onychomycosis treated with a fractional carbon dioxide laser and topical antifungal cream. J Am Acad Dermatol 2014;70:918–923.

10.22214/IJRASET

45.98

IMPACT FACTOR:
7.129

IMPACT FACTOR:
7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)