iJRASET

International Journal For Research in
Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: II Month of publication: February

DOI: <http://doi.org/10.22214/ijraset.2019.2077>

www.ijraset.com

Call: 08813907089

E-mail ID: ijraset@gmail.com

The Effect and Clinical Relevance of Superoxide Dismutase and RBC Glutathione Levels in Type-2 Diabetic Male and Female in Kolkata, West Bengal, India

Rajarshi Kayal¹, Krishanu Banik², Saheli Kayal³, Debojyoti Bhattacharjee⁴, Sudip K Banerjee⁵

¹Researcher, Techno India University, EM-4/1, Sector-V, Salt Lake, Kolkata-700091, India

²Assistant Professor, Department of Medicine, College of Medicine and JNM Hospital, The WBUHS, Kalyani, Nadia, West Bengal 741235, India.

³Assistant Teacher, Birla Bharati, Taratolla, Santoshpore New Road, Near Nature Park, P. O.-Bidhangarh, Kolkata, West Bengal 700066, India

⁴Dept of Biochemistry, Institute of Post Graduate Medical Education & Research (S.S.K.M. Hospital) Kolkata West Bengal India.

⁵Professor, Techno India University, Department of Biochemistry, EM -4 Saltlake Sector -5, Kolkata -700091

Abstract: Diabetes mellitus (DM) is a group of metabolic diseases characterized by hyperglycemia resulting from defects in insulin secretion, insulin action or both. This study reveals the role of antioxidant enzymes SOD and RBC Glutathione as major source of antioxidant status in type-2 diabetes. A total 360 subjects were involved in the study of which 90 were male diabetics (cases), 90 were female diabetics (cases) and 90 male non diabetics (control), 90 females non diabetics (control). The mean values SOD and RBC Glutathione were significantly lower ($p < 0.05$) among cases for both sexes compared to control. The mean values for FBS, PPBS, HbA1c%, were significantly higher ($p < 0.05$) among cases for both sexes compared to controls. In DM there is increased level of oxidative stress in humans.

Keywords: Diabetes mellitus, Superoxide dismutase, RBC Glutathione, Antioxidants, Hyperglycemia

I. INTRODUCTION

Diabetes mellitus may be recognized as a group of metabolic disease which might be characterized by increased level hyperglycemia which is a result of the defects in insulin secretion and its consecutive action or both. (1) Chronic hyperglycemia is associated with the damages in long run, of various organ in the body which includes eyes, kidneys, nerves, heart and blood vessels etc. It has also been seen that there is altered metabolism of carbohydrates lipids, proteins with increased risk of vascular disease. Regardless of the exact mechanism the reactive oxygen species (ROS) has been implicated to gear these diseases in the humans. Reactive oxygen species or free radicals has a very important role either in-vivo or in-vitro of lipids resulting in tissue damage and death of affected cells.

The primary target of ROS attack are PUFA (poly unsaturated fatty acid) in a membrane lipids causing lipid peroxidation which leads to disorganization of cells structure and function. (2) The harmful effect of ROS are counter attacked by class of protective agents termed as antioxidants which prevents the oxidative damage by reaction of free radicals before any other molecule can become a target.

The major types of antioxidant enzymes are superoxide dismutase (SOD), catalase, RBC Glutathione which are found to play a very important role in the protection of cells and tissues. (3,4). The evidence suggests that the oxidative stress plays a role in the diabetes mellitus pathogenesis and complication (5).

Hyperglycemia always increase oxidative stress which contribute to the main process of impairment that is the insulin action and insulin secretion.

Thus to protect the cell and the organ of the body against ROS humans have a very complex antioxidative protection (6,7,8). Free radicals has been found to effect the insulin signaling cascade (9). In certain case oxidative stress caused by hyperglycemia in diabetes patients impair insulin signaling which leads to insulin resistance and certain other mechanism which have not been completely established. (10,11).

Several studies have also shown that the diabetes mellitus is accompanied by increased formation of free radicals and decreased antioxidant activity which leads to oxidative damage of cell component.(12). Under diabetic condition the reactive oxygen species are produced through the glycation reaction which occurs in various tissues playing the major role in the development of diabetic complication (13). Although the induction of glycation reaction in diabetes has been found in neural cells and lens crystal which are actually known as target diabetic complication. In this background a study has been formulated to estimate the antioxidant status in chronic type 2 Diabetes mellitus (T2DM) patients.

II. MATERIALS AND METHODS

The study was carried out at Department of Biochemistry, Calcutta National Medical college and Hospital, Kolkata, West Bengal. Total three hundred and sixty (360) subjects age between 18 to 45 years constituted the study population. One hundred and eighty (180) apparently healthy and non diabetic were used as control and one hundred and eighty (180) diabetes used to test subjects. A total one hundred and eighty (180) males and one hundred and eighty (180) females were involved in the study. These subjects of both the sexes were categorized under three age groups such as 18 to 26 years, 27 to 36 years and 37 to 45 years. In females samples are collected in between 1-4 days of menstrual cycle. The research protocol was approved by ethics committee of the Institution. Sample collected aseptically from cases and control was subjected to serum analysis. Proper vein puncture technique was employed in the collection of the 5ml blood sample from the subjects. Blood samples were centrifuged at 3000rpm for 10 minutes. Fasting blood glucose and post prandial blood glucose were measured by GOD-POD method(14) and HbA1c% was measured by Ion exchange resin method by coral clinical system kit. The superoxide dismutase (SOD) was performed by Kakkar's et. al method (15) and Glutathione peroxidase was performed by DTNB method(16).

Data generated were analyzed using statistical package for social science (SPSS) version 20.00 and Microsoft excel 2007.

Comparison mean and standard deviation values were made for the various parameters for test and control subjects using student-t test. Results were considered statistically significant 95% confidence interval ($p < 0.05$).

III. RESULTS

Age group (years)	18 to 26		27 to 36		37 to 45	
Male	Case (Mean and SD)	Control (Mean and SD)	Case (Mean and SD)	Control (Mean and SD)	Case (Mean and SD)	Control (Mean and SD)
SOD(milliuunits/g of hb)	3.51±0.65	5.15±0.87	3.44±0.57	4.18±0.42	2.99±0.75	4.10±0.81
Glutathione(miuM/g of hb)	11.65±1.39	16.68±3.22	8.35±0.39	15.86±2.06	7.58±0.59	15.59±3.80
FBS (mg/dl)	175±5.45	78±5.63	201±7.40	88±5.34	167±8.58	98±7.01
PPBS(mg/dl)	254±16.99	102±13.51	236±8.23	110±8.44	210±9.35	115±6.09
HbA1C%	9.10±0.71	4.15±0.57	7.4±0.66	4.15±0.65	7.53±1.10	4.25±0.66
P value	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05

Table:-1 Age based comparison between case & controls for serum SOD, RBC Glutathione, FBS, PPBS and HbA1C% parameters among males.

Age group (years)	18 to 26		27 to 36		37 to 45	
Female	Case (Mean and SD)	Control (Mean and SD)	Case (Mean and SD)	Control (Mean and SD)	Case (Mean and SD)	Control (Mean and SD)
SOD(milliuunits/g of hb)	2.50±0.57	5.32±1.17	3.21±0.88	4.58±0.50	2.83±0.34	4.09±0.45
Glutathione(miuM/g of hb)	10.21±0.98	15.94±2.61	7.24±1.40	14.67±1.37	8.25±0.27	15.4±0.65
FBS (mg/dl)	179.96±16.70	82.33±8.35	159.33±10.01	92.89±3.92	158.43±9.29	105.10±3.18
PPBS(mg/dl)	232.76±36.77	105.16±9.30	222.80±25.60	111.58±5.68	210.96±13.57	118±5.73
HbA1C%	9.64±1.00	4.67±0.67	8.13±0.77	4.54±0.49	7.53±0.26	5.17±0.32
P value	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05

Table:-2 Table:-Age based comparison between case & controls for serum SOD, RBC Glutathione, FBS, PPBS and HbA1C% parameters among females

IV. DISCUSSION

The result of this study showed that the mean and Sd values of FBS, PPBS, HbA1c%, SOD and RBC Glutathione were calculated for all Type 2 Diabetic cases and controls. The mean and Sd values of FBS, PPBS, HbA1c%, were significantly higher among cases ($p \leq 0.05$) for both sexes compared to controls as shown in Table 1, 2

The mean and Sd values of SOD and RBC Glutathione for type -2 diabetics were significantly lower ($p \leq 0.05$) when compared with that of the control subjects as shown in Table 1 and 2. These findings show that type-2 diabetes is associated with decrease in antioxidant enzyme resulting in increase oxidative stress. These findings do agree with Djordjević et. al. findings which discovered a reduced antioxidative defense in patients with type-2 diabetes mellitus. Similarly in Table -1 and Table -2, the level of antioxidant SOD and glutathione were found to be significantly lower in diabetic patients than that of control subjects. These findings also show that type-2 diabetes is associated with decrease in antioxidant enzyme resulting in increase oxidative stress. These findings also do agree with Djordjević et. al. findings which discovered a reduced antioxidative defense in patients with type-2 diabetes mellitus (17).

Our finding also show that the level of antioxidant enzymes SOD and RBC Glutathione decreases with ages of diabetic subjects. This finding is similar with that of Chandankhede et.al and Bohr et. al.(18,19) With increase in mitochondrial ROS production endogenous antioxidant activity get diminished in aging individuals. Excessive production of ROS and reduced antioxidant defence with age significantly contribute to aging. This findings agrees with the literature of Poljsak and Milisav, 2013) (20)

V. CONCLUSION

From this research paper it has been found that type-2 diabetes is associated with decreased antioxidative status as the levels of antioxidant enzymes like SOD and RBC Glutathione were found to be significantly reduced in diabetic patients. This is because hyperglycemia is the cause of type-2 diabetes which increases the generation of ROS and that might depress the antioxidant defense system exposing the cells to damage from oxidative stress which leads to the development of diabetic complication. Moreover with the increase in ROS production of the antioxidant enzymes can suppressed the antioxidant system as well as its adaptive responses.

It can be concluded that T2DM is related to decreased antioxidative status as result in decrease in level of antioxidant enzymes SOD and RBC Glutathione where reduce significantly in diabetic subjects. The antioxidative system can be depressed if the subject is suffered from the illness for a prolonged period of time. Progression of disease is inversely proportional to antioxidative parameters, that is, antioxidative parameters are depleted with the progress in the condition of the disease.

It can be suggested that the antioxidant therapy is to be incorporated in the management of type-2 diabetes to supplement the antioxidative system as this prevent the progression of disease and other complication in diabetes. These imbalances lead to the tissue damage and inflammation in disease thus it can be said the antioxidant therapy along with conventional drugs should be used to prevent the disease progression.

REFERENCES

- [1] Brown Holy, Briggs Ojoye Ngoye. Clinical relevance of superoxide dismutase and glutathione peroxidase levels in management of diabetes type2. International Journal of Contemporary Medical Research 2016;3(5):1380-1382.
- [2] Asmat U, Diabetes mellitus and oxidative stress-A concise review, Saudi Pharmaceutical Journal, 2015
- [3] Lipinski B, Pathophysiology of oxidative stress in diabetes mellitus, J Diabetes its complications 2001.
- [4] Kumari M.Kusuma, Sankaranarayana. T, Evaluation of Oxidative Stress in Type 2 Diabetes Mellitus Patients, IOSR Journal of Dental and Medical Sciences (IOSR-JDMS) e-ISSN: 2279-0853, p-ISSN: 2279-0861, Volume 13, Issue 5 Ver. III. (May. 2014), PP 46-50 www.iosrjournals.org
- [5] Ojoye Ngoye Briggs, Holy Brown, Kemzi Elechi-amadi, Ferdinand Ezeiruaku, Nsirim Nduka, Superoxide Dismutase and Glutathione Peroxidase Levels in Patients with Long Standing Type 2 Diabetes in Port Harcourt, Rivers State, Nigeria, International Journal of Science and Research (IJSR). Volume 5 Issue 3, March 2016
- [6] Kunwar Amit, Priyadarsini KI, Free radicals, Oxidative Stress And Importance Of Antioxidants In Human Health, J Med Allied Sci 2011;1(2):53-60.
- [7] Giacco, F., & Brownlee, M. Oxidative stress and diabetic complications. Circulation Research. 2010;107: 1058-1070.
- [8] Montonen, J., Knekt, P., Jarvinen, R., & Reunanen, A. (2004). Dietary antioxidant intake and risk of type 2 diabetes. Diabetes Care, 27, 362-366.
- [9] Rains, J. L., & Jain, S. K. (2011). Oxidative stress, insulin signaling, and diabetes. Free Radical Biology and Medicine, 50(5), 567-575.
- [10] Mahadev, K., Motoshima, H., Wu, X., Ruddy, J. M., Arnold, R. S., & Cheng, G. (2004). The NAD(P)H oxidase homolog Nox4 modulates insulin-stimulated generation of H₂O₂ and plays an integral role in insulin signal transduction. Molecular and Cell Biology, 24(5), 1844-1854.
- [11] Bashan, N., Kovsan, J., Kachko, I., Ovadia, H., & Rudich, A. (2009). Positive and negative regulation of insulin signaling by reactive oxygen and nitrogen species. Physiological Reviews, 89, 27-71.
- [12] Brownlee, M. (2001). Biochemistry and molecular cell biology of diabetic complications. Nature, 414(6865), 813-820.
- [13] Memisogullari R, Taysi S, Bakan E, Capogly I., Antioxidant status and lipid peroxidation in type II diabetes mellitus, Cell Biochem 2003; 21(3): 291- 296
- [14] Kaplan L.A. Clinical Chemistry. The C.V. Mosby Co. St. Louis. Baltimore. Philadelphia. Toronto. 854-856, 1989.

- [15] Kakkar Poonam, Das Ballabh, Viswanathan P N, A modified Spectrophometric assay of Superoxide Dismutase, Indian journal of Biochemistry and Biophysics, Vol 21, April 1984, pp-130-132.
- [16] Beuder E, The Glutathione instability of drug sensitive red cells: a new method for the in vitro detection of drug sensitivity, J.Lab. Clin. Med. 49:84-94. 1957..
- [17] Djordjević, G. M., Djurić, S. S., Djordjević, V. B., Apostolski, S., & Živković, M. (2011), The role of oxidative stress in pathogenesis of diabetic neuropathy: erythrocyte superoxide dismutase, catalase and glutathione peroxidase level in relation to peripheral nerve conduction in diabetic neuropathy patients, In: C. Croniger (Ed.) , Role of the Adipocyte in Development of Type 2 Diabetes. (pp 153-172), Rijeka, Croatia: InTech.
- [18] Chandankhede Manju S. & Gupta Madhur M Oxidative stress and antioxidant status in patients with rheumatoid arthritis Int J Biol Med Res. 2013; 4(2): 3088-3090
- [19] Bohr, V., Anson, S., Mazur, R. M., & Dianov, G. (1998). Oxidative DNA damage processing and changes with aging. Toxicology Letters, 102-103, 47-52.
- [20] Poljsak, B., & Milisav, I. (2013). Aging, oxidative Stress and antioxidants. In: J.A. Morales-Gonzalez (Ed.), Oxidative stress and chronic degenerative diseases: A role for antioxidants (pp. 345). Rijeka, Croatia: InTech

10.22214/IJRASET

45.98

IMPACT FACTOR:
7.129

IMPACT FACTOR:
7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)