iJRASET

International Journal For Research in
Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: VIII Month of publication: August 2019

DOI: <http://doi.org/10.22214/ijraset.2019.8145>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Effect of Swimming Exercise on Lipid Profile in Type 2 Diabetes Mellitus

Raut B.U.¹, Dr. Kolkhede P.V.²

¹Reserch Scholar Department Of Zoology, J.D. Patil Sangludkar Mahavidyalay Daryapur Dist Amravati M.S. India

²Assosiate professor & Head of Zoology, Department. J.D. Patil Sangludkar Mahavidyalay Daryapur Dist Amravati M.S India

Abstract: *The aim of this study was to observe the effect of swimming exercise on lipid profile type 2 diabetic patients we all known swimming exercise may well be the perfect exercise for diabetes type 1 or type 2 diabetes for everyone with diabetes swimming can keep active and healthy swimming exercise mobilize and improve metabolic body function. The diabetes disease development of hypertension in diabetic patients so keeping of this point of view twenty(N=20) diabetic patients aged 45 to 55 years were selected for experimentation purpose and the variable such as systolic diastolic blood pressure body weight and lipid profile including TG,TC, HDLC, LDLC, VLDLC, They were randomly divided into two groups such as controle group experimentation groups. The blood sample collected before swimming exercise serves as control groups while sample collected after swimming exercise serve as experimental group of diabetic patients. Patients were suggested for swimming exercise two hour per daily for a period of 12 weeks. The systolic, diastolic blood pressure, body weight and lipid profile including TG,TC,LDLC,VLDLC,(except HDLC) were significantly low in swimming exercise groups ($P<0.005$) compared to the control groups. The present studies show that lipid profile value can be effectively control in diabetic patients type 2 through the swimming exercise daily in the morning .*

Keywords: *Diabetise mellitus, lipid profile, diabetic patients, swimming exercise.*

I. INTRODUCTION

Diabetes is a chronic metabolic syndrome caused by body's inability to produce enough insulin or to use insulin that is available as a result there is an increase in the concentration of glucose metabolic abnormalities (1). Diabetes has become a wide spread epidemic primarily because lack of exercise and awareness and stress diet exercise has been considered a cornerstone of diabetes management (5). Swimming keeps you moving for a length of time rather than having you expend energy then rest. This important for cardiovascular fitness in that it allow one-two pump blood and oxygen more efficiently Exercise burns 350 to 420 calories per hour (2). Physical exercise has been conceded as one of the cornerstone in the treatment of diabetes mellitus along with nutrition and medication since from the past 100 year ago (14). Swimming exercise to improve blood glucose control in type 2 diabetes, reduces cardiovascular risk factor contribute to weight loss and improve well being (riddle) (7). It has long known that exercise have substantial benefits for people with both the type of diabetes (T1DM) and type (T2DM) It can increase insulin sensitivity, improve cardiovascular fitness and help sustain weight loss (10). Guidelines recommended that patients with T2DM should perform atleast 150 min resistance exercise minimum 3 times per week (8). Scientific pedestal and popularize at among the general public we planned to undertake study of effect of swimming exercise on lipid profile (VLDLC) cholesterol (TC) VLDLC, HDLC and triglycerides (TG) in type 2 diabetic patients.

II. MATERIAL AND METHOD

A 3 month exercise two groups of twenty (N=20) diabetic patients (age 45 to 65 years) have been selected from single center in diabetic camp in Amravati district we conducted a study involving the swimming at selected adult patients of T2DM. The systolic, diastolic blood pressure and body weight were measured at visit. The sample were collected from people involved in the experiment before and after 3 month. The physiological parameter like systolic and diastolic blood pressure measured by using standard Sphygmomanometer (ISI mark) and body weight by weighing machine. Experiment for testing lipid profile parameters such as TC, LDLC, VLDLC, HDLC and TG by using kit method (Foosati 1982 and Richmond 1973 (3-6). and obtained data compared with control readings. The mean of the difference of values and standard deviation obtained of these two groups were calculated and statistically analyzed by utilizing the student "s" "t" test.

III. RESULT AND DISCUSSION

Table No.1- Mean values of lipid profile systolic, diastolic blood pressure and body weight(wt) in subjects participated before and after swimming exercise.

	Biochemical parameter	Before	After
1	Systolic blood pressure	158.5 $\pm$ 9.39	127.36 $\pm$ 10.43***
2	Diastolic blood pressure	108.80 $\pm$ 10.11	86.00 $\pm$ 79.32***
3	Body weight	55.84 $\pm$ 10.43	98.26 $\pm$ 09.32****
4	Total Cholesterol mg/dl(TC)	181.2 $\pm$ 39.92	168.0 $\pm$ 37.92*
5	Serum Triglycerides mg/dl (TG)	75.29 $\pm$ 28.26	81.02 $\pm$ 33.89
6	Very low density lipoproteins mg/dl (VLDL)	15.32 $\pm$ 05.42	17.52 $\pm$ 10.02
7	Low density lipoproteins mg/dl(LDL)	108.70 $\pm$ 38.22	95.61 $\pm$ 28.42***
8	High density lipoproteins(HDL)	50.00 $\pm$ 9.92	55.52 $\pm$ 8.62

***P<0.001, P<0.005, Non significant as compare to swimming exercise

Table No.1 – Show that systolic, diastolic blood pressure, body weight and lipid profile's level in control groups did not show any significant variation (P<0.005) Whereas the experimental groups which underwent swimming exercise (P<0.001) show reduction in systolic and diastolic blood pressure (Table1) which is quite significant after swimming exercise. The data was analyzed by using student paired 't' test and P values of less than 0.005 were accepted as significant difference between the compared values. The TC (P,0.005) and VLDL (P,0.001) was significantly decreased.

Graph- Show the percentage (%) of lipid profile in subjects participated before and after swimming exercise.

Figure no 1- Percentage change (%) of mean values of lipid profile blood pressure and body weight in subjects participated before and after swimming exercise.

The review shows that swimming exercise had a beneficial effect on body weight, blood pressure, blood glucose level, and cholesterol (13). Aerobic exercise improves the physiological parameters including glycemic control, fasting blood glucose level, and lipid profile (11). Regarding management of T2DM, researches have highlighted the use of modern medicine, alternative or herbal medicine, and exercise management therapy. The entire management proves to have a positive impact on the disease. Therefore, physical activity or exercise is considered as the beneficial treatment regimen for the treatment of T2DM. The aim of the present systematic review was to focus on the positive effect of exercise towards T2DM (12). However, most of the randomized trial studies showed that high volume aerobic exercise produced weight loss with significant improvement in insulin sensitivity (9). The practice of exercise was associated with a significant decrease in cholesterol among subjects with cardiovascular disease, hypertension, angina, arteriosclerosis, and type 2 diabetes at different durations of practice of exercise (4). The present investigation confirms that swimming practices control the lipid profile in T2DM.

REFERENCES

- [1] Anna chuddy k; Robert J; Petrella E;(2011):Effect of exercise on cardiovascular risk factor in type2 diabetes Vol 34 no.spp-1228-1237
- [2] Daniel Bubnis ,M.S(SEPT-2017):Ashley Marcine (Na.J.Mt-Pt).
- [3] Foosati P (1982) serum triglycerides determine colorimetricallay with an enzyme that produces hydrogen peroxide clin chem. 28:2077-2080
- [4] Gorden L.A.M,Morinson E.Y,Mcgrowder D.A,Young R.,Fraser Y.T . Zamora E.M.etal.(2008): Effect of exercise therapy on lipid profile and oxidative stress indicators in patients with Type 2 diabetes.B.M.CJ.Alt comple.Med 8:21
- [5] Quinn Philips (2011):Exercise and diet diabetes self management.
- [6] Richmond W.(1973):Preparation and properties of a cholesterol oxidase from nocardia sp.and its application to the enzymatic assay of total cholesterol in serum clin chem. 19: 1350-1356
- [7] Riddled M. Perkins B.A.(2009): Exercise and glucose metabolism in persons with diabetes mellitus perspectives on the role for continous glucose monitoring J. diabetes sci.technol 1914-1923.
- [8] Ronald J. sigal, Glen F, Kenny G.P, David H.,Wasserman,Carman Cost aned as sceppas,Russel D.white(2006):-Physical exercise and type2 diabetes Vol.29,No.614,pp33-143.
- [9] Ross R.,Dagnone D.,Jones P.J ,Smith H, Paddagase A.,et .al(2000): Reduction in obesity and related comorbid condition after diet induced weight loss in men randomized controlled trial Ann.Inter.med 133-92-103
- [10] Siga R.J,Kenny G.P.Wasserman,D.H Castaned(2004): Physical activity exercise and type2 diabetes: Diabetes care 27(10)PP 2518-2539
- [11] Steward KJ (2002): Exercise training the cardiovascular consequences of type 2 diabetes and hypertension plausible mechanism for improving cardiovascular health TAMA 288: 1622-1631.
- [12] Tuomilehto J, Lindstrom J,Errikson J.U, Valle J.J, Hamalainen H.et. al(2001):Prevention of type2 diabetes mellitus by changes in life style among subject with impaired glucose tolerance N.Eng J. Med 344:1343-1350.
- [13] Yang K.A (2007): Review of yoga/physical exercise programmes for leading risk factor of chronic diseases.Evidences based compte. Alt medicine 4(4) 487-491.
- [14] Zarchi Thent ; Srist Daj; Leonard Joseph; Henry (2014):Role of exercise in the management of diabetes mellitus.

10.22214/IJRASET

45.98

IMPACT FACTOR:
7.129

IMPACT FACTOR:
7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)