iJRASET

International Journal For Research in
Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 9 Issue: II Month of publication: February 2021

DOI: <https://doi.org/10.22214/ijraset.2021.33072>

www.ijraset.com

Call: 08813907089

E-mail ID: ijraset@gmail.com

Antimicrobial and Antioxidant Activities of some Newly Synthesized Pyrazolo Thiazolo Pyrimidines

A. V. Pawde

Department of Chemistry, A.E.S's Arts, commerce and Science College, Hingoli-431513, MS, India

Abstract: synthesis of 3-amino -4-imino-7-methyl -2-N-(substituted) pyrazolo [3,4-b] thiazolo [1,2-a] pyrimidine (5A.17a-i) have been reported by condensing 6-cyano-5-imino-2-methyl-7-(methylthio)-5H-thiazolo [3,2-a] pyrimidine with hydrazine and its various derivatives. The chemical structure of the product was proved on the basis of their spectral IR, ¹H-NMR, ¹³C-NMR, Mass and analytical studies. Some of the newly synthesized compounds have been screened for antimicrobial and antioxidant activity.

Keywords: Pyrazolothiazolo pyrimidine, Bis (methylthio) methylene malononitrile, N,N-dimethyl formamideanhydrous potassium carbonate, Hydrazine.

I. INTRODUCTION

Extensive investigation has been done in the area of preparation of the pyrazolothiazolo pyrimidines. These compounds have wide range of biological activities¹⁻⁸. Some of these series of compounds show effective calcium channel blockers⁹ and few compounds show a strong antiproliferative activity¹⁰. With an intention of preparing the compounds possessing better therapeutic potential, we have undertaken the synthesis of pyrazolothiazolo pyrimidine derivatives. In this paper we reported synthesis of 6-cyano-5-imino-2-methyl-7-(methylthio)-5H-thiazolo [3,2-a] pyrimidine and their hydrazine derivatives which shows promising biological activities.

II. MATERIALS AND METHODS

Melting point were determined by open capillary tubes and were uncorrected. Progress of reaction was monitored by thin layer chromatography carried out on aluminium silica plates using UV Chamber for detection. Infrared spectra were recorded in potassium bromide pellets on an infrared spectrophotometer, nuclear magnetic spectra were obtained on Bruker advance spectrophotometer; 400MHz mass spectra were recorded on ET-VC7070H mass spectrophotometer with the use of EI technique at 70eV. All the reactions were carried out under ambient atmosphere.

III. RESULT AND DISCUSSION

In the present investigation, we have reported synthesis of 3-amino -4-imino-7-methyl -2-N-(substituted) pyrazolo [3,4-b] thiazolo [1,2-a] pyrimidine (5a-i). The reaction started by preparation of 6-cyano-5-imino-2-methyl-7-(methylthio)-5H-thiazolo[3,2-a] pyrimidine (3) by condensing 2-amino-5-methyl thiazole (1) with bis(methylthio) methylene malononitrile (2) in presence of DMF and anhydrous K₂CO₃ as catalyst. (Scheme -1).

Scheme 1

The compound (3) has replaceable active methylthio group at 2-position which is activated by ring 1-nitrogen atom and electron withdrawing cyano group at 3-position. Hence, the susceptibility of the compound (3) towards cyclization with hydrazine hydrate and their different substituted derivatives (4a-i) have been investigated. These reactions result into formation of 3-amino -4-imino-7-methyl -2-N-(substituted) pyrazolo [3,4-b] thiazolo [1,2-a] pyrimidine (5a-i). According to these reactions, the compound (3) independently reacted with hydrazine hydrate (80%) (4a), phenyl hydrazine (4b), 4-nitro phenyl hydrazine (4c), 2,4-dinitro phenyl hydrazine (4d), 2-hydrazino benzothiazole (4e), 6-methyl-2-hydrazino benzothiazole (4f), 6-methoxy-2-hydrazino benzothiazole (4g), 6-chloro-2-hydrazino benzothiazole (4h), and 6-nitro-2-hydrazino benzothiazole (4i) in DMF and anhydrous K₂CO₃ to obtain, 3-amino -4-imino-7-methyl -2-N-(substituted) pyrazolo [3,4-b] thiazolo [1,2-a] pyrimidine (5a-i).

(Scheme 2).

Compound No.	R	Compound No.	R
4a	H	4f
4b		4g
4c		4h
4d		4i
4e

IV. ANTIOXIDANT ACTIVITY OF NEWLY SYNTHESIZED COMPOUNDS

A. DPPH Assay

DPPH (2, 2, diphenyl-1-picrylhydrazyl) radical scavenging assay was carried out as per reported methods with slight modification (Kato et al., 1998). Briefly, 1ml of test solution (Test compound) added to equal quantity of 0.1mmol solution of DPPH in ethanol. After 20 min incubation at room temperature, the DPPH reductions were measured by reading the absorbance at 517 nm. Ascorbic acid used as reference compound.

Table No. 1: Antioxidant potential of pyrazolo [3,4-*b*] thiazolo [1, 2-*a*] pyrimidines

Sr. No	Compound	DPPH radical scavenging activity (%)
1	5a	63±0.44
2	5b	54 ± 0.98
3	5d	75 ± 0.41
4	5e	68 ± 0.57
5	5g	44±0.54
6	Ascorbic Acid	86 ± 0.88

V. ANTIMICROBIAL ACTIVITY

A. Disc Diffusion Method

Kirby-Bauer Method was followed for disc diffusion assay. In vitro antimicrobial activity was screened by using Mueller Hinton Agar (MHA) obtained from Himedia (Mumbai). The MHA plates were prepared by pouring 15 mL of molten media into sterile petriplates. The plates were allowed to solidify for 5 min and 0.1 % inoculum suspension was swabbed uniformly and the inoculum was allowed to dry for 5 min. The concentration of compounds were set at (10 µg disc-1) were loaded on 5 mm sterile individual discs. The loaded discs were placed on the surface of medium and the compound was allowed to diffuse for 5 min and the plates were kept for incubation at 37°C for 24 h. Penicillin (10 µg disc-1) was used as positive control. At the end of incubation, inhibition zones formed around the disc were measured with transparent ruler in millimeter.

Table No. 2: Antimicrobial potential of pyrazolo [3,4-*b*] thiazolo [1, 2-*a*]

Sr.no.	Name of compound	Zone of inhibition in mm	
		<i>E. coli</i>	<i>B. subtilis</i>
1	5a	NR	NR
2	5b	NR	NR
3	5d	14	07
4	5e	NR	18
5	5g	12	08
6	penicillin	26	28

NR: No Respose

VI. CONCLUSION

Simple and efficient synthesis of 3-amino -4-imino-7-methyl -2-N-(substituted) pyrazolo [3,4-*d*] thiazolo [3,2-*a*] pyrimidine (5a-i) has been reported . Among these synthesized compounds (5d) showed remarkable antioxidant and antimicrobial activity and compound (5e) exhibit promising antimicrobial activity against *B. subtilis*. The result of the present work demonstrated that pyrazolothiazolo pyrimidines are potent antioxidant and antimicrobial agents and it will attract researchers to design new potent pharmacological pyrazolothiazolo pyrimidines.

REFERENCES

- [1] Chandrahas N. Khobragade, Ragini G., Journal of Enzyme Inhibition and Medicinal Chemistry., 2010, Vol.25, pp. 615-621.
- [2] A.V. Bentya., Russian Journal of Organic Chemistry., 2007, Vol.44, pp. 1362-13.
- [3] B.A.Bilal., Cheminform., 2010, Vol. 24, pp. 345-351.
- [4] Pier Giovanni Baraldi., Drug Development research., 2001, pp. 406-415.
- [5] Abheshek kumar Singh,Cheminform, 2005, Vol. 36, pp. 367-371.
- [6] Johansson., Current topics in Medical Chemistry., 2003, Vol. 3.
- [7] Hayam H. Sayed, Acta Pharmaceutica, 2006,Vol. 56.
- [8] A. Mobinikhaledi, Cheminform, 2008, Vol. 36.
- [9] Guccione, S., Modica, M., Longmore, J., Show, D., Barretta,G.U., Santagati, A., Santagati, M., and Russo, F., Bioorg. Med. Chem. Lett., 1996, Vol. 6, pp. 59-64.
- [10] Russo, F., Guccione, S., Romeo, G., Baretta, G.U.,Pueci, S., Caruso, A., Amico-Roxas, V., and Cutuli, V., Eur. J. Med. Chem., 1993, vol. 28, pp. 363

10.22214/IJRASET

45.98

IMPACT FACTOR:
7.129

IMPACT FACTOR:
7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)