KABACHNIK-FILDS REACTIONS WITH 2,3-TRIMETHYLENE-1,2,3,4-TETRAHYDROPYRIDO[2,3-D]PYRIMIDIN-4-ONE

KH. I. AKBAROV¹, Z.U. SAMAROV², D. E. TURGUNOV³, D. SH. KIYAMOVA³

¹National University of Uzbekistan named after Mirzo Ulugbek, Uzbekistan

²Samarkand State University, Uzbekistan

³Samarkand State Medical Institute, Uzbekistan

ABSTRACT

In the article synthesized 2,3-trimethylene-3,4-dihydropyrido[2,3-d]pyrimidin-4-one from 2-Aminonicotinic acid and pyrrolidone-2 in the presence different agents, such as PCl₅, POCl₃. Its reduction reaction with NaBH₄ carrying out. Obtained 2,3-trimethylene-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-4-one – this three-component coupling of a carbonyl, an amine and a hydrophosphoryl compound leads to α-aminophosphonates, phosphorous acid-formaldehyde; aldehydes in three component system; aminomethylphosphonic acid synthesis based on kabachnik-filds reaction. 2,3-trimethylene-3,4-dihydropyrido[2,3-d]pyrimidin-4-one 35% da product obtained. The reason of low reaction yield was studied. Structures were confirmed using IR and 1H NMR spectroscopies. Obtained product reduced using NaBH₄ and 2,3-trimethylene-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-4-one synthesis was carried out with high yield. 2,3-trimethylene-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-4-one and respective aminophosphonic acids were synthesized in three component system with high yield.

KEYWORDS: 2,3-trimethylene-3,4-dihydropyrido[2,3-d]pyrimidin-4-one, 2,3-trimethylene-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-4-one, 2-Aminonicotinic acid, pyrrolidone-2, Kabachnik-Filds reactions, IR spectra

Received: Jun 08, 2020; **Accepted:** Jun 28, 2020; **Published:** Sep 24, 2020; **Paper Id.:** IJMPERDJUN20201401

INTRODUCTION

In the world synthesis of new physiologically active derivatives of pyrido[2,3-d]pyrimidin and creation on their basis of modern medicinal facilities come with the use of high-tech. It is known that the anti-cancer drugs applied to date, destroying malignant cancer cells, simultaneously damage healthy cells.

Representatives of pyrido[2,3-d]pyrimidins is preferentially operating anti-cancer drugs of palbocyclik, preparations of antibacterial action of pipemidic and piromid acids are worked out by the world scientists. These medical facilities promote practical interest in derivatives.

Organophosphorus compounds are ubiquitous in nature and find applications in the fields of agriculture, medicine, and industry [1-3]. Some organophosphorus compounds are important pesticides [4], bactericides [5-7], and antibiotics [5]. Phosphorus analogues of α-pyrones act as HIV protease inhibitors [8]. α-aminophosphonic acids constitute important motifs among the organophosphorus compounds in medicinal chemistry due to their obvious structural similarities to α-amino acids [9,10]. Many natural and synthetic aminophosphonic acids and their ester and peptide derivatives display a wide range of biological activities [11,12], act as herbicides [13], enzyme inhibitors [14], and antibacterial [15,16] antiviral [10], and antitumor [17] agents, and may even be

peptide mimics [18].

The most common synthetic route to α -aminophosphonic acids is via chemical manipulation of the corresponding α -aminophosphonates [19-21]. The hydrophosphonylation of imines is a widely used method for the synthesis of α -aminophosphonates [19-27]. This is achieved by one of two pathways: (I) in a two-component fashion known as the Pudovik reaction [28,29] or (II) by the Kabachnik-Filds reaction [22, 23, 30, 31] which combines in situ formation of imine by condensation of amines with an aldehyde or ketone and an hydrophosphonylation step [32].

One-pot Kabachnik-Filds reaction can be promoted by acidic or basic catalysts, microwave irradiation, or by heating [33]. Due to the above-mentioned factors, in this paper we reported the synthesis of α -aminophosphonates with high yield using a recyclable catalyst for applications in medicine and industry.

MATERIALS AND METHODS

Threecyclic 2,3-thremetilenpyrido[2,3-d]pyrimidin-4-one (1) was synthesized coming from 2-aminonicotinic (2-amino-3-pyridin carboxylic) acid that was exposed to condensation with pyrrolidone-2 in the presence of POCl_3 .

Scheme 1: Synthesis of 2,3-trimethylene-3,4-dihydro-pyrido[2,3-d]pyrimidin-4-one

Yield of products of condensation with lactam row 2-aminotiophen-3-carboxylic, 2-aminobenzoyic(antranilic) and 2-aminopyridin-3-carboxylic acids go down. So, if 2,3-threemetylentieno[2,3-d]-pyridin-4-one is got with 81% yield, and 2,3-polymetylen-hinazalon - with 56-70% yield, in the cases of 2,3-threemetylenpyrido-[2,3-d]pyrimidin-4-one (1) he made a 35%. This fact explained by that π -electron a surplus tiophenic ring facilitates, and π -deficit pyridinic ring hampers flowing of this condensation.

In infrared spectrum 1 there are absorptions bands characteristic for amido carbonyl groups ($\text{N}-\text{C}=\text{O}$) in area of 1683 cm^{-1} , and absorption bands of $\text{C}2=\text{N}1$ double bond show up in area of 1625 cm^{-1} .

In ^1H NMR spectrum 1 (CDCl_3) aromatic protons are observed in the weak fields at 7.31-7.34 ppm in a kind duplicate of duplicates (1H, dd, $J=4.6, 7.9$, H-6), 8.48-8.53 (1H, ppm $J=2.0, 7.9$, H-5) and 8.86-8.89 (1H, ppm, $J=2.0, 4.6$, H-7).

Selective reduction of $\text{N}1=\text{C}2$ bonds. Reduction of 2,3-three-metylenopyrido[2,3-d] by the pyrimidin-4-one (1) borhydrid of natrium in a spirit solution at boiling results in formation of 2,3-threemytilen-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-4-on (2). In this case, selective reduction goes $\text{N}1=\text{C}2$ to double bonds of pyrimidinic ring, and carbonyl group $\text{C}=\text{O}$ in position 4 remains.

Scheme 2. Reactions of 1 with NaBH₄

ESI-MS the spectrum of product of reaction gives a strong band of molecular [M+H] ion at *m/z* 190, that specifies on molecular mass of this fragment 189, and along with ¹H NMR results spectral data leads to the molecular formula of C₁₀H₁₁N₃O.

After reduction a 1 absorption band at 1625 cm⁻¹ (C=N) disappears in IR spectrum (2). The absorption band of carbonyl group at C₄, that reveals at 1718 cm⁻¹ (CO) in initial connection, is displaced toward low-frequency areas (1654 cm⁻¹). Saturated N1 appears as a result of selective restoration-C₂ bond (product of reduction 2).

In ¹H NMR spectrum 2 signals of protons α- and β- CH₂ of groups appear in area of 1.85-2.40 (4H) as multiplet, unlike initial bonds [3.20 (2H, trip, J=7.9, α-CH₂), 2.26 (2H, quar., β-CH₂)

Signals of protons-CH₂ groups showing up in areas 3.59 (¹H, td, J=8.6; 2.9, γ a-CH₂) and 3.70 (¹H, m, γ e-CH₂) also differ from the signals of analogical protons in initial connection [4.16 (2H, t, J=7.3, γ-CH₂)].

Here becomes obvious circumstance that recovery of N1=C2 is connections to HN-CH results in noticeable displacement of chemical changes of protons of methylene groups toward the strong fields. Found out also substantial displacement of signals of protons of pyridinic ring after reduction. So, in initial step they appear at ppm: 7.34 (¹H, dd, J=4.6, 7.9, C6-H), 8.53 (¹H, dd, J=2.0, 7.9, C5-H), 8.89 (¹H, dd, J=2.0, 4.6, C7-H), accordingly; and in the product of renewal the same protons are observed in areas: 6.74 (¹H, dd, J = 5.0; 7.6, C6-H), 8.06 (¹H, dd, J=1.8; 7.4, C5 - H) and 8.10 (¹H, dd, J=1.8; 5.0, C7-H), accordingly; i.e. and here is displacement of chemical changes of protons toward the strong fields. Proton of asymmetric carbon of C2-H found out at 5.10.

In the initial experiments, the one-pot, three-component reaction of 2,3-trimethylene-1,2,3,4-dihydropyrido[2,3-d]pyrimidin-4-one, *p* and *m*-nitro-benzaldehyde, and phosphorous acid was chosen as the model reaction to optimize the reaction conditions. In the present work, the procedures followed for the synthesis of α-aminophosphonates are conventional shown in Table-1, entries 3-5. The products α-aminophosphonates were obtained by solvent-free microwave irradiation of aldehyde, amine, and phosphorous acid for 1 min. In toluene, without any catalyst, the product formed was in a good yield.

Thin layer chromatography (TLC) was employed to monitor reaction progress and to determine the purity of the products. The reaction was carried out using catalytic amount of HCl, in toluene for 30 min. All the title compounds are readily soluble in polar organic solvents.

The IR spectra of compounds (3-5) showed the N-CH₂ band in the range of 1550 cm⁻¹. The sharp band observed in the range 1240–1291 cm⁻¹ is due to the ν_{P=O}, and a band for P–C stretching occurred in the range 740–770 cm⁻¹. All the stretching frequencies are compiled in Table 2. ¹H NMR spectra of the compounds (3-5) were recorded in the DMSO-d₆ solvent. The aromatic protons of α-aminophosphonic acids appeared as a multiplet in the region δ 6.15–8.69. The P–C–H group resonated as a multiplet in the range δ 3.77–4.86.

Scheme 3: Reaction with Paraformum

Scheme 4: Reaction with Aromatic Aldehyde

Table 1: Reaction time and percentage yield of 2 in different reaction condition

Entry	Catalyst used	Reaction condition	Reaction time	% yield
3	HCl	Toluene/reflux	5	80
4	HCl	Toluene/reflux	5	87
5	HCl	Toluene/reflux	5	88

In toluene, without any catalyst, the product formed was in a good yield, but the time taken was 4 to 5 h, which is considerably longer.

Table 2: Elemental analysis and IR data of compounds (1-5)

	Mol. formula	Elemental analysis found (Calc)			IR spectral data in cm ⁻¹				
		C	H	N	C=N	C=O	P=O	P-O	P-C
1	C ₁₀ H ₉ N ₃ O	64.16	4.85	22.45	1625	1718			
2	C ₁₀ H ₁₁ N ₃ O	63.48	5.86	22.21		1760			
3	C ₁₁ H ₁₄ N ₃ O ₄ P	46.65	4.98	14.84	3135	1690	1210	1050	751
4	C ₁₈ H ₂₀ N ₄ O ₆	51.56	4.81	13.36	3381	1650	1268	1060	745
5	C ₁₈ H ₂₀ N ₄ O ₆	51.56	4.81	13.36	3340	1680	1263	1040	763

EXPERIMENTAL DETAILS

Synthesis of hydrochloride of 2,3-trimethylene-3,4-dihydropyrido[2,3-d]pyrimidin-4-one (1).

POCl₃ (90 ml) was dropped to mixture of 2-aminonicotinic acid (65.3 g, 0.47 mol) and pyrrolidone-2 (47.94 g, 0.564 mol) by stirring at room temperature. Reaction mixture heated at 80-90°C for 4 hours, cooled to room temperature and added 100 ml water. Aqueous solution worked up by 5% NaOH solution to pH 7.5-8. Extracted by chloroform (3x100 ml) and organic phase dried with Na₂SO₄, filtrated and evaporated solvent. Yield of compound 1 is 30.76 g (35%), m.p. 139-140°C. R_f 0.48 (10:1 chloroform-methanol):

To solution of a 28,05 g (0,15 mol) 2,3 - trimethylene - 3,4 - dihydropyrido[2,3 - d]pyrimidin - 4 - one (1) in 350 mls of alcohol added a 30 g (0,78 mol) of sodium borohydride. Reaction mixture was boiled on water bath during 3H and left on a 1 twenty-four hours. A solvent was driven away, remaining product was dissolved in water and left for 1 hour. Falling out sediment was filtered washed 3 times water, dried and recrystallized from a hexane. Yield of compound 2 is 28.35 g (91%), m.p. 140-142°C. R_f 0.58 (10:1 chloroform-methanol).

Synthesis of compound 3. A mixture of paraphorm (0.005 mol), 2,3-trimethylene-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-4-one (0.005 mol) and phosphide acid in dry toluene was stirred for 10 min at room temperature. Then the temperature was raised to reflux for 5H. The reaction was monitored by TLC. After completion of the reaction, toluene was removed by distillation and the residue was purified using column chromatography (5:4, benzene : hexane). Yield of compound 3 is 1.124g (80%), m.p. 167-169°C. R_f 0.4 (5:4, benzene:hexane)

Synthesis of compound 4. A mixture of m-nitrobenzaldehyde (0.005 mol), 2,3-trimethylene-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-4-one (0.005 mol) and phosphide acid in dry toluene was stirred for 10 min at room temperature. Then the temperature was raised to reflux for 5H. the reaction was monitored by TLC. After completion of the reaction, toluene was removed by distillation and the residue was purified using column chromatography (5:4, benzene : hexane). Yield of compound 4 is 1.76 g (87%), m.p. 167-169°C. R_f 0.52 (5:4, benzene:hexane).

Synthesis of compound 5. A mixture of p-nitrobenzaldehyde (0.01 mol), 2,3-trimethylene-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-4-one (0.01 mol) and phosphide acid in dry toluene was stirred for 10 min at room temperature. Then the temperature was raised to reflux for 5H. the reaction was monitored by TLC. After completion of the reaction, toluene was removed by distillation and the residue was purified using column chromatography (5:4, benzene: hexane). Yield of compound 5 is 1.78g (88%), m.p. 167-169°C. R_f 0.4 (5:4, benzene:hexane)

CONCLUSIONS

The new method of preparation of 2,3-threemytilenpyrido[2,3-d]-pyrimidin-4-one condensation of 2-aminonicotinic acid with lactam.

Carries out selective renewal of N1=C2 double bonds of 2,3-threemytilenpyrido[2,3-d]-pyrimidin-4-one.

The synthesis of new α -aminophosphonic acid was achieved in high yields through a one-pot three-component reaction process, a Kabachnik-Filds reaction. It involves the reactions among 2,3-trimethylene-1,2,3,4-

tetrahydropyrido[2,3-d]pyrimidin-4-one, substituted aromatic aldehydes, and phosphorous acid in dry toluene at reflux temperature. Their structures were determined by elemental analysis IR, ¹H-NMR, and mass spectral data.

REFERENCES

1. E. Breuer, *The Chemistry of Organophosphorus Compounds*, vol. 4, John Wiley & Sons, New York, NY, USA, 1996.
2. K. Srinivasulu, M. Anilkumar, C. Nagaraju, and C.S. Reddy "Synthesis and bioactivity of some new 2 substituted 3,4-dihydro-1-(9H-carbazol-4-yloxy)methyl-3-[2-(2-methoxyphenoxy)ethyl]-1.3.2λ5-oxazaphosphole 2-oxides, sulfides and selenides," *Arkivoc*, vol. 2007, no. 14, pp. 100-109.
3. T.K. Prakasha, R.O. Day, and R.R. Holmes, "New class of bicyclic oxyphosphoranes with an oxaphosphorinane ring: Molecular structures and activation energies for ligand exchange," *Journal of the American Chemical Society*, vol. 116, no. 18, pp. 8095-8104, 1994.
4. C. Fest and K.J. Schmidt, *The Chemistry of Organophosphorus Pesticides*, vol. 12, Springer, 1982.
5. M.S. Bhatia and P. Pawanjit, "Phosphorus containing heterocycles as fungicides: synthesis of 2,2' diphenylene chlorophosphonate and 2,2' diphenylene chlorothiophosphonate", *Experientia*, vol. 32, no. 9, p. 1111, 1976.
6. P.N. Manne, S.D. Deshmukh, N.G.V. Rao, H.G. Dodale, and S.N. Tikar, "Efficacy of some insecticide against *Helicoverpa armigera*", *Pestology*, vol. 34, p 65, 2000.
7. D. Hendlin, E. O. Stapley, M. Jackson et al., "Phosphonomycin, a new antibiotic produced by strains of *Streptomyces*", *Science*, vol. 166, no. 3901, pp. 122-123, 1969.
8. A.M. Polozov and S.E. Cremer, "Synthesis of 2H-1,2-oxaphosphorin 2-oxides", *Journal of Organometallic Chemistry*, vol. 646, no. 1-2, pp. 153-160, 2002.
9. V.P. Kukhar and H.R. Hundson, *Aminophosphonic and Aminophosphinic Acids: Chemistry and Biological Activity*, John Wiley & Sons, New York, NY, USA, 2000.
10. J. Huang and R. Chen, "HYPERLINK an overview of recent advances on the synthesis and biological activity of α-aminophosphonic acid derivatives", *Heteroatom Chemistry*, vol. 11, pp.480-492, 2000.
11. L.D. Quin, *A Guide to Organophosphorus Chemistry*, vol. 11, Wiley, New York, NY, USA, 2000.
12. J. Hiratake and J. ODA, "Aminophosphonic and aminoboronic acids as key elements of a transition state analogue inhibitor of enzymes," *Bioscience, Biotechnology, and Biochemistry*, vol. 61, no. 2, pp. 211-218, 1997.
13. L. X. Xiao, K. Li, and D. Q. Shi, "A Convenient Synthesis and Herbicidal Activity of N-phosphonoalkylpyrazolo[4,3-e][1,2,4]-triazolo[1,5-d]pyrimidines," *Phosphorus, Sulfur, and Silicon and the Related Elements*, vol. 183, no. 12, pp. 3156-3165, 2008.
14. M. C. Allen, W. Fuhrer, B. Tuck, R. Wade, and J. M. Wood, "Renin inhibitors. Synthesis of transition-state analogue inhibitiros containing phosphorus acid derivatives at the scissile bond," *Journal of Medicinal Chemistry*, vol. 32, no. 7, pp. 1652-1661, 1989.
15. F. R. Atherton, C. H. Hassall, and R. W. Lambert, "Synthesis and structure-activity relationships of antibacterial phosphonpeptides incorporating (1-aminoethyl) phosphonic acid and (aminomethyl) phosphonic acid," *journal of medicinal chemistry*, vol. 29, no. 1, pp. 29-40, 1986.

16. R. F. Pratt, "Inhibition of a class C β -lactamase by a specific phosphonate monoester," *Science*, vol. 246, no. 4932, pp. 917-919, 1989.
17. G. lavielle, P. Hautefaye, C. Schaffer, J. A. Boutin, C. A. Cudennec, and A. Pierre, "New α -amino phosphonic acid derivatives of vinblastine: chemistry and antitumor activity," *Journal of medicinal chemistry*, vol. 34, no. 7, pp. 1998-2003, 1991.
18. P. Kafarski and B. Lejczak, "Biological activity of aminophosphonic acids," *phosphorus, Sulfur, and Silicon and the Related Elements*, vol. 63, no. 1-2, pp. 193-215, 1991.
19. V. Ria and I. N. N. Namboothiri, "Enantioselective conjugate addition of dialkyl phosphites to nitroalkenes," *Tetrahedron Asymmetry*, vol. 19, 20, pp. 2335-2338, 2008.
20. E. Kuliszewska, M. Hanbauer, and F. Hammerschmidt, "preparation of α -aminobenzylphosphonic acids with a stereogenic quaternary carbon atom via microscopically configurationally stable α -aminobenzylolithiums," *Chemistry A*, vol. 14, no. 28, pp. 8603-8614, 2008.
21. V. Coeffard, I. Beaudet, M. Evain, E. Le Grogne, and J. P. Quintard, "Preparation and transmetallation of enantioenriched α -aminoorganostannanes derived from N-boc phenylglycinol: Organic Chemistry, vol. 2008, no. 19, pp. 3344-3351, 2008.
22. R. A. Cherkasov and V. I. Galkin, "The Kabachnik-fields reaction: synthetic potential and the problem of the mechanism," *Russian Chemical Reviews*, vol. 67, no. 10, pp. 587-882, 1998.
23. N. S. Zefirov and E. D. Matveeva, "Catalytic Kabachnik-Fields reaction: new horizons for old reaction," *ARKIVOC*, vol. 1, no. 11, pp. 1-17, 2008.
24. R. Jacquier, M. L. Hassani, C. Petrus, and F. Petrus, "Asymmetric synthesis of 1-aminoalkylphosphonic acids," *Phosphorus, Sulfur, and Silicon and the Related Elements*, vol. 81, no. 1-4, pp. 83-87, 1993.
25. G. Jommi, G. Miglierini, R. Pagliarin, g. Sello, and M. Sisti, "Studies toward a model for predicting the diastereoselectivity in the electrophilic amination of chiral 1,2,3-oxazaphospholanes," *Tetrahedron*, vol. 48, no. 35, pp. 7275-7288, 1992.
26. S. Hannesian and Y. Bennani, "Electrophilic amination and azidation of chiral α -alkyl phosphonamides: asymmetric syntheses of α -amino α -alkyl phosphonic acids," *Synthesis*, vol. 1994, no. 12, pp. 1272-1276, 1994.
27. S. E. Denmark, N. Chatani, and S. V. Pansare, "Asymmetric electrophilic amination of chiral phosphorus-stabilized anions," *Tetrahedron*, vol. 48, no. 11, pp. 2191-2208, 1992.
28. N. Pudovik, "New method of synthesis of esters of phosphonocarboxylic acids and their derivatives," *Doklady Akademii Nauk*, vol. 85, pp. 349-351, 1952.
29. N. Pudovik and I.V. Konovalova, "Addition reactions of esters of phosphorus (III) acids with unsaturated systems," *Synthesis*, vol. 1979, no. 2, pp. 81-96, 1979.
30. M. I. Kabachnik and T. Y. Medvel, "New synthesis of aminophosphonic acids," *Doklady Akademii Nauk*, vol. 83, pp. 689-692, 1952.
31. E. K. Fields, "The synthesis of esters of substituted amino phosphonic acids," *Journal of the American Chemical Society*, vol. 74, no. 6, pp. 1528-1531, 1952.
32. D. F. Wiemer, "Synthesis of nonracemic phosphonates," *Tetrahedron*, vol. 53, no. 49, pp. 16609-16644, 1997.

33. C. Ranu and A. Hajra, "A simple and green procedure for the synthesis of α -aminophosphonate by a one-pot threecomponent condensation of carbonyl compound, amine and diethyl phosphite without solvent and catalyst," *Green Chemistry*, vol. 4, no. 6, pp. 551-554, 2002.