Effects of interval training in patients with coronary artery disease and positive exercise stress test: a pilot study

Iwona Korzeniowska-Kubacka, Barbara Dobraszkiewicz-Wasilewska, Maria Bilińska Ewa Rydzewska, Stanisław Rudnicki, Ryszard Piotrowicz.

Department of Cardiac Rehabilitation and Noninvasive Electrocardiology
Institute of Cardiology, Warsaw, Poland

Summary

Background: Ischemia during exercise stress test (EST) in patients (pts) with stable CAD is not a contraindication to physical training. There are only few publications on training effects and safety in patients (pts) with positive stress test. The aim of the study was to assess the effects and safety of interval training in patients with stable CAD and exercise-induced myocardial ischemia.

Material and Methods: Thirty male pts, aged 55 ± 10 years, after MI or/and CABG with positive EST underwent a three-month interval training program on cycloergometer, which included 2 cycles of 16 trainings each, three times a week. EST was performed before and at the end of each cycle of trainings. The training HR limit was set 10 beats below the HR at which ST segment depressed by 1 mm during baseline EST. The following variables were assessed during EST: walking distance (m), duration (min), time to 1 mm ST depression (min), double product (DP, mmHg/min) at 1 mm ST depression, maxVO₂ (ml/kg/min), metabolic equivalent (METs), and complications during training.

Results: After a training program all analyzed variables improved significantly: walking distance (565.6 ± 142.6 vs 686.1 ± 150.5 , $p < 0.001$), EST duration (11.1 ± 2.07 vs 12.7 ± 2.1 , $p < 0.001$), time to 1 mm ST depression (10.2 ± 2.3 vs 11.8 ± 2.3 , $p = 0.001$), DP at 1 mm ST depression (186.5 ± 51.3 vs 208 ± 48.6 , $p < 0.03$), maxVO₂ (21.4 ± 4.6 vs 24.6 ± 5.1 , $p < 0.001$), and maximal workload (6.35 ± 1.6 vs 7.49 ± 1.5 , $p = 0.002$). There were no complications during training.

Conclusion: Interval training below the threshold of ischemia is effective and safe in patients with stable CAD and exercise-induced myocardial ischemia.

Key words

coronary artery disease, physical training, positive exercise stress test.

Ocena efektów treningu interwałowego u pacjentów z chorobą niedokrwienną serca i dodatnią próbą wysiłkową

Streszczenie

Wstęp. Dodatnia próba wysiłkowa u pacjentów ze stabilną chorobą wieńcową nie stanowi przeciwwskazań do treningu fizycznego. Jednak w piśmiennictwie jest niewiele prac oryginalnych dotyczących efektów i bezpieczeństwa takiego treningu.

Cel pracy. Celem pracy była ocena efektów i bezpieczeństwa treningu fizycznego u pacjentów ze stabilną chorobą wieńcową i dodatnią próbą wysiłkową.

Metody. Badaniami objęto 30 pacjentów (mężczyźni, wiek 55 ± 10 lat) po zawale serca i/lub operacyjnym leczeniu choroby wieńcowej z dodatnią próbą wysiłkową. Wszyscy badani uczestniczyli w 3 miesięcznym programie treningowym, który zawierał 2 cykle po 16 treningów interwałowych na cycloergometrze 3 razy w tygodniu. Na wstępie i po zakończeniu cyklu treningów wykonywano ergospirometryczną próbę wysiłkową (EPW). Limit tętna treningowego wyznaczono do częstotliwości rytmu serca o 10 uderzeń/min mniejszy od tej, przy której pojawiało się obniżenie odcinka ST o 1 mm we wstępnej próbie wysiłkowej. Ocenie poddano: czas trwania próby (minuty), czas do obniżenia odcinka ST o 1 mm (minuty), DP do obniżenia odcinka ST o 1 mm (mmHg/min), maxVO₂ (ml/kg/min), przebytą drogę (metry), maksymalne obciążenie (MET) oraz powikłania w czasie treningu.

Wyniki. Po odbyciu programu treningowego nastąpiło istotne wydłużenie czasu próby z $11,1 \pm 2$ do $12,7 \pm 2,1$ ($p < 0,001$), wydłużenie czasu do obniżenia ST o 1 mm z $10,2 \pm 2,3$ do $11,8 \pm 2,3$ ($p = 0,001$), wzrost DP przy obniżeniu ST o 1 mm z $186,5 \pm 51,3$ do $208 \pm 48,6$ ($p < 0,03$), wzrost maxVO_2 z $21,4 \pm 4,6$ do $24,6 \pm 5,1$ ($p < 0,001$), przebytej drogi z $565,6 \pm 142,6$ do $686,1 \pm 150,5$ ($p < 0,001$) i maksymalnego obciążenia z $6,35 \pm 1,6$ do $7,49 \pm 1,5$ ($p = 0,002$). Nie stwierdzono żadnych powikłań w czasie trwania treningów.

Wnioski. Trening interwałowy poniżej progu niedokrwienia jest efektywny i bezpieczny u pacjentów ze stabilną chorobą wieńcową i dodatnią próbą wysiłkową.

Słowa kluczowe

trening fizyczny, dodatnia próba wysiłkowa

Background

Patients after myocardial infarction (MI) referred for cardiac rehabilitation on the out-patient basis include those whose exercise stress test (EST) has been positive.

In our population of rehabilitated patients with proved coronary artery disease (CAD) exercise stress test appeared to be negative in a majority (87%) and positive in only 13% of patients (1).

According to the present standards, myocardial ischemia during an exercise stress test displayed as horizontal or downsloping ST segment depression on ECG and/or anginal pain, needs to be analysed thoroughly before a training program is designed.

The intensity of exercise is derived from the results of the exercise stress test. The limit of the training heart rate (HR) should be 10 beats per minute (bpm) below the ischemia threshold, which results from angina and/or ischemic ST segment depression by ≥ 1 mm during an exercise stress test (2). Moreover during training, patients after MI with positive exercise stress test should be carefully monitored by experienced personnel.

To date, the literature provides only few original studies of this issue, and the answers to the question if ischemia during exercise should be avoided are inconsistent (2-4).

Therefore, the objective of the study was to assess the effects and safety of physical training in patients with stable coronary artery disease and positive exercise stress test.

Material and Methods

The study comprised 30 male patients, aged 55 ± 10 years, who entered the cardiac rehabilitation program 32.8 ± 20.8 days after MI and/or after coronary artery bypass grafting (CABG), with positive exercise stress test. All the patients had preserved left ventricular function (ejection fraction $> 50\%$) and no evidence of right ventricular dysfunction.

Patients were excluded if they had unstable angina, congestive heart failure, uncontrolled hypertension (blood pressure $> 180/120$ mmHg at rest), cardiac rhythm disturbances, valvular heart disease, mitral valve prolapse, left bundle branch block (at rest or during exercise), echocardiographically defined left ventricular hypertrophy, diabetes, impaired renal or hepatic function.

All the patients underwent a three-month interval training program on cycloergometer. Before the program and after the first and second cycles of trainings, patients underwent an exercise stress test.

The study protocol was approved by the Institutional Ethics Committee on human research and all the participants gave their written informed consent.

Exercise stress test

The exercise stress test was performed on a treadmill according to a modified Bruce protocol (ZAN 600 USB CPX equipment by Reynolds) before the program and after the first and second training cycles. A 3-lead ECG, usually aVF, V2, V5 was monitored continuously before, during, and for 10 min after the exercise test. EST was discontinued when a subject became tired, his blood pressure exceeded 230/130 mmHg, in the case of ST segment depression by at least 2 mm, anginal pain, blood pressure drop by more than 10 mmHg compared to the baseline value. An exercise stress test was considered positive when ST segment depression of at least 1 mm was horizontal or downsloping, 80 ms beyond the J point. The following parameters were analyzed: walking distance (m), duration (min), time to 1 mm ST segment depression (min), maximal oxygen consumption (maxVO_2 , ml/kg/min), metabolic equivalents (METs), heart rate (HR, bpm), double product, i.e. product of heart rate and systolic blood pressure (DP, mmHg/min $\times 100$) at rest, at peak exercise and at 1 mm ST segment depression and maximal ST segment depression (mm).

Interval training

All subjects underwent a interval training program on cycloergometer (Ergoline) three times a week. The training program consisted of two training cycles of 16 trainings each. Each session lasted 40 minutes. The training heart rate limit was set at 10 bpm below the heart rate when 1 mm ST segment depression had occurred during the baseline exercise stress test. During the study period all patients were carefully monitored whether complications such as unstable angina, MI, heart failure and / or serious cardiac rhythm disturbances had occurred.

Statistics

All results are expressed as mean $\pm$ SD. Mean values of quantitative variables were compared with one-way analysis of variance followed by Student's t-test for matched pairs to assess the significance of differences between means. Non-normally distributed variables were compared with Kruskal-Wallis test followed either by the Mann-Whitney or the Wilcoxon rank test, as appropriate. The value of $p < 0.05$ was considered statistically significant.

Statistical analysis was performed using SAS statistical software (version 8.02; Cary, North Carolina, USA).

Results

We studied 30 male patients, aged 55 ± 10 years, with stable angina pectoris (Canadian Cardiovascular Society class 0 or I) and reproducible positive exercise tests performed previously. The distribution of coronary risk factors (included body mass index, hypertension, smoking habits, hyperlipidemia, diabetes) and baseline clinical characteristics of the studied patients are presented in Table 1.

Table 1. Baseline clinical characteristics of the study group.

Characteristics	Patients n=30
Mean age (years)	55 ± 10
Risk factors:	
BMI (kg/m ²)	27.6 ± 2.7
Hypertension, n (%)	19 (63.4)
Smokers:	
- past, n (%)	14 (46.6)
- current, n (%)	5 (16.6)
Hyperlipidemia, n (%)	28 (93.4)
Diabetes, n (%)	7 (23.4)
History of MI, n (%)	21 (70)
History of CABG, n (%)	15 (50)
CCS Classification:	
- Class 0, n (%)	18 (60)
- Class I, n (%)	12 (40)
Vessel disease, n (%)	
one-vessel	14 (46.6)
two-vessel	6 (20)
three-vessel	10 (33.4)
Start of rehabilitation (days after MI or CABG)	32.8 ± 20.8
LVEF (%)	53.4 ± 3.2
Physical capacity (max VO ₂ ml/kg/min)	21.45 ± 4.61
Medications:	
Beta-blockers (%)	93
ACE-I (%)	93
Antiplatelet agents (%)	100
Statins (%)	100

All values are presented as mean $\pm$ SD or percentages; BMI, body mass index; MI, myocardial infarction; CABG, coronary artery bypass grafting;

CCS, Classification of angina according to the Canadian Cardiovascular Society;

LVEF, left ventricular ejection fraction; max VO₂, maximal oxygen uptake;

ACE-I, angiotensin-converting enzyme inhibitors.

All patients had proved coronary artery disease (CAD), and the majority of them had history of myocardial infarction (MI). With respect to angiographic findings, almost two

in third of our patients had either left main stenosis $\geq 50\%$ or three vessel disease. Mean time between MI or surgery (CABG) and the start of rehabilitation was 32.8 ± 20.8 days. Moreover, at entry all the patients had preserved left ventricular ejection fraction (ejection fraction $>50\%$) and no evidence of right ventricular dysfunction on echocardiography and good physical capacity. In addition, all patients had prescribed antiplatelet agents (aspirin and/or clopidogrel) and statins, whereas prescription of beta-blockers and ACE inhibitors were slightly lower. None of the patients received nitrates or diuretics.

Exercise stress testing results before (I), after the first (II) and the second (III) cycle of trainings presented Table 2.

Table 2. Exercise stress testing results before (I), after the first (II) and the second (III) cycle of trainings.

	Before trainings (I)	After trainings (II)	After trainings (III)	p		
				I vs II	II vs III	I vs III
Walking distance (m)	565.6 ± 142.67	678.33 ± 118.06	686.11 ± 150.52	<0.001	ns	<0.001
Duration (min)	11.12 ± 2.07	12.54 ± 1.38	12.70 ± 2.14	<0.001	ns	<0.001
maxVO ₂	21.45 ± 4.61	24.32 ± 4.71	24.67 ± 5.11	<0.001	ns	<0.001
Workload (METs)	6.35 ± 1.61	7.49 ± 1.12	7.49 ± 1.59	0.001	ns	0.002
HR at rest	65.6 ± 10.5	65.27 ± 9.68	64.86 ± 9.9	ns	ns	ns
HR at peak	119.07 ± 12.54	124.4 ± 15.76	124.71 ± 12.78	<0.02	ns	0.02
DP at rest	77.47 ± 16.76	78.90 ± 14.64	82.54 ± 15.79	ns	ns	ns
DP at peak	199.43 ± 49.05	211.80 ± 42.99	222.89 ± 46.4	ns	ns	<0.02
DP1mm	186.57 ± 51.35	189.50 ± 45.95	208.04 ± 48.68	ns	ns	<0.03
Time to 1 mm ST depression (min)	10.23 ± 2.38	11.20 ± 2.31	11.82 ± 2.35	<0.03	0.02	0.001
Max ST depression (mm)	1.24 ± 0.29	1.35 ± 0.47	1.17 ± 0.32	ns	ns	ns

Data are expressed as mean $\pm$ SD; ns, non significant; maxVO₂, maximal oxygen uptake (ml/kg/min); HR, heart rate (beats per minute, bpm);

DP, double product (heart rate-systolic blood pressure product, mmHg $\times$ 100);

DP 1mm, double product at 1 mm ST segment depression

The subjects' physical capacity improved significantly after the interval training. The improvement was visible already after the first cycle of trainings, as longer test duration, walking distance, higher max VO₂ and maximal workload (Table 2).

Moreover, the training program resulted in a significant lengthening of the time until 1 mm ST segment depression after the first and the second cycle of trainings (Table 2).

Of note, physical trainings also caused a significant increase in DP at 1 mm ST segment depression, which represents an objective index of myocardial oxygen demand at the onset of ischemia. In addition, ST segment depression was observed to have decreased from 1.24 ± 0.29 mm to 1.17 ± 0.32 mm, yet these differences were not statistically significant.

None of the subjects developed complications while performing the training program.

Discussion

Incorporating the interval training into the rehabilitation program resulted in an improvement in physical capacity assessed from a significant increase in maxVO₂, maximal workload, lengthening of the test duration and the distance covered. The fact has been confirmed in numerous papers reporting a beneficial impact of training on physical capacity in CAD patients (5-7). In our material, the physical capacity improved significantly already after the first cycle of the training program. Continuing the training program did not result in further improvement, it only kept up the physical capacity achieved. The study focused on subjects whose stress test was positive. Following the training, the time to 1 mm ST segment depression during exercise stress test prolonged significantly and more importantly, DP at 1 mm ST segment depression increased, which may be associated with the myocardium lower demand for oxygen at a higher workload.

A beneficial impact of the training on the myocardial oxygen demand was observed already after the first training cycle. When the training was continued, the time to 1 mm ST segment depression during exercise stress test prolonged even more.

It has been established that myocardial oxygen consumption during exercise stress test can be estimated from the product of heart rate and systolic pressure, called double product (DP) (2). There is a linear dependence between myocardial oxygen consumption and coronary blood flow. In our study, even the training below the ischemia threshold improved the balance between oxygen supply and myocardial demand, thus increasing the threshold of ischemia and displaying an anti-ischemic activity.

To date, there are only few publications examined the influence of exercise training on myocardial perfusion using, beyond standard ECG exercise stress test, nuclear techniques.

Belardinelli et al. observed an increase in physical capacity and myocardial perfusion assessed scintigraphically as soon as after 8 weeks of medium effort at 60% peak VO₂ in patients with ischemic cardiomyopathy. Prolonging the training to 12 months did not result in changes in myocardial perfusion (8). Nishikawa et al. trained CAD patients with preserved left ventricular function and scintigraphically confirmed ischemia during exercise stress test, whose physical capacity and cardiac perfusion improved after a 12-week training (9).

In our study, performing only standard ECG exercise stress

test, we were able to assess only myocardial oxygen demand. As we mentioned above, DP at 1 mm ST segment depression, which represents an objective index of myocardial oxygen demand at the onset of ischemia, as well as time to 1 mm ST depression increased significantly in all patients after the training program. On the contrary, Digenio et al., after a 6-month physical training, observed an improvement in physical capacity, however, the time to 1 mm ST segment depression during exercise stress test did not prolong. The authors suggest it might have resulted from the fact that the focus had been on physical capacity improvement rather than on modifying the risk factors which would have impact on endothelium function and myocardial perfusion (slowed down the atherosclerotic process) (10). Similarly, the Heidelberg Study findings also showed only an increase in physical capacity by 28%, but DP at 1 mm ST depression had not changed. These authors believed the training might not have been intensive enough to provoke an anti-ischemic effect (11).

Importantly, Ehsani et al., training CAD patients for 12 months at 50% to 80% maxVO₂, regularly provoked uncomplicated myocardial ischemia during trainings (12). They observed not only a significant increase in physical capacity. The 12-month training resulted in decreased maximal ST segment depression from 0.2 ± 0.04 mm to 0.16 ± 0.08 and a 22% increase in DP at 1 mm ST segment depression. The authors suggest that ischemia should be provoked during training, because it may trigger the phenomenon of myocardial preconditioning, consequently protecting it from another ischemic stress.

It is noteworthy, that according to the standards, only training below ischemic threshold is recommended as safe (2).

Recently, Noël et al. studied the safety of ischemia-provoking training (13). The study comprised 22 patients with CAD and positive exercise stress test. Half of them trained above ischemia threshold, and half below. None of the patients developed such complications as dangerous arrhythmia, troponin level increase or myocardial dysfunction. Training resulted in a significant increase of maximal workload (Watts) in both groups, but maxVO₂, DP and ST segment depression during exercise stress test did not change significantly.

The results of the training below ischemia threshold presented in the paper show that even such training has an anti-ischemic influence. Recommending training which provokes ischemia needs to be investigated further.

In conclusion, interval training below ischemia threshold is effective and safe in patients with stable CAD and exercise-induced myocardial ischemia. However, all patients with positive exercise stress test should be carefully monitored by experienced personnel.

References

1. Korzeniowska-Kubacka I, Wolszakiewicz J, Smolis-Bąk E, Kongres Europrevent Ateny 2006. *Kardiologia Polska* 2006;64: 1043-1044
2. Fletcher G F, Balady GJ, Amsterdam EA et al. Exercise standards for testing and training: A statement for health care professionals from the American Heart Association. *Circulation* 2001; 104 :1694-1740.

3. Bertelinek R, Oh P, Exercise training and the ischemic patient. *Curr Issues Card Rehabil Prev* 2006; 14: 17-20.
4. Nigam A, Cardiff JC. The place of exercise in the patient with chronic stable angina. *Heart Metab* 2008; 38: 34-37.
5. Laslett LJ, Paumer L, Amsterdam EA. Increase in myocardial oxygen consumption indexes by exercise training at onset of ischemia in patients with coronary artery disease. *Circulation* 1985; 71: 958-962.
6. Hofman-Bang C, Lisspers J, Nordlander R et al. Two-year results of a controlled study of residential rehabilitation for patients treated with percutaneous transluminal coronary angioplasty: a randomised study of a multifactorial programme. *Eur Heart J* 1999; 20:1448-50.
7. Maines TY, Lavie CJ, Milani RV et al. Effect of cardiac rehabilitation and exercise programs on exercise capacity, coronary risk factors, behaviour, and quality of life in patients with coronary artery disease. *South Med J* 1997;90: 43-49.
8. Belardinelli R, Georgiou D, Ginzton L, Cianci G, Purcaro A. Effects of moderate exercise training on thallium uptake and contractile response to low dose dobutamine of dysfunctional myocardium in patients with ischemic cardiomyopathy. *Circulation* 1998; 97: 553-561
9. Nishikawa E, Tomiyama H, Inagaki M et al. Safety and effectiveness of exercise training in patients with silent myocardial ischemia. *J Cardiol* 1996; 28: 321-7
10. Digenio AG, Noakes TD, Joughin H, Daly L. Effect of myocardial ischaemia on left ventricular function and adaptability to exercise training. *Med Sci Sports Exerc* 1999; 31: 1094-1101.
11. Gohlke H. Exercise training in Coronary Heart Disease. In; Perk J .Gohlke H, Monpere C, Hellemans J, McGea H, Sellier P, Saner H (eds). *Cardiovascular Prevention and Rehabilitation*, London: Springer; 2007, 126-127.
12. Ehsani AA, Heath GW, Hagberg JM, Sobel BE, Holloszy JO. Effects of 12 months of intense exercise training on ischemic ST-segment depression in patients with coronary artery disease. *Circulation* 1981; 64: 1116-1124.
13. Noël M, Jobin J, Marcoux A, Poirier P, Dagenais GR, Bogaty P. Can prolonged exercise-induced myocardial ischemia be innocuous? *Eur Heart J* 2007;28: 1559-1565.