ON A CHARACTERIZATION OF HILBERTIAN NORMS

BY

TEODOR PRECUPANU

Abstract. In this paper we prove that a characterization of hilbertian norms established in smooth normed spaces X with $\dim X \geq 3$ also holds in two dimensional spaces.

Mathematics Subject Classification 2000: 46B20, 46C15.

Key words: smooth linear normed space, hilbertian norm, height of a triangle.

1. Introduction

Let X be a real smooth linear normed space, that is there exists

$$\lim_{t \rightarrow 0} \frac{\|x + ty\|^2 - \|x\|^2}{2t}$$

for all $x, y \in X$. Usually, this limit is denoted by $n'(x; y)$. It is well known, that $n(x; \cdot)$ is a linear continuous functional on X and $n'(x; x) = \|x\|^2$ for every $x \in X$. Therefore, X is a smooth space (see [1], [2]). Using an euclidean property of heights of a triangle, in [4] it is established the following characterization of hilbertian norms:

Theorem 1. *In a smooth real linear normed space X with $\dim X \geq 3$ the norm is hilbertian if and only*

$$(1) \quad \begin{aligned} & n'(y; x)[n'(x; y)n'(x - y; x) - \|x\|^2 n'(x - y; y)] \\ & = n'(x; y)[\|y\|^2 n'(x - y; x) - n'(y; x)n'(x - y; y)], \quad \forall x, y \in X. \end{aligned}$$

In the present paper we show that this result is also true if $\dim X = 2$. Our goal is to find the smooth norms of a two dimensional space X which have the property (1). For this we use an idea of [3] taking into account that $n'(x; \cdot)$ is a linear functional for any $x \in X$. Thus, if $x = (x_1, x_2)$, $y = (y_1, y_2)$, $x_1, x_2, y_1, y_2 \in \mathbb{R}$, then

$$(2) \quad n'(x; y) = A(x_1, x_2)y_1 + B(x_1, x_2)y_2,$$

and

$$(3) \quad \|x\|^2 = A(x_1, x_2)x_1 + B(x_1, x_2)x_2,$$

where A, B are two real functions on $\mathbb{R}^2$. Moreover, since $n'(tx; y) = tn'(x; y)$, for all $t \in \mathbb{R}$, we also have

$$(4) \quad \begin{aligned} A(tx_1, tx_2) &= tA(x_1, x_2), \\ B(tx_1, tx_2) &= tB(x_1, x_2), \quad \text{for all } t \in \mathbb{R}. \end{aligned}$$

2. Proof of Theorem 1 in two dimensional spaces

Taking into account the equalities (2) and (3), the property (1) becomes

$$\begin{aligned} &(A(y_1, y_2)x_1 + B(y_1, y_2)x_2)[(A(x_1, x_2)y_1 + B(x_1, x_2)y_2) \\ &(A(x_1 - y_1, x_2 - y_2)x_1 + B(x_1 - y_1, x_2 - y_2)x_2) - (A(x_1, x_2)x_1 \\ &+ B(x_1, x_2)x_2)(A(x_1 - y_1, x_2 - y_2)y_1 + B(x_1 - y_1, x_2 - y_2)y_2)] \\ &= (A(x_1, x_2)y_1 + B(x_1, x_2)y_2)[(A(y_1, y_2)y_1 + B(y_1, y_2)y_2) \\ &(A(x_1 - y_1, x_2 - y_2)x_1 + B(x_1 - y_1, x_2 - y_2)x_2) - (A(y_1, y_2)x_1 \\ &+ B(y_1, y_2)x_2)(A(x_1 - y_1, x_2 - y_2)y_1 + B(x_1 - y_1, x_2 - y_2)y_2)]. \end{aligned}$$

If $x = (1, t)$, $y = (0, t)$, $t \in \mathbb{R}$, we obtain

$$\begin{aligned} &(A(0, t) + tB(0, t))[tB(1, t)(A(1, 0) + tB(1, 0)) - tB(1, 0)(A(1, t) + tB(1, t))] \\ &= tB(1, t)[tB(0, t)(A(1, 0) + tB(1, 0)) - tB(1, 0)(A(0, t) + tB(0, t))]. \end{aligned}$$

By (4) we get

$$\begin{aligned} &(A(0, 1) + tB(0, 1))[B(1, t)(A(1, 0) + tB(1, 0)) - B(1, 0)(A(1, t) + tB(1, t))] \\ &= tB(1, t)[B(0, 1)(A(1, 0) + tB(1, 0)) - B(1, 0)(A(0, 1) + tB(0, 1))], \end{aligned}$$

that is,

$$(5) \quad (A(0,1) + tB(0,1))(A(1,0)B(1,t) - B(1,0)A(1,t)) \\ = tB(1,t)(A(1,0)B(0,1) - A(0,1)B(1,0)).$$

Therefore, if $B(1,t) \neq 0$, we obtain

$$\frac{A(1,t)}{B(1,t)} = \frac{A(0,1)(tB(1,0) + A(1,0))}{B(1,0)(tB(0,1) + A(0,1))},$$

whenever $t \neq -\frac{A(0,1)}{B(0,1)}$, $B(0,1) \neq 0$. Hence, the function $t \rightarrow \frac{A(1,t)}{B(1,t)}$, $t \in \mathbb{R}$, is a real function of homographic type, i.e.

$$\frac{A(1,t)}{B(1,t)} = \frac{\alpha t + \beta}{\gamma t + \alpha}, \quad \text{if } B(1,t) \neq 0, \quad t \neq -\frac{\alpha}{\gamma},$$

where $\alpha = A(0,1)A(1,0)$, $\beta = A(0,1)B(1,0)$, $\gamma = B(0,1)B(1,0)$.

On the other hand if $x_1 \neq 0$, $x_2 \neq 0$ and $B(1, \frac{x_2}{x_1}) \neq 0$ we have

$$n'(x,y) = x_1 B\left(1, \frac{x_2}{x_1}\right) \left[\frac{A(1, \frac{x_2}{x_1})}{B(1, \frac{x_2}{x_1})} y_1 + y_2 \right] = x_1 B\left(1, \frac{x_2}{x_1}\right) \left[\frac{\alpha x_2 + \beta x_1}{\gamma x_2 + \alpha x_1} y_1 + y_2 \right].$$

By (3) it follows

$$(6) \quad \|x\|^2 = x_1 B\left(1, \frac{x_2}{x_1}\right) \frac{\beta x_1^2 + 2\alpha x_1 x_2 + \gamma x_2^2}{\alpha x_1 + \gamma x_2}.$$

Since the norm is differentiable it is well known that

$$A(x_1, x_2) = \frac{\partial}{\partial x_1} \left(\frac{1}{2} \|x\|^2 \right), \quad B(x_1, x_2) = \frac{\partial}{\partial x_2} \left(\frac{1}{2} \|x\|^2 \right),$$

and so, denoting $B(1,t) = f(t)$, $t \in \mathbb{R}$, by usual computations we find

$$x_1 f\left(\frac{x_2}{x_1}\right) = \frac{\partial}{\partial x_2} \left(\frac{1}{2} \|x\|^2 \right) = \frac{1}{2} x_1 f\left(\frac{x_2}{x_1}\right) \frac{(2\alpha^2 - \beta\gamma)x_1^2 + 2\alpha\gamma x_1 x_2 + \gamma^2 x_2^2}{(\alpha x_1 + \gamma x_2)^2} \\ + \frac{1}{2} f'\left(\frac{x_2}{x_1}\right) \frac{\beta x_1^2 + 2\alpha x_1 x_2 + \gamma x_2^2}{\alpha x_1 + \gamma x_2},$$

i.e.

$$\begin{aligned} x_1 f\left(\frac{x_2}{x_1}\right) & \frac{2(\alpha x_1 + \gamma x_2)^2 - (2\alpha^2 - \beta\gamma)x_1^2 - 2\alpha\gamma x_1 x_2 - \gamma x_2^2}{\alpha x_1 + \gamma x_2} \\ & = f'\left(\frac{x_2}{x_1}\right) (\beta x_1^2 + 2\alpha x_1 x_2 + \gamma x_2^2). \end{aligned}$$

Dividing by $\beta x_1^2 + 2\alpha x_1 x_2 + \gamma x_2^2$ we obtain the following simple differential equation $\frac{f'(t)}{f(t)} = \frac{\gamma}{\gamma t + \alpha}$. Thus, it follows that $f(t) = (\gamma t + \alpha)k$, where k is a real constant. In fact, $k = \|(1, 0)\|^{-2}$.

Now, we return to the relation (6) and so we get

$$(7) \quad \|x\|^2 = k(\beta x_1^2 + 2\alpha x_1 x_2 + \gamma x_2^2).$$

If $B(1, t) = 0$, then from (3) it follows that $\|(1, t)\|^2 = A(1, t) > 0$. In this case we consider the function $g(t) = A(1, t)$, $t \in \mathbb{R}$. Similarly, we obtain the same equality (7). Also, since the norm is a continuous function, the equality (7) also holds if $x_1 = 0$ or $x_2 = 0$ or $\frac{x_2}{x_1} = -\frac{\alpha}{\gamma}$.

Finally, we established (7) for all $(x_1, x_2) \in \mathbb{R}^2$.

Therefore, the norm is obviously hilbertian.

Remark. Since a norm is hilbertian if its restriction to every two dimensional linear subspace is hilbertian, the above proof can be considered as a new proof of Theorem 4 in [4] for any linear normed space.

REFERENCES

1. BARBU, V.; PRECUPANU, T. – *Convexity and Optimization in Banach Spaces*, D. Reidel Publ. Comp., Dordrecht, 1986.
2. HOLMES, R.B. – *Geometric Functional Analysis and Its Applications*, Graduate Texts in Mathematics, Springer-Verlag, New York-Heidelberg-Berlin, 1975.
3. Precupanu, T. – *Ortogonalitate in spații liniare*, (in Romanian), St. Cerc. Mat., 6 (1972), 881-889.
4. PRECUPANU, T.; IONICA, I. – *Heights of a triangle in a linear normed space and Hilbertian norms*, An. Științ. Univ. "Al.I. Cuza" Iași. Mat. (N.S.), 55 (2009), 35-47.

Received: 15.III.2009

*Department of Mathematics,
University "Al.I. Cuza",
11, Bd. Carol I, 700506, Iași,
ROMANIA
tprecup@uaic.ro*