NEW EXAMPLES OF BIHARMONIC SUBMANIFOLDS IN $\mathbb{C}P^n$ AND $\mathbb{S}^{2n+1}$ *

BY

WEI ZHANG

Abstract. We construct biharmonic real hypersurfaces and biharmonic Lagrangian submanifolds of Clifford torus type in $\mathbb{C}P^n$ via the Hopf fibration; and get new examples of biharmonic submanifolds in $\mathbb{S}^{2n+1}$ as byproducts.

Mathematics Subject Classification 2000: 58E20.

Key words: biharmonic maps, real hypersurfaces, Lagrangian submanifolds, Hopf fibration, Clifford torus.

1. Introduction

Let $\phi : (M, g) \rightarrow (N, h)$ be a smooth map between two Riemannian manifolds and define its first tension field as $\tau(\phi) = \text{trace } \nabla d\phi$. The map ϕ is called harmonic if it is a critical point of the energy functional

$$E(\phi) = \frac{1}{2} \int_M |d\phi|^2 v_g.$$

This amounts to $\tau(\phi) = 0$. In the isometric immersions case, harmonic is equivalent to minimal.

Considering the bienergy functional $E_2(\phi) = \frac{1}{2} \int_M |\tau(\phi)|^2 v_g$, its critical points are called biharmonic maps ([10],[7]). The associated Euler-Lagrange equation is given by the vanishing of the bitension field

$$\tau_2(\phi) = -\Delta^\phi \tau(\phi) - \text{trace } R^N(d\phi, \tau(\phi))d\phi,$$

*Supported by Tian Yuan Math. Fund. and the Fundamental Research Funds for the Central Universities

where $R^N(X, Y)Z = \nabla_X^N \nabla_Y^N Z - \nabla_Y^N \nabla_X^N Z - \nabla_{[X, Y]}^N Z$.

Obviously, any harmonic map is biharmonic. The non-harmonic biharmonic maps are called proper biharmonic, and a submanifold is proper biharmonic if its inclusion map is proper biharmonic.

Concerning the proper biharmonic maps, there are several non-existence results for the non-positive sectional curvature codomains([10, 3, 7]), for instance

Theorem 1 ([3, 4]). *Let $\phi : (M, g) \rightarrow (N, h)$ be a smooth map. If M is compact, orientable and $\text{Riem}^N \leq 0$, then ϕ is biharmonic if and only if it is harmonic.*

Suggested by similar results

Generalized Chen's Conjecture. *Biharmonic submanifolds of a manifold N with $\text{Riem}^N \leq 0$ are minimal.*

This conjecture has been proved to be false recently by OU and TANG ([8]), while it is still sensible to focus on the proper biharmonic submanifolds in sphere or other non-negatively curved spaces. Although some partial classification results had been obtained ([1]), the known examples are still relatively rare. In this article, we will construct series of new examples of proper biharmonic submanifolds in $\mathbb{S}^{2n+1}$ and $\mathbb{C}P^n$.

Theorem 2. *Denote by $M_{p,q}^C(r, s)$ the image of the generalized Clifford torus $M_{p,q}(r, s) = \mathbb{S}^{2p+1}(r) \times \mathbb{S}^{2q+1}(s)$ in $\mathbb{S}^{2n+1}$ via the Hopf map, where $p + q = n - 1$. Then it is biharmonic in $\mathbb{C}P^n$ if and only if*

$$(1) \quad \left(\frac{s}{r}\right)^2 (2p+1) + \left(\frac{r}{s}\right)^2 (2q+1) = 2(n+2).$$

And

Theorem 3. *Identify $\mathbb{R}^{2n+2}$ with $\mathbb{C}^{n+1}$. Define the Clifford torus $\mathbb{T}^{n+1} = \{z = (z_1, \dots, z_i, \dots, z_{n+1}) \in \mathbb{C}^{n+1} \mid |z_i| = a_i, \sum_{i=1}^{n+1} a_i^2 = 1\}$ in $\mathbb{S}^{2n+1}$. Then it is biharmonic if and only if*

$$(2) \quad a_i d - \frac{1}{a_i^3} = 2(n+1) \left((n+1)a_i - \frac{1}{a_i} \right), \quad i = 1, \dots, n+1,$$

where $d = \sum \frac{1}{a_i^2}$.

Followed by

Theorem 4. *Let $T_{\mathbb{C}}^n$ be the quotient of $\mathbb{T}^{n+1}$ by the $\mathbb{S}^1$ action. Then it is biharmonic in $\mathbb{C}P^n$ if and only if*

$$(3) \quad a_i d - \frac{1}{a_i^3} = 2(n+3) \left((n+1)a_i - \frac{1}{a_i} \right).$$

The explicit expressions of r , s and a_i will be solved out in section 3.

Among these tori, we can pick out plenty proper biharmonic ones. As this is a routine construction, we will emphasise on the method rather than the concrete examples.

2. Preliminary

2.1. Hopf fibration

There are already many known examples of biharmonic submanifolds in spheres. To find examples in $\mathbb{C}P^n$, the Hopf fibration $\pi : \mathbb{S}^{2n+1} \xrightarrow{\mathbb{S}^1} \mathbb{C}P^n$ is a natural candidate, where π is a Riemannian submersion with totally geodesic fibres $\mathbb{S}^1$ and $\mathbb{C}P^n$ has constant holomorphic sectional curvature 4.

This submersion has lots of good properties, such as

Lemma 5 ([9]). *If $\tilde{M}$, M are submanifolds with respect to this Riemannian submersion, i.e. the following diagram commutes*

$$\begin{array}{ccc} \tilde{M} & \longrightarrow & \mathbb{S}^{2n+1} \\ \pi \downarrow & & \downarrow \pi \\ M & \longrightarrow & \mathbb{C}P^n \end{array}$$

then $\tilde{M}$ minimal (totally geodesic) is equivalent to M minimal (totally geodesic). More precisely, the mean curvature vector field $\tilde{H}$ of $\tilde{M}$ is the horizontal lift of H of M .

In this paper, to simplify the notations, we make the convention that H is non-normalized, i.e. $H = \tau(i)$.

Unfortunately, this submersion does not preserve the biharmonicity. See [6], consider the Hopf map $\mathbb{S}^3 \rightarrow \mathbb{S}^2(\frac{1}{2})$. Lifting the biharmonic submanifold

$\mathbb{S}^1(\frac{\sqrt{2}}{4})$, we get the Clifford torus $\mathbb{S}^1(\frac{\sqrt{2+\sqrt{2}}}{2}) \times \mathbb{S}^1(\frac{\sqrt{2-\sqrt{2}}}{2})$, which is not biharmonic in $\mathbb{S}^3$.

But it provides us the stereotype of constructing biharmonic submanifolds in $\mathbb{C}P^n$. If we modify the radii of the standard Clifford torus properly, its image will be biharmonic in $\mathbb{C}P^n$.

2.2. The biharmonic equations in $\mathbb{S}^n$ and $\mathbb{C}P^n$

When the submanifold lies in $\mathbb{S}^n$, it is convenient to split the bitension field in its normal and tangent components.

Theorem 6 ([2]). *If M^m is a submanifold of $\mathbb{S}^n$, then it is biharmonic if and only if*

$$(4) \quad \begin{cases} -\Delta^\perp H - \text{trace } B(\cdot, A_H \cdot) + mH = 0 \\ 2 \text{trace } A_{\nabla_{(\cdot)}^\perp H}(\cdot) + \frac{1}{2} \text{grad}(|H|^2) = 0 \end{cases}.$$

Moreover, if we assume M has parallel mean curvature vector field, the equation turns to

$$(5) \quad \text{trace } B(\cdot, A_H \cdot) = \sum_{i,j=1}^m \langle B_{ij}, H \rangle B_{ij} = mH,$$

where $B_{ij} = B(e_i, e_j)$ and $\{e_i\}_{i=1}^m$ is an orthogonal basis on M ; $\nabla^\perp$ and $\Delta^\perp$ are the connection and the Laplacian in the normal bundle, respectively.

When the ambient space is $\mathbb{C}P^n$, considering the real hypersurfaces or the Lagrangian submanifolds, the biharmonic equation has a similar form

Proposition 7. *If M^{2n-1} is a real hypersurface in $\mathbb{C}P^n$, then it is biharmonic if and only if*

$$(6) \quad \begin{cases} -\Delta^\perp H - \text{trace } B(\cdot, A_H \cdot) + 2(n+1)H = 0 \\ 2 \text{trace } A_{\nabla_{(\cdot)}^\perp H}(\cdot) + \frac{1}{2} \text{grad}(|H|^2) = 0 \end{cases}.$$

If, in addition, M has constant mean curvature the biharmonic equation becomes

$$(7) \quad |B|^2 = 2(n+1).$$

And

Proposition 8. *If M^n is a Lagrangian submanifold of $\mathbb{C}P^n$, then it is biharmonic if and only if*

$$(8) \quad \begin{cases} -\Delta^\perp H - \text{trace } B(\cdot, A_H \cdot) + (n+3)H = 0 \\ 2 \text{ trace } A_{\nabla_{(\cdot)}^\perp H}(\cdot) + \frac{1}{2} \text{grad}(|H|^2) = 0 \end{cases}.$$

When M has parallel mean curvature vector field, the biharmonic equation becomes

$$(9) \quad \text{trace } B(\cdot, A_H \cdot) = \sum_{i,j} \langle B_{ij}, H \rangle B_{ij} = (n+3)H.$$

Proof of Proposition 7. Denote the canonical inclusion map as i , then $\tau(i) = H$.

Since $\text{trace } R^{\mathbb{C}P^n}(di, \tau(i))di = -2(n+1)\tau(i) = -2(n+1)H$, the inclusion map i is biharmonic if and only if

$$\tau_2(i) = \text{trace } \nabla dH + 2(n+1)H = 0.$$

For this is a tensor equation, we can choose orthonormal frame $\{e_i\}$ around $p \in M$, where $1 \leq i \leq 2n-1$, s.t. $\nabla_{e_i} e_j|_p = 0$ and do all the computation restricted to the point p

$$\begin{aligned} \text{trace } \nabla dH &= \nabla_{e_i}^{\mathbb{C}P^n} \nabla_{e_i}^{\mathbb{C}P^n} H = \nabla_{e_i}^{\mathbb{C}P^n} (\nabla_{e_i}^\perp H - A_H(e_i)) \\ &= \nabla_{e_i}^\perp \nabla_{e_i}^\perp H - A_{\nabla_{e_i}^\perp H}(e_i) - \nabla_{e_i} A_H(e_i) - B(e_i, A_H(e_i)) \\ &= -\Delta^\perp H - \text{trace } B(\cdot, A_H \cdot) - (A_{\nabla_{e_i}^\perp H}(e_i) + \nabla_{e_i} A_H(e_i)). \end{aligned}$$

Then, rewrite A_H and use Codazzi equation

$$\begin{aligned} \nabla_{e_i} A_H(e_i) &= (\nabla_{e_i} \langle B(e_j, e_i), H \rangle) e_j \\ &= \langle B(e_j, e_i), \nabla_{e_i}^\perp H \rangle e_j + \langle (\nabla_{e_i}^\perp B)(e_j, e_i), H \rangle e_j \\ &= A_{\nabla_{e_i}^\perp H}(e_i) + \langle (\nabla_{e_j}^\perp B)(e_i, e_i), H \rangle e_j \\ &\quad + \langle R^{\mathbb{C}P^n}(e_i, e_j)e_i, H \rangle e_j \\ &= A_{\nabla_{e_i}^\perp H}(e_i) + \frac{1}{2} \text{grad}(|H|^2), \end{aligned}$$

where we use the fact that $\mathbb{C}P^n$ is Einstein to deduce that

$$\langle R^{\mathbb{C}P^n}(e_i, e_j)e_i, H \rangle = -\text{Ricci}^{\mathbb{C}P^n}(e_j, H) = -\langle e_j, H \rangle_{\mathbb{C}P^n} = 0$$

for any j .

Replacing trace ∇dH in the identity and arranging the terms in tangent and normal components, we get what desired. $\square$

The proof of Proposition 8 is similar.

Our general modus operandi is to adjust the radii of the Clifford torus until its second fundamental form satisfies the biharmonic equation. The fact that the Clifford tori we concern about all have parallel mean curvature vector field simplifies the computation.

3. The examples

3.1. Generalized equators $M_{p,q}^{\mathbb{C}}(r, s)$ in $\mathbb{C}P^n$

In [5], LAWSON introduced the concept of generalized equator $M_{p,q}^{\mathbb{C}}$ which is minimal in $\mathbb{C}P^n$. Thus we call $M_{p,q}^{\mathbb{C}}(r, s)$ the generalized equator in the biharmonic category, which is not always minimal.

Since the generalized Clifford torus in $\mathbb{S}^{2n+1}$ has constant mean curvature, so does $M_{p,q}^{\mathbb{C}}(r, s)$.

By (7), $M_{p,q}^{\mathbb{C}}(r, s)$ is biharmonic if $|B|^2 = 2(n+1)$, while from [5], we know $|B|^2 = |\tilde{B}|^2 - 2$, where $\tilde{B}$ is the second fundamental form of the generalized Clifford torus in $\mathbb{S}^{2n+1}$. Combining with the fact $|\tilde{B}|^2 = (\frac{s}{r})^2(2p+1) + (\frac{r}{s})^2(2q+1)$, we get (1). It is easy to check the solution is never minimal.

This is a table of biharmonic real hypersurfaces in $\mathbb{C}P^n$ (We only list the case $n = 5$).

n	p	q	r	s
5	0	4	$\sqrt{(3 * (5 - \sqrt{3}))/22}$	$\sqrt{(7 + (3 * \sqrt{3}))/22}$
5	0	4	$\sqrt{(3 * (5 + \sqrt{3}))/22}$	$\sqrt{(7 - (3 * \sqrt{3}))/22}$
5	1	3	$\sqrt{(13 - \sqrt{15})/22}$	$\sqrt{(9 + \sqrt{15})/22}$
5	1	3	$\sqrt{(13 + \sqrt{15})/22}$	$\sqrt{(9 - \sqrt{15})/22}$
5	2	2	$\sqrt{(11 + \sqrt{11})/22}$	$\sqrt{(11 - \sqrt{11})/22}$

3.2. Clifford torus $\mathbb{T}^{n+1}$ in $\mathbb{S}^{2n+1}$

Decompose $\mathbb{C}^{n+1}$ into $V_1 \oplus \cdots \oplus V_{n+1}$, where each V_i is the standard one dimensional complex subspace. Identify V_i with $\mathbb{R}^2$, let x_i be the unit vector s.t. $|x_i| = 1$, i.e. all the unit vectors constitute $\mathbb{S}^1 \subset \mathbb{R}^2$. Denote the position vector of $\mathbb{T}^{n+1}$ by $x = a_i x_i$, where a_i are non-negative real number, then $\mathbb{T}^{n+1}$ is included in $\mathbb{S}^{2n+1}$ if and only if $\sum_{i=1}^{n+1} a_i^2 = 1$. For $\mathbb{T}^{n+1}$ is a product manifold, restrict to $\mathbb{S}^1(a_i)$, we can choose frame e_i of the tangent space s.t. $Je_i = x_i$ where J is the complex structure. Hence $\{e_i\}_{i=1}^{n+1}$ is a normal frame s.t. $Je_i = x_i$ where J is the complex structure of $\mathbb{C}^{n+1}$. Direct computation shows that $B_{ij} = \delta_{ij}(-\frac{x_i}{a_i} + x)$ and $H = \sum(a_i(n+1) - \frac{1}{a_i})x_i$.

Thus

$$\sum_{i,j=1}^{n+1} \langle B_{ij}, H \rangle B_{ij} = \left(\sum (a_i d - \frac{1}{a_i^3}) x_i \right) - (n+1)H,$$

where $d = \sum_{i=1}^{n+1} \frac{1}{a_i^2}$.

Furthermore, $\mathbb{T}^{n+1}$ has parallel mean curvature vector field. If it is biharmonic, by (5)

$$\left(\sum (a_i d - \frac{1}{a_i^3}) x_i \right) - (n+1)H = (n+1)H.$$

By comparing each component of H , we have (2).

The equation system (2) is some how funny. Denote $b_i = a_i^2$, rewrite it as

$$(10) \quad (2(n+1)^2 - d)b_i^2 - 2(n+1)b_i + 1 = 0,$$

where $d = \sum_{i=1}^{n+1} \frac{1}{b_i}$.

There are $n+2$ equations and $n+1$ variables, while the equations are not independent. It is always solvable. To see this, view d as a known number. Let r, s be the two different roots of the quadratic (10), and p, q be their multiplicities in b_i respectively. Then $pr + qs = 1$, and

$$d = \frac{p}{r} + \frac{q}{s} = \frac{p}{r} + \frac{p}{s} + \frac{q-p}{s} = p \frac{r+s}{rs} + \frac{q-p}{\frac{1-p(r+s)}{q-p}}.$$

Let $t = 2(n+1)^2 - d$, use Vieta's Theorem

$$(2q(n+1) - t)(t - 2p(n+1)) = (q-p)^2 t,$$

i.e.

$$t^2 - (2(n+1)^2 - (p-q)^2)t + 4pq(n+1)^2 = 0.$$

This equation's discriminant is

$$\begin{aligned}\Delta &= (p-q)^4 + 4(n+1)^4 - (4(p-q)^2(n+1)^2 + 16pq(n+1)^2) \\ &= (p-q)^4 + 4(n+1)^4 - 4(p+q)^2(n+1)^2 = (p-q)^4.\end{aligned}$$

The root $t = (n+1)^2$ leads to minimal torus, so we pick the one $t = (n+1)^2 - (p-q)^2$. The original quadratic becomes

$$((n+1)^2 - (p-q)^2)b_i^2 - 2(n+1)b_i + 1 = 0.$$

Its two roots are $\frac{1}{(n+1) \pm (p-q)}$ with multiplicities p and q .

Thus the torus is with elegant form $T_{p,q}^{n+1}(\frac{1}{\sqrt{2p}}, \frac{1}{\sqrt{2q}})$. It can not be minimal unless $p = q$.

3.3. Clifford torus $T_{\mathbb{C}}^n$ in $\mathbb{C}P^n$

Denote by $T_{\mathbb{C}}^n$ the quotient of $\mathbb{T}^{n+1}$ by the $\mathbb{S}^1$ action. It is Lagrangian in $\mathbb{C}P^n$, and has parallel mean curvature vector field.

Choose an orthogonal frame $\{e_i\}_{i=1}^n$ on $T_{\mathbb{C}}^n$, $i = 1, \dots, n$. To compute $\sum_{i,j} \langle B_{ij}, H \rangle B_{ij}$, as the Hopf map preserve the horizontal part of second fundamental form([9]) and the mean curvature vector field, we lift it to $\mathbb{S}^{2n+1}$. Denote by f_i the horizontal lift of e_i , when $i \leq n$, and $f_{n+1} = \nu$, where $\nu = Jx$, x the position vector, is the vertical vector field. Then $\{f_\alpha\}_{\alpha=1}^{n+1}$ is an orthogonal frame of $\mathbb{T}^{n+1}$.

$$\begin{aligned}\sum_{\alpha,\beta} \langle \tilde{B}_{\alpha\beta}, \tilde{H} \rangle \tilde{B}_{\alpha\beta} - 2 \sum_i \langle \tilde{B}_{i\nu}, \tilde{H} \rangle \tilde{B}_{i\nu} - \langle \tilde{B}_{\nu\nu}, \tilde{H} \rangle \tilde{B}_{\nu\nu} \\ = \sum_{i,j} \langle B_{ij}, H \rangle \tilde{B}_{ij}.\end{aligned}$$

It is not hard to show

$$\sum_i \langle \tilde{B}_{i\nu}, \tilde{H} \rangle \tilde{B}_{i\nu} = \sum_i -Jf_i \langle -Jf_i, \tilde{H} \rangle = \tilde{H},$$

and $\tilde{B}_{\nu\nu} = 0$.

Follow the biharmonic equation (9)

$$\sum_{\alpha,\beta} \langle \tilde{B}_{\alpha\beta}, \tilde{H} \rangle \tilde{B}_{\alpha\beta} = (n+5)\tilde{H}.$$

This is nothing but (3). Notice that $\sum_{\alpha,\beta} \langle \tilde{B}_{\alpha\beta}, \tilde{H} \rangle \tilde{B}_{\alpha\beta}$ is independent of the choice of basis, thus we solve it in the same way as in Section 3.2.

$$d = \frac{p}{r} + \frac{p}{s} + \frac{q-p}{s} = p \frac{r+s}{rs} + \frac{q-p}{\frac{1-p(r+s)}{q-p}}.$$

Let $t = 2(n+1)(n+3) - d$, we have

$$(11) \quad t^2 - (2(n+1)(n+3) - (p-q)^2)t + 4pq(n+3)^2 = 0.$$

Its discriminant is

$$\Delta = (p-q)^4 + 8(n+3)(p-q)^2.$$

Both roots of t make the original quadratic about r and s solvable. As in the generalized equators $M_{p,q}^{\mathbb{C}}(r, s)$ case, the biharmonic torus $T_{\mathbb{C}_{p,q}}^n(r, s)$ could never be minimal.

The software “Mathematica” is helpful to get the explicit expressions. The following is the table of $T_{\mathbb{C}_{p,q}}^n(r, s)$ when $n = 4$

n	p	q	r	s
4	1	4	$\frac{1}{2}\sqrt{(11 - \sqrt{65})/7}$	$\frac{1}{4}\sqrt{(17 + \sqrt{65})/7}$
4	1	4	$\frac{1}{2}\sqrt{(11 + \sqrt{65})/7}$	$\frac{1}{4}\sqrt{(17 - \sqrt{65})/7}$
4	2	3	$\frac{1}{2}\sqrt{(13 - \sqrt{57})/14}$	$\frac{1}{2}\sqrt{(15 + \sqrt{57})/21}$
4	2	3	$\frac{1}{2}\sqrt{(13 + \sqrt{57})/14}$	$\frac{1}{2}\sqrt{(15 - \sqrt{57})/21}$

4. Stability

We add a few remarks on the biharmonic stability of such tori.

A biharmonic submanifold is called stable if for any variation of the inclusion map $\{i_t\}$, its second derivative of E_2 is non-negative.

In [4], JIANG had proved that there is no stable proper biharmonic submanifold in sphere, so we only need to check the stabilities of $M_{p,q}^{\mathbb{C}}(r, s)$ and $T_{\mathbb{C}_{p,q}}^n(r, s)$.

We will deal with these two cases simultaneously. Assume that M is a proper biharmonic submanifold in $\mathbb{C}P^n$, being either a hypersurface or a Lagrangian submanifold, with an orthonormal frame $\{e_i\}_{i=1}^m$, where $m = 2n-1$ for the hypersurface and $m = n$ for the Lagrangian submanifold. Using $H = \tau(i)$ as the variation vector fields, by the second variation formula in [4]

$$\begin{aligned} \frac{1}{2} \frac{d^2}{dt^2} E_2(i_t)|_{t=0} &= 4 \int_M \langle R^{\mathbb{C}P^n}(e_k, H) \nabla_{e_k}^{\mathbb{C}P^n} H, H \rangle v_g \\ &= 4 \int_M (-|H|^4 + 3 \langle e_k, JH \rangle \langle \nabla_{e_k}^{\mathbb{C}P^n} H, JH \rangle) v_g. \end{aligned}$$

Notice that $JH = \langle JH, e_l \rangle e_l$ holds both in the hypersurface case and the Lagrangian submanifold case, and

$$\langle \nabla_{e_k}^{\mathbb{C}P^n} H, e_l \rangle = - \langle \nabla_{e_k}^{\mathbb{C}P^n} e_l, H \rangle = - \langle B_{kl}, H \rangle,$$

the above expression becomes

$$(12) \quad -4 \int_M (|H|^4 + 3 \langle J e_k, H \rangle \langle B_{kl}, H \rangle \langle J e_l, H \rangle) v_g.$$

Still lift them to $\mathbb{S}^{2n+1} \subset \mathbb{C}^{n+1}$ for computation. Let $\{f_\alpha\}_{\alpha=1}^{m+1}$ be the orthogonal frame on $\tilde{M}$ s.t. f_i is the horizontal lift of e_i and $f_{m+1} = \nu$, then

$$\langle J e_k, H \rangle \langle B_{kl}, H \rangle \langle J e_l, H \rangle = \langle J f_k, \tilde{H} \rangle \langle \tilde{B}_{kl}, \tilde{H} \rangle \langle J f_l, \tilde{H} \rangle,$$

and $\langle J \nu, \tilde{H} \rangle = \langle x, \tilde{H} \rangle = 0$. Where x is the position vector in $\mathbb{C}^{n+1}$ and without any ambiguity, J can be the complex structure in $\mathbb{C}P^n$ or $\mathbb{C}^{n+1}$ respectively. Thus we have

$$\begin{aligned} \langle J e_k, H \rangle \langle B_{kl}, H \rangle \langle J e_l, H \rangle &= \langle J f_\alpha, \tilde{H} \rangle \langle \tilde{B}_{\alpha\beta}, \tilde{H} \rangle \langle J f_\beta, \tilde{H} \rangle \\ &= \langle J f_\alpha, \tilde{H} \rangle^2 \langle \tilde{B}_{\alpha\alpha}, \tilde{H} \rangle, \end{aligned}$$

if $\tilde{B}$ can be diagonalized by $\{f_\alpha\}$.

In the $M_{p,q}^{\mathbb{C}}(r, s)$ case, one can compute

$$\begin{aligned} \frac{1}{2} \frac{d^2}{dt^2} E_2(i_t)|_{t=0} &= -4 \int \left[\frac{s}{r}(2p+1) - \frac{r}{s}(2q+1) \right]^4 \\ &+ 3 \left[\frac{s}{r}(2p+1) - \frac{r}{s}(2q+1) \right]^2 \left[\frac{s^4}{r^2}(2p+1) + \frac{r^4}{s^2}(2q+1) - 2n \right] v_g \end{aligned}$$

Combining Eq.(1), after tedious computation, we get

$$\frac{1}{2} \frac{d^2}{dt^2} E_2(i_t)|_{t=0} < 0.$$

In the $T_{\mathbb{C}_{p,q}}^n(r, s)$ case, every term in Eq.(12) can also be computed explicitly,

$$\frac{1}{2} \frac{d^2}{dt^2} E_2(i_t)|_{t=0} = -4 \int [d - (n+1)^2]^2 + 3[2(n+1)^3 + (\sum \frac{1}{a_i^4}) - 3(n+1)d]v_g.$$

Recall (11) and $t = 2(n+1)(n+3) - d$, we can show

$$\frac{1}{2} \frac{d^2}{dt^2} E_2(i_t)|_{t=0} < 0.$$

In conclusion, all the examples appearing in this paper are unstable.

At the same time, by (12) the expression of second variation, we easily get

Proposition 9. *Let M be a compact proper biharmonic submanifold of $\mathbb{C}P^n$. If the eigenvalues of its Weingarten operator are all nonnegative, then it cannot be stable.*

Although it can't apply to our examples.

All the computations in above sections are valid in $\mathbb{Q}P^n$ and $\mathbb{S}^{4n+3}$, so the same method can generate lots of examples of biharmonic submanifolds in $\mathbb{Q}P^n$ as well.

Finally, the author wants to thank his advisor Prof. Yuxin Dong for the instructions, Prof. Qingchun Ji and Prof. Xiaoli Chao for valuable conversations, and the referee for kind comments.

REFERENCES

1. BALMUŞ, A.; MONTALDO, S.; ONICIUC, C. – *Classification results for biharmonic submanifolds in spheres*, Israel J. Math., 168 (2008), 201–220.
2. CADDEO, R.; MONTALDO, S.; ONICIUC, C. – *Biharmonic submanifolds of S^3* , Internat. J. Math., 12 (2001), 867–876.

3. JIANG, G.Y. – *2-harmonic isometric immersions between Riemannian manifolds*, Chinese Ann. Math. Ser. A, 7 (1986), 130-144.
4. JIANG, G.Y. – *2-harmonic maps and their first and second variational formulas*, Chinese Ann. Math. Ser. A, 7 (1986), 389-402.
5. LAWSON, H.B. – *Rigidity theorems in rank-1 symmetric spaces*, J. Differential Geometry, 4 (1970), 349-357.
6. LOUBEAU, E.; ONICIUC, C. – *On the biharmonic and harmonic indices of the Hopf map*, Trans. Amer. Math. Soc., 359 (2007), 5239-5256.
7. MONTALDO, S.; ONICIUC, C. – *A short survey on biharmonic maps between Riemannian manifolds*, Rev. Un. Mat. Argentina, 47 (2006), 1-22.
8. OU, YE-LIN; TANG, LIANG – *The generalized Chen's conjecture on biharmonic submanifolds is false*, preprint, arXiv:math.DG/1006.1838v1.
9. WU, CHUANXI; LI, GUANGHAN – *Geometry of Submanifolds*, (Chinese) Science Press, Beijing, 2002.
10. LOUBEAU, E.; MONTALDO, S.; ONICIUC, C. (Editors) – *The Bibliography of Biharmonic Maps*, <http://people.unica.it/biharmonic/>.

Received: 19.VI.2010

Department of Mathematics,
University of Science and Technology of China,
Hefei, 230026,
P.R. CHINA
zhangw81@ustc.edu.cn