REFINEMENT OF HARDY'S INEQUALITIES INVOLVING MANY FUNCTIONS VIA SUPERQUADRATIC FUNCTIONS

BY

J.A. OGUNTUASE and B.A. POPOOLA

Abstract. Some new refined Hardy type integral inequalities involving n functions ($n \in \mathbb{Z}_+$) via superquadratic functions are established for $p \geq 2$ and their dual inequalities are also derived. In particular, the results obtained complement and improve some recent results of OGUNTUASE and PERSSON.

Mathematics Subject Classification 2000: 26D15, 26A51.

Key words: Hardy's inequality, refinement of Hardy's inequalities, dual inequalities, superquadratic functions, subquadratic functions.

1. Introduction

In 1920, HARDY [3] announced and proved in [4] the following integral inequality (see also [5, Theorem 330, p. 245])

$$(1.1) \quad \int_0^\infty \left(\frac{1}{x} \int_0^x f(t) dt \right)^p dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty f^p(x) dx, \quad p > 1,$$

where f is a nonnegative measurable function and the constant $(\frac{p}{p-1})^p$ is the best possible. The prehistory of (1.1) up to the time when Hardy finally proved (1.1) in 1925 can be found in [5], [6] and [7]. Concerning the history and development of inequality (1.1) we refer interested reader to the books [7], [8] and [10] devoted to this subject and also the recent historical article [6] and the references given therein.

A well-known simple fact is that (1.1) can equivalently (via the substitution $f(x) = h(x^{(1-\frac{1}{p})}x^{-\frac{1}{p}})$, be rewritten in the form

$$(1.2) \quad \int_0^\infty \left(\frac{1}{x} \int_0^x h(t) dt \right)^p \frac{dx}{x} \leq \int_0^\infty h^p(x) \frac{dx}{x}$$

and in this form it even holds with equality when $p = 1$. In this form we see that Hardy's inequality is a simple consequence of Jensen's inequality but this was not discovered in the dramatic period when Hardy discovered and finally proved this inequality in [4].

In a recent paper, OGUNTUASE and PERSSON [9] using mainly the notion of superquadratic and subquadratic functions proved for $p \geq 2$ the refined form of Hardy's inequality (1.1), namely

$$\begin{aligned}
 & \int_0^b x^{-m} \left(\int_0^x f(t) dt \right)^p dx \\
 & + \frac{m-1}{p} \int_0^b \int_t^b \left| \frac{p}{m-1} \left(\frac{t}{x} \right)^{1-\frac{m-1}{p}} f(t) - \frac{1}{x} \int_0^x f(t) dt \right|^p \\
 (1.3) \quad & \times x^{p-m-\frac{m-1}{p}} dx t^{\frac{m-1}{p}-1} dt \\
 & \leq \left(\frac{p}{m-1} \right)^p \int_0^b \left[1 - \left(\frac{x}{b} \right)^{\frac{m-1}{p}} \right] x^{p-m} f^p(x) dx
 \end{aligned}$$

for $m > 1$, $0 < b \leq \infty$, and

$$\begin{aligned}
 & \int_b^\infty x^{-m} \left(\int_x^\infty f(t) dt \right)^p dx \\
 & + \frac{1-m}{p} \int_b^\infty \int_b^t \left| \frac{p}{1-m} \left(\frac{t}{x} \right)^{\frac{1-m}{p}+1} f(t) - \frac{1}{x} \int_x^\infty f(t) dt \right|^p \\
 (1.4) \quad & \times x^{\frac{1-m}{p}+p-m} dx t^{\frac{m-1}{p}-1} dt \\
 & \leq \left(\frac{p}{1-m} \right)^p \int_b^\infty \left[1 - \left(\frac{b}{x} \right)^{\frac{1-m}{p}} \right] x^{p-m} f^p(x) dx
 \end{aligned}$$

for $m < 1$, $0 \leq b < \infty$.

Furthermore, it was shown in [9] that (1.3)-(1.4) hold in the reversed direction if $1 < p \leq 2$. In particular, this means that for $p = 2$ we have equality in (1.3)-(1.4), respectively.

In this paper we generalize and unify some of these results by proving some new refined Hardy type integral inequalities involving n functions ($n \in \mathbb{Z}_+$) via superquadratic functions. Moreover the corresponding dual inequalities are also derived. In particular, our results further complement and improve some recent results in [9].

Our main tool in the proofs is to use the notion of superquadratic functions introduced by ABRAMOVICH, JAMESON and SINNAMON in [1] (see also [2]).

Definition 1.1 ([1, Definition 2.1]). A function $\varphi : [0, \infty) \rightarrow \mathbb{R}$ is superquadratic provided that for all $x \geq 0$ there exists a constant $C_x \in \mathbb{R}$ such that

$$\varphi(y) - \varphi(x) - \varphi(|y - x|) \geq C_x (y - x)$$

for all $y \geq 0$. We say that f is subquadratic if $-f$ is superquadratic.

The paper is organized as follows: In Section 2 we present some preliminaries, including some inequalities involving superquadratic and subquadratic functions involving n functions ($n \in \mathbb{Z}_+$) of independent interest. In Section 3 we present some new refined Hardy type inequalities involving n functions and their proofs. Our final Section 4 is devoted to some concluding remarks and examples.

Notations. Throughout this paper, all functions are assumed to be measurable and expressions of the form $0 \cdot \infty$, $\frac{\infty}{\infty}$, and $\frac{0}{0}$ are taken to be equal to zero. In addition, by a weight function u we mean a nonnegative measurable function on the actual interval.

2. Preliminaries

First, we state the following refinement of Jensen's inequality in [1], which is very useful in the proofs of our results.

Lemma 2.1 ([1, Theorem 2.3]). *Let (Ω, μ) be a probability measure space. The inequality*

$$(2.1) \quad \begin{aligned} \varphi \left(\int_{\Omega} f(s) d\mu(s) \right) &\leq \int_{\Omega} \varphi(f(s)) d\mu(s) \\ &- \int_{\Omega} \varphi \left(\left| f(s) - \int_{\Omega} f(s) d\mu(s) \right| \right) d\mu(s) \end{aligned}$$

holds for all probability measures μ and all nonnegative μ -integrable functions f if and only if φ is superquadratic. Moreover, (2.1) holds in the reversed direction if and only if φ is subquadratic.

Proposition 2.1. *Let $f_1, f_2, \dots, f_n$ be nonnegative integrable functions and define*

$$(2.2) \quad F_k(x) := \frac{1}{x} \int_0^x f_k(t) dt, \quad k = 1, 2, \dots, n.$$

Suppose that $0 < b \leq \infty$, let $u : (0, \infty) \rightarrow \mathbb{R}$ be a nonnegative weight function such that the function $x \rightarrow \frac{u(x)}{x^2}$ is locally integrable on $(0, \infty)$, and define the weight function v by

$$v(t) = t \int_t^b \frac{u(x)}{x^2} dx, \quad t \in (0, b).$$

If the real-valued function φ is superquadratic and nondecreasing on (a, c) , $0 \leq a < c \leq \infty$, then

$$(2.3) \quad \begin{aligned} & \int_0^b u(x) \varphi \left(\left[\prod_{k=1}^n F_k(x) \right]^{\frac{1}{n}} \right) \frac{dx}{x} \\ & + \int_0^b \int_t^b \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right| \right) \frac{u(x)}{x^2} dx dt \\ & \leq \int_0^b v(x) \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(x) \right) \frac{dx}{x} \end{aligned}$$

holds for all f_k with $a < f_k(x) < c$, $0 < x \leq b$, $k \in \{1, 2, \dots, n\}$.

Proof. First we apply the Arithmetic-Geometric Mean inequality

$$(2.4) \quad \left(\prod_{k=1}^n F_k(x) \right)^{\frac{1}{n}} \leq \frac{1}{n} \sum_{k=1}^n F_k(x),$$

to the first term on the left hand side of inequality (2.3) and thereafter, by using Lemma 2.1 and Fubini's theorem we have that

$$\int_0^b u(x) \varphi \left(\left[\prod_{k=1}^n F_k(x) \right]^{\frac{1}{n}} \right) \frac{dx}{x} \leq \int_0^b u(x) \varphi \left(\frac{1}{n} \sum_{k=1}^n F_k(x) \right) \frac{dx}{x}$$

$$\begin{aligned}
&= \int_0^b u(x) \varphi \left(\frac{1}{x} \int_0^x \left(\frac{1}{n} \sum_{k=1}^n f_k(t) \right) dt \right) \frac{dx}{x} \\
&\leq \int_0^b \frac{u(x)}{x^2} \int_0^x \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(t) \right) dt dx \\
&\quad - \int_0^b \frac{u(x)}{x^2} \int_0^x \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right| \right) dt dx \\
&= \int_0^b \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(t) \right) \int_t^b \frac{u(x)}{x^2} dx dt \\
&\quad - \int_0^b \int_t^b \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right| \right) \frac{u(x)}{x^2} dx dt \\
&= \int_0^b v(t) \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(t) \right) \frac{dt}{t} \\
&\quad - \int_0^b \int_t^b \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right| \right) \frac{u(x)}{x^2} dx dt.
\end{aligned}$$

from which (2.3) follows and the proof is complete. $\square$

Remark 2.1. For the case $n = 1$ Proposition 2.1 coincides with Proposition 2.1 (a) in [9].

Proposition 2.2. Let $0 \leq b < \infty$, $u : (b, \infty) \rightarrow \mathbb{R}$ be a nonnegative locally integrable function on (b, ∞) , and define the function v by

$$v(t) = \frac{1}{t} \int_b^t u(x) dx, \quad t \in (b, \infty).$$

If the real-valued function φ is superquadratic and nondecreasing on (a, c) , $0 \leq a < c \leq \infty$, then the inequality

$$\int_b^\infty u(x) \varphi \left(\left[\prod_{k=1}^n G_k(x) \right]^{\frac{1}{n}} \right) \frac{dx}{x}$$

$$\begin{aligned}
(2.5) \quad & + \int_b^\infty \int_b^t \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - x \int_x^\infty \frac{1}{n} \sum_{k=1}^n f_k(t) \frac{dt}{t^2} \right| \right) u(x) dx \frac{dt}{t^2} \\
& \leq \int_b^\infty v(x) \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(x) \right) \frac{dx}{x}
\end{aligned}$$

holds for all f_k with $a < f_k(x) < c$, $0 < x \leq b$, $k \in \{1, 2, \dots, n\}$, where

$$G_k(x) := x \int_x^\infty f_k(t) \frac{dt}{t^2}.$$

Proof. The proof is similar to that of Proposition 2.1 so we omit the details. $\square$

Remark 2.2. For the case $n = 1$ Proposition 2.2 coincides with Proposition 2.2 (a) in [9].

3. Refinement of Hardy-type inequalities involving many functions

Our first result reads:

Theorem 3.1. Let $p > 1$, $m > 1$, $0 < b \leq \infty$, $n \in \mathbb{Z}_+$ and let the functions f_k , $k \in \{1, 2, \dots, n\}$, be nonnegative and locally integrable on $(0, b)$ such that

$$0 < \int_0^b x^{p-m} \left(\sum_{k=1}^n f_k(x) \right)^p dx < \infty.$$

(i) If $p \geq 2$, then

$$\begin{aligned}
(3.1) \quad & \int_0^b x^{-m} \left(\prod_{k=1}^n \int_0^x f_k(t) dt \right)^{\frac{p}{n}} dx \\
& + \frac{m-1}{p} \int_0^b \int_t^b \left| \frac{p}{m-1} \sum_{k=1}^n \frac{f_k(t)}{n} \left(\frac{t}{x} \right)^{1-\frac{m-1}{p}} \right. \\
& \left. - \frac{1}{x} \int_0^x \sum_{k=1}^n \frac{f_k(t)}{n} dt \right|^p x^{p-m-\frac{m-1}{p}} dx t^{\frac{m-1}{p}-1} dt \\
& \leq \left(\frac{p}{nm-n} \right)^p \int_0^b \left[1 - \left(\frac{x}{b} \right)^{\frac{m-1}{p}} \right] x^{p-m} \left(\sum_{k=1}^n f_k(x) \right)^p dx.
\end{aligned}$$

(ii) If $1 < p \leq 2$, then inequality (3.1) holds in the reversed direction.

Proof. This follows directly from the proof of Theorem 3.1 in [9] by using the substitution $f(x) = \sum_{k=1}^n \frac{f_k(x)}{n}$ and the arithmetic-geometric inequality (2.3). $\square$

Remark 3.1. In the special case $n = 1$ Theorem 3.1 coincides with Theorem 3.1 in [9]. For the case $p = 2$ and $n = 1$ we even have equality in (3.1).

Next, we formulate the dual version of Theorem 3.1.

Theorem 3.2. Let $p > 1$, $m < 1$, $0 \leq b < \infty$, $n \in \mathbb{Z}_+$ and let the functions f_k , $k \in \{1, 2, \dots, n\}$, be nonnegative and locally integrable on (b, ∞) such that $0 < \int_b^\infty x^{p-m} (\sum_{k=1}^n f_k(x))^p dx < \infty$.

(iii) If $p \geq 2$, then

$$\begin{aligned}
 & \int_b^\infty x^{-m} \left(\prod_{k=1}^n \int_x^\infty f_k(t) dt \right)^{\frac{p}{n}} dx \\
 & + \frac{1-m}{p} \int_b^\infty \int_b^t \left| \frac{p}{1-m} \sum_{k=1}^n \frac{f_k(t)}{n} \left(\frac{t}{x} \right)^{\frac{1-m}{p}+1} \right. \\
 (3.2) \quad & \left. - \frac{1}{x} \int_x^\infty \sum_{k=1}^n \frac{f_k(t)}{n} dt \right|^p \times x^{\frac{1-m}{p}+p-m} dx t^{\frac{m-1}{p}-1} dt \\
 & \leq \left(\frac{p}{n-nm} \right)^p \int_b^\infty \left[1 - \left(\frac{b}{x} \right)^{\frac{1-m}{p}} \right] x^{p-m} \left(\sum_{k=1}^n f_k(x) \right)^p dx.
 \end{aligned}$$

(iv) If $1 < p \leq 2$, then inequality (3.2) holds in the reversed direction.

Proof. This follows directly from the proof of Theorem 3.2 in [9] by using the substitution $f(x) = \sum_{k=1}^n \frac{f_k(x)}{n}$ and the arithmetic-geometric inequality (2.3). $\square$

Remark 3.2. If we set $n = 1$, then Theorem 3.2 reduces to inequality Theorem 3.2 in [9]. For the case $p = 2$ and $n = 1$ we even have equality in (3.2).

4. Concluding remarks and examples

Next, we obtain the following generalizations of our results in Theorems 3.1 and 3.2 respectively.

Theorem 4.1. Let $p > 1, m > 1, 0 < b \leq \infty$, and $n \in \mathbb{Z}_+$. Let $\{\alpha_k\}_{k=1}^n$ be a positive sequence such that $\sum_{k=1}^n \alpha_k = 1$ and $\{f_k\}_{k=1}^n$ be a sequence of integrable functions on $(0, b)$ such that

$$0 < \int_0^b x^{p-m} \left(\sum_{k=1}^n f_k(x) \right)^p dx < \infty.$$

(v) If $p \geq 2$, then

$$\begin{aligned} & \int_0^b x^{(1-m)n\alpha_k-1} \left(\prod_{k=1}^n \int_0^x f_k(t) dt \right)^{p\alpha_k} dx \\ (4.1) \quad & + \left(\left| \frac{m-1}{p} \right| \right)^{p+1-np\alpha_k} \int_0^b \int_t^b \left| \frac{p}{m-1} \sum_{k=1}^n \frac{f_k(t)}{n} \left(\frac{t}{x} \right)^{1-\frac{m-1}{p}} \right. \\ & \left. - \frac{1}{x} \int_0^x \sum_{k=1}^n \frac{f_k(t)}{n} dt \right|^p x^{p-m-\frac{m-1}{p}} dx t^{\frac{m-1}{p}-1} dt \\ & \leq \alpha_k^p \left(\left| \frac{p}{m-1} \right| \right)^{np\alpha_k} \int_0^b \left[1 - \left(\frac{x}{b} \right)^{\frac{m-1}{p}} \right] x^{p-m} \left(\sum_{k=1}^n f_k(x) \right)^p dx. \end{aligned}$$

(vi) If $1 < p \leq 2$, then inequality (4.1) holds in the reversed direction.

Proof. Let $p \geq 2$. Then, by the more general Arithmetic-Geometric Mean inequality

$$\prod_{k=1}^n g_k^{\alpha_k}(x) \leq \sum_{k=1}^n \alpha_k g_k(x),$$

we have that

$$(4.2) \quad \left(\prod_{k=1}^n F_k^{\alpha_k}(x) \right)^p \leq \left(\sum_{k=1}^n \alpha_k F_k(x) \right)^p = \left(\frac{1}{x} \int_0^x \sum_{k=1}^n \alpha_k f_k(t) dt \right)^p.$$

The rest of the proof is just formal modification of the proof of Theorem 3.1 by using inequality (4.2) together with the substitution $f(x) = \sum_{k=1}^n \alpha_k f_k(x)$. So we omit the details. $\square$

Remark 4.1. If $\alpha_k = \frac{1}{k}$, $k \in \{1, 2, \dots, n\}$, and $m > 1$, then Theorem 4.1 reduces to Theorem 3.1.

Theorem 4.2. Let $p > 1, m < 1, 0 \leq b < \infty$, and $n \in \mathbb{Z}_+$. Let $\{\alpha_k\}_{k=1}^n$ be a positive sequence such that $\sum_{k=1}^n \alpha_k = 1$ and $\{f_k\}_{k=1}^n$ be a sequence of integrable functions on (b, ∞) such that

$$0 < \int_b^\infty x^{p-m} \left(\sum_{k=1}^n f_k(x) \right)^p dx < \infty.$$

(vii) If $p \geq 2$, then

$$\begin{aligned} & \int_b^\infty x^{(1-m)n\alpha_k-1} \left(\prod_{k=1}^n \int_x^\infty f_k(t) dt \right)^{p\alpha_k} dx \\ (4.3) \quad & + \left(\frac{1-m}{p} \right)^{p+1-np\alpha_k} \int_b^\infty \int_b^t \left| \frac{p}{1-m} \sum_{k=1}^n \frac{f_k(t)}{n} \left(\frac{t}{x} \right)^{\frac{1-m}{p}+1} \right. \\ & \left. - \frac{1}{x} \int_x^\infty \sum_{k=1}^n \frac{f_k(t)}{n} dt \right|^p x^{\frac{1-m}{p}+p-m} dx t^{\frac{m-1}{p}-1} dt \\ & \leq \alpha_k^p \left(\frac{p}{1-m} \right)^{np\alpha_k} \int_b^\infty \left[1 - \left(\frac{x}{b} \right)^{\frac{1-m}{p}} \right] x^{p-m} \left(\sum_{k=1}^n f_k(x) \right)^p dx. \end{aligned}$$

(viii) If $1 < p \leq 2$, then inequality (4.3) holds in the reversed direction.

Proof. The proof of Theorem 4.2 is similar to the proof of Theorems 4.1 and so we omit the details. $\square$

Remark 4.2. If $\alpha_k = \frac{1}{k}$, $k \in \{1, 2, \dots, n\}$, and $m > 1$, then Theorem 4.2 reduces to Theorem 3.2.

Example 4.1. In Proposition 2.1, by putting $u(x) \equiv 1$ we obtain the weight function v to be equal to

$$v(x) = \begin{cases} 1 - \frac{x}{b}, & b < \infty \\ 1, & b = \infty \end{cases}.$$

Hence for $b < \infty$, inequality (2.3) takes the form

$$\int_0^b \varphi \left(\left[\prod_{k=1}^n F_k(x) \right]^{\frac{1}{n}} \right) \frac{dx}{x}$$

$$\begin{aligned}
& + \int_0^b \int_t^b \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right| \right) \frac{dx}{x^2} dt \\
& \leq \int_0^b \left(1 - \frac{x}{b} \right) \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(t) \right) \frac{dx}{x}
\end{aligned}$$

and

$$\begin{aligned}
& \int_0^\infty \varphi \left(\left[\prod_{k=1}^n F_k(x) \right]^{\frac{1}{n}} \right) \frac{dx}{x} \\
& + \int_0^\infty \int_t^\infty \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right| \right) \frac{dx}{x^2} dt \\
& \leq \int_0^\infty \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(t) \right) \frac{dx}{x}
\end{aligned}$$

when $b = \infty$.

Example 4.2. Also by putting $u(x) \equiv 1$ in Proposition 2.2 yields

$$v(x) = \begin{cases} 1 - \frac{b}{x}, & b > 0 \\ 1, & b = 0 \end{cases}$$

and so for the case $b > 0$, inequality (2.5) becomes

$$\begin{aligned}
& \int_b^\infty \varphi \left(\left[\prod_{k=1}^n G_k(x) \right]^{\frac{1}{n}} \right) \frac{dx}{x} \\
& + \int_b^\infty \int_b^t \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - x \int_x^\infty \frac{1}{n} \sum_{k=1}^n f_k(t) \frac{dt}{t^2} \right| \right) dx \frac{dt}{t^2} \\
& \leq \int_b^\infty \left(1 - \frac{b}{x} \right) \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(x) \right) \frac{dx}{x}
\end{aligned}$$

while for $b = 0$ it reads

$$\int_0^\infty \varphi \left(\left[\prod_{k=1}^n G_k(x) \right]^{\frac{1}{n}} \right) \frac{dx}{x}$$

$$\begin{aligned}
& + \int_0^\infty \int_0^t \varphi \left(\left| \frac{1}{n} \sum_{k=1}^n f_k(t) - x \int_x^\infty \frac{1}{n} \sum_{k=1}^n f_k(t) \frac{dt}{t^2} \right| \right) dx \frac{dt}{t^2} \\
& \leq \int_0^\infty \varphi \left(\frac{1}{n} \sum_{k=1}^n f_k(x) \right) \frac{dx}{x}.
\end{aligned}$$

Remark 4.3. For the case $n = 1$ Examples 4.1 and 4.2 coincide with Examples 4.1 and 4.2 in [9].

By using Theorem 3.1 with $m = p$ we obtain:

Example 4.3. Let $0 < b \leq \infty$ and let the function f_k , $k \in \{1, 2, \dots, n\}$ be locally integrable on $(0, b)$ such that $0 < \int_0^b x^{p-m} (\sum_{k=1}^n f_k(x))^p dx < \infty$. If $p \geq 2$, then

$$\begin{aligned}
(4.4) \quad & \int_0^b \left(\frac{1}{x^n} \prod_{k=1}^n \int_0^x f_k(t) dt \right)^{\frac{p}{n}} dx \\
& + \frac{p-1}{p} \int_0^b \int_t^b \left| \frac{p}{p-1} \frac{1}{n} \sum_{k=1}^n f_k(t) \left(\frac{t}{x} \right)^{\frac{1}{p}} \right. \\
& \left. - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right|^p x^{-\frac{p-1}{p}} dx t^{-\frac{1}{p}} dt \\
& \leq \left(\frac{p}{np-n} \right)^p \int_0^b \left[1 - \left(\frac{x}{b} \right)^{\frac{p-1}{p}} \right] \left(\sum_{k=1}^n f_k(x) \right)^p dx.
\end{aligned}$$

Remark 4.4. For $p = 2$ and $b = \infty$ we obtain the following identity for all $f_k \in L_2(0, \infty)$, $k \in \{1, 2, \dots, n\}$:

$$\begin{aligned}
(4.5) \quad & \int_0^\infty \left(\frac{1}{x^n} \prod_{k=1}^n \int_0^x f_k(t) dt \right)^{\frac{2}{n}} dx \\
& + \frac{1}{2} \int_0^\infty \int_t^b \left(2\sqrt{\frac{t}{x}} \frac{1}{n} \sum_{k=1}^n f_k(t) - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right)^2 \frac{dx}{\sqrt{x}} \frac{dt}{\sqrt{t}} \\
& = \left(\frac{2}{n} \right)^2 \int_0^\infty \left(\sum_{k=1}^n f_k(x) \right)^2 dx.
\end{aligned}$$

In particular, if $n = 1$ inequality (4.5) coincides with Remark 4.2 in [9].

Remark 4.5. For the case $p = n$ in Theorem 3.1, inequality (3.1) reads:

$$\begin{aligned}
 & \int_0^b x^{-m} \left(\prod_{k=1}^n \int_0^x f_k(t) dt \right) dx \\
 (4.6) \quad & + \frac{m-1}{n} \int_0^b \int_t^b \left| \frac{1}{m-1} \sum_{k=1}^n f_k(t) \left(\frac{t}{x} \right)^{1-\frac{m-1}{n}} \right. \\
 & \left. - \frac{1}{x} \int_0^x \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right|^n \times x^{n-m-\frac{m-1}{n}} dx t^{\frac{m-1}{n}-1} dt \\
 & \leq \left(\frac{1}{m-1} \right)^n \int_0^b \left[1 - \left(\frac{x}{b} \right)^{\frac{m-1}{n}} \right] x^{n-m} \left(\sum_{k=1}^n f_k(x) \right)^n dx,
 \end{aligned}$$

$m > 1, n = 2, 3, 4, \dots$

Example 4.4. For the case $n = 1$ and $b = 0$ in Theorem 3.2 we obtain that

$$\begin{aligned}
 & \int_0^\infty x^{-m} \left(\int_x^\infty f(t) dt \right)^p dx + \frac{1-m}{p} \int_0^\infty \int_0^t \left| \frac{p}{1-m} f(t) \left(\frac{t}{x} \right)^{\frac{1-m}{p}+1} \right. \\
 (4.7) \quad & \left. - \frac{1}{x} \int_x^\infty f(t) dt \right|^p x^{\frac{1-m}{p}+p-m} dx t^{\frac{m-1}{p}-1} dt \\
 & \leq \left(\frac{p}{1-m} \right)^p \int_0^\infty x^{p-m} f^p(x) dx.
 \end{aligned}$$

Remark 4.6. For the case $n = p$ in Theorem 3.2 the inequality (3.2) reads:

$$\begin{aligned}
 & \int_b^\infty x^{-m} \left(\prod_{k=1}^n \int_x^\infty f_k(t) dt \right) dx \\
 & + \frac{1-m}{n} \int_b^\infty \int_b^t \left| \frac{1}{1-m} \sum_{k=1}^n f_k(t) \left(\frac{t}{x} \right)^{\frac{1-m}{n}+1} - \frac{1}{x} \int_x^\infty \frac{1}{n} \sum_{k=1}^n f_k(t) dt \right|^n \\
 (4.8) \quad & \times x^{\frac{1-m}{n}+n-m} dx t^{\frac{m-1}{n}-1} dt \\
 & \leq \left(\frac{1}{1-m} \right)^n \int_b^\infty \left[1 - \left(\frac{b}{x} \right)^{\frac{1-m}{n}} \right] x^{n-m} \left(\sum_{k=1}^n f_k(x) \right)^n dx,
 \end{aligned}$$

$m < 1, n = 2, 3, 4, \dots$

Acknowledgement. We thank the referee for some good advices, which have improved the final version of this paper.

REFERENCES

1. ABRAMOVICH, S.; JAMESON, G.; SINNAMON, G. – *Refining Jensen's inequality*, Bull. Math. Soc. Sci. Math. Roumanie (N.S.), 47 (2004), 3–14.
2. ABRAMOVICH, S.; JAMESON, G.; SINNAMON, G. – *Inequalities for averages of convex and superquadratic functions*, JIPAM. J. Inequal. Pure Appl. Math., 5 (2004), Article 91, 14 pp.
3. HARDY, G.H. – *Note on a theorem of Hilbert*, Math. Z., 6 (1920), 314–317.
4. HARDY, G.H. – *Notes on some points in the integral calculus, LX, An inequality between integrals*, Messenger of Math., 54 (1925), 150–156.
5. HARDY, G.H.; LITTLEWOOD, J.E.; PÓLYA, G. – *Inequalities*, Reprint of the 1952 edition, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1988.
6. KUFNER, A.; MALIGRANDA, L.; PERSSON, L.-E. – *The prehistory of the Hardy inequality*, Amer. Math. Monthly, 113 (2006), 715–732.
7. KUFNER, A.; MALIGRANDA, L.; PERSSON, L.-E. – *The Hardy Inequality. About its History and Some Related Results*, Vydavatelský Servis, Plzeň, 2007.
8. KUFNER, A.; PERSSON, L.-E. – *Weighted Inequalities of Hardy Type*, World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
9. OGUNTUASE, J.A.; PERSSON, L.-E. – *Refinement of Hardy's inequalities via superquadratic and subquadratic functions*, J. Math. Anal. Appl., 339 (2008), 1305–1312.
10. OPIC, B.; KUFNER, A. – *Hardy-Type Inequalities*, Pitman Research Notes in Mathematics Series, 219, Longman Scientific & Technical, Harlow, 1990.

Received: 4.XI.2009

Revised: 7.IV.2010

Department of Mathematics,
University of Agriculture,
Abeokuta, Ogun State,
NIGERIA
oguntuase@yahoo.com

Department of Mathematics and Statistics,
Federal College of Education,
Osiele, Abeokuta, Ogun State,
NIGERIA
bolajipopoola2002@yahoo.com