ON LINEAR OPERATORS PRESERVING ORTHOGONALITY

BY

I. IONICĂ

Abstract. In this paper we present some relationships among Birkhoff orthogonality and other concepts in convex analysis. In this way we obtain a result of Blanco and Turnšek concerning the linear operators that preserve Birkhoff orthogonality.

Mathematics Subject Classification 2010: 46B20, 46B04, 46C50.

Key words: orthogonality (in Birkhoff sense), directional derivative, linear operator preserving orthogonality.

1. Introduction

Let X be a real linear normed space and let us denote by $\perp$ the orthogonality relation (in Birkhoff sense) on X as:

$$(1) \quad x \perp y \Leftrightarrow \|x + ty\| \leq \|x\|, \text{ for all } t \in \mathbb{R}$$

(see, for example, [1], [4], [9], [11], [12]).

In general, this orthogonality is neither symmetric nor additive, but it is always homogeneous. Because the function $t \rightarrow \|x + ty\|$, $t \in \mathbb{R}$, is convex, the orthogonality can be easily characterized using the directional derivatives of the norm (see [1]).

In 1992, KOLDOBSKY [13] has proved that a linear operator $T : X \rightarrow X$ preserves orthogonality if and only if T is an isometry multiplied by a positive constant. In 2006, BLANCO and TURNŠEK [5], extend this result to the case of the linear operators between two linear normed spaces. Moreover, the linear normed spaces X, Y can be considered over C , the field of the complex numbers.

Obviously, if in Koldobsky's result the operator T is the identical operator on a linear space we get that if the orthogonality generated by a norm implies the orthogonality generated by another norm, then that two norms are proportional. A straight demonstration of this result has been given by SCHÖPF [15] using the norm's derivatives.

In the next paragraph we recall a few concepts and basic results in convex analysis (see, for instance [3], [9], [10]), which will be used in the next sections. In the third section we establish some simple properties which characterize the linear operators preserving the orthogonality. Shortly, using a Schöpf's idea, the result of Blanco and Turnšek can be obtained taking into account the monotonicity property of the function $t \rightarrow \|Tx + tTy\|_1^2 \cdot \|x + ty\|_2^{-2}$, where T is a linear operator between the real linear normed spaces $(X, \|\cdot\|_1), (Y, \|\cdot\|_2)$ and x, y are two fixed elements in X .

2. Preliminaries

We recall some basic properties of directional derivatives (see, for example, [2], [3], [9], [14]). Let X be a real linear normed space and X^* its dual. If we consider a proper function (that is, there exist $\tilde{x} \in X$ such that $f(\tilde{x}) \in \mathbb{R}$) $f : X \rightarrow]-\infty, +\infty]$, then the *directional derivatives* (right and left) of f are defined as:

$$(2) \quad f'_\pm(x; y) = \lim_{t \rightarrow \pm 0} \frac{f(x + ty) - f(x)}{t},$$

where $x \in Df = \{x \in X; f(x) < \infty\}$, called the *domain* of the function f and $y \in X$.

If f is convex, then $f'_\pm(x; y)$ exist for all $x \in Df, y \in X$. We also have the next well known properties:

$$(3) \quad f'_\pm(x; -y) = -f'_\mp(x; y), \quad x \in D(f), \quad y \in X;$$

$$(4) \quad f'_\pm(x; \lambda y) = \lambda f'_\pm(x; y), \quad \lambda \geq 0, x \in D(f), y \in X;$$

$$(5) \quad f'_-(x; y) \leq f'_+(x; y), \quad x \in D(f), \quad y \in X;$$

$$(6) \quad f'_+(x; y_1 + y_2) \leq f'_+(x; y_1) + f'_+(x; y_2), \quad x \in D(f), \quad y_1, y_2 \in X.$$

The function f is said to be (Gâteaux) *differentiable* at x if $y \rightarrow f'_+(x, y)$, $y \in X$, is a linear continuous functional on X . If f is a convex function, then it is differentiable at x whenever f is continuous at x and $f'_-(x, y) = f'_+(x, y)$ for all $y \in X$.

If $f(x) = \frac{1}{2}\|x\|^2$, $x \in X$, we denote its directional derivatives by $n'_\pm(x; y)$. The basic properties of $n'_\pm$ to be frequently used are the following:

$$(7) \quad n'_\pm(x; x) = \|x\|^2, \quad x \in X;$$

$$(8) \quad n'_\pm(x; y) \leq \|x\| \cdot \|y\|, \quad x, y \in X;$$

$$(9) \quad n'_+(x; \alpha x + y) = n'_+(x; y) + \alpha \|x\|^2, \quad x \in X, \alpha \in \mathbb{R}.$$

If $n'_-(x, y) = n'_+(x, y)$ for all $x, y \in X$, then the linear normed space X is said to be *smooth*. Obviously, a linear normed space is smooth if and only if its norm $(\frac{1}{2}\|\cdot\|^2)$ is differentiable on X .

3. Characterizations of linear operators that preserve orthogonality

If we consider the definition (1) of orthogonality relation, we observe that $t = 0$ is a minimum point of the convex function $t \rightarrow \|x + ty\|$, $t \in \mathbb{R}$. Taking into account the characterization of minimum points for convex functions on $\mathbb{R}$ and using the directional derivatives [1], [14], we get that:

$$(10) \quad x \perp y \text{ if and only if } n'_-(x; y) \leq 0 \leq n'_+(x; y).$$

Particularly, it follows:

$$(11) \quad \text{if } n'_+(x; y) = 0 \text{ or } n'_-(x; y) = 0, \text{ then } x \perp y.$$

Taking into account the relations (3) and (9), we can easily obtain from (11) that:

$$(12) \quad x \perp \frac{n'_+(x; y)}{\|x\|^2} x - y \text{ for any } x \in X \setminus \{0\}, y \in Y.$$

Let us denote by $T : X_1 \rightarrow X_2$ a linear operator, where X_1 and X_2 are two real linear normed spaces and by $\perp_1, \perp_2$ the orthogonality generated on X_1 , respectively X_2 .

In the next paragraph we will specify some properties using the directional derivatives of the norms on the spaces X_1, X_2 . We denote these derivatives by $n'_{1\pm}$, respectively $n'_{2\pm}$. These properties will prove the property that preserves the orthogonality, namely:

$$(13) \quad \text{if } x \perp_1 y, \text{ then } Tx \perp_2 Ty, \quad x, y \in X_1.$$

If we have two linearly independent vectors $x, y \in X_1$, let us denote by:

$$(14) \quad F_{x,y}(t) = \|Tx + tTy\|_2^2 \cdot \|x + ty\|_1^{-2}, \quad t \in \mathbb{R}.$$

Theorem 1. *If $T : X_1 \rightarrow X_2$ is a linear operator, then the following properties are equivalent:*

- (i) *T preserves the orthogonality;*
- (ii) *if $n'_{1+}(x; y) = 0$, $x, y \in X_1$, then $n'_{2+}(Tx; Ty) \geq 0$;*
- (iii) *the function $F_{x,y}$ is non-decreasing on $\mathbb{R}$;*
- (iv) *the function $F_{x,y}$ is constant on $\mathbb{R}$;*
- (v) *there exist $k > 0$ such that $\|Tx\|_2 = k\|x\|_1$, for all $x \in X_1$ (that is T is an isometry multiplied by a positive constant).*

Proof. (i)→(ii). Let us consider the linear operator T with property (13). If $n'_{1+}(x; y) = 0$, and taking into account the property (11), we obtain that $x \perp_1 y$. Therefore, $Tx \perp_2 Ty$ and so, using (10) we get $n'_{2+}(Tx; Ty) \geq 0$, that is (ii) is fulfilled.

(ii)→(iii). The function $F_{x,y}$ is right derivable on $\mathbb{R}$ because the norms of the linear normed spaces X_1, X_2 have directional derivatives. More than this, after an usual computation, we obtain:

$$(15) \quad (F_{x,y})'_r(t) = [\|x + ty\|_1^2 \cdot n'_{2+}(Tx + tTy; Ty) - \|Tx + tTy\|_2^2 \cdot n'_{1+}(x + ty; y)] \cdot \|x + ty\|_1^{-2}.$$

On the other hand, using (9), we have:

$$(16) \quad n'_{1+} \left(u; \frac{n'_{1+}(u; v)}{\|u\|^2} u - v \right) = 0, \text{ for all } u \in X_1 \setminus \{0\}, v \in X_2.$$

Thus, by hypothesis (ii) it follows:

$$(17) \quad n'_{2+} \left(Tu; \frac{n'_{1+}(u;v)}{\|u\|^2} Tu - Tv \right) \geq 0,$$

that is,

$$(18) \quad \frac{n'_{1+}(u;v)}{\|u\|^2} \|Tu\|_2^2 - n'_{2+}(Tu;Tv) \geq 0.$$

Now, taking $u = x + ty$, $v = y$ we obtain that $(F_{x,y})'_r(t) \leq 0$, for all $t \in \mathbb{R}$. This means that $F_{x,y}$ is a decreasing function on $\mathbb{R}$ (see, for example, [6], p.21).

(iii)→(iv). We observe that $F_{x,y}(-t) = F_{x,-y}(t)$ and so $t \rightarrow F_{x,y}(-t)$, $t \in \mathbb{R}$ is also a decreasing function. On the other hand, this function is non-decreasing because $t \rightarrow F_{x,y}(t)$, $t \in \mathbb{R}$ is decreasing. Finally $F_{x,y}$ is constant on $\mathbb{R}$, that is (iv) is satisfied.

(iv)→(v). As $\lim_{t \rightarrow \infty} F_{x,y}(t) = \frac{\|Ty\|_2^2}{\|y\|_1^2}$ and $F_{x,y}(0) = \frac{\|Tx\|_2^2}{\|x\|_1^2}$ we get that

$$(19) \quad \frac{\|Ty\|_2}{\|y\|_1} = \frac{\|Tx\|_2}{\|x\|_1}, \text{ for all } x, y \in X_1 \setminus \{0\}$$

and this means (v) is satisfied. The implication (v)→(i) is obvious. Thus, the proof of theorem is complete.

Corollary 1. *Two orthogonalities on the same real linear normed space X are equivalent if and only if the corresponding norms are proportional.*

Proof. We take $X_1 = X_2 = X$ and T the identical operator.

Remark 1. The statement (ii) is obviously equivalent to the following statement:

(ii') $n'_-(x, y) = 0$, $x, y \in X_1$, then $n'_-(Tx, Ty) \leq 0$.

Indeed, if $x \perp y$, then $x \perp -y$.

Remark 2. Let us consider the next special orthogonalities:

$$(20) \quad x \perp^s y \text{ if } n'_+(x; y) = 0 \text{ (called Birkhoff strong orthogonality),}$$

respectively

$$(21) \quad x \perp^w y \text{ if } n'_+(x; y) \geq 0 \text{ (called Birkhoff weak orthogonality).}$$

Obviously, $\perp^s$ is stronger than $\perp$, while $\perp^w$ is weaker than $\perp$. So, taking into account the hypothesis (ii) a linear operator preserves the Birkhoff's orthogonality if and only if satisfies the next weaker condition:

$$(22) \quad Tx \perp_2^w Ty \text{ if } x \perp_1^s y, \quad x, y \in X.$$

Particularly, if $\perp_1$ and $\perp_2$ are two orthogonality relations on the same linear real space X , then $\perp_1 \rightarrow \perp_2$ if and only if $\perp_1^s \rightarrow \perp_2^w$.

Other special orthogonalities of Birkhoff type were defined by CHMIELIŃSKI in [7], [8]. In these papers, he establishes some results concerning mappings preserving orthogonality.

Remark 3. The proof of Theorem 1 can be considered as a new proof of Blanco and Turnšek's result in the real case. Their proof use some special properties of support functionals.

Now, we consider the following three statements:

- (i) a linear operator $T : X_1 \rightarrow X_2$, where X_1, X_2 are two real linear normed spaces, preserves the orthogonality relation if and only if is an isometry multiplied by a positive constant;
- (ii) a linear operator $T : X \rightarrow X$, where X is a real linear normed space, preserves the orthogonality relation if and only if is an isometry multiplied by a positive constant;
- (iii) two orthogonality relations on the same real linear normed space X , generated by two norms on X , are equivalent if and only if these two norms are proportional.

Theorem 2. The statements (i), (ii), (iii) are equivalent.

Proof. Obviously (i) $\rightarrow$ (ii) $\rightarrow$ (iii). Now, we suppose the statement (iii) is true and let us consider an arbitrary linear operator $T : X_1 \rightarrow X_2$ that preserves the orthogonality relation. Exactly as BLANCO and TURNŠEK proved (see [5], Theorem 3.1), the operator must be injective. Indeed, if we suppose by contrary that exists an element $\tilde{x} \in X_1$, $\|\tilde{x}\| = 1$, such that $T\tilde{x} = 0$, then taking an arbitrary element $y \in X_1$, $\|y\| = 1$, by (1), it follows that the elements $\alpha\tilde{x} + y$ and $\tilde{x}$ are not orthogonal, whenever $\alpha > 2$. Now, using (9) and (11) we obtain that the elements $\alpha\tilde{x} + y$ and $\beta(\alpha\tilde{x} + \tilde{y}) + \tilde{x}$ are orthogonal if $\beta = -n'_+(\alpha\tilde{x} + y, \tilde{x})\|\alpha\tilde{x} + y\|^{-2}$, $\alpha > 2$. Moreover, we

have $\beta \neq 0$. Since, by hypothesis, the linear operator T preserves the orthogonality relation, we also have $T(\tilde{\alpha}\tilde{x} + y) \perp_2 (T\tilde{x} + \beta T(\tilde{\alpha}\tilde{x} + y))$, that is $Ty \perp_2 Ty$. Hence, $Ty = 0$, for all $y \in X$, that is T is the null operator, and so, we arrive at a contradiction with the property (13) of T . Therefore, T is necessarily injective. Thus, if we denote

$$\|x\|'_1 = \|Tx\|_2, \quad x \in X_1,$$

we obtain that $\|\cdot\|'_1$ is also a norm on X_1 and $\perp_1$ implies $\perp'_1$. Using (iii), we get that the norms $\|\cdot\|_1$, $\|\cdot\|'_1$ are proportional and this proves that T is a scalar multiple of a linear isometry, meaning (i) is satisfied, so the proof is finished.

REFERENCES

1. ALONSO, J.; MARTINI, H.; WU, S. – *On Birkhoff orthogonality and isosceles orthogonality in normed linear spaces*, Aequationes Math., 83 (2012), 153–189.
2. AMIR, D. – *Characterizations of Inner Product Spaces*, Operator Theory: Advances and Applications, 20, Birkhäuser Verlag, Basel, 1986.
3. BARBU, V.; PRECUPANU, T. – *Convexity and Optimization in Banach Spaces*, Fourth Edition, Springer, 2012.
4. BIRKHOFF, G. – *Orthogonality in linear metric spaces*, Duke Math. J., 1 (1935), 169–172.
5. BLANCO, A.; TURNŠEK, A. – *On maps that preserve orthogonality in normed spaces*, Proc. Roy. Soc. Edinburgh Sect. A, 136 (2006), 709–716.
6. BOURBAKI, N. – *Fonctions d'une variable réelle*, Chap. 1, 2 et 3, Hermann, Paris, 1958.
7. CHMIELIŃSKI, J. – *Linear mappings approximately preserving orthogonality*, J. Math. Anal. Appl., 304 (2005), 158–169.
8. CHMIELIŃSKI, J. – *Remarks on orthogonality preserving mappings in normed spaces and some stability problems*, Banach J. Math. Anal., 1 (2007), 117–124.
9. DAY, M.M. – *Normed Linear Spaces*, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1958.
10. HOLMES, R.B. – *Geometric Functional Analysis and its Applications*, Graduate Texts in Mathematics, No. 24, Springer-Verlag, New York-Heidelberg, 1975.
11. JAMES, R.C. – *Orthogonality in normed linear spaces*, Duke Math. J., 12 (1945), 291–302.

12. JAMES, R.C. – *Orthogonality and linear functionals in normed linear spaces*, Trans. Amer. Math. Soc., 61 (1947), 265–292.
13. KOLDOBSKY, A. – *Operators preserving orthogonality are isometries*, Proc. Roy. Soc. Edinburgh Sect. A, 123 (1993), 835–837.
14. PRECUPANU, T.; IONICĂ, I. – *Heights of a triangle in a linear normed space and Hilbertian norms*, An. Ştiinţ. Univ. "Al.I. Cuza" Iaşi. Mat. (N.S.), 55 (2009), 35–47.
15. SCHÖPF, P. – *Orthogonality and proportional norms*, Anz. Österreich. Akad. Wiss. Math.-Natur. Kl., 133 (1996), 11–16 (1997).

Received: 15.V.2011

*Department of Mathematics,
University "Al.I. Cuza",
11, Bd. Carol I, 700506, Iaşi,
ROMANIA
ionica_iuliana@yahoo.com*