A multi-component reaction approach to the synthesis of potent antidiabetic agents five-membered iminosugars analogues

Agustono Wibowo ^{1,*}, Zurina Shaameri ^{2,3}, Mohd Fazli Mohammat ^{2,3}
and Ahmad Sazali Hamzah ^{2,3*}

¹Faculty Applied of Sciences, Universiti Teknologi MARA (UiTM) Pahang Branch, Jengka Campus,
26400 Bandar Tun Abdul Razak Jengka, Pahang Darul Makmur, Malaysia

²Faculty Applied of Sciences, Universiti Teknologi MARA (UiTM), 40450 Shah Alam, Selangor Darul
Ehsan, Malaysia

³Organic Synthesis Laboratory, Institute of Science, Universiti Teknologi MARA (UiTM)42300
Bandar-Puncak Alam, Selangor Darul Ehsan, Malaysia

(Received August 08, 2020; Revised September 01, 2020; Accepted September 05, 2020)

Abstract: Some novel five-membered iminosugar analogues were efficiently synthesized via the MCRs of three components; amines, aldehydes and oxaloacetate to give 4-hydroxymethylpyrrolidines. Reduction of 3-hydroxy-4-carbomethoxy-3-pyrroliden-2-ones with Pd/C-catalyzed hydrogenation gave the corresponding *cis*-hydrogenated products at C₃/C₄ stereoselectively. Then following reductions of ester and amide functionalities with LiAlH₄ afforded 4-(hydroxymethyl)pyrrolidin-3-ols. Among the synthesized compounds 4-hydrazineyl-3-hydroxymethyl-2-(4-methoxyphenyl)-N-methylpyrrolidine and 3-hydrazineyl-4-hydroxymethyl-1-methylpyrrolidine were found to be the most promising α -glucosidase inhibitor with IC₅₀ values of 1.12 and 1.17 mM, respectively.

Keywords: Iminosugar; pyrrolidine; one-pot reaction; stereoselective hydrogenation; antidiabetic. ©2020 ACG Publications. All right reserved.

1. Introduction

Polyhydroxylated piperidines and pyrrolidines, also known as iminosugars, are important class of compounds with interesting structures and immense biological significance, especially as glycosidase inhibitors,^{1,2} thus making them important targets for organic synthesis. Among them, naturally occurring five-membered iminosugars such as 1,4-dideoxy-1,4-iminoheptitol (iminoheptitol) and 1-butyl-1,4-dideoxy-1,4-iminoarabinitol (iminoarabinitol). The synthetic analogues of iminoheptitol and iminoarabinitol are known to show powerful glycosidase inhibition though stereochemical similarity between these iminosugars and the structure of hexoses or enzyme-bound intermediates.³⁻⁷ As a result, several synthetic routes towards these pyrrolidine-based iminosugars have been reported in the literature.¹

* Corresponding author: E-Mail: agustono@uitm.edu.my and asazali@uitm.edu.my

Most of the synthetic routes to the skeleton of iminosugar analogues began with their carbohydrate analogues by taking advantage of the carbohydrates' many stereocenters. However, non-carbohydrate routes such as ring closing metathesis and consecutive reductive aminations are becoming popular,⁸⁻¹¹ since they exhibit an increase in stereoselectivity and are more efficient in introducing the amine moiety.¹⁰ Recently, some synthetic approaches to these biologically active pyrrolidine-based iminosugars have been reported.^{12,13} However, very few reports are known on the preparation of five-membered iminosugars through multi-component reactions (MCRs). In continuation to our ongoing research projects in our laboratory on pyrrolidine-type compounds,¹⁴⁻¹⁹ we herein report on the synthesis and antidiabetic evaluation of 11 new pyrrolidine-based iminosugar analogues in search of new potential glucosidase inhibitors. In addition to the syntheses, all synthesized iminosugars were evaluated for α -glucosidase inhibitory activity.

2. Experimental

2.1. Chemical Material and Apparatus

High-resolution mass spectra were obtained from Agilent 1290 Infinity LC system coupled to Agilent 6520 Accurate-Mass Q-TOF. The ^1H and ^{13}C NMR spectra were registered in CDCl_3 with Joel Resonance ECZ400S [400 MHz (^1H) and 100 MHz (^{13}C)] using TMS as the internal standard. Analytical TLC was performed on silica gel 60 F₂₅₄, Merck (layer thickness 0.25 mm, Merck) and visualized with UV light and KMnO_4 as the detecting agent. All reagents and starting materials such as sodium diethyl oxaloacetate, amine and aldehyde were purchased from Sigma-Aldrich Co. and Merck Chemical Co.

2.2. General Procedure for the Synthesis of Pyrrolidine Intermediates **2a-e** (one-pot reaction)

A suspension of sodium diethyl oxaloacetate (10 mmol), amine (10 mmol) and aldehyde (10 mmol) dissolved in absolute EtOH (40 mL) were heated at reflux for 0.5-2 h. After cooling, the mixture was added onto ice-water and then acidified with 12M HCl until pH 2. The precipitate was filtered, washed with H_2O and Et_2O for removal of unreacted aldehyde to yield pyrrolidine intermediates (**a-e**).

Ethyl 2-oxo-3-hydroxypyrrole-4-carboxylate (2a): Compound **2a** was previously reported (See Scheme 1)²⁰ White solid; 58%, ^1H -NMR (CDCl_3 , 400 MHz) δ_{H} 4.89 (2H, s, CH_2), 4.36 (2H, q, $J=7.2$ Hz, CH_2), 1.35 (3H, t, $J=7.1$ Hz, CH_3). ^{13}C -NMR (CDCl_3 , 100 MHz), δ_{C} 166.5, 164.2, 151.3, 107.6, 66.1, 62.1, 14.2.

Ethyl 2-oxo-3-hydroxy-1-methylpyrrole-4-carboxylate (2b): Compound **2b** was previously reported (See Scheme 1)²⁰ Yellowish solid; 57%, ^1H -NMR (CDCl_3 , 400 MHz) δ_{H} 4.31 (2H, q, $J=7.2$ Hz, OCH_2), 3.96 (2H, s, CH_2), 3.07 (3H, s, NCH_3), 1.33 (3H, t, $J=7.2$ Hz, CH_3). ^{13}C -NMR (CDCl_3 , 100 MHz), δ_{C} 165.2, 164.1, 157.4, 107.6, 61.2, 48.1, 30.1, 14.3.

Ethyl 2-oxo-3-hydroxy-1,5-dimethylpyrrole-4-carboxylate (2c): Yellowish solid; 68%, ^1H -NMR (CDCl_3 , 400 MHz) δ_{H} 4.28 (1H, m, NCH), 4.22 (2H, m, OCH_2), 2.90 (3H, s, CH_3), 2.09 (1H, m, CH_2 -b), 1.95 (1H, m, CH_2 -a), 1.27 (3H, t, $J=6.8$ Hz, CH_3), 0.48 (3H, d, $J=7.6$ Hz, CH_3). ^{13}C -NMR (CDCl_3 , 100 MHz), δ_{C} 188.4, 163.8, 156.5, 109.7, 60.2, 57.9, 26.4, 20.7, 13.8, 4.9.

Ethyl 2-oxo-5-ethyl-3-hydroxy-1-methylpyrrole-4-carboxylate (2d): Yellowish solid; 53%, ^1H -NMR (CDCl_3 , 400 MHz) δ_{H} 4.28 (2H, q, $J=7.2$ Hz, OCH_2), 4.10 (1H, q, $J=6.4$ Hz, NCH), 2.93 (3H, s, CH_3), 1.37 (3H, d, $J=6.4$ Hz, CH_3), 1.28 (3H, t, $J=7.2$ Hz, CH_3). ^{13}C -NMR (CDCl_3 , 100 MHz), δ_{C} 164.4, 164.0, 156.0, 112.0, 60.2, 54.1, 26.2, 16.6, 13.8.

Ethyl 2-oxo-3-hydroxy-1-methyl-5-(4-methoxyphenyl)pyrrole-4-carboxylate (2e): Compound **2e** was previously reported (See Scheme 1)²⁰ Yellow solid; 58%, ^1H -NMR (CDCl_3 , 400 MHz) δ_{H} 7.08 (2H, d,

$J = 9.6$ Hz, Ar), 6.88 (2H, *d*, $J = 9.6$ Hz, Ar), 4.98 (1H, *s*, CH), 4.18 (2H, *q*, $J = 7.2$ Hz, OCH₂), 3.82 (3H, *s*, OCH₃), 2.81 (3H, *s*, NCH₃), 1.17 (3H, *t*, $J = 7.2$ Hz, CH₃). ¹³C-NMR (CDCl₃, 100 MHz), δ_c 165.3, 163.7, 159.9, 157.8, 128.8, 126.5, 114.3, 112.9, 62.2, 61.0, 55.4, 27.6, 14.0.

2.3. Synthesis of Ethyl 2-oxo-3-hydroxyaminopyrrolidine-4-carboxylate (**5**)

To a solution of pyrrolidine **2a** (1.5 g, 8.82 mmol) in CH₂Cl₂ (100 mL) was added NH₂OH.HCl (0.74 g, 10.59 mmol), NaHCO₃ (1.07 g, 12.71 mmol) and Na₂SO₄ (1.8 g, 12.71 mmol). The mixture was stirred at room temperature for 24 h, filtered and evaporated to furnish the hydroxyamino pyrrolidine **5** as white solid in a yield of 95%. White solid. ¹H-NMR (D₂O, 400 MHz) δ_H 3.95 (2H, *m*, OCH₂), 4.62 (2H, *s*, NCH₂), 1.05 (3H, *t*, $J = 6.8$ Hz, CH₃). ¹³C-NMR (CDCl₃, 100 MHz) δ_c 175.6, 166.2, 158.0, 104.3, 67.1, 60.0, 48.5, 13.3.

2.4. General Procedure for the Synthesis of Pyrrolidine Intermediates (**8a-e**):

NH₂NH₂.H₂O (5.8 mmol) was added to a mixture of ethyl 2-oxo-3-hydroxy-5(substituted)-pyrrole-4-carboxylate (**2a-e**) (2.9 mmol), AcOH (2.9 mmol) and EtOH (15 mL). The mixture was heated at reflux for 60 minutes. For **8a-d**, the reaction mixtures were evaporated under reduced pressure, extracted with EtOAc, washed with H₂O, dried over MgSO₄ and evaporated to afford the products. For **8e**, the resulting precipitate was filtered and washed with EtOH to give a white precipitate product **8e**.

Ethyl 2-oxo-3-hydrazinylpyrrolidine-4-carboxylate (8a): Yield: 95%; ¹H NMR (400 MHz, DMSO-*d*₆) δ_H 4.66 (2H, *s*, NCH₂), 3.98 (2H, *q*, $J = 7.2$ Hz, OCH₂), 3.12-3.96 (3H, *brs*, NHNH₂), 1.12 (3H, *t*, $J = 7.2$ Hz, CH₃). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_c 173.8, 164.7, 160.3, 98.3, 67.0, 58.2, 15.2.

Ethyl 2-oxo-3-hydrazinyl-1-methylpyrrolidine-4-carboxylate (8b): Yield: 74%; ¹H NMR (400 MHz, DMSO-*d*₆) δ_H 3.96 (2H, *q*, $J = 7.0$ Hz, OCH₂), 3.74 (2H, *s*, CH₂), 2.86 (3H, *s*, NCH₃), 1.11 (3H, *t*, $J = 7.1$ Hz, CH₃). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_c 170.08, 166.69, 164.90, 93.90, 57.74, 48.63, 30.24, 15.37.

Ethyl 2-oxo-3-hydrazinyl-1,5-dimethylpyrrolidine-4-carboxylate (8c): Yield: 70%; ¹H NMR (400 MHz, CDCl₃) δ_H 4.16 (2H, *q*, $J = 7.0$ Hz, OCH₂), 3.99 (1H, *q*, $J = 6.4$ Hz, CH), 2.90 (3H, *s*, NCH₃), 1.30 (3H, *d*, $J = 6.4$ Hz, CH₃), 1.24 (3H, *t*, $J = 7.1$ Hz, CH₃). ¹³C NMR (100 MHz, CDCl₃) δ_c 165.83, 163.57, 157.18, 112.96, 59.63, 55.39, 27.26, 18.01, 14.53.

Ethyl 2-oxo-5-ethyl-3-hydrazinyl-1-methylpyrrolidine-4-carboxylate (8d): Yield: 91%; ¹H NMR (400 MHz, CD₃OD) δ_H 4.21-4.09 (2H, *m*, OCH₂), 4.15 (1H, *t*, $J = 7.2$ Hz, CH), 2.91 (3H, *s*, NCH₃), 2.12-2.08 (1H, *m*, CH₂-a), 1.87-1.84 (1H, *m*, CH₂-b), 1.26 (3H, *t*, $J = 7.1$ Hz, CH₃), 0.47 (3H, *t*, $J = 7.3$ Hz, CH₃). ¹³C NMR (100 MHz, CD₃OD) δ_c 171.65, 164.23, 163.87, 109.74, 58.53, 58.27, 26.33, 21.03, 14.16, 13.64.

Ethyl 2-oxo-3-hydrazinyl-1-methyl-5-(4-methoxyphenyl)pyrrolidine-4-carboxylate (8e): Yield: 86%; ¹H NMR (400 MHz, CDCl₃) δ_H 7.03 (2H, *d*, $J = 8.5$ Hz, CHAr), 6.81 (2H, *d*, $J = 8.2$ Hz, CHAr), 4.93 (1H, *s*, NCH), 4.07 (2H, *q*, $J = 7.0$ Hz, OCH₂), 3.74 (3H, *s*, OCH₃), 2.74 (3H, *s*, NCH₃), 1.08 (3H, *t*, $J = 7.1$ Hz, CH₃). ¹³C NMR (100 MHz, CDCl₃) δ_c 164.90, 164.08, 159.89, 157.03, 128.82, 126.13, 114.22, 112.92, 62.29, 60.92, 55.34, 27.64, 14.03.

2.5. Synthesis of 2-oxo-5(4-methoxyphenyl)-1-methylpyrrolidine-2,3-dione (**11**)

Pyrrolidine **2e** (0.20 g, 0.77 mmol) was dispersed in 10% aq. HCl solution (10 mL) and heated under reflux for 7 hours which was gradually dissolved to give a brownish solution. The reaction mixture was then cooled and concentrated to dryness. The crude product was triturated with Et₂O and extracted with CH₂Cl₂ and water. The organic layer was dried over MgSO₄, filtered and concentrated in vacuo to give the pure product pyrrolidinone intermediate **11** in a yield of 75%. ¹H-NMR (CDCl₃, 400 MHz) δ_H 7.11 (2H, *d*, $J = 8.8$ Hz, Ar), 6.93 (2H, *d*, $J = 8.8$ Hz, Ar), 4.74 (1H, *dd*, $J = 7.4$ Hz, 3.2 Hz, NCH), 3.79 (3H, *s*, OCH₃), 3.17-3.10 (1H, *dd*, $J = 20.4$ Hz; $J = 7.4$ Hz, H_{4a}), 2.87 (3H, *s*, NCH₃), 2.53

(1H, *dd*, $J = 20.4$ Hz, $J = 3.2$ Hz, H_{4b}). ^{13}C -NMR (CDCl_3 , 100 MHz), δ_{C} 198.07, 160.04, 159.3, 130.0, 127.6, 114.8, 57.8, 55.3, 40.9, 29.6,

2.6. General procedure for the Synthesis of Iminosugars (**4a-e**, **7**, **10a-e**, **12**):

To a stirred solutions of **2a-e**, **5** and **8a-e** (6 mmol) in EtOH (40 mL) were added catalytic amount of Pd-C (10% wt.) (0.90 g) and AcOH (0.2 mL). The reactions were stirred vigorously under H_2 atmosphere for 24 h and then filtered through Celite. After removal of the solvent, the crude products were submitted to the next steps without purification. The crude products were dissolved in dry THF and were added slowly into solution of LiAlH_4 (excess) in a 50 mL round bottom flask. The mixtures were heated at 90 °C for 4-6 h and then cooled to 0 °C. The reaction mixtures were quenched by the addition of distilled H_2O , and the mixtures were filtered through Celite and concentrated in vacuo to give the pure products iminosugar analogues (**4a-e**, **7**, **10a-e**, **12**) as solids after recrystallization or purification by column chromatography.

3-hydroxy-4-hydroxymethylpyrrolidine (4a): Compound **4a** was previously reported (See Scheme 2)²⁰ White solid; 90%. HRESI-MS m/z : $[\text{M}+\text{H}]^+$ 118.0674 (Calc. for $\text{C}_5\text{H}_{12}\text{NO}_2$, 118.0862). ^1H -NMR ($\text{DMSO}-d_6$, 400 MHz) δ_{H} 3.52-3.40 (3H, *m*, CHOH and CH_2OH), 3.41-3.25 (4H, *m*, 2 x NHCH_2), 1.53 (1H, *m*, CH). ^{13}C -NMR ($\text{DMSO}-d_6$, 100 MHz) δ_{C} 70.6, 64.8, 60.3, 59.8, 46.2.

3-hydroxy-4-hydroxymethyl-1-methylpyrrolidine (4b): Compound **4b** was previously reported (See Scheme 2)²⁰ Brown solid; 90%. HRESI-MS m/z : $[\text{M}]^+$ 131.0908 (Calc. for $\text{C}_6\text{H}_{13}\text{NO}_2$, 131.0941). ^1H -NMR (CD_3OD , 400 MHz) δ_{H} 4.35 (1H, *dt*, $J = 5.2, 4.4$ Hz, CHOH), 3.77 (1H, *dd*, $J = 8.0, 11.2$ Hz, $\text{OCH}_2\text{-a}$), 3.60 (1H, *dd*, $J = 6.8, 11.2$ Hz, $\text{OCH}_2\text{-b}$), 3.52-3.40 (3H, *m*, CHOH and CH_2OH), 3.01 (1H, *dd*, $J = 5.6, 10.4$ Hz, $\text{NCH}_2\text{-a}$ CHOH), 2.79 (1H, *dd*, $J = 8.0, 9.2$ Hz, $\text{NCH}_2\text{-a}$), 2.44 (1H, *m*, $\text{NCH}_2\text{-b}$ CHOH), 2.41 (1H, *m*, $\text{NCH}_2\text{-b}$), 2.34 (3H, *s*, NCH_3) and 1.61 (1H, *m*, CH). ^{13}C -NMR (CD_3OD , 100 MHz) δ_{C} 71.3, 64.2, 60.3, 57.5, 45.3, 41.6.

5-ethyl-3-hydroxy-4-hydroxymethyl-1-methylpyrrolidine (4d): Brown solid; 85%. HRESI-MS m/z : $[\text{M}]^+$ 159.9179 (Calc. for $\text{C}_8\text{H}_{17}\text{NO}_2$, 159.1254). IR (ATR) ν/cm^{-1} : 3539 (stretching, OH), 3124-3024 (stretching, CH aliphatic), 1476-1347 (bending, CH aliphatic), 1043 (stretching, C-O). ^1H -NMR (CD_3OD , 400 MHz) δ_{H} 4.53 (1H, *dd*, $J = 5.6, 9.2$ Hz, CHOH), 3.83 (1H, *dd*, $J = 6.0, 10.8$ Hz, $\text{OCH}_2\text{-b}$), 3.79 (1H, *dd*, $J = 6.0, 10.8$ Hz, $\text{OCH}_2\text{-a}$), 3.46 (1H, *dd*, $J = 3.2, 11.6$ Hz, $\text{NCH}_2\text{-b}$), 3.13 (1H, *dd*, $J = 6.0, 12.0$ Hz, $\text{NCH}_2\text{-a}$), 2.80 (3H, *s*, NCH_3), 2.54 (1H, *dt*, $J = 7.6, 13.2$ Hz, NCH), 1.86 (2H, *sx*, $J = 7.2$ Hz, CH_2), 1.07 (3H, *t*, $J = 7.2$ Hz, CH_3). ^{13}C -NMR (CD_3OD , 100 MHz) δ_{C} 71.4, 69.2, 62.9, 56.2, 46.1, 41.2, 21.1, 10.4.

3-hydroxy-4-hydroxymethyl-1-methyl-5(4-methoxyphenenyl)pyrrolidine (4e): IR (ATR) ν/cm^{-1} : 3652 (stretching, OH), 3211-3179 (stretching, CH Ar), 3110-2967 (stretching, CH aliphatic), 1658 and 1559 (stretching, C=C Ar), 1505-1226 (bending, CH aliphatic), 1282 and 1070 (stretching, C-O-C), 1054 (stretching, C-O). Brown solid; 78%. HRESI-MS m/z : $[\text{M}]^+$ 237.1249 (Calc. for $\text{C}_{13}\text{H}_{19}\text{NO}_3$, 237.1359). ^1H -NMR (CD_3OD , 400 MHz) δ_{H} 7.24 (2H, *d*, $J = 8.4$ Hz, Ar-CH), 6.87 (2H, *d*, $J = 8.4$ Hz, Ar-CH), 3.80 (1H, *m*, CHOH), 3.76 (3H, *s*, OCH_3), 3.47 (1H, *dd*, $J = 4.0, 10.8$ Hz, $\text{OCH}_2\text{-b}$), 3.36 (1H, *dd*, $J = 6.4, 10.8$ Hz, $\text{OCH}_2\text{-a}$), 3.32 (1H, *m*, NCH), 3.13 (1H, *t*, $J = 7.6$ Hz, $\text{NCH}_2\text{-b}$), 2.71 (1H, *d*, $J = 9.2$ Hz, $\text{NCH}_2\text{-a}$), 2.22 (1H, *m*, CH), 2.07 (3H, *s*, NCH_3). ^{13}C -NMR (CD_3OD , 100 MHz) δ_{C} 159.5, 132.3, 129.0, 113.6, 73.5, 62.6, 55.6, 54.3, 49.4, 39.2.

3-hydroxyamino-4-hydroxymethylpyrrolidine (7): Brown solid; 70.4%. HRESI-MS m/z : $[\text{M}]^+$ 132.0428 (Calc. for $\text{C}_5\text{H}_{12}\text{N}_2\text{O}_2$, 132.0893). IR (ATR) ν/cm^{-1} : 3556 (stretching, OH), 3426 (stretching, NH), 3097-2991 (stretching, CH aliphatic), 1460-1367 (bending, CH aliphatic), 1069 (stretching, C-O). ^1H -NMR ($\text{DMSO}-d_6$, 400 MHz) δ_{H} 3.52-3.40 (2H, *m*, CH_2OH), 3.41-3.25 (4H, *m*, 2x NHCH_2), 2.6 (1H, *m*, HONHCH), 1.35 (1H, *m*, CH). ^{13}C -NMR ($\text{DMSO}-d_6$, 100 MHz) δ_{C} 61.2, 33.0, 31.8, 29.2, 26.0, 22.6.

3-hydrazinyl-4-hydroxymethylpyrrolidine (10a): Pale yellow solid; 93%. HRESI-MS m/z : $[\text{M}]^+$ 131.0853 (Calc. for $\text{C}_5\text{H}_{13}\text{N}_3\text{O}$, 131.1053). IR (ATR) ν/cm^{-1} : 3545 (stretching, OH), 3394 (stretching,

NH), 3097-2976 (stretching, CH aliphatic), 1463-1333 (bending, CH aliphatic), 1074 (stretching, C-O). ¹H-NMR (D₂O, 400 MHz) δ_{H} 3.35-3.19 (2H, *m*, CH₂OH), 3.21-3.01 (4H, *m*, 2 x NCH₂), 1.28 (1H, *m*, CH). ¹³C-NMR (D₂O, 100 MHz) δ_{C} 70.2, 63.4, 59.2, 58.7, 44.5.

3-hydrazinyl-4-hydroxymethyl-1-methylpyrrolidine (10b): White solid; 90%. HRESI-MS (negative mode) *m/z*: [M-H]⁻ 144.0791 (Calc. for C₆H₁₄N₃O, 144.1142). IR (ATR) ν/cm^{-1} : 3547 (stretching, OH), 3394 (stretching, NH), 3083-2974 (stretching, CH aliphatic), 1462-1330 (bending, CH aliphatic), 1059 (stretching, C-O). ¹H-NMR (CD₃OD, 400 MHz) δ_{H} 3.85-3.90 (2H, *m*, CH₂OH), 3.80 (1H, *m*, NH₂NHCH), 3.70-3.50 (4H, *m*, 2 x NCH₂), 2.80 (3H, *s*, NCH₃), 1.80 (1H, *m*, CH). ¹³C-NMR (D₂O, 100 MHz) δ_{C} 71.2, 68.8, 62.1, 59.4, 55.9, 53.4.

3-hydrazinyl-4-hydroxymethyl-1,5-dimethylpyrrolidine (10c): White solid; 95%. HRESI-MS (negative mode) *m/z*: [M+H]⁺ 159.0902 (Calc. for C₇H₁₇N₃O, 159.1377). IR (ATR) ν/cm^{-1} : 3531 (stretching, OH), 3392 (stretching, NH), 3103-2981 (stretching, CH aliphatic), 1468-2981 (bending, CH aliphatic), 1073 (stretching, C-O). ¹H-NMR (CD₃OD, 400 MHz) δ_{H} 3.84 (1H, *dd*, *J* = 7.2, 10.8 OCH₂-b), 3.72 (1H, *dd*, *J* = 5.2, 10.8 Hz, OCH₂-a), 3.58 (1H, *q*, *J* = 7.2 Hz, NH₂NHCH), 3.55 (1H, *m*, NCH₂-b), 3.41 (1H, *dd*, *J* = 4.8, 11.6 Hz, NCH₂-a), 2.33 (1H, *m*, NCH), 2.27 (3H, *s*, NCH₃), 1.70 (1H, *m*, CH), 0.94 (1H, *d*, *J* = 7.2 Hz, CH₃). ¹³C-NMR (CD₃OD, 100 MHz) δ_{C} 72.4, 69.2, 63.9, 60.4, 55.9, 46.2, 9.24.

5-ethyl-3-hydrazinyl-4-hydroxymethyl-1-methylpyrrolidine (10d): White solid; 90%. HRESI-MS (negative mode) *m/z*: [M+K-2H]⁻ 210.9579 (Calc. for C₈H₁₇N₃OK, 210.1014). IR (ATR) ν/cm^{-1} : 3531 (stretching, OH), 3393 (stretching, NH), 3113-2970 (stretching, CH aliphatic), 1466-1285 (bending, CH aliphatic), 1058 (stretching, C-O). ¹H-NMR (DMSO-*d*₆, 400 MHz) δ_{H} 3.66 (1H, *dd*, *J* = 4.8, 10.8 OCH₂-b), 3.52 (1H, *dd*, *J* = 5.2, 10.8 OCH₂-a), 3.44 (1H, *m*, NH₂NHCH), 3.24 (1H, *m*, NCH₂-b), 3.10 (1H, *m*, NCH₂-a), 2.90 (3H, *s*, NCH₃), 2.01 (1H, *m*, CH), 2.19 (1H, *q*, *J* = 5.6, NCH), 1.75 (2H, *qt*, *J* = 7.2 Hz, CH₂), 1.06 (1H, *t*, *J* = 7.2 Hz, CH₃). ¹³C-NMR (DMSO-*d*₆, 100 MHz) δ_{C} 72.4, 69.2, 63.9, 60.4, 55.9, 39.8, 25.1, 9.29.

3-hydrazinyl-4-hydroxymethyl-1-methyl-5-(4-methoxyphenyl)pyrrolidine (10e): Brown solid; 80%. HRESI-MS (negative mode) *m/z*: [M]⁻ 251.1182 (Calc. for C₁₃H₂₁N₃O₂, 251.1639). IR (ATR) ν/cm^{-1} : 3532 (stretching, OH), 3391 (stretching, NH), 3211-3174 (stretching, CH Ar), 3119-2957 (stretching, CH aliphatic), 1659 and 1559 (stretching, C=C Ar), 1469-1286 (bending, CH aliphatic), 1282 and 1073 (stretching, C-O-C), 1055 (stretching, C-O). ¹H-NMR (CD₃OD, 400 MHz) δ_{H} 7.31 (2H, *d*, *J* = 8.8 Hz, Ar-CH), 6.87 (2H, *d*, *J* = 8.8 Hz, Ar-CH), 4.21 (1H, *m*, NHCH), 3.75 (3H, *s*, OCH₃), 3.60 (1H, *dd*, *J* = 4.4, 11.6 Hz, OCH₂-b), 3.39 (1H, *dd*, *J* = 6.0, 11.6 Hz, OCH₂-a), 3.01 (1H, *d*, *J* = 10.8 Hz, NCH₂-b), 2.91 (1H, *d*, *J* = 9.6 Hz, NCH₂-a), 2.50 (1H, *dd*, *J* = 6.0, 11.2 Hz, CH), 2.06 (3H, *s*, NCH₃). ¹³C-NMR (CD₃OD, 100 MHz) δ_{C} 159.5, 131.8, 129.2, 113.6, 72.9, 71.5, 64.5, 60.0, 59.7, 54.4, 39.2.

3-hydroxy-5-(4-methoxyphenyl)pyrrolidine (12): Pale brown solid; 85%. HRESI-MS (negative mode) *m/z*: [M]⁻ 207.0912 (Calc. for C₁₂H₁₇NO₂, 207.1265). IR (ATR) ν/cm^{-1} : 3670 (stretching, OH), 3211-3173 (stretching, CH Ar), 3109-2980 (stretching, CH aliphatic), 1659 and 1560 (stretching, C=C Ar), 1497-1237 (bending, CH aliphatic), 1283 and 1072 (stretching, C-O-C), 1068 (stretching, C-O). ¹H-NMR (CD₃OD, 400 MHz) δ_{H} 7.27 (2H, *d*, *J* = 8.8 Hz, Ar-CH), 6.84 (2H, *d*, *J* = 8.8 Hz, Ar-CH), 4.39 (1H, *m*, CHOH), 4.29 (1H, *m*, NCH), 3.74 (3H, *s*, OCH₃), 3.50 (1H, *dd*, *J* = 6.8, 10.4 Hz, NCH₂-b), 3.50 (1H, *dd*, *J* = 7.2, 9.6 Hz, NCH₂-a), 2.52 (1H, *dt*, *J* = 7.6, 14.0 Hz, CH₂-b), 2.38 (1H, *m*, CH₂-a), 2.04 (3H, *s*, NCH₃). ¹³C-NMR (CD₃OD, 100 MHz) δ_{C} 159.5, 131.2, 128.7, 113.5, 70.9, 68.6, 65.4, 54.4, 44.8, 39.2.

2.7. Biological Assay

α -Glucosidase enzyme inhibitory activity assay was adopted from method as described by Elya *et al.* (2012) with slight modifications.²⁵

3. Results and Discussion

3.1. Chemistry

Synthesis of five-membered iminosugars analogues in this study are based on the modification of the core structures of glucosidase inhibitor iminoheitol and iminoarabinitol. Conventionally, synthesis of hydroxyprolines and pyrrolidines utilize carbohydrates as starting materials, due to the similarity of their structure.²¹ In this work, we prepared all starting substances for targeted iminosugar analogues via multicomponent reaction strategy (MCRs) using readily accessible oxaloacetate, aldehyde and amine as starting material.²²

In this study, five key intermediate ethyl 2-oxo-3-hydroxy-5(substituted)-pyrrole-4-carboxylates (**2a-e**) were prepared from MCRs of three components including diethyl oxaloacetate (ammonia and methylamine), aldehyde (formaldehyde, acetaldehyde, ethanaldehyde and *p*-methoxybenzaldehyde) and amine precursors (Scheme 1). The reaction was performed in the EtOH under reflux condition as reported previously by Mohammad et al. (2009).¹⁷ The MCRs begins with condensation of amine and aldehyde *via* nucleophilic addition to give a hemiaminal intermediate, followed by an elimination of water to yield the imine intermediate. Then the reaction continues with the formation of pyrrolidine ring *via* one-step addition and cyclization reaction of imine and diethyl oxaloacetate. In this route, the racemic mixture is formed, due to the direction of nucleophile attack can be achieved from both side of imine intermediate (Scheme 1). The product yield of this step is only moderate (57-68%) due to the presence of byproducts and side-products during the reactions.²³ The ¹³C NMR spectral data show the presence of four carbon signals in the downfield region belong to amide and ester (around δ_c 163.0 – 167.0) and enolic alkene part (around δ_c 157.0, 106.0). The racemic excess at C₍₅₎ position was recognized from the presence of replication signals on their ¹H NMR spectrum.

Scheme 1. Preparation of starting compounds (**2a-e**) and a proposed formation mechanism

Furthermore, Pd/C-catalyzed hydrogenation of **2a,b,d,e** in the presence of AcOH gave ethyl 2-oxo-3-hydrazinyl-5(substituted)-pyrrole-4-carboxylate (**3a,b,d,e**). Further reductions of amide and ester at using excess of LiAlH_4 in dry Et_2O lead to our desired products 3-hydroxy-4-hydroxymethyl-5(substituted)-pyrrolidines (**4a,b,d,e**) in good yield (75-90%). In the Pd/C-catalysis hydrogenation, hydrogen atom is always delivered in the same face of the alkene (“syn” addition), therefore the orientation of methine protons $\text{H}_{(3)}/\text{H}_{(4)}$ at **4a,b,d,e** are always in *cis*-configuration as show in Scheme 2. From this products, the important proton signals of **4a,b,d,e** were observed at around δ_{H} 1.28 (methine), δ_{H} 4.53 (oxygenated methine), δ_{H} 2.71 (methylene) and δ_{H} 3.47 (oxygenated methylene). The *cis*-configuration of protons at C₍₃₎ and C₍₄₎ was confirmed by the *J* coupling value of protons $\text{H}_{(3)}/\text{H}_{(4)}$ ($^3J=5.6$ Hz).²⁴ Those evident was supported by HRESI-MS experiments.

Scheme 2. Synthesis of iminosugar analogues (**4a,b,d,e**)

Synthesis of compound **7** was carried out by treatment of intermediate **2a** with $\text{NH}_2\text{OH} \cdot \text{HCl}$ in the presence of a weak base NaHCO_3 at room temperature gave enamine-imine tautomerism products of ethyl 2-oxo-3-hydroxyamino-pyrrole-4-carboxylate and 2-oxo-3-hydroxyimino-pyrrole-4-carboxylate (**5**) (95%). This tautomerization was confirmed from the presence of additional signals belong to aliphatic methine (δ_{C} 48.5) and quaternary sp^2 carbon (δ_{C} 158.0), in which oxime product only exist as minor products. Further catalytic hydrogenations of **5** using in the presence of Pd/C and then reduction with LiAlH_4 lead to the targeting compound 3-hydroxyamino-4-hydroxymethyl-pyrrolidine (**7**) in the good yield (70.4%) (Scheme 3). In these reductions, the *cis*-orientation of methine protons $\text{H}_{(3)}/\text{H}_{(4)}$ was anticipated due to the utilization of stereoselective hydrogenation catalyst H_2 Pd/C. The structures of **5** and **7** were confirmed by NMR and HRESI-MS spectral data.

Scheme 3. Synthesis of 3-hydroxyamino-4-hydroxymethylpyrrolidine **7**.

For synthesis of iminosugars (**10a-e**), the transformation of hydroxyl group of **2a-e** into hydrazinyl functionalities have been done through the treatment with hydrazine hydrate in EtOH under reflux condition.²⁵ In this step, the appropriate products ethyl 2-oxo-3-hydrazinyl-5(substituted)-pyrrole-4-carboxylate (**8a-e**) were formed in moderate to good yields (70-95%). Further steps are continued with catalytic hydrogenations in the presence of Pd/C and reduction with LiAlH_4 respectively to give targeting 3-hydrazinyl-4-hydroxymethyl-5(substituted)-pyrrolidine (**10a-e**) in the excellent yields (80-95%) (Scheme 4). The characteristic signals in the skeleton of intermediates (**8a-e**) and targeting iminosugars (**10a-e**) have been confirmed by NMR and HRESI-MS.

Scheme 4. Synthesis of targeting iminosugar analogues (**10a-e**)

Synthesis of targeting iminosugar **12** was started with decarboxylation of key intermediate **2e** using 10% aqueous HCl under reflux condition which gave pyrrolidinone intermediate **11** in good yield (75%). Furthermore, direct reduction of ketone and amide groups in **11** using LiAlH_4 in the dry Et_2O at 60°C lead to 3-hydroxy-5-(4-methoxyphenyl)-pyrrolidine (**12**). The structure of **12** was confirmed by the presence of two methylene groups and absence proton signals for ethyl ester group. The structure of **12** was also supported by HRESI-MS spectral data.

Scheme 5. Synthesis of 3-hydroxy-5-(4-methoxyphenyl)-pyrrolidine (**12**)

All synthesized iminosugars have been evaluated for their inhibitory potential against α -glucosidase as reported by Elya et al. (2012) with slight modification.²⁶ The tested compounds showed a variable degree of α -glucosidase inhibition with IC_{50} values ranging between 1.116 ± 1.010 mM and 2.938 ± 1.076 mM as compared with 1-deoxynojirimycin (DNJ) ($\text{IC}_{50} = 0.558 \pm 1.030$ mM). Compound **10e** is found to be the most potent compound in the series (Table 2). This compound has hydrazine group at C-4 position and methoxybenzene group at C-2 position which seem to play an important role in the inhibition activity. Besides that, among of the 4-hydroxypyrrolidine analogues (**4a-e**), **4a** has the highest inhibitory activity because the compound highly mimics the structure of substrate, thus it fits well in the active site.

Table 2. α -Glucosidase inhibition studies of five-membered iminosugars

Compound	IC ₅₀ (mM)*
4a	1.404 $\pm$ 1.005
4b	2.938 $\pm$ 1.076
4d	1.800 $\pm$ 1.016
4e	1.660 $\pm$ 1.017
7	2.171 $\pm$ 1.365
10a	1.751 $\pm$ 1.006
10b	1.311 $\pm$ 1.003
10c	1.271 $\pm$ 1.859
10d	1.523 $\pm$ 1.006
10e	1.116 $\pm$ 1.010
12	1.169 $\pm$ 1.012
DNJ (control)	0.558 $\pm$ 1.030

*Data was expressed as mean $\pm$ S.D., n = 3.

As a conclusion, a diverse framework of pyrrolidine-based iminosugars analogues was successfully synthesized from MCRs strategy with simple procedure under mild conditions, short steps and good yields. The utilization of non-carbohydrate precursor for the synthesis of iminosugar have opened the diversity of the iminosugar skeleton for the variety of biological activity study. This study also demonstrated that the newly synthesized pyrrolidine-based iminosugar analogues had promising effects in inhibiting α -glucosidase and further optimization can be done to improve their activity.

Acknowledgements

The author would like to thank the generous support of Institute of Science (IOS), UiTM Shah Alam, Malaysia, Prof. Dr Mohd Zaki Bin Salleh from Integrative Pharmacogenomics Institute (iPROMISE), UiTM Selangor, Malaysia for mass spectrometer measurement of samples. The Ministry of Higher Education (MOHE), Malaysia for the financial support under the Fundamental Research Grant Scheme (Grants No.: 600-RMI/FRGS 5/3 (109/2019).

ORCID

Agustono Wibowo: [0000-0001-6588-9507](https://orcid.org/0000-0001-6588-9507)

Zurina Shaameri: [0000-0002-0314-8532](https://orcid.org/0000-0002-0314-8532)

Mohd Fazli Mohammat: [0000-0002-6926-0447](https://orcid.org/0000-0002-6926-0447)

Ahmad Sazali Hamzah: [0000-0002-6826-1659](https://orcid.org/0000-0002-6826-1659)

References

- [1] Compain, P.; Martin, O.R. Iminosugars: From Synthesis to Therapeutic Applications. Wiley 2007.
- [2] Stütz, A.E. Iminosugars as Glycosidase Inhibitors: Nojirimycin and Beyond. Wiley-VCH 1999.
- [3] Doddi, V.R.; Kokatla, H.P.; Pal, A.P.J.; Basak, R.K.; Vankar, Y.D. Synthesis of hybrids of D-glucose and D-galactose with pyrrolidine-based iminosugars as glycosidase inhibitors. *Eur. J. Org. Chem.* **2008**, *34*, 5731-5739.
- [4] Donohoe, T.J.; Freestone, G.C.; Headley, C.E.; Rigby, C.L.; Cousins, R.P.C.; Bhalay, G. Enantioselective partial reduction of 2,5-disubstituted pyrroles via a chiral protonation approach. *Org. Lett.* **2004**, *6*(18), 3055-3058.
- [5] Cren, S.; Wilson, C.; Thomas, N.R. A rapid synthesis of hexofuranose-like iminosugars using ring-closing metathesis. *Org. Lett.* **2005**, *7*(16), 3521-3523.
- [6] Carmona, A.T.; Fuentes, J.; Robina, I.; García, E.R.; Demange, R.; Vogel, P.; Winters, A.L. Stereoselective syntheses of 1,4-dideoxy-1,4-imino-octitols and novel tetrahydroxyindolizidines. *J. Org. Chem.* **2003**, *68*(10), 3874-3883.
- [7] Karaveg, K.; Siriwardena, A.; Tempel, W.; Liu, Z.-J.; Glushka, J.; Wang, B.-C.; Moremen, K.W.; Mechanism of class 1 (Glycosylhydrolase Family 47) α -mannosidases involved in N-glycan processing and endoplasmic reticulum quality control. *J. Biol. Chem.* **2005**, *280*(16), 16197-16207.

- [8] Karl Dieter, R.; Lu, K. α -(N carbamoyl)alkylcuprate chemistry in the synthesis of nitrogen heterocycles. *J. Org. Chem.* **2002**, 67(3), 847–855.
- [9] White, J. D.; Hrcniar, P. Synthesis of polyhydroxylated pyrrolizidine alkaloids of the alexine family by tandem ring-closing metathesis–transannular cyclization. (+)-australine. 2001. *J. Org. Chem.* **2000**, 65(26), 9129–9142
- [10] Zhao, H.; Hans, S.; Cheng, X.; Mootoo, D. R. Allylated monosaccharides as precursors in triple reductive amination strategies: synthesis of castanospermine and swainsonine. *J. Org. Chem.* **2001**, 66(5), 1761–1767
- [11] Back, T.G.; Nakajima, K.; A convenient new route to piperidines, pyrrolizidines, indolizidines, and quinolizidines by cyclization of acetylenic sulfones with β - and γ -chloroamines. Enantioselective total synthesis of indolizidines (–)-167B, (–)-209D, (–)-209B, and (–)-207A. *J. Org. Chem.* **2000**, 65(15), 4543–4552.
- [12] Lindstrom, U.M.; Ding, R.; Hidestøl, O. Efficient asymmetric synthesis of an azasugar in water, *Chem. Commun.* **2005**, 13, 1773–1774.
- [13] Ayad, T.; Génisson, Y.; Broussy, S.; Baltas, M.; Gorrichon, L. A flexible route towards five-membered ring imino sugars and their novel 2-deoxy-2-fluoro analogues. *Eur. J. Org. Chem.* **2003**, 15, 2903–2910.
- [14] Shaameri, Z.; Azib, N.A.; Mohamat, M.F.; Hamzah, A.S. Synthesis of 3,4-fused γ -lactone- γ -lactam bicyclic moieties as multifunctional synthons for bioactive molecules. *J. Heterocycl. Chem.* **2016**, 53(4), 1059–1064.
- [15] Mohammad, M.F.; Mansor, N.S.; Shaameri, Z.; Hamzah, A.S. Diastereoselective reduction of 2,3-dioxo-4-carboxy-5-substituted pyrrolidines using $\text{NaBH}_4/\text{AcOH}$ and heterogeneous hydrogenation reactions. *J. Korean. Chem. Soc.* **2015**, 59(1), 31–35.
- [16] Shaameri, Z.; Ali, S.H.S.; Mohamat, M.F.; Yamin, B.M.; Hamzah, A.S. A short and elegant synthesis of ($\pm$)-streptopyrrolidine. *J. Heterocycl. Chem.* **2013**, 50(2), 320–325.
- [17] Mohammad, M.; Shaameri, Z.; Hamzah, A. Synthesis of 2,3-dioxo-5-(substituted)arylpyrroles and their 2-oxo-5-aryl-3-hydrazone pyrrolidine derivatives, *Molecules* (Basel, Switzerland), **2009**, 14(1), 250–256.
- [18] Mohalid, N.; Hamzah, A.S.; Shaameri, Z.; Chemical exploration of 4-hydroxybenzylated 3-substituted tetramic acid. *Malaysian J. Anal. Sci.* **2015**, 19(2), 359–368.
- [19] Wibowo, A.; Shaameri, Z.; Mohammad, F.; Hamzah, A.S. Asymmetric reduction of 3-ketoproline ethyl ester by modified borohydrides and various vegetables. *J. Korean Chem. Soc.* **2017**, 61(5), 244–250.
- [20] Bacho M.Z.; Mohammad M.F.; Shaameri Z.; Wibowo A.; Kamarulzaman F.; Hamzah A.S.A. Facile synthesis of pyrrolidine-based iminosugars as potential α -glucosidase inhibitors. *Orient. J. Chem.* **2020**, 36(2), 309–319
- [21] Compain, P.; Chagnault, V.; Martin, O.R. Tactics and strategies for the synthesis of iminosugar C-glycosides: a review. *Tetrahedron Asymm.* **2009**, 20(6), 672–711.
- [22] Manta, S.; Kollatos, N.; Mitsos, C.; Chatziefraimidi, G.-A.; Papanastasiou, I.; Gallos, J.K.; Komiotis, D. Multicomponent reaction of aldehydes, amines and oxalacetate analogues leading to biologically attractive pyrrole derivatives. *Mini-Rev. Med. Chem.* **2020**, 20(10), 818–830.
- [23] Hayashi, Y. Pot economy and one-pot synthesis. *Chem. Sci.* **2016**, 7(2), 866–880.
- [24] Ansari, A.A.; Vankar, Y.D. Synthesis of pyrrolidine iminosugars, (–)-lentiginosine, (–)-swainsonine and their 8a-epimers from d-glycals. *RSC Adv.* **2014**, 4(24), 12555–12567.
- [25] Dabideen, D.R.; Cheng, K.F.; Aljabari, B.; Miller, E.J.; Pavlov, V.A.; Al-Abed, Y. Phenolic hydrazones are potent inhibitors of macrophage migration inhibitory factor proinflammatory activity and survival improving agents in sepsis. *J. Med. Chem.* **2007**, 50(8), 1993–1997.
- [26] Elya, B.; Basah, K.; Mun'im, A.; Yuliastuti, W.; Bangun, A.; Septiana, E.K. Screening of α -glucosidase inhibitory activity from some plants of Apocynaceae, Clusiaceae, Euphorbiaceae, and Rubiaceae. *J. Biomed. Biotechnol.* **2012**, 281078.