Phytoextraction potential of different grasses for the uptake of cadmium and lead from industrial wastewater

Sana Ullah^{*1}, Zafar Iqbal¹, Sajid Mahmood¹, Kalsoom Akhtar² and Rehmat Ali¹

¹Soil and Environmental Sciences Division, Nuclear Institute for Agriculture (NIA), Tandojam Tando Jam 70050, Sindh, Pakistan

¹Soil and Environmental Sciences Division, Nuclear Institute for Agriculture and Biology (NIAB), Faisalabad 38000, Pakistan

²National Institute for Biotechnology and Genetic Engineering (NIBGE), Faisalabad 38000, Pakistan

[Received: June 03, 2019 Accepted: January 04, 2020 Published Online: March 21, 2020]

Abstract

Industrial wastewater contains a variety of contaminants like salts, organic carbon, and heavy metals. Among the heavy metals, cadmium (Cd) and lead (Pb) are considered highly toxic even at low concentration. These metals could enter the food chain through a process of phytoassimilation, hence, lethal for living beings. The present study aimed to investigate Cd and Pb phytoextraction in four grass species viz. Dhab (*Desmostachya bipinnata*), Sporobolus (*Sporobolus arabicus*), Kallar (*Leptochloa fusca*) and Para grass (*Brachiaria mutica*) from industrial wastewater. The grasses were grown hydroponically in plastic pots in industrial wastewater as growth medium under greenhouse conditions. The experiment trial was arranged following completely randomized design (CRD) with three replicates. Results showed that *B. mutica* had maximum shoot metal content (Pb = 21, Cd = 0.66 mg kg⁻¹ dry matter), shoot metal uptake (Pb = 201.8, Cd = 6.39 µg plant⁻¹), translocation factor (Pb = 0.73, Cd = 0.55), shoot dry matter (28.8 g pot⁻¹), and root dry matter (4.2 g pot⁻¹). Root Pb concentration was highest in *B. mutica* followed by *D. bipinnata* and *L. fusca*. Whereas *S. arabicus* showed depressed growth, minimum shoot metal accumulation and uptake potential. Thereby among the tested grass species *B. mutica* could be suitable option to remediate industrial wastewater contaminated with moderate levels of Pb and Cd.

Keywords: Metal uptake, para grass, kallar grass, dry biomass, metal concentration

Introduction

Heavy metals usually indicate environmentally bad metals which have higher atomic number, or atomic weight, density (3.5-7.0 g cm⁻³ or above), properties of metallic substance at room temperature, and are extremely toxic for living organisms even at lower concentration (Duffus, 2002; Mitra, 2019). These metals include lead (Pb), cadmium (Cd), chromium, nickel, arsenic and mercury which are well-known environmental contaminants due to their toxic nature, bioaccumulation and persistence in the environment (Latif *et al.*, 2019; Ali *et al.*, 2019). Heavy metal contamination distresses the environment and food security ultimately posing serious health concerns (Rai *et al.*, 2019; Mushtaq *et al.*, 2019). With rampant inflation in anthropogenic activities, the release of toxic metals into the environment is increasing day by day. According to an estimate, global annual release of Pb and Cd into the environment is 783,000 and 22,000 metric tons, respectively

(Singh *et al.*, 2003). Wastewater from different sources (domestic, municipal and industries) plays significant role in release of heavy metals like Pb and Cd. About 2 million tons of sewage sludge and effluents are polluting the world's water per day (Azizullah *et al.*, 2011). Issue is more serious in developing countries due to improper treatment of domestic, municipal and industrial waste. Such as in India 24% of wastewater (industry and domestic) is treated, whereas in Pakistan only 2% of total sewerage or wastewater is treated properly (IWMI, 2003; Minhas and Samra, 2004; Majeed *et al.*, 2018). Wastewater released from different industries contains a variety of toxic metals such as Pb, Cd, chromium and arsenic (Rehman *et al.*, 2018, 2019; Hussain *et al.* 2019; Afzal *et al.* 2019). Use of industrial effluents/wastewater for cultivation of crops has created the problem of heavy metal contamination in soil (Hussain *et al.*, 2019). Moreover, in developing countries, like Pakistan, shortage of good quality irrigation water also urges the farmers to use wastewater for irrigation purpose

*Email: msunny9887@gmail.com

(Qadir *et al.*, 2010). Use of contaminated wastewater to irrigate crops and vegetables increases the risk of entry of toxic metal into our food chain. For example, Ahmad *et al.* (2019) observed Pb and Cd contamination in wheat grain up to 0.2 and 1.6 mg kg⁻¹ respectively, due to irrigation with wastewater. Hussain *et al.* (2019) also observed Pb and Cd contamination in different vegetables (carrot, radish, and spinach) irrigated with wastewater.

In human beings, heavy metals cause severe health problems such as Pb poisoning especially in children and kidney disease due to Cd toxicity (WHO, 1997; Padmavathiamma and Li, 2007; Rehman *et al.*, 2018). Intake of Pb and Cd above their permissible limits could leave carcinogenic impacts in human body (Salman *et al.*, 2019). Cadmium and Pb could easily cause acute and chronic risk even at their lower concentrations, causing renal damage, headache, hypertension and malformation in fetuses (Shi and Chatt, 2014; Zhou *et al.*, 2019). Svetlana *et al.* (2019) observed that higher levels of Pb and Cd caused chronic sinusitis in children. In plants, heavy metals retard shoot and root growth, reduce mineral uptake (Fe, Cu, Zn, Mn), and stimulate the production of reactive oxygen species (H₂O₂, O₂⁻) that damage plasma membrane (Zou *et al.*, 2019). Improper management to dispose-off wastewater can cause ground and/or drinking water contamination with toxic metals, thereby, it is imperative to strictly monitor the wastewater management (Hussain *et al.*, 2019; Rehman *et al.*, 2019).

Recently, much consideration has been paid to various techniques/ approaches in order to remove toxic metals

from effluents discharged in to water bodies such as membrane filtration, reverse osmosis, ion exchange, solvent extraction, absorption/adsorption, precipitation and electrochemical treatments (Miretzky *et al.*, 2006; Singh *et al.*, 2012; Liu *et al.*, 2018). But these techniques cover higher cost, technical expertise and are difficult to apply at large scale (Olguin and Sanchez-Galvan, 2012), thereby, phytoremediation is a well-suited technique. It is cost effective, easy to apply, and is environment friendly (Cheng, 2003; Sarwar *et al.*, 2017; Liu *et al.*, 2018). A hyperaccumulator to be used to extract water soluble heavy metals, should have fast growth, ample root system, higher biomass, resistant to higher concentration of metals, and retain higher metal concentration (Garbisu and Alkorta, 2001; Soda *et al.*, 2012; Rev *et al.*, 2017). In this regard, use of different grass species is gaining promising importance due to their tolerance to phytotoxicity, hinder natural succession of weeds, suitable for grazing, bulk biomass, rapid growth and compact root system. Well-developed root system in grasses triggers phytostabilization by retarding metal mobility and improves metal uptake and their accumulation in plant body hence make them more exploring phytoremediant agent as compared to shrubs, herbs, and trees (Ghosh *et al.*, 2017). Literature has confirmed the potential of different grass species (*Cynodon dactylon* L, *Chrysopogon zizanioides*, *Imperata cylindrical*) for the uptake of heavy metals like Pb and Cd from wastewater and contaminated soil (Pongthornpruek, 2017; Chen *et al.*, 2019; Kiiskila *et al.*, 2019; Zheng *et al.*, 2019).

Table 1: Some salient characteristics of grasses used in the experiment

Grass species	Origin	Salient feature	Salt tolerance potential [Root zone salinity (EC, dS m ⁻¹) causing 50% decrease in yield]	Reference
Dhab (<i>Desmostachya bipinnata</i>)	China, India, Pakistan and Africa	C4 perennial grass, used as fodder source in dry-saline areas	9.0	NIAB (2007); Gulzar <i>et al.</i> (2007); Ahmad <i>et al.</i> (2009); Asrar <i>et al.</i> (2017)
Kallar grass (<i>Leptochloa fusca</i>)	Australia, now well adapted to Pakistan and India	C4 perennial grass, used as animal fodder and reclamation agent for saline-sodic soils	22.0-14.6	Qureshi <i>et al.</i> (1982); Ashraf <i>et al.</i> (2012); Adabnejad <i>et al.</i> (2015)
Para grass (<i>Brachiaria mutica</i>)	African countries	C4 plant, known as buffalo grass, used as pasture, fodder	Salt tolerant	Ashraf <i>et al.</i> (2012); Mohanty and Patra, (2012)
<i>Sporobolus</i> (<i>Sporobolus arabicus</i>)		C4 perennial grass, used as desalinization tool for saline-sodic soils	21.7	Ashraf <i>et al.</i> (2012); Falla <i>et al.</i> (2017); Yobi <i>et al.</i> (2017)

Therefore, present experiment was conducted to investigate the growth response of four different salt tolerant grass species in wastewater and to assess the phytoextraction potential of four different salt tolerant grass species for Pb and Cd from wastewater in order to identify suitable grass species for the remediation of contaminated wastewater.

Materials and Methods

Plant material

Grass species viz. Kallar grass (*Leptochloa fusca*), Para grass (*Brachiaria mutica*), *Sporobolus* (*Sporobolus arabicus*) and Dhab (*Desmostachya bipinnata*) were brought from the Biosaline Research Station Pakka Anna, a substation of Nuclear Institute for Agriculture and Biology (NIAB), Faisalabad. The silent features of the grasses are presented in Table 1. Each grass was grown in plastic pots (three plants/cuttings per pot) containing gravel and three liters of wastewater which was aerated manually by placing pots outside on wooden sticks for two hours twice a day (morning and evening). Each grass was grown in three independent pots, representing three replicates. The pots were placed in the greenhouse of NIAB under natural conditions/ photoperiod, with mean temperatures of 23-26 °C (night-day) and relative humidity of 60-65%. Wastewater was replaced twice a week with its fresh counterpart till plants were harvested after three months. After harvesting root and shoot samples were dried at 70 °C in oven until constant weight and dry matter was recorded.

Table 2: Characteristics of industrial wastewater used in the experiment

Parameter	Value	Unit
pH	7.42	
SAR	254	
EC	3.57	dsm ⁻²
RSC	7	me L ⁻¹
TSS	3390	mg L ⁻¹
CO ₃ ⁻	ND*	mg L ⁻¹
HCO ₃ ⁻	17	mg L ⁻¹
Na ⁺	570	mg L ⁻¹
K ⁺	56	mg L ⁻¹
Total N	2.372	mg L ⁻¹
NH ₄ -N	0.07	mg L ⁻¹
NO ₃ -N	0.0006	mg L ⁻¹
Phosphorus	0.278	mg L ⁻¹
Organic carbon	42.3	mg L ⁻¹
Ca+Mg	10	me L ⁻¹

Cl ⁻	20	me L ⁻¹
Pb	0.92	mg L ⁻¹
Cd	0.03	mg L ⁻¹

ND*: Not detected

Water quality analysis

The wastewater used in this study was collected from industrial wastewater collecting system of village 'Chakaira 217 R.B.' Faisalabad, Punjab, Pakistan. The collected wastewater was stored in the plastic tanks once and analyzed for different quality indicators which are presented in Table 2. The pH was measured by pH meter (HI-8520, Hanna Instruments, UK), EC and total soluble salts (TSS) by conductivity meter (LF-538 WTW, Germany), carbonates (CO₃⁻), bicarbonates (HCO₃⁻), chlorides (Cl⁻), calcium (Ca⁺²) and magnesium (Mg⁺²) were determined by titration method as described by Richards (1954). Sodium (Na⁺) and potassium (K⁺) were determined by Flame Photometer (PFP7-Jenway, UK). Phosphorus was measured by Watanabe and Olsen (1965) method, organic carbon by Riehm and Ulrich (1954) method. Nitrogen (NH₄⁺-N, NO₃⁻-N, total N) was determined by following Kroon (1993) and Kjeldahl method with some modification. Wastewater was concentrated in water bath and the organic bound N was digested by adding the mixture of conc. H₂SO₄, K₂SO₄ and Se as catalyst.

Metal concentration, uptake and translocation factor (TF)

Root and shoot samples were ground and digested following the method described by Wolf (1982). After digestion, extracts of samples were used to determine Pb and Cd concentration by atomic absorption spectrometer (AA240FS, Varian, Australia). Shoot metal uptake was calculated by multiplying shoot dry matter with shoot metal concentration:

$$\text{Shoot metal uptake } (\mu\text{g/plant}) = \text{shoot dry matter (g)} \times \text{shoot metal concentration } (\mu\text{g/g})$$

Translocation factor (TF) was calculated by following formula (Soda *et al.*, 2012):

$$TF = \frac{\text{shoot metal concentration}}{\text{root metal concentration}}$$

Statistical analysis

Statistical differences among different grass species were assessed by one-way analysis of variance (ANOVA) at $p \leq 0.05$ and least significant difference (LSD) test was used to separate the significant means of treatments using Statistix 8.1.

Results

Wastewater characteristics

Different physicochemical parameters of wastewater

(Table 2) revealed that wastewater contained significant amount of essential plant nutrients (N, P, K, Ca, Mg), organic C while high levels of Na^+ (570 mg L^{-1}), SAR (254), RSC (7 mg L^{-1}), and TSS (3390 mg L^{-1}) were recorded. Heavy

Figure 1: Effect of industrial wastewater on shoot and root dry matter yield of various grasses. Data represent the means of three replicates. The bars sharing similar letters do not differ significantly at $p \leq 0.05$.

Figure 2: Effect of industrial wastewater on shoot and root Pb concentration of various grasses. Data represent the means of three replicates. The bars sharing similar letters do not differ significantly at $p \leq 0.05$.

Figure 3: Effect of industrial wastewater on shoot and root Cd concentration of various grasses. Data represent the means of three replicates. The bars sharing similar letters do not differ significantly at $p \leq 0.05$.

metals Pb and Cd were found to be 0.92 and 0.03 mg L⁻¹, respectively.

Dry matter yield

Results regarding shoot dry matter yield are presented in Figure 1. *B. mutica* had maximum shoot dry matter yield (28.8 g pot⁻¹) as compared to other grass species, following *L. fusca* (20.6 g per pot) and *D. bipinnata* (12.6 g pot⁻¹), while the *S. arabicus* showed lowest dry matter yield of 3.4 g pot⁻¹. Similarly, in case of root dry matter yield (Figure 1), application of Chakaira-wastewater significantly affected the root dry matter of all grass species at $p \leq 0.05$. Root dry matter was observed in the sequence of *B. mutica* > *L. fusca* > *D. bipinnata* > *S. arabicus* ranging between 1.1 to 4.21 g pot⁻¹.

Shoot and Root metal concentration

Being statistically at par with each other, *B. mutica* and *L. fusca* had more Pb concentration in shoot, i.e. 21.05 and 17.14 mg kg⁻¹ dry weight, respectively, than *D. bipinnata* and *S. arabicus* with corresponding values of 9.55 and 8.19 mg kg⁻¹ dry weight, respectively (Figure 2). Data regarding root Pb concentration (Figure 2) revealed that highest root Pb concentration was recorded in *B. mutica* (28.82 mg kg⁻¹ dry weight) while the lowest was in *S. arabicus* (8.19 mg kg⁻¹ dry weight). Root Pb concentration in *D. bipinnata* and *L. fusca* was 23.61 and 20.06 mg kg⁻¹ dry weight, respectively. In case of shoot Cd concentration (Figure 3), high values were recorded for *B. mutica* (0.66 mg kg⁻¹ dry weight) followed by *L. fusca* (0.38 mg kg⁻¹ dry weight). The *S. arabicus* and *D. bipinnata* showed lower shoot Cd concentration, 0.23 and 0.27 mg kg⁻¹ dry weight

Figure 4: Effect of industrial wastewater on Pb and Cd shoot uptake of various grasses. Data represent the means of three replicates. The bars sharing similar letters do not differ significantly at $p \leq 0.05$.

Figure 5: Effect of industrial wastewater on Pb and Cd translocation factor (TF) of various grasses. Data represent the means of three replicates. The bars sharing similar letters do not differ significantly at $p \leq 0.05$.

respectively, with non-significant difference among them. However, no significance difference was observed in root Cd concentration of different grass species (Figure 3) and it ranged from 1.24-1.57 mg kg⁻¹ dry weight.

Metal uptake and Translocation factor

Shoot metal uptake varied with different grass species and was in the range of 9.4-201.8 and 0.26-6.39 µg plant⁻¹ for Pb and Cd (Figure 4), respectively. Uptake for both the metals by different grass species was in the order *B. mutica* > *L. fusca* > *D. bipinnata* > *S. arabicus*. However, *B. mutica* indicated better efficiency with significant uptake of Pb and Cd, 201.8 and 6.39 µg plant⁻¹ respectively, in comparison to other grass species. The translocation factor (TF) for Pb and Cd did not exceed the unity (Figure 5). *B. mutica* and *L. fusca* gave higher TF-values for Pb 0.73 and 0.85, respectively, compared to other grasses. Moreover, TF did not differ significantly for *D. bipinnata*, *L. fusca* and *S. arabicus* except for *B. mutica* which showed highest TF-value 0.55 ($p \leq 0.05$).

Discussion

The pollution status is becoming worse especially in the developing countries due to modernization, such as Pakistan where industries share the major portion in water pollution (Sial *et al.*, 2006; Majeed *et al.*, 2018; Kadam *et al.*, 2018; Hussain *et al.*, 2018). These industries are generating over 435 million gallons of sewage and effluents daily (Ijaz *et al.*, 2016). A large quantity of this wastewater is discharged in the outer environment, and farmers near urban areas are likely to use this water for irrigation purpose due to shortage of good quality irrigation water. It contains toxic metals i.e. Cd (0.03 mg L⁻¹) and Pb (0.92 mg L⁻¹) (Table 2). Amount of Pb present in wastewater was above the permissible limit set by National Environmental Quality Standards (NEQS), Pakistan. Thereby, there is dire need for appropriate treatment of wastewater in order to prevent the entry of toxic metals into food chain.

Growth potential of grasses in wastewater showed variable influence on shoot and root dry biomass of grass species, especially *B. mutica* indicated good growth which might be due to the presence of organic matter and inorganic mineral ions like N, P, K, Ca, and Mg in wastewater which are necessary for plant growth and development (Adrover *et al.*, 2012; Rev *et al.*, 2017). The results of present study are consistent to the findings that use of urban wastewater as irrigation enhanced the photosynthesis, growth and dry matter yield of chickpea (Tak *et al.*, 2013). *S. arabicus* showed least growth potential compared to other grass species which might be

due to the suppressive effects of toxic elements (Cd, Pb) present in the wastewater (Nair *et al.*, 2008; Adrover *et al.*, 2012). These toxic metals may reduce plant growth by inhibiting photosynthesis, respiration and enzymatic activity, decreasing chlorophyll and nitrogen contents, increasing the production of reactive oxygen species, disequilibrium in nutrients, water and hormonal balance of plants (Li *et al.*, 2012; Lou *et al.*, 2017; Akhtar *et al.*, 2018). It was observed that grass species showed variable response regarding accumulation of Pb and Cd in roots and shoots in wastewater growth medium. However, the Pb and Cd concentration in roots of all grass species was more than shoot. This might be due to negative charges present on root surface capable to bind the metal cations and decrease root to shoot transport (Zhivotovsky *et al.*, 2011). These results are in line with the findings of Roongtanakiat *et al.* (2007) who found more concentration of heavy metals (Pb, Mn, Cu, Fe, Zn) in roots as compared to shoot of *Vetiveria zizanioides* growing in industrial wastewater. Moreover, Silva *et al.* (2016) also observed high Cd concentration in roots compared to above ground parts of forage grasses.

Results regarding shoot metal concentration and shoot metal uptake showed that *B. mutica* had maximum metal concentration (Pb 21 and Cd 0.66 mg kg⁻¹ dry matter) and uptake (Pb 201.8 and Cd 6.4 µg plant⁻¹) followed by *L. fusca*. Good response of *B. mutica* for uptake and accumulation of heavy metals (Pb, Cd) could be due to different genetically controlled mechanisms responsible for varying uptake and transportation potential of metal cations from soil/ solution to aerial parts of the plant such as composition and quantity of organic acids released through roots to mobilize/ chelate the metal cations (Hall, 2002; Sarwar *et al.*, 2010; Najafi *et al.*, 2015; Alves *et al.*, 2016). These results are in agreement with our recent study in which *B. mutica* showed good potential regarding the shoot metal concentration (Cd > 150 and Pb > 1000 mg kg⁻¹ dry matter) and shoot metal uptake (Pb 6000 and Cd 1200 µg pot⁻¹) under contaminated nutrient solution hydroponic culture (Ullah *et al.*, 2019). Mohanty and Patra (2012) observed significant Cr uptake (i.e. total accumulation rate 8.2 mg kg⁻¹ day⁻¹ and transportation index 6.16) with luxuriant growth and biomass, thereby, suggesting para grass an excellent plant for remediation of heavy metal contaminated wastewater and soil. The TF value refers to the efficiency of plant to transport the metal from root to above ground parts i.e. shoot, leaves. In our experiment, for both metals all grass species showed TF value less than one which shows the lower rate of metal transfer from root to shoot. Similar results were observed by Aran *et al.* (2017) who observed TF values less than

one for different heavy metals (Pb, Cr, Ni, Zn) in *L. laevigatum* growing under wastewater.

Conclusion

It can be concluded on the basis of current findings, among the test grass species that *B. mutica* showed better efficacy regarding metal transfer, accumulation, and uptake with luxuriant growth or biomass. Thereby, *B. mutica* could be a better option to filter out the toxic metals from wastewater.

References

- Adabnejad, H., H.R. Kavousi, H. Hamidi and T. Avassollian. 2015. Assessment of the vacuolar Na⁺/H⁺ antiporter (NHX1) transcriptional changes in *Leptochloa fusca* L. in response to salt and cadmium stresses. *Molecular Biology Research Communications* 4: 133-142.
- Adrover, M., G. Moya and J. Vadell. 2012. Effect of treated wastewater irrigation on plant growth and biological activity in three soil types. *Communications in Soil Science and Plant Analysis* 43:1163-1180.
- Afzal, M., K. Rehman, G. Shabir, R. Tahseen, A. Ijaz, A.J. Hashmat and H. Brix. 2019. Large scale remediation of oil-contaminated water using floating treatment wetlands. *npj Clean Water* doi:10.1038/s41545-018-0025-7.
- Ahmad, F., M.A. Khan, M. Ahmad, M. Zafar, A. Nazir and S.K. Marwat. 2009. Taxonomic studies of grasses and their indigenous uses in the salt range area of Pakistan. *African Journal of Biotechnology* 8: 231-249.
- Ahmad, K., K. Wajid, Z.I. Khan, I. Ugulu, H. Memoona, M. Sana, K. Nawaz, I.S. Malik, H. Bashir and M. Sher. 2019. Evaluation of potential toxic metals accumulation in wheat irrigated with wastewater. *Bulletin of Environmental Contamination and Toxicology* 102: 822-828.
- Akhtar, M.J., S. Ullah, I. Ahmad, A. Rauf, S.M. Nadeem, M.Y. Khan, S. Hussain and L. Bulgariu. 2018. Nickel phytoextraction through bacterial inoculation in *Raphanus sativus*. *Chemosphere* 190: 234-242.
- Ali, H., E. Khan and I. Ilahi. 2019. Environmental chemistry and ecotoxicology of hazardous heavy metals: Environmental persistence, toxicity, and bioaccumulation. *Journal of Chemistry* doi.org/10.1155/2019/6730305.
- Alves, J.C., A.P. Souza, M.L.A. Porto, R.L.F. Fontes, J. Arruda and L.F. Marques. 2016. Potential of sunflower, castor bean, common buckwheat and vetiver as lead phytoaccumulators. *Revista Brasileira de Engenharia Agrícola e Ambiental*. 20: 243-249.
- Aran, D.S., C.A. Harguinteguy, A. Fernandez-Cirelli and M.L. Pignata. 2017. Phytoextraction of Pb, Cr, Ni, and Zn using the aquatic plant *Limnobium laevigatum* and its potential use in the treatment of wastewater. *Environmental Science and Pollution Research* 24:18295-18308.
- Ashraf, M.Y., A.R. Awan and K. Mahmood. 2012. Rehabilitation of saline ecosystems through cultivation of salt tolerant plants. *Pakistan Journal of Botany* 44: 69-75.
- Asrar, H., T. Hussaina, S.M.S. Hadi, B. Gul, B.L. Nielsen and M.A. Khan. 2017. Salinity induced changes in light harvesting and carbon assimilating complexes of *Desmostachya bipinnata* (L.) Staph. *Environmental and Experimental Botany* 135: 86-95.
- Azizullah, A., M.N.K. Khattak, P. Richter and D.P. Hader. 2011. Water pollution in Pakistan and its impact on public health, A review. *Environment International* 37: 479-497.
- Chen, B., S. Tan, Q. Zeng, A. Wang and H. Zheng. 2019. Soil nutrient heterogeneity affects the accumulation and transfer of cadmium in Bermuda grass (*Cynodon dactylon* (L.) pers.). *Chemosphere* 221: 342-348.
- Cheng, S. 2003. Heavy metals in plants and phytoremediation. *Environmental Science and Pollution Research International* 10:335-340.
- Duffus J. H. 2002. "Heavy metals"-a meaningless term. *Pure App. Chemistry*: 74: 793-807.
- Falla, F., D. Dioufa, D. Fall, N. Bakhouma, B. Thioyea, A. Kanea, C. Ndiaye, I. Ndoyea and A.M. Bae. 2017. Growth and physiological responses of *Sporobolus robustus* kunth seedlings to salt stress. *Arid Land Research and Management* 31: 46-56.
- Garbisu, C. and I. Alkorta. 2001. Phytoextraction: A cost-effective plant-based technology for the removal of metals from the environment. *Bioresource Technology* 77: 229-236.
- Ghosh, I., M. Ghosh and A. Mukherjee. 2017. doi.org/10.1007/978-3-319-55426-6_7. Remediation of mine tailings and fly ash dumpsites: Role of poaceae family members and aromatic grasses. p.1-50. In: Enhancing cleanup of environmental pollutants. N. Anjum, S. Gill and N. Tuteja. (eds). Springer International Publishing.
- Gulzar, S., M. Khan and X. Liu. 2007. Seed germination strategies of *Desmostachya bipinnata*: a fodder crop for saline soils. *Rangeland Ecology and Management* 60: 401-407.
- Hall, J.L. 2002. Cellular mechanisms for heavy metal detoxification and tolerance. *Journal of Experimental Botany* 53: 1-11.

- Hussain, A., M. Priyadarshi and S. Dubey. 2019. Experimental study on accumulation of heavy metals in vegetables irrigated with treated wastewater. *Applied Water Science* 9:122.
- Hussain, S., M. Habib-Ur-Rehman, T. Khanam, A. Sheer, Z. Kebin and Y. Jianjun. 2019. Health risk assessment of different heavy metals dissolved in drinking water. *International Journal of Environmental Research and Public Health* 16: 1737.
- Hussain, Z., M. Arslan, M.H. Malik, M. Mohsin, S. Iqbal and M. Afzal. 2018. Treatment of the textile industry effluent in a pilot-scale vertical flow constructed wetland system augmented with bacterial endophytes. *Science of the Total Environment* 645: 966-973.
- Ijaz, A., Z. Iqbal and M. Afzal. 2016. Remediation of sewage and industrial effluent using bacterially assisted floating treatment wetlands vegetated with *Typha domingensis*. *Water Science and Technology* 74: 2192-2201.
- IWMI (International Water Management Institute), 2003. Confronting the realities of wastewater use in agriculture. Water Policy Briefing 9. IWMI, Colombo, Sri Lanka.
- Kadam, S.K., V.V. Chandanshive, N.R. Rane, S.M. Patil, A.R. Gholave, R.V. Khandare, A.R. Bhosale, B.H. Jeon and S.P. Govindwar. 2018. Phytobeds with *Fimbristylis dichotoma* and *Ammannia baccifera* for treatment of real textile effluent: An in-situ treatment, anatomical studies and toxicity evaluation. *Environmental Research* 160: 1-11.
- Kiiskila, J.D., D. Sarkar, S. Panja, S.V. Sahi and R. Datta. 2019. Remediation of acid mine drainage-impacted water by vetiver grass (*Chrysopogon zizanioides*): A multiscale long-term study. *Ecological Engineering* 129: 97-108.
- Kroon, H. 1993. Determination of nitrogen in water: comparison of a continuous-flow method with on-line UV digestion with the original Kjeldahl method. *Analytica Chimica Acta* 276: 287-293.
- Latif, S.A., S. Sharif, S.M. Hossain, M.A. Islam, I.M. Mehedi and M.S. Sultana. 2019. Heavy metal contamination of surface soils in southern part of Bangladesh. *Soil and Environment* 38(1): 112-118.
- Li, H., H. Hao, X. Yang, L. Xiang, F. Zhao, H. Jiang and Z. He. 2012. Purification of refinery wastewater by different perennial grasses growing in a floating bed. *Journal of Plant Nutrition* 35: 93-110.
- Liu, L., W. Li, W. Song, and M. Gou. 2018. Remediation techniques for heavy metal-contaminated soils: Principles and applicability. *Science of the Total Environment* 63: 206-219.
- Lou, Y., P. Zhao, D. Wang, E. Amombo, X. Sun, H. Wang, et al. 2017. Germination, physiological responses and gene expression of tall fescue (*Festuca arundinacea* Schreb.) growing under Pb and Cd. *PLoS ONE* 12: e0169495.
- Majeed, S., S. Rashid, A. Qadir, C. Mackay and F. Hayat. 2018. Spatial patterns of pollutants in water of metropolitan drain in Lahore, Pakistan, using multivariate statistical techniques. *Environmental Monitoring and Assessment* 190: 128.
- Minhas, P.S. and J.S. Samra. 2004. Wastewater use in peri-urban agriculture: impacts and opportunities. Central Soil Salinity Research Institute, Karnal, India. 75 p.
- Miretzky, P., A. Saralegui and A.F. Cirelli. 2006. Simultaneous heavy metal removal mechanism by dead macrophytes. *Chemosphere* 62: 247-254.
- Mitra, A. 2019. Heavy metal status in the lower Gangetic Delta. P. 113-156. In: Estuarine Pollution in the Lower Gangetic Delta. A. Mitra (ed.). Springer, Cham.
- Mohanty, M. and H.K. Patra. 2012. Phytoremediation potential of paragrass-an in-situ approach for chromium contaminated soil. *International Journal of Phytoremediation* 14: 796-805.
- Mushtaq, J. and R. Batool. 2019. Cr (VI) resistant *Bacillus* and *Acinetobacter* isolated from soil of Narran valley. *Soil and Environment* 38(1): 57-65.
- Nair, J., J. Levitan and N. Oyama. 2008. Zinc and copper uptake by silver beet grown in secondary treated effluent. *Bioresource Technology* 99: 2537-2543.
- Najafi, S. and M. Jalali. 2015. Effects of organic acids on cadmium and copper sorption and desorption by two calcareous soils. *Environmental Monitoring and Assessment* 9: 187-585.
- NIAB 2007. Activities at Biosaline Research Station, Pakka Anna. In: Thirty-Five Years of NIAB. Nuclear Institute for Agriculture and Biology (NIAB), Faisalabad, Pakistan.
- Olguin, E.J. and G. Sanchez-Galvan. 2012. Heavy metal removal in phytoremediation and phycoremediation: the need to differentiate between bioadsorption and bioaccumulation. *New Biotechnology*. 30: 3-8.
- Padmavathiamma, P.K. and L.Y. Li. 2007. Phytoremediation technology: Hyper-accumulation. metals in plants. *Water Air Soil Pollution* 184: 105-126.
- Pongthornpruek, S. 2017. Swine farm wastewater treatment by constructed wetland planted with vetiver grass. *Environment and Natural Resources Journal* 15: 13-20.
- Qadir, M., D. Wichelns, L. Raschid-Sally, P.G. McCornick, P. Drechsel, A. Bahri and P.S. Minhas.

2010. The challenges of wastewater irrigation in developing countries. *Agricultural Water Management* 97: 561-568.
- Qureshi, R.H., M. Salim, M. Abdullah and M.G. Pitman. 1982. *Diplachne fusca*: An Australian salt tolerant grass used in Pakistan agriculture. *Journal of Australian Institute and Agriculture Science* 48: 195-199.
- Rai, P.K., S.S. Lee, M. Zhang, Y.F. Tsang and K.H. Kim. 2019. Heavy metals in food crops: Health risks, fate, mechanisms, and management. *Environment International* 125: 365-385.
- Rehman, K., A. Ijaz, M. Arslan and M. Afzal. 2019. Floating treatment wetlands as biological buoyant filters for wastewater reclamation. *International Journal of Phytoremediation* doi.org/10.1080/15226514.2019.1633253.
- Rehman, K., A. Imrana, I. Amina and M. Afzal. 2018. Inoculation with bacteria in floating treatment wetlands positively modulates the phytoremediation of oil field wastewater. *Journal of Hazardous Materials* 349: 242-251.
- Rev, A., B. Toth, A. Solti, G. Sipos and F. Fodor. 2017. Responses of Szarvasi-1 energy grass to sewage sludge treatments in hydroponics. *Plant Physiology and Biochemistry* 118: 627-633.
- Richards, L.A. 1954. U.S. Salinity Laboratory Staff. Diagnosis and Improvement of Saline and Alkali Soils. Agricultural handbook 60. U.S. Dept. of Agriculture, Washington D.C., 160 p.
- Riehm, H. and B. Ulrich. 1954. Quantitative colorimetrische Bestimmung der organischen Substanz im Boden. *Landwirtsch Forsch* 6: 173-176.
- Roongtanakiat, N., S. Tangruangkiat and R. Meesat. 2007. Utilization of vetiver grass (*Vetiveria zizanioides*) for removal of heavy metals from industrial wastewaters. *Science Asia* 33: 397-403.
- Salman, S.A., A.A. Asmoay, A. El-Gohary and H. Sabet. 2019. Evaluation of human risks of surface water and groundwater contaminated with Cd and Pb in the southern El-Minya Governorate, Egypt. *Drinking Water Engineering and Science* 12: 23-30.
- Sarwar, N., Saifullah, S.S. Malhi, M.H. Zia^{d,e}, A. Naeem, S. Bibia and G. Farid. 2010. Role of mineral nutrition in minimizing cadmium accumulation by plants. *Journal of Science Food and Agriculture* 90: 925-937.
- Sarwar, N., M. Imran, M.R. Shaheen, W. Ishaque, M.A. Kamran, A. Matloob, A. Rehim and S. Hussain. 2017. Phytoremediation strategies for soils contaminated with heavy metals: Modifications and future perspectives. *Chemosphere* 171:710-721.
- Shi, Y. and A. Chatt. 2014. Simultaneous determination of inorganic, As (III), As (V), Sb (III), Sb(V), and Se (IV) species in natural waters by APDC/MIBK-NAA. *Journal of Radioanalytical Nuclear Chemistry* 299: 867-877.
- Sial, R.A., M.F. Chaudhary, S.T. Abbas, M.I. Latif and A.G. Khan. 2006. Quality of effluents from Hattar Industrial Estate. *Journal of Zhejiang University Science B* 7: 974-80.
- Silvaa, E.B., F.G. Fonseca, L.R.F. Alleoni, S.S. Nascimento, P.H. Graziottia and B.O. Nardisa. 2016. Availability and toxicity of cadmium to forage grasses grown in contaminated soil. *International Journal of Phytoremediation* 18: 847-852.
- Singh, D., A. Tiwari and R. Gupta. 2012. Phytoremediation of lead from wastewater using aquatic plants. *Journal of Agricultural Technology* 8: 1-11.
- Singh, O.V., S. Labana, G. Pandey, R. Budhiraja and R.K. Jain. 2003. Phytoremediation: An overview of metallic ion decontamination from soil. *Applied Microbiology and Biotechnology* 61: 405-412.
- Soda, S., T. Hamada, Y. Yamaoka, M. Ike, H. Nakazato, Y. Saeki, T. Kasamatsu and Y. Sakurai. 2012. Constructed wetlands for advanced treatment of wastewater with a complex matrix from a metal-processing plant: Bioconcentration and translocation factors of various metals in *Acorus gramineus* and *Cyperus alternifolius*. *Ecological Engineering* 39: 63-70.
- Svetlana, S., A.A. Tinkov and A.V. Skalny. 2019. Hair toxic metal and metalloid levels in children with chronic sinusitis. *Journal of Elementology* 24: 1091-1100.
- Tak, H.I., O.O. Babalola, M.H. Huyser and A. Inam. 2013. Urban wastewater irrigation and its effect on growth, photosynthesis and yield of chickpea under different doses of potassium. *Soil Science and Plant Nutrition* 59: 156-167.
- Ullah, S., T. Mahmood, Z. Iqbal, A. Naeem, R. Ali and S. Mahmood. 2019. Phytoremediative potential of salt-tolerant grass species for cadmium and lead under contaminated nutrient solution. *International Journal of Phytoremediation* doi.10.1080/15226514.2019.1594683.
- Watanabe, F.S. and S.R. Olsen. 1965. Test of ascorbic acid method for determining phosphorus in water and NaHCO₃ extract from soils. *Soil Science Society of American Journal* 29: 677-678.
- WHO. 1997. Health and environment in sustainable development. Geneva: WHO.

- Wolf, B. 1982. A comprehensive system of leaf analyses and its use for diagnosing crop nutrient status. *Communications in Soil Science and Plant Analysis* 13:1035-59.
- Yobi, A., K.A. Schlauch, R.L. Tillett, W.C. Yim, C. Espinoza, B.W.M. Wone, J.C. Cushman and M.J. Oliver. 2017. *Sporobolus stapfianus*: Insights into desiccation tolerance in the resurrection grasses from linking transcriptomics to metabolomics. *BMC Plant Biology* 17:67.
- Zheng, H., Z. Zhang, X. Xing, T. Hu, C. Qu, W. Chen and J. Zhang. 2019. Potentially toxic metals in soil and dominant plants from tonglushan Cu-Fe deposit, Central China. *Bulletin of Environmental Contamination and Toxicology* 102: 92-97.
- Zhivotovsky, O.P., J.A. Kuzovkina, C.P. Schulthess, T. Morris, D. Pettinelli, D. and M. Ge. 2011. Hydroponic screening of willows (*Salix* L.) for lead tolerance and accumulation. *International Journal of Phytoremediation* 13: 75-94.
- Zhou, M, W. Tian, J. Zhang, X. Chen, Y. Wu and S. Wang. 2019. A rapid on-site analysis method for the simultaneous extraction and determination of Pb^{2+} and Cd^{2+} in cereals. *RSC Advances* doi: 10.1039/c9ra05587h.
- Zou, J., X. Shang, C. Li, J. Ouyang, B. Li and X. Liu. 2019. Effects of cadmium on mineral metabolism and antioxidant enzyme activities in *Salix matsudana* Koidz. *Polish Journal of Environmental Studies* 28: 989-999.