RESEARCH ARTICLE

VERNALIZATION AND GIBBERELIC ACID RESPONSE IN SUMMER ONION'S (*ALLIUM CEPA* L.) REPRODUCTIVE PHASES

Lamia Khatun^a, Md. Rezaul Karim^b, Fakhar Uddin Talukder^c, Md. Sohanur Rahman^{c*}, Md. Israfil Jahan^d

^a Department of Horticulture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

^b Professor, Department of Horticulture, Bangladesh Agricultural University, Mymensingh – 2202, Bangladesh.

^c Scientific Officer, Pest Management Division, Bangladesh Jute Research Institute, Manik Mia Avenue, Dhaka-1207, Bangladesh.

^d Department of Seed Science and Technology, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

*Corresponding author email: sohanbau2010@gmail.com

This is an open access article distributed under the Creative Commons Attribution License CC BY 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ARTICLE DETAILS

Article History:

Received 22 May 2020

Accepted 23 June 2020

Available online 06 July 2020

ABSTRACT

The present study was aimed at determining the effect of vernalization and Gibberellic Acid (GA₃) on seed yield and quality of summer onion. There were three vernalization treatments viz., no vernalization (control), vernalization at 5°C for 14 days and vernalization at 10°C for 14 days and four GA₃ treatment viz., 0, 50, 100 and 150 ppm. The two-factor experiment was laid out in the Randomized Complete Block Design with three replications. The results of the experiment showed that the vernalization and GA₃ had significant influence on all the parameters studied. Solitary Effect of vernalization and GA₃ was significant on the parameters such as plant height, number of leaves plant⁻¹, the highest number of flowering stalk, number of umbels plant⁻¹, number of bud umbel⁻¹, percent flowering at 45 and 60 DAP, number of seeds umbel⁻¹, weight of seeds umbel⁻¹, weight of seeds plant⁻¹, weight of seeds plot⁻¹, 1000 seed weight, seed yield, number of fruits umbel⁻¹, percent of fruit set umbel⁻¹ and percent germination. The highest seed yield (248.16 kg ha⁻¹) was obtained from vernalization at 5°C for 14 days. The lowest values of all the parameters were recorded at control treatment. The highest seed yield (191.77 kg ha⁻¹) was obtained from 100 ppm GA₃. The lowest values of all the parameters were recorded in the 0 ppm GA₃ treatment. Use of suitable vernalization of mother bulb and proper concentration of gibberellic acid can be a possible way to expand onion production at the time of summer.

KEYWORDS

vernalization, gibberellic acid, onion, reproductive phases.

1. INTRODUCTION

Onion (*Allium cepa* L.) is one of the most important spice crops in Bangladesh. Usually onion is grown in Bangladesh in the winter season. One possibility to expand onion production is to cultivate onion during the summer. There are still some problems to cultivate and store summer onion. One of the most important problems is that the unavailability of good quality seeds due to lack of information of producing summer onion seed production. Many crops e.g. onion grown in temperate climates require vernalization and must introduce a period of low winter temperature to initiate or accelerate the flowering process. This assures that reproductive development and seed production happens in spring and winters, rather than in autumn. Typical vernalization temperatures are between 1 to 7 degrees Celsius (34- and 45-degrees Fahrenheit) (Taiz and Murphy, 2015). Vernalization of bulb are required for flower initiation in onion (Brewster, 1994).

Suitable vernalization temperature of mother bulb incites early flowering and produces better yield of seed (Jones and Mann, 1963). Different growth phases of onion have different environmental requirements. Onion bulbs require a certain period of cool treatment (vernalization) for maximum flowering of the stalks. All the plants raised from bulbs do not

produce flower stalks due to improper vernalization. Vernalization of bulb prior to planting ensures effective flowering of seed crop (Brewster, 1994). Vernalization is a suitable process required for some definite plant species, e.g. onion (*Allium cepa* L.), to start the reproductive stage, through an introduce to low but non-freezing temperatures (Streck, 2003). Onion is herbaceous biennial plant, and temperature is a major environmental factor that control its development (Brewster, 1983, 1987; Rabinowitch, 1985). Onion developmental models has a temperature function (Brewster, 1987).

The expression of vegetative plants (after a juvenile stage that corresponds to a minimum 7-10 initiated leaves) or bulbs to low temperatures conduces flowering, which terminates the growth of the main axis (Rabinowitch, 1985). The outpouring to low temperatures either in natural winter or in artificial cold treatment that brings about induction of flowering in plant species (e.g. onion) is called vernalization (Pinthus, 1985; Flood and Halloran, 1986). Vernalization in onion is an important research interest because of the need to induce flowering for seed production for breeding purposes in the shortest period of time (Brewster, 1987). Lack of cold temperatures to induce flowering is the prime constrain to onion seed production in tropical countries (Kimani et al., 1994). The reaction of plants to vernalization is occurred by the

Quick Response Code

Access this article online

Website:
www.taec.com.my

DOI:
10.26480/taec.01.2020.07.14

combination of 2 factors, the duration of the vernalization period and the temperature during the vernalization period (Hodges and Ritchie, 1991; Ritchie, 1991).

Brewster processed data every day vernalization rate to temperature with a stepwise linear function (Brewster, 1987). A step-wise function is not the most suitable response work as a temperature response function because biological systems seems to respond to temperature, in a continuous and smooth fashion rather than in a combination of linear functions which introduce sudden changes in the response (Shaykewich, 1995; Streck, 2002a). In case of onion, short duration cold treatment of 10 °C for 10 days enhances the seed yield (Yalamalle, 2016). The inoculation of mycorrhizae and the application of humic substances can enhance bulb growth and quality (Sekara et al., 2017). Proper vernalization temperature of mother onion bulbs enhance early flowering and produces a higher yield of onion seed (Khokhar, 2014).

Plant growth regulators are considered as key factors in vegetative growth, flowering, fruit setting and seed production in plant. It is revealed from experiments conducted in various part of the world that IAA, NAA, GA₃, BA, 2, 4-D, IBA and ethrel at various concentrations have remarkable effects on number of flowers per umbel, umbel size and quality seed production of onion (El-Habbasha et al., 1985). GA₃ application brings metabolic changes that affect both quality and quantity of the desired product. It stimulates the synthesis of hydrolic enzymes which are secreted and act on starchy endosperm in turn affecting physiology of seed germination and establishment of seedlings. Gibberellins are one of the major regulators of plant growth and development which suppress the growth and promoting cell elongation and division (Olszewski et al., 2002; Tomas et al., 2009).

It is well known that GA₃ enhances plant growth and its accompanying metabolite production (Jones et al., 2009). The applications of gibberellic acid (GA₃) on plants promotes plant height and, secondarily, increase dry weight (Sharma et al., 1999). The effect of GA₃ on onion seed production is characterized to increasing in number of inflorescences per plant, and synchronization of pistillate & staminate patterns (Corgan and Montano, 1975). Medium or large bulbs sprayed with Gibberellic acid at 50ppm provide significantly higher seed yield per hectare, germination and vigor (Shaikh et al., 2002). Spraying of GA₃ on plants increased seed yield and quality parameters compared with controls (Rizk et al., 1996). Research work on effects of vernalization and GA₃ on seed production is scanty in Bangladesh specially in summer season. The present research work was undertaken with the following objectives:

1. To find out optimum temperature for vernalization for shortening its life cycle as well as increasing production of high-quality summer onion seed.
2. To identify optimum dose of GA₃ for the production of summer onion seed.

2. MATERIALS AND METHODS

2.1 Experimental site, Soil, Climate

The present research work was conducted at the Horticultural Farm, Bangladesh Agricultural University, Mymensingh during the period from November 2017 to April 2018. The experimental area is located at 24°46'N latitude and 90°24'E longitude. The experimental area belongs to Old Brahmaputra Flood Plain under the AEZ number-9 of Bangladesh. The field was above inundation level and texturally silty loam with a pH value 6.8. The climate of the experimental area was sub-tropical, characterized by heavy and scanty rainfall.

2.2 Experimental material

The bulbs (Different size graded) of summer onion variety BARI Piaz-3 were used in this study. These seed bulbs were collected from Bangladesh Agricultural Research Institute, Gazipur.

2.3 Experimental treatments

Vernalization: the cooling of crop seed during germination for accelerating flowering when it is planted.

Gibberellic acid (GA₃): it is a hormone found in plants and fungi. Its chemical formula is C₁₉H₂₂O₆. When purified, it is a white to pale-yellow solid.

The experiment had two factors. The levels of each factor were Factor A: Vernalization

V₀ =No vernalization, V₁ =Vernalization at 5°C for 14 days, V₂ =Vernalization at 10°C for 14 days

Factor B: Concentrations of GA₃

G₀ = 0 ppm, G₁ = 50 ppm, G₂ = 100 ppm, G₃ = 150 ppm

2.4 Design and layout of the experiment

The two-factor experiment consisting of twelve treatment combinations was laid out in the Randomized Complete Block Design (RCBD) with three replications. Thus, in total 36-unit plots were obtained. The total area of this experiment was divided into three blocks and each blocks contained twelve plots. The treatment combinations were randomly placed to unit plots in each blocks. The size of each unit plot was 1m × 1m. The distance between the blocks was 75 cm and between the plots was 50 cm.

2.5 Vernalization of mother bulbs

Selected bulbs of the same size were put in white cotton cloth bags. Bulbs were then vernalized for 14 days in two separate refrigerators calibrated at 5 °C and 10 °C.

2.6 Planting of bulbs in the experimental plot

Bulbs were taken out from the refrigerators four hours prior to planting and then kept under ceiling fan for surface drying. The distance maintained between the row was 25 cm and between the plant was 20 cm.

2.7 Preparation of GA₃ solution and its application

Laboratory grade reagents were used to prepare 50, 100 and 150 ppm of GA₃ solution. The stock solution of 1000 ppm of GA₃ was prepared by protein 1.0 gm of GA₃ into a conical flask. Then a little amount of ethyl alcohol (C₂H₅OH) was added into the conical flask. Then the flask was stirred for some time and exact volume in the volumetric flask was made up by adding distilled water with frequent stirring to have the desired 1L stock solution. The stock solution was then preserved in glass jars at low temperature in a refrigerator for preparing solutions of desired concentrations. To prepare three concentrations of GA₃ solution of 50,100 and 150 ppm; at first 50,100- and 150-ml solution had taken from GA₃ stock solution (1000 ppm) and then were diluted to prepare 1 liter volume with the required amount of distilled water. The prepared solutions were sprayed over the plants in two times one at 30 and 50 DAP. The solutions were sprayed as foliar spray with the help of hand sprayer.

2.8 Harvesting

Matured seed umbels were harvested in several installments when 10-20% of the capsules were split and exposed their black seeds. Umbels were harvested with a small portion of flowering stalk in the morning to prevent shattering of seed.

2.9 Postharvest operation

Harvested umbels were dried in open sunlight on brown paper for 4-5 days. Threshing was done manually, seed were cleaned and were dried again until they reached safe moisture content (Brewster, 1994). Seeds were then kept in polythene bag, which were kept in refrigerator at 8±10°C.

2.10 Germination test

Germination percentage of onion seed was recorded in the Postgraduate Laboratory of the Department of Horticulture, BAU, Mymensingh at ambient room temperature immediately after drying seeds with 3 replications. For each treatment 75 seeds were taken for each replication. It was performed following TP (top of paper) method and in sterile petridish (90 mm diameter). In each petridish 25 seeds were placed on three layer of tissue paper saturated with water.

2.11 Data collection

To avoid border effect with the highest precision six plants were selected randomly from each plot out of 30 discarding the outer two rows and outer plants of the middle lines.

2.11.1 Plant height

The height of plants was measured from ground level to the tip of the longest leaf. Average plant height was determined from selected plants at 20, 40 and 60 DAP in centimeter (cm).

2.11.2 Number of leaves plant⁻¹

Number of leaves including green and dry from six selected plants were recorded at 20, 40 and 60 DAP.

2.11.3 Height of flowering stalk

It was measured from ground level to the tip of the scape before harvest, and the average was recorded in centimeter (cm) from selected plants.

2.11.4 Number of umbels plant⁻¹

Average number of umbels per plant was recorded from randomly selected plants at full bloom stage.

2.11.5 Number of fruits umbel⁻¹

Average number of seeded fruits per umbel was computed from randomly selected six sampled umbels, after harvesting.

2.11.6 Number of bud umbel⁻¹

Average number of buds per umbel was computed from randomly selected six sampled umbels at maximum flowering stage and the average number was recorded.

2.11.7 Percent flowering plants

The number of plants flowered was conducted at 5 days interval from initiation to completion.

$$\text{Percent flowering plants} = \frac{\text{No. of plants flowered}}{\text{Total no. of plants in the plot flowered}} \times 100$$

2.11.8 Number of seeds umbel⁻¹

Average number of seeds per umbel was computed from randomly selected six sampled umbels at maximum flowering stage and the average number was recorded.

2.11.9 Weight of seeds umbel⁻¹

Average weight of seed per umbel was determined in gram (g) from randomly selected 6 umbels.

2.11.10 Weight of seeds plant⁻¹

It was recorded from randomly selected 6 plants and was expressed in gram (g).

2.11.11 Weight of seeds plot⁻¹

Total amount of seeds for each plot were bulked and weighted in gram (g).

2.11.12 Percent of fruit set umbel⁻¹

The average number of seeded fruits per umbel was recorded from randomly selected 6 sampled umbels after harvesting. Then it was counted per plot. Seeded fruits divided by no. of flowers per umbel were calculated as % fruit set.

2.11.13 Number of umbels plot⁻¹

Average number of umbels per plot was determined from previously selected six plants.

2.11.14 Yield of seeds plant⁻¹

The seed were collected, weighed and recorded from randomly selected six plants and the mean weight was expressed in gram (g).

2.11.15 Yield of seeds plot⁻¹

Total amount of seeds for each plot was bulked, was weighed and was expressed in gram (g).

2.11.16 Seed yield (kg ha⁻¹)

It was determined from conversion of the respective seed yield per plot to ha and was expressed in kg per ha.

2.11.17 1000 seed weight

For each plot 1000 seeds were counted and weighed by electric balance and was recorded in gram (g).

2.11.18 Germination percentage

The number of seeds germinated was recorded respectively for each petridish daily. Seeds were considered to have germinated when radical emerged about 1 mm from the seed.

$$\text{Germination percentage} = \frac{\text{No. of seeds germinated}}{\text{No. of seeds set for germination}} \times 100$$

2.12 Statistical analysis

The collected data were statistically analyzed by the computer using statistical package programme MSTAT-C. The means for all the treatments were calculated and the analysis of variance was performed by F variance test. The pair of comparisons were performed by Least Significant Difference (LSD) test at 5% and 1% level of probability (Gomez and Gomez, 1984).

3. RESULTS

The effects of vernalization and GA₃ on seed production of onion have been presented in Tables and Figures. Results of different parameters studied in this experiment have been presented and discussed under the following headlines.

3.1 Main effect of vernalization on plant height

Vernalization significantly affected plant height at 20 DAP, 40 DAP and 60 DAP. At 60 DAP the tallest plant (47.72 cm) was obtained from the bulbs that were vernalized at 5°C for 14 days (V₁) period to planting (Figure 1). The shortest plant (45.82 cm) for the same was recorded from the bulbs having no vernalization treatment (V₀).

Figure 1: Effect of vernalization on plant height of onion. Vertical bars indicate LSD at 1% level of probability. V₀, V₁, and V₂ indicates no vernalization, vernalization at 5°C and 10°C for 14 days respectively.

3.2 Main effect of GA₃ on plant height

GA₃ had significantly affected plant height at 20 DAP, 40 DAP and 60 DAP. At 60 DAP the tallest plant (50.78 cm) was obtained from the bulbs that were GA₃ at 100 ppm (G₂) (Figure 2). The shortest plant (41.66 cm) for the same was recorded from the bulbs having no GA₃ treatment (G₀).

Figure 2: Effect of GA₃ on plant height of onion. Vertical bars indicate LSD at 1% level of probability. G₀, G₁, G₂, G₃ indicates 0, 50, 100 and 150 ppm

respectively.

3.3 Main effect of vernalization on number of leaves plant⁻¹

Duration of vernalization significantly influenced the number of green leaves at 20 DAP, 40 DAP and 60 DAP. Plant produced from the bulbs vernalized at 5°C for 14 days prior to planting gave maximum leaves at 60 DAP (30.38 leaves plant⁻¹). The minimum number (20.79 green leaves plant⁻¹) were produced by the control treatment (Figure 3).

Figure 3: Effect of vernalization on number of leaves of onion. Vertical bars indicate LSD at 1% level of probability. V₀, V₁, and V₂ indicate no vernalization, vernalization at 5°C and 10°C for 14 days respectively

3.4 Main effect of GA₃ on number of leaves plant⁻¹

Significant variation was found in increased number of leaves plant⁻¹ due to the effect of GA₃. The maximum number (29.67) of leaves plant⁻¹ was obtained from 100 ppm (G₂) which was identically followed by 150 ppm (28.28) at 60 DAP. The control treatment gave the minimum number of leaves (19.50) plant⁻¹ at the same DAP (Figure 4).

Figure 4: Effect of GA₃ on number of leaves of onion. Vertical bars indicate LSD at 1% level of probability. G₀, G₁, G₂, G₃ indicates 0, 50, 100 and 150 ppm respectively.

3.5 Main effect of vernalization on percent germination of summer onion

Duration of vernalization significantly influenced the percent germination of summer onion. Plant produced from the bulbs vernalized at 5°C for 14 days prior to planting gave maximum germination at (61.50 percent germination). The minimum germination (54.50 percent germination) were produced by the control treatment (Table 1).

Vernalization	Germination (%)
V ₀	54.50
V ₁	61.50
V ₂	56.50
LSD _{0.05}	0.69
LSD _{0.01}	0.94
Level of significance	156.00**

** = Significant at 1% level of probability. Here V₀, V₁, and V₂ indicates no vernalization, vernalization at 5°C and 10°C for 14 days respectively.

3.6 Main effect of GA₃ on percent germination of summer onion

Significant variation was found in increased percent germination due to the effect of GA₃. The maximum germination (65.33) was obtained from 100 ppm (G₂) which was identically followed by 150 ppm (58.00). The control treatment gave the minimum germination (50.67) at the same DAP (Table 2).

GA ₃ concentration	Germination (%)
G ₀	50.67
G ₁	56.00
G ₂	65.33
G ₃	58.00
LSD _{0.05}	0.80
LSD _{0.01}	1.09
Level of significance	331.67**

** = Significant at 1% level of probability. G₀, G₁, G₂, G₃ indicates 0, 50, 100 and 150 ppm respectively

3.7 Effect of vernalization on flowering characteristics at 60 DAP

There was a significant variation in respect of height of flowering stalk, highest percent flowering, number of umbels plant⁻¹, number of flowers plant⁻¹, number of seeds umbel⁻¹, number of fruits umbel⁻¹ at 60 DAP due to different levels of vernalization. The longest flowering stalk (78.95cm) was found in the plots which received 5°C vernalization for 14 days while the shortest (66.95cm) was recorded in the plant that received no vernalization. The highest percent flowering (79.60) was found in the plots which received 5°C vernalization for 14 days while the lowest (73.52) was recorded in the plant that received no vernalization. The maximum number of umbels plant⁻¹, number of flowers umbel⁻¹ (6.71, 236.89) was found in the plant that received vernalization at 5°C for 14 days and minimum (5.10, 197.56) was observed respectively in the plant that received no vernalization. The maximum number of seeds umbel⁻¹, number of fruits umbel⁻¹ (273.39, 88.24) was found in the plant that received vernalization at 5°C for 14 days and minimum (214.62, 77.46) was observed respectively in the plant that received no vernalization (Table 3).

Vernalization	Height of flowering stalk	Highest percent flowering	No. of umbels/plant	No. of flowers/umbel	No. of seeds/umbel	No. of fruits/umbel
V ₀	66.95	73.52	5.10	197.56	214.62	77.46
V ₁	78.95	79.60	6.71	236.89	273.39	88.24
V ₂	74.99	76.21	5.74	211.59	246.09	82.37
LSD _{0.05}	0.429	0.276	0.089	3.45	4.45	0.94
LSD _{0.01}	0.583	0.375	0.121	4.69	6.05	1.27
Level of significance	448.41**	111.09**	7.87**	4766.09**	10377.28**	349.21**

** = Significant at 1% level of probability. V₀, V₁, and V₂ indicates no vernalization, vernalization at 5°C and 10°C for 14 days respectively.

3.8 Effect of GA₃ on flowering characteristics at 60 DAP

There was a significant variation in respect of height of flowering stalk, highest percent flowering, number of umbels plant⁻¹, number of flowers plant⁻¹, number of seeds umbel⁻¹, number of fruits umbel⁻¹ at 60 DAP due to different levels of GA₃. The longest flowering stalk (79.97cm) was found

in the plots which received 100 ppm GA₃ while the shortest (67.67cm) was recorded in the plant that received no GA₃. The highest percent flowering (79.20) was found in the plots which received 100 ppm GA₃ while the lowest (74.03) was recorded in the plant that received no GA₃. The maximum number of umbels plant⁻¹, number of flowers umbel⁻¹ (6.87, 241.27) was found in the plant that received 100 ppm GA₃ and minimum

(4.67, 186.72) was observed respectively in the plant that received no GA₃. The maximum number of seeds umbel⁻¹, number of fruits umbel⁻¹ (275.47, 92.06) was found in the plant that received 100 ppm GA₃ and minimum

(216.75, 71.73) was observed respectively in the plant that received no GA₃ (Table 4).

Table 4: Effect of GA₃ concentration on flowering characteristics of summer onion

GA ₃ concentration	Height of flowering stalk	Highest percent flowering	No. of umbels/plant	No. of flowers/umbel	No. of seeds/umbel	No. of fruits/umbel
G ₀	67.67	74.03	4.67	186.72	216.75	71.73
G ₁	71.93	75.68	5.54	209.89	233.93	81.00
G ₂	79.97	79.20	6.87	241.27	275.47	92.06
G ₃	74.94	76.85	6.33	223.50	252.64	85.96
LSD _{0.05}	0.496	0.318	0.103	3.98	5.14	1.08
LSD _{0.01}	0.674	0.433	0.139	5.41	6.98	1.47
Level of significance	241.21**	42.46**	8.28**	4763.27**	5720.88**	664.36**

** = Significant at 1% level of probability. G₀, G₁, G₂, G₃ indicate 0, 50, 100 and 150 ppm respectively.

3.9 Effect of vernalization on fruit & seed production of summer onion at 60 DAP

There was a significant variation in respect of number of fruits plant⁻¹, percent fruits, weight of seeds umbel⁻¹, 1000 seeds weight, weight of seeds plant⁻¹, weight of seeds plot⁻¹ and seed yield due to different levels of vernalization. The highest number of fruits plant⁻¹ (600.03) was found in the plots which received 5°C vernalization for 14 days while the shortest (399.62) was recorded in the plant that received no vernalization. The highest percent fruits set (40.84) was found in the plots which received

5°C vernalization for 14 days while the lowest (38.94) was recorded in the plant that received no vernalization. The maximum weight of seeds umbel⁻¹, 1000 seeds weight (1.08, 2.67) was found in the plant that received vernalization at 5°C for 14 days and minimum (0.42, 2.33) was observed respectively in the plant that received no vernalization. The maximum weight of seeds plot⁻¹ and seed yield (59.01, 248.16) was found in the plant that received vernalization at 5°C for 14 days and minimum (23.10, 97.15) was observed respectively in the plant that received no vernalization (Table 5).

Table 5: Effect of vernalization on fruit & seed production of summer onion

Vernalization	No. of fruits/ plant	% Fruit set	Weight of seeds/ umbel	1000 seed wt (g)	Weight of seeds/ plant (g)	Weight of seeds/ plot (g)
V ₀	399.62	38.94	0.42	2.33	2.53	23.10
V ₁	600.03	40.84	1.08	2.67	6.45	59.01
V ₂	478.73	39.42	0.67	2.43	4.04	36.91
LSD _{0.05}	10.31	0.235	0.027	0.038	0.076	1.48
LSD _{0.01}	14.01	0.319	0.036	0.051	0.103	2.01
Level of significance	122278.27**	11.617**	1.313**	0.363**	47.09**	18838.52**

** = Significant at 1% level of probability. V₀, V₁, and V₂ indicates no vernalization, vernalization at 5°C and 10°C for 14 days respectively.

3.10 Effect of GA₃ on fruit & seed production at 60 DAP

There was a significant variation in respect of number of fruits plant⁻¹, percent fruits, weight of seeds umbel⁻¹, 1000 seeds weight, weight of seeds plant⁻¹, weight of seeds plot⁻¹ and seed yield due to different levels of GA₃. The highest number of fruits plant⁻¹ (636.17) was found in the plots which received 100 ppm GA₃ while the shortest (335.96) was recorded in the plant that received no GA₃. The highest percent fruits set (40.54) was found in the plots which received 100 ppm GA₃ while the lowest (38.81) was recorded in the plant that received no GA₃. The maximum weight of seeds umbel⁻¹, 1000 seeds weight (0.80, 2.82) was found in the plant that received 100 ppm GA₃ and minimum (0.61, 2.18) was observed respectively in the plant that received no GA₃. The maximum weight of seeds plot⁻¹ and seed yield (45.60, 191.77) was found in the plant that received 100 ppm GA₃ and minimum (33.58, 141.24) was observed respectively in the plant that received no GA₃ (Table 6).

Table 6: Effect of GA₃ concentration on fruit & seed production of summer onion

GA ₃ concentration	No. of fruits/ plant	% Fruit set	Weight of seeds/ umbel	1000 seed wt (g)	Weight of seeds/ plant (g)	Weight of seeds/ plot (g)
G ₀	335.96	38.81	0.61	2.18	3.67	33.58
G ₁	452.27	39.42	0.69	2.36	4.11	37.61
G ₂	636.17	40.54	0.80	2.82	4.99	45.60
G ₃	546.78	40.16	0.79	2.54	4.58	41.90
LSD _{0.05}	11.90	0.271	0.031	0.044	0.087	1.71
LSD _{0.01}	16.18	0.369	0.042	0.059	0.119	2.32
Level of significance	149133.08**	5.29**	0.073*	0.666*	2.68**	1073.28**

** = Significant at 1% level of probability. G₀, G₁, G₂, G₃ indicates 0 ppm, 50 ppm, 100 ppm and 150 ppm respectively.

3.11 Main effect of vernalization on seed yield

Vernalization had significant effect on seed yield. The highest seed yield (248.16 kg ha⁻¹) was found in the plots which received 5°C vernalization for 14 days while the lowest seed yield (97.15 kg ha⁻¹) was recorded in the plant that received no vernalization. (Figure 5).

Figure 5: Effect of vernalization on seed yield of onion. Vertical bar indicates LSD at 1% level of probability. V₀, V₁, and V₂ indicates no vernalization, vernalization at 5°C and 10°C for 14 days respectively.

3.12 Main effect of GA₃ on seed yield

GA₃ had significant effect on seed yield. The highest seed yield (191.77 kg ha⁻¹) was found in the plots which received 100 ppm GA₃ while the lowest seed yield (141.24 kg ha⁻¹) was recorded in the plant that received no GA₃. (Figure 6).

Figure 6: Effect of GA₃ on seed yield of onion. Vertical bar indicates LSD at 1% level of probability. G₀, G₁, G₂, G₃ indicate 0 ppm, 50 ppm, 100 ppm and 150 ppm respectively.

4. DISCUSSION

Vernalization treatment of 10 °C for 10 days is effective in enhancing the seed yield in onion (Yalamalle, 2016). Low temperatures exposure (in natural or artificial conditions) that causes induction of flowering is called vernalization (Lee et al., 2013; Thomas and Vince-Prue, 1997). Due to incomplete vernalization seed yield will be poor (Voss et al., 2013). A group researchers noticed that the higher activity of soluble invertase, glucose, sucrose and fructose levels in onion bulbs stored at 10 °C than those stored at 20 °C (Benkeblia et al., 2004). Vernalization helped in conversion of complex sugars in the bulbs to simple sugars which were deliberately available for transition to the reproductive growth. These resulted in increase in number of flowers and increase of seed yield subsequently (Yalamalle, 2016). Optimum duration to completely vernalize onion cultivars at low temperatures (3 to 11 °C) are in the range of 7 to 90 days (Woodbury, 1950; Kruzhilin and Shvedskaya, 1962; Rabinowitch, 1985; Pike, 1986; Bertaud, 1988; Peters, 1990; Rabinowitch, 1990).

In case of onion, a comparison of the cardinal temperatures for vernalization (0 °C, 10 °C, 16 °C) with other species gives some similarities and differences. The cardinal temperatures for vernalization in carrot are -1, 6.5, and 16 °C, in lily are 0, 5, and 21 °C, in calabrese are -2.8, 15.8, and 23.6 °C, and in winter wheat are -1.3, 4.9, and 15.7 °C (Porter and Gawith, 1999; Atherthon et al., 1990; Roh and Wilkins, 1977; Roh, 1985; Holcomb and Berghage, 2001; Wurr et al., 1995). All the treatments show maximum plant height at 60 DAP. Vernalization at 5 °C for 14 days (V₁) gave the tallest plant (47.72 cm) while vernalization treatment V₀ (no vernalization) gave the shortest (45.82 cm) plant. On the other hand, the tallest plant height (50.78 cm) was obtained with the treatment of G₂ (100 ppm) and the shortest plant height (41.66 cm) was observed with the treatment of G₀ (control). Streck, Nereu A 2003 observed a maximum vernalization response of various onion cultivars in the range of 9-12 °C and a reduction when temperature departs from optimum temperature.

The number of leaves plant⁻¹ gradually increased with the increasing time after planting. Single effect of vernalization on leaf production was significantly affected by the selected treatments. Vernalization at 5 °C for 14 days treatment produced the maximum number of leaves (30.38) at 60 DAP while the minimum (20.79) number of leaves was produced with control treatment. On the other hand, maximum number of leaves (29.67) was observed in the treatment of 100 ppm (G₂) and minimum number of leaves (19.50) was produced with control treatment. The emergence and differentiation of scape depends on temperature (Zemah et al., 2001). Duration of vernalization significantly influenced the number of green leaves at 20 DAP, 40 DAP and 60 DAP. Plant produced from the bulbs vernalized at 5 °C for 14 days prior to planting gave maximum germination at (61.50 percent germination). The minimum germination (54.50 percent germination) was produced by the control treatment.

Significant variation was found in increased percent germination due to the effect of GA₃. The maximum germination (65.33) was obtained from 100 ppm (G₂) which was identically followed by 150 ppm (58.00) at 60 DAP. The control treatment gave the minimum germination (50.67) at the same DAP. Some researchers prove that the cold treatment enhances the

scape emergence (Muthamia, 1994; Khokhar, 2009). Vernalization treatment mostly alters the source sink relation in favor of reproductive growth (Linwattana et al., 1997). According to a study at the time of thermo phase the onion seed crop apart from temperature, high concentration of soluble carbohydrates is an important factor enhancing flowering in onion and bulbing of *Alliums* (Brewster, 1994). There was a significant variation in respect of height of flowering stalk, highest percent flowering, number of umbels plant⁻¹, number of flowers plant⁻¹, number of seeds umbel⁻¹, number of fruits umbel⁻¹ at 60 DAP due to different levels of vernalization.

The longest flowering stalk (78.95 cm) was found in the plots which received 5 °C vernalization for 14 days while the shortest (66.95 cm) was recorded in the plant that received no vernalization. The highest percent flowering (79.60) was found in the plots which received 5 °C vernalization for 14 days while the lowest (73.52) was recorded in the plant that received no vernalization. Relevant results were published by who observed earlier and higher flowering for vernalized bulb planting (Behairy and Habbasha, 1979; Aguiar et al., 1983; Ramos and Ramos, 1993; Msika et al., 1997). High temperature (28-30 °C) during storage inhibits inflorescence initiation and prevents floral initiation (Rabinowitch, 1990). On the other hand, 8-12 °C storage temperatures were optimum for the occurrence of such events (Brewster, 1994). The maximum number of umbels plant⁻¹, number of flowers umbel⁻¹ (6.71, 236.89) was found in the plant that received vernalization at 5 °C for 14 days and minimum (5.10, 197.56) was observed respectively in the plant that received no vernalization.

The maximum number of seeds umbel⁻¹, number of fruits umbel⁻¹ (273.39, 88.24) was found in the plant that received vernalization at 5 °C for 14 days and minimum (214.62, 77.46) was observed respectively in the plant that received no vernalization. Larger mother plant produced larger umbel, therefore, conditions which lead to the improved growth would usually result also in larger flower heads. Usually a flowering stalk may consist of 50 to 2000 flowers (Jones and Mann, 1963). A higher proportion of scape emergence and greater uniformity in floral initiation were observed when at 10 °C increased as the length of vernalization (D'Angelo and Goldman, 2019). Flowering in bolting resistant cultivars can be particularly difficult and large bulb size or long treatment at low temperature may be required (Shishido and Saito, 1977). The increase in yield could be characterized to the greater number of flowers in vernalized plants. The results are similar with the previous findings (Msika et al., 1997; Khokhar, 2009). Some researchers noticed that 100 ppm GA₃ spray increased the flowering by 47% (Geetharani et al., 2008).

There was a significant variation in respect of height of flowering stalk, highest percent flowering, number of umbels plant⁻¹, number of flowers plant⁻¹, number of seeds umbel⁻¹, number of fruits umbel⁻¹ at 60 DAP due to different levels of GA₃. The longest flowering stalk (79.97 cm) was found in the plots which received 100 ppm GA₃ while the shortest (67.67 cm) was recorded in the plant that received no GA₃. The highest percent flowering (79.20) was found in the plots which received 100 ppm GA₃ while the lowest (74.03) was recorded in the plant that received no GA₃. The maximum number of umbels plant⁻¹, number of flowers umbel⁻¹ (6.87, 241.27) was found in the plant that received 100 ppm GA₃ and minimum (4.67, 186.72) was observed respectively in the plant that received no GA₃. The maximum number of seeds umbel⁻¹, number of fruits umbel⁻¹ (275.47, 92.06) was found in the plant that received 100 ppm GA₃ and minimum (216.75, 71.73) was observed respectively in the plant that received no GA₃. A group researchers investigated that 30-40% increase in seed yield due to GA₃ application (Naamni et al., 1980; Looper and Waller, 1982). The variation in the results of seed stalk length during both seasons could be characterized to the effect of the environment, because the genotypes used were identical (Elsiddig et al., 2015).

There was a significant variation in respect of number of fruits plant⁻¹, percent fruits, weight of seeds umbel⁻¹, 1000 seeds weight, weight of seeds plant⁻¹, weight of seeds plot⁻¹ and seed yield due to different levels of vernalization. The highest number of fruits plant⁻¹ (600.03) was found in the plots which received 5 °C vernalization for 14 days while the shortest (399.62) was recorded in the plant that received no vernalization. The highest percent fruit set (40.84) was found in the plots which received 5 °C vernalization for 14 days while the lowest (38.94) was recorded in the plant that received no vernalization. The maximum weight of seeds umbel⁻¹, 1000 seeds weight (1.08, 2.67) was found in the plant that received vernalization at 5 °C for 14 days and minimum (0.42, 2.33) was observed respectively in the plant that received no vernalization. Size of mother bulb and storage of mother bulbs are also major factors that affect flowering. Larger mother plant produced larger umbel, therefore,

conditions which lead to the improved growth would usually result also in larger flower heads.

Usually a flowering stalk may consist of 50 to 2000 flowers (Jones and Mann, 1963). The maximum weight of seeds plot⁻¹ and seed yield (59.01, 248.16) was found in the plant that received vernalization at 5°C for 14 days and minimum (23.10, 97.15) was observed respectively in the plant that received no vernalization. Increasing seed yield with vernalization treatment was observed (Aguiar et al., 1983). Early flowering shortens the life cycle of onion. Because of early flowering onion plants are exposed to different adverse condition (High temperature, Disease incidence, early rainfall etc.) for a short period than the no vernalized plants. This might have contributed to obtain higher seed yield due to vernalization. Better seed yield was obtained when bulbs were stored at low temperature. Onion, being a biennial crop needs vernalization of bulbs before flowering. The vernalization treatment depends upon the varieties and existing climatic condition. It might be that 5±1°C to 10±1°C cold treatment of bulbs were enough to meet up the vernalization requirement of the cultivar, which ultimately resulted in higher seed yield. Vernalization significantly increased the number of seeded fruits per umbel compared to the control treatment. The seed stalk elongation rate was increased by higher temperatures and long photoperiods (Brewster, 1982; Brewster, 1987). Better part of the plants with GA₃ treatment exhibited stunted growth and were chlorotic in appearance shows the toxic effect mostly due to high dosage (Yalamalle, 2016). There was a significant variation in respect of number of fruits plant⁻¹, percent fruits, weight of seeds umbel⁻¹, 1000 seeds weight, weight of seeds plant⁻¹, weight of seeds plot⁻¹ and seed yield due to different levels of GA₃.

The highest number of fruits plant⁻¹ (636.17) was found in the plots which received 100 ppm GA₃ while the shortest (335.96) was recorded in the plant that received no GA₃. The highest percent fruit set (40.54) was found in the plots which received 100 ppm GA₃ while the lowest (38.81) was recorded in the plant that received no GA₃. The maximum weight of seeds umbel⁻¹, 1000 seeds weight (0.80, 2.82) was found in the plant that received 100 ppm GA₃ and minimum (0.61, 2.18) was observed respectively in the plant that received no GA₃. The maximum weight of seeds plot⁻¹ and seed yield (45.60, 191.77) was found in the plant that received 100 ppm GA₃ and minimum (33.58, 141.24) was observed respectively in the plant that received no GA₃. Vernalization of bulbs (cultivar -Texas Early Grano) at 4-5°C for 180 days with the application of GA₃ at the rate of 600 ppm for seed production is suitable under Gezira state (Sudan) conditions (Elsiddig et al., 2015). GA₃ increased number of flowers per plant, length of flower stalks and seed yield per plant. The highest seed yield was achieved with the application of GA₃ at 75 ppm (Bhopale et al., 1999). Without vernalization, application of GA₃ singly, all the concentrations tested failed to induce flowering of onion (cultivar -Texas Early Grano) when planted off season but it accelerated the effect of vernalization on bolting and seed production (Elsiddig et al., 2015).

5. CONCLUSION

The results of present research indicated that application of vernalization and GA₃ had significant effect on yield and quality of summer onion production. For maximum seed yield, percentage of germination and for other good quality reproductive phases, vernalization at 5°C for 14 days and 100 ppm GA₃ responded independently. Therefore, to obtain the highest seed yield and quality seeds of summer onion the use of vernalization at 5°C for 14 days or 100 ppm GA₃ may be suggested to Mymensingh region. This experiment may be repeated under different agro-climatic regions of Bangladesh.

ACKNOWLEDGEMENT

Authors are grateful to the ministry of science and technology, Bangladesh for providing financial support to conduct the research work.

CONFLICT OF INTEREST

The authors have no conflict of interest to report.

REFERENCES

- Aguiar, P.A.A., Oliveira, L.O.B.D., Assuncao, M.V., 1983. Bulb vernalization in onion seed production in the lower middle Sao Francisco region. *Peaquis Agropecuaria Brasileira* 18(7)741-746. [Cited from Seed Abstr., 7 (4), Pp. 612 1984.
- Atherton, J.G., Craigon, J., Basher, E.A., 1990. Flowering and bolting in carrot. I. Juvenility, cardinal temperatures and thermal times for vernalization. *Journal of Horticultural Science*, Ashford, 65 (4), Pp. 423-429. <https://doi.org/10.1080/00221589.1990.11516075>
- Behairy, A.G., Habbasha, K.M.E., 1979. Onion (*Allium cepa* L.) seed production as affected by vernalization of bulbs. *Zeitschrift fur Acher and Pflanzenbau*, 148 (2), Pp. 109-114. [Cited from Hort. Abstr.50(1)270, 1980].
- Benkeblia, N., Onodera, S., Yoshihira, T., Kosaka, S., Shiomi, N., 2004. Effect of temperature on soluble invertase activity, and glucose, fructose and sucrose status of onion bulbs (*Allium cepa* L.) in store. *International Journal of Food Science and Nutrition*, 55 (4), Pp. 325-331. <https://doi.org/10.1080/09637480412331290512>
- Bertaud, D.S., 1988. Effects of chilling duration photoperiod and temperature on floral initiation and development in sprouted and unsprouted onion bulbs. In: *Proceedings of the fourth EUCARPIA Allium Symposium*, Institute of Horticultural Research, Wellesbourne, UK, Pp. 245-261.
- Bhopale, S.R., Dod, V.N., Bharad, S.G., Gholap, S.V., Jadhao, B., 1999. Seed Production of Onion as Influenced by the Application of Growth Regulators. *Journal of Soils and Crops*, 9 (1), Pp. 78-79.
- Brewster, J. L., 1982. Flowering and seed production in over wintered cultivars of bulb onion, I. Effects of different raising environments, temperatures and day lengths. *Journal of Horticultural Sciences*, 57, Pp. 93 - 96. <https://doi.org/10.1080/00221589.1982.11515028>
- Brewster, J.L., 1983. Effects of photoperiod, nitrogen nutrition and temperature on inflorescence initiation and development in onion (*Allium cepa* L.). *Annals of Botany*, London, 51 (4), Pp. 429-440. <https://doi.org/10.1093/oxfordjournals.aob.a086487>
- Brewster, J.L., 1987. Vernalization in the onion - a quantitative approach. In: *ATHERTON, J.G. Manipulation of flowering*. London: Butterworths, Cap., 13, Pp. 171-183. <https://doi.org/10.1016/b978-0-407-00570-9.50018-1>
- Brewster, J.L., 1994. *Onion and Other Vegetable Alliums*. CAB International, UK, Pp. 236.
- Corgan, J.N., Montano, T.M., 1975. Bolting and other responses of onion (*Allium cepa* L.) to growth regulating chemicals. *J. Amer. Soc. Hort. Sci.*, 100 (3), Pp. 273-227.
- D'Angelo, C.J., Goldman, I.L., 2019. Annualization of the Long Day Onion Breeding Cycle through Threshold Vernalization and Dormancy Disruption. *Crop Breed Genom.*, 1, Pp. e190009. <https://doi.org/10.20900/cbgg20190009>
- El-Habbasha, K.M., Mahmoud, H.A., Thabet, N.G., Abdou, F.E., 1985. Effect of GA₃ and IAA Application on the flowering and seed production of onion (*Allium cepa* L.). *Iraq Journal of Agricultural Water Resource Research*, 4 (2), Pp. 13-26.
- Elsiddig, E.A.M., Elamin, O.M., Elkishif, M.E., 2015. Induction of Flowering in Texas Early Grano Onion Cultivar Using Vernalization and Gibberellic Acid under Gezira State Conditions, Sudan. *International Journal of Scientific and Research Publications*, 5 (9).
- Flood, R.G., Halloran, G.M., 1986. Genetics and physiology of vernalization response in wheat. *Advances in Agronomy*, 39 (1), Pp. 87-125. [https://doi.org/10.1016/s0065-2113\(08\)60466-6](https://doi.org/10.1016/s0065-2113(08)60466-6)
- Geetharani, P., Manivannan, M.I., Ponnuswamy, A.S., 2008. Seed production of onion as influenced by the application of growth regulators and nutrients. *Asian Journal of Horticulture*, 3 (2), Pp. 301-303.
- Gomez, K.A., Gomez, A.A., 1984. *Statistical Procedures for Agricultural Research*. New York: John Wiley and Sons Inc. New York, Pp. 67-215.
- Hodges, T., Ritchie, J.T., 1991. The Ceres-Wheat phenology model. In: *Hodges, T. Predicting crop phenology*. Boston: CRC Press, Cap., 12, Pp. 133-141.
- Holcomb, E.J., Berghage, R., 2001. Photoperiod, chilling, and light quality during daylight extension affect growth and flowering of tissue-cultured Easter lily plants. *HortScience*, Mt. Vernon, 36 (1), Pp. 53-55. <https://doi.org/10.21273/hortsci.36.1.53>
- Jones, A., Saxena, P.K., Murch, S.J., 2009. Elicitation of Secondary Metabolism in *Echinacea purpurea* L. by Gibberellic Acid and Triazoles. *Eng Life*; <https://doi.org/10.1002/elsc.200800104>
- Jones, H.A., Mann, L.K., 1963. *Onions and their Allies*. Leonard Hill, (Books) Ltd., London. Pp. 1-169.

- Khokhar, K.M., 2009. Effect of set-size and storage temperature on bolting, bulbing and seed yield in two onion cultivars. *Scientia Horticulturae*, 122, Pp. 187–194. <https://doi.org/10.1016/j.scienta.2009.05.008>
- Khokhar, K.M., 2014. Flowering and Seed Development in Onion—A Review. *Open Access Library Journal*, 1: e1049. <http://dx.doi.org/10.4236/oalib.1101049>
- Kimani, P.M., Peters, R., Rabinowitch, H.D., 1994. Potential of onion seed production in a tropical environment. *Acta Horticulturae*, Wageningen, 358 (1), Pp. 341–348. <https://doi.org/10.17660/actahortic.1994.358.56>
- Kruzhilin, A.S., Shvedskaya, Z.M., 1962. Peculiarities of phasic development and morphogenesis in onion (*Allium cepa* L.). *Fiziologia Rastenii*, 9, Pp. 466–475.
- Lee, R., Baldwin, S., Kenel, F., McCallum, J., Macknight, R., 2013. Flowering Locus T genes control onion bulb formation and flowering. *Nature Communications*, 4, Pp. 2884. <https://doi.org/10.1038/ncomms3884>
- Linwattana, G., Protacio, C.M., Mabesa, R.C., 1997. Tropical low land seed production of non-heading Chinese cabbage (*Brassica rapa* L. *pekinensis* group) using vernalization and gibberellic acid. *Philippine Journal of Crop Science*, 22 (3), Pp. 161–166.
- Loper, G.M., Waller, G.D., 1982. GA3-increased bolting and seed production in late-planted onions. *Hortscience*, 17 (6), Pp. 922–923.
- Msika, R.L., Jackson, J.E., Currah, L., 1997. Onion Seed Production Trials in the Highveld of Zimbabwe. *Acta Horticulturae*, 433, Pp. 337–346. <https://doi.org/10.17660/actahortic.1997.433.36>
- Muthamia, K.W., 1994. Effect of bulb size and duration of vernalization on onion (*Allium cepa* L.) seed yield. MSc thesis University of Nairobi. Available from <http://erepository.uonbi.ac.ke/handle/11295/23061>.
- Naamni, F., Rabinowitch, H.D., Kedar, N., 1980. The effect of GA3 application on flowering and seed production in onion. *Journal of the American Society for Horticultural Science*, 105 (2), Pp. 164–167.
- Olszewski, N., Sun, T.P., Gubler, F., 2002. Gibberellin signaling biosynthesis, catabolism, and response pathways. *Plant Cell*, (14), Pp. S61–S80. <https://doi.org/10.1105/tpc.010476>
- Peters, R., 1990. Seed production in onions and some other *Allium* species. In: Rabinowitch, H.D., Brewster, J.L. (Eds.), *Onions and Allied Crops*. Vol. I. Botany, Physiology and Genetics. CRC Press, Boca Raton, Florida, Pp. 161–176. <https://doi.org/10.1201/9781351075169-8>
- Pike, L.M., 1986. Onion breeding. In: Basset, M.J. (Ed.), *Breeding Vegetable Crops*. AVI Publishing Co., Connecticut, Pp. 357–394.
- Pinthus, M.J., 1985. *Triticum*. In: HALEY, H.C. *CRC Handbook of flowering*. Volume IV. Boca Raton: CRC Press, Pp. 418–443. <https://doi.org/10.1201/9781351072564-59>
- Porter, J.R., Gawith, M., 1999. Temperatures and the growth and development of wheat: a review. *European Journal of Agronomy*, Montrouge Cedex, 10 (1), Pp. 23–36.
- Rabinowitch, H.D., 1990. Physiology of flowering. In: Rabinowitch, H.D., Brewster, J.L. (Eds.), *Onions and Allied Crops*. Vol. I. Botany, Physiology and Genetics. CRC Press, Boca Raton, Florida, Pp. 133–134. <https://doi.org/10.1201/9781351075169-5>
- Rabinowitch, H.D., 1985. Onions and other edible *Alliums*. In: Halevy, A.H. (Ed.), *CRC Handbook of Flowering*. Vol. I. CRC Press, Boca Raton, Florida, Pp. 398–405.
- Ramos, D.S.G., Ramos, A.A., 1993. Preliminary observations of the production of onion seed in the state of Merida, Venezuela. *Onion Newsletter for the Tropics*, 5, Pp. 44–48.
- Ritchie, J.T., 1991. *Modeling Plant and Soil Systems*. Madison: ASA, CSSA, and SSSA, Cap., 3, Pp. 31–54.
- Rizk, F.A., Shaheen, A.M., El-Habbasha, K.M., 1996. Flowering and seed yield of onion (*Allium cepa* L.) plants as affected by dates of planting and some growth regulators. *Egypt J Hort.*, 23, Pp. 113–127.
- Roh, M.S., 1985. Flowering response of mid-century hybrid lilies to bulb vernalization and shoot photoperiod treatment. *Hort. Science*, Mt. Vernon, 20 (4), Pp. 710–713.
- Roh, M.S., Wilkins, H.F., 1977. Comparison of continuous and alternating bulb temperature treatments on the growth and flowering in *Lilium longiflorum* Thunb. *Journal of the American Society for Horticultural Science*, Boston, 102 (3), Pp. 242–247.
- Sekara, A., Pokluda, R., Del Vacchio, L., Somma, S., Caruso, G., 2017. Interactions among genotype, environment and agronomic practices on production and quality of storage onion (*Allium cepa* L.) – A review. *Hort. Sci. (Prague)*, 44, Pp. 21–42. <https://doi.org/10.17221/92/2015-hortsci>
- Shaikh, A.M., Vyakaranahal, B.S., Shekhargouda, M., Dharmatti, P.R., 2002. Influence of Bulb Size and Growth Regulators on Growth, Seed Yield and Quality of Onion cv. Nasik Red. *Seed Research*, 30 (2), Pp. 223–229.
- Sharma, N., Kaur, N., Gupta, A.K., 1999. Effects of gibberellic acid and chlorocholine chloride on tuberisation and growth of potato (*Solanum tuberosum* L.). *J. Sci. Food Agric.*, 78 (4), Pp. 466–470.
- Shaykewich, C.F., 1995. An appraisal of cereal crop phenology modeling. *Canadian Journal of Plant Science*, Ottawa, 75 (2), Pp. 329–341.
- Shishido, Y., Saito, T., 1977. Studies on the flower bud formation in onion plants. III. Effects of physiological conditions on the low temperature induction of flower buds in bulbs. *J. Jpn. Soc. Hortic. Sci.*, 46, Pp. 310–316.
- Streck, N.A., 2003. A Vernalization Model in Onion (*Allium cepa* L.). *Modelo De Vernalização Em Cebola (Allium cepa L.)*. Departamento de Fitotecnia, Centro de Ciências Rurais, Universidade Federal de Santa Maria. 97105-900 – Santa Maria, RS, Brasil. <https://doi.org/10.21475/ajcs.18.12.12.p1434>
- Streck, N.A., 2002a. A generalized nonlinear air temperature response function for node appearance rate in muskmelon (*Cucumis melo* L.). *Revista Brasileira de Agrometeorologia*, Santa Maria, 10 (1), Pp. 105–111.
- Taiz, L., Murphy, A., 2015. *Plant Physiology and Development*. Sunderland, Massachusetts (USA): Sinauer Associates. Pp. 605. ISBN 978-1-60535-255-8. <https://doi.org/10.3119/0035-4902-117.971.397>
- Thomas, B., Vince-Prue, D., 1997. *Photoperiodism in Plants*. Academic Press, San Diego.
- Ubeda Tomas, S., Federici, F., Casimiro, I., 2009. Gibberellin signaling in the endodermis controls arabidopsis root meristem size. *Curr Biol.*, 19 (14), Pp. 1194–1199. <https://doi.org/10.1016/j.cub.2009.06.023>
- Voss, R.E., Murray, M., Bradford, K., Mayberry, K., Miller, I., Long, R., Gillespie, S., 2013. Onion seed production in California. Available from <http://anrcatalog.ucdavis.edu/pdf/8008.pdf> <https://doi.org/10.3733/ucanr.8008>
- Woodbury, G.W., 1950. A study of factors influencing floral initiation and seed stalk development in the onion, *Allium cepa* L. Idaho Agricultural Experiment Station Research Bulletin No. 18.
- Wurr, D.C.E., Fellows, J.R., Phelps, K., 1995. Vernalization in calabrese (*Brassica oleracea* var. *italica*) – a model for apex development. *Journal of Experimental Botany*, Oxford, 46 (291), Pp. 1487–1496.
- Yalamalle, V.R., 2016. Vernalization Responses in Onion (*Allium cepa*) Pre-flowering and Reproductive Phases. ICAR-Directorate of Onion and Garlic Research, Rajgurunagar, Pune (412 105), India. *International Journal of Bio-resource and Stress Management*, 7 (6), Pp. 1413–1417. <https://doi.org/10.23910/ijbsm/2016.7.6.1728a>
- Zemah, H., Rabinowitch, H.D., Kamenetsky, R., 2001. Florogenesis and the effect of temperatures on the development of *Allium* *aflatunense*. *Journal of Horticultural Science & Biotechnology*, 76, Pp. 507–513. <https://doi.org/10.1080/14620316.2001.11511401>