ISSN: 2735-0274 (Online)
CODEN: TARGDB

Tropical Agroecosystems (TAEC)

DOI: <http://doi.org/10.26480/taec.02.2021.79.81>

RESEARCH ARTICLE

INTERCROPPING CHILLI WITH GROUNDNUT IN CHAR LAND

M Kadir^a, Jubaidur Rahman^b, MI Riad^c, MK Alam^d

^aPrinciple Scientific Officer, Oilseed Research Centre, Bangladesh Agricultural Research Institute, Jamalpur-2000, Bangladesh

^bScientific Officer, Agronomy Division, Bangladesh Agricultural Research Institute, Jamalpur-2000, Bangladesh

^cSenior Scientific Officer, Plant Genetic Resources Centre, Bangladesh Agricultural Research Institute, Jamalpur-2000, Bangladesh

^dSenior Scientific Officer, Oilseed Research Centre, Bangladesh Agricultural Research Institute, Jamalpur-2000, Bangladesh

*Corresponding Author E-mail: mkadirbari@yahoo.com

This is an open access article distributed under the Creative Commons Attribution License CC BY 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ARTICLE DETAILS

Article History:

Received 01 June 2021

Accepted 03 July 2021

Available online 22 July 2021

ABSTRACT

Intercropping is a farming practice involving two or more crop species, or genotypes, growing together and coexisting for a time. On the fringes of modern intensive agriculture, intercropping is important in many subsistence or low-input/resource-limited agricultural systems. Intercropping in order to improve land use efficiency, take advantage of intercropping facilitation and improve economic benefits. Also, linked with the higher yield, the associated higher amount of vegetables-legume by-product is preferred for animal feed or human consumption. The objective of this study is to find out the suitable groundnut - chilli intercropping system and increase productivity, economic and nutritional development of charland farmers of Bangladesh. In the experiment, intercropping system such as One row of chilli in between two normal rows of groundnut, two rows of chilli in between two normal rows of groundnut which was compared with sole groundnut and sole chilli production. Total yields of chilli /groundnut intercropping were higher than that of sole chilli and groundnut. chilli /groundnut intercropping is the efficient use of light because of complementary use of space between the chilli plants and the groundnut plants. Chilli /groundnut intercrop gross return and BCR per hectare were found higher than from sole crop system. The total yield of intercropped crops was greater than sole cropping shown by LER>1. The overall advantage of intercropping ranged from 71 to 81%. The highest land equivalent value of 81% was recorded from paired row chilli in between four normal rows of groundnut sowing arrangements indicated a yield advantage of 81% over sole crop.

KEYWORDS

Intercropping, chilli, groundnut and Charland.

1. INTRODUCTION

Intercropping is the cultivation of two or more crop species in close intermingling on the same field for the whole or a part of their growing period (Vandermeer, 1989; Willey, 1990). Intercropping generally produces more yield per unit of land than sole cropping (Yu et al., 2015; Zhang and Li, 2003) and reduces the risk of crop failure due to pests and diseases (Boudreau, 2013; Horwith, 1985; Lithourgidis et al., 2011; Zhang et al., 2019). These advantages mainly arise from complementary patterns of resource uptake (light, water and nutrients) between crop species, better nutrient cycling, and through suppression of pests, weeds and diseases (Vandermeer, 1989). Intercropping also has the potential to counteract resource degradation (Feike et al., 2012; Lichtfouse, 2009), it can contribute to increases in soil carbon due to increased root biomass input, and it can increase organic soil nitrogen as a result of better nitrogen cycling (Cong et al., 2015). Intercropping is receiving increasing global interest as an agricultural practice as farmers strive to be more sustainable and maintain soil health (Glaze-Corcoran et al., 2020). Groundnut (*Arachis hypogaea* L.) is an important leguminous oilseed crop which is commonly known as poor man's nut as it is a cheaper source of protein when comparable to other nuts like cashew nut. It is also called as peanut, monkey nut and goober nut. Groundnut seed contains 44 to 56% oil and 22 to 30% protein on dry seed basis and is a rich wellspring of minerals

(Phosphorus, Calcium, Magnesium and Potassium) and vitamins (Kaba et al., 2014). Chilli (*Capsicum*) is an important crop and is considered an almost indispensable spice in daily life (W. B. Paul, 2013). The trace elements, minerals, and nutrients in their fruits have good health properties (R. Arimboor, 2015; G. Subha, 2017; Z. Shi, 2018 and A. Srivastava, 2019). This crop is grown year-round, seasonally independent (R. D. R. Elizanilda, 2016). In Bangladesh, there are about 0.82 million hectares of char land (Ahmed et al., 1987). "Charland" is the Bengali term, its English meaning is "Riverine Island" for mid-channel island that emerges periodically from riverbed as a consequence of accretion (Elahi, 1991). In Bangladesh the char lands can be divided into five sub areas which has highly potential for groundnut production (The Jamuna, the Ganges, The Padma, The upper Meghna and the lower Meghna River) where Tista and old Brahmaputra also constitute some char land areas (Islam et al., 2012). Due to decreasing cultivable land, farmers of char areas (Riverine Island) in Bangladesh have been practicing intercropped garden pea with onion (Rahman et al., 2015), coriander with onion (Talukder et al. 2015), sweet gourd with onion (Talukder et al. 2015), vegetables, pulse and oilseed crops with wheat (Talukder et al. 2016), different gourds with brinjal (Rahman et al., 2020) are common practice to the farmers of char areas. Groundnut and brinjal are the most important oil/vegetable crop production in Bangladesh especially in char land and cultivated all over the country in Rabi and Kharif seasons. Groundnut

Quick Response Code

Access this article online

Website:
www.taec.com.my

DOI:
10.26480/taec.02.2021.79.81

(87131 acres) chilli (239203 acres) of land with an annual production of groundnut (62832 M Ton) chilli (149473 M Ton) in Rabi and Kharif seasons respectively (BBS, 2020). Farmers of charland area cultivate sole groundnut or sole chilli as a result productivity also reduced for this situation intercropped practice may increase productivity, economic and nutritional development of charland farmers. In this case the experiment is undertaken in charland.

2. MATERIALS AND METHODS

The experiment was conducted at Nawvanger char, sadar, Jamalpur, Bangladesh 24°57' north latitudes and 89°55' east longitudes. The annual average temperature of this district varies from maximum 36.63°C to minimum 9.4°C. Annual average rainfall is 1549.45mm (Regional Research Report 2019-2020). The experimental site was of medium high land belonging to the agro-ecological zone Old Brahmaputra Floodplain under Agro-Ecological Zone 9 (UNDP & FAO, 1988). The experiment was conducted at the during *rabi* 2019-2020 and 2020-2021 to find out the suitable groundnut - brinjal intercropping system for increase productivity, economic and nutritional development of charland farmers. Design of the experiment was RCB with 03 (three) replications having the unit of plot 3.6m × 3.2m. BARI Badam-9 was used as a variety in the experiment. Treatments included in the experiment were: T₁= Sole chilli, T₂= Sole groundnut, T₃= One row of chilli in between two normal rows of groundnut, T₄= Paired row chilli in between four normal rows of groundnut. ½ N and all other fertilizers as basal. Rest N will be applied at 30-35 and 55-60 DAS after irrigation (FRG, 2018). Crops were sown on October 29, 2020 and harvested from January 28, 2021 to April 25, 2021. Intercultural operations like watering, weeding and spraying insecticides were followed as and when necessary. Collected data were analyzed statistically with the help of STAR software and mean separation was done as per LSD test at 5% level of significance.

2.1 Nutrient status of initial soil

Location	pH	OC (%)	OM (%)	N (%)	K Meq 100g ⁻¹	P	S	Zn	B
Naovangar char, Jamalpur	6.3		0.98	0.049	0.22	μg g ⁻¹			
						4.56	1.41	0.71	0.39

3. RESULTS AND DISCUSSION

Although total grain yields of chilli/groundnut intercropping were higher than that of sole chilli and groundnut. Another important aspect for chilli/groundnut intercropping is the efficient use of light because of complementary use of space between the chilli plants and the groundnut plants. The complementary effect is also shown for the length of the growth period because their life cycles are different, with the maturity of intercropped. The yield, gross return, variable cost and BCR of chilli/groundnut are presented in Table 1. From the Table, it was revealed that chilli/groundnut intercrop gross return and BCR per hectare were found higher than from sole crop system. In sole crop system which delivered less BCR than intercropped system which range 1.6-2.68. Groundnut Equivalent Yield (GEY) was highest from paired row chilli in between four normal rows of groundnut (4.89 t/ha) system which was statistically similar to other intercropped system.

Table 1: Two year polled data of intercropping chilli with groundnut at 2019-2020 and 2020-2021

Treatment combination	Yield (t/ha)		Gross return (Tk.ha ⁻¹)	TVC (Tk.ha ⁻¹)	BCR	GEY	LER
	Groundnut	Chilli					
Sole chilli	-	8.0	240000	150000	1.6	3.43	-
Sole groundnut	2.01	-	140700	112500	1.25	2.01	-
One row of chilli in between two normal rows of groundnut	1.80	6.5	321000	120000	2.68	4.59	1.71
Paired row chilli in between four normal rows of groundnut	1.85	7.1	342500	132000	2.59	4.89	1.81

Chilli (Green) = 30 Tk per kg and Groundnut= 70 Tk per kg

4. CONCLUSION

The increased yield with the intercropping was largely due to improved interspecies interaction and facilitation. Intercropping in order to improve land use efficiency, take advantage of intercropping facilitation and improve economic benefits. Also, linked with the higher yield, the associated higher amount of vegetables-legume by-product is preferred for animal feed or human consumption. Therefore, chilli /groundnut intercropping systems increase productivity, economic and nutritional development of charland farmers.

ACKNOWLEDGEMENT

The authors are too much grateful to 'Enhancement of nutritional status and economic condition of charland farmers through development of modern groundnut varieties and intercropping technologies' for funding this programme.

REFERENCES

- Ahmed, M.M., Alam, N., Kar, N.K., Maniruzzaman, A.F.M., Abedin, Z., Jasimuddin, G. 1987. Crop production in saline and charlands-existing situation and potentials. *Advances in Agronomic Research in Bangladesh*, vol. 2, pp. 1-27.
- Arimboor, R., Natarajan, R.B., Menon, K.R., Chandrasekhar, L.P., Moorkoth, V. 2015. Red pepper (*Capsicum annum*) carotenoids as a source of natural food colors: analysis and stability a review, *Journal of Food Science and Technology*, vol. 52, no. 3, pp. 1258-1271.
- Bangladesh Bureau of Statistics (BBS). 2020. Statistical Year Book of Bangladesh, Ministry of Planning, Government of the People's Republic of Bangladesh, Dhaka.
- Boudreau, M.A. 2013. Diseases in intercropping systems. *Annu. Rev. Phytopathol.*, vol. 51, pp. 499-519.
- Cong, W., Hoffland, E., Li, L., Six, J., Sun, J., Bao, X., Zhang, F., Van der Werf, W. 2015. Intercropping enhances soil carbon and nitrogen. *Glob. Chang. Biol.*, vol. 21, pp. 1715-1726.
- Elahi, K.M. 1991. Impacts of riverbank erosion and flood in Bangladesh: An introduction. *Riverbank Erosion Impact Study*.
- Elizanilda, R.D.R., Mailson, M.D.R., Fernando, L.F. 2016. Production and Breeding of Chilli Peppers (*Capsicum* spp.), Springer International Publishing.
- FAP 16/19, 1993a. Charland Study Overview: Summary Report, Flood Plain Coordination Organization, Ministry of Irrigation Water Development and Flood Control, Dhaka.
- Feike, T., Doluschitz, R., Chen, Q., Graeff-Honninger, S., Claupein, W. 2012. How to overcome the slow death of intercropping in the North China plain. *Sustainability*, vol. 4, pp. 2550-2565.
- FRG. 2018. Fertilizer Recommendation Guide, Bangladesh Agricultural Research Council (BARC), Farmgate, Dhaka, pp. 1215.
- Glaze-Corcoran, S., Hashemi, M., Sadeghpour, A., Jahanzad, E., Afshar, R.K., Liu, X., et al. 2020. Understanding intercropping to improve agricultural resiliency and environmental sustainability. *Adv. Agron.*, vol. 162, pp. 199-256. Doi: 10.1016/bs.agron.2020.02.004
- Islam, M.S., Hasan, T., Chowdhury, M.S.I.R., Rahaman, M.H., Tusher, T.R. 2012. Coping Techniques of Local People to Flood and River Erosion in Char Areas of Bangladesh, *J. Environ. Sci. & Natural Resources*, vol. 5, no. 2, pp. 251 - 261. Jahangirnagar University, Dhaka, pp. 11-67.
- Jubaidur, R., Begum, A.A., Fouzia, S.S., Akter, A., Saha, R.R. 2020. Relay Intercropping of Different Gourds with Brinjal in Charland Area. *Acta Scientifica Malaysia*, vol. 4, no. 1, pp. 10-12.
- Kaba, J.S., Kumaga, F.K., Ofori, K. 2014. Effect of flower production and time of flowering on pod yield of peanut (*Arachis hypogaea* L) genotypes. *Journal of Agriculture and Veterinary Science*, vol. 7, no. 4, pp. 44-9.
- Lichtfouse, E. 2009. Sustainable Agriculture Reviews: Climate Change, Intercropping, Pest Control and Beneficial Microorganisms. Springer Netherlands, Dordrecht.
- Lithourgidis, A., Dordas, C., Damalas, C., Vlachostergios, D. 2011. Annual

- intercrops: an alternative pathway for sustainable agriculture. *Aust. J. Crop. Sci.*, vol. 5, pp. 396–410.
- Motiur Rahman Talukder, A.H.M., Bulbul, A., Lutfun, N., Fahid Hossain, K.M., Jubaidur, R., Santosh, K.P. 2016. Enhancement of farm productivity through intercropping of vegetables, pulse and oilseed crops with wheat at Jamuna Char area of Islampur in Jamalpur District. *International Journal of Applied Research*, vol. 2, no. 2, pp. 71-76
- Paul, W.B., Eric, J.V. 2013. *Peppers: Vegetable and Spice Sapsicums*, Centre for Agriculture and Bioscience International, 2nd edition.
- Rahman, J., Talukder, A.H.M.M.R., Nahar, L., Rahman, M.M., Kayser, N. 2015. *International Journal of Applied Research*, vol. 1, no. 2, pp. 79-82
- Regional Research Report. 2019-20. Regional Agricultural Research Station, BARI, Jamalpur-2000
- Saini, D.K., Kaushik, P. 2019. Visiting eggplant from a biotechnological perspective: A review. *Sci. Hortic.*, vol. 253, pp. 327–340.
- Shi, Z. 2018. Chilli intake is inversely associated with hypertension among adults,” *Clinical Nutrition ESPEN*, vol. 23, pp. 67–72.
- Srivastava, A., Sharma, V.R., Dishri, M., Dikshit, A., Mangal, M., Kalia, P. 2019. Inheritance of fruit attributes in chilli pepper, *Indian Journal of Horticulture*, vol. 76, no. 1, pp. 86–93.
- Subha, G., Praveen, K.P., Parveez, A.P., Vijay, S. 2017. Medicinal Properties of Chilli Pepper in Human Diet: An Editorial,” *ARC Journal of Public Health and Community Medicine*, vol. 2, no. 1, pp. 6-7.
- Talukder, A.H.M.M.R., Rahman, J., Nahar, L., Rahman, M.M., Kayser, N. 2015. Mixed cropping onion with different plant population of sweet gourd. *IOSR Journal of Agriculture and Veterinary Science*, pp. 45-50
- Talukder, A.H.M.M.R., Rahman, J., Nahar, L., Rahman, M.M., Kayser, N. 2015. Mixed cropping onion with different plant population of sweet gourd. *IOSR Journal of Agriculture and Veterinary Science*, pp. 45-50
- UNDP-FAO. 1988. Land resources appraisal of Bangladesh for agricultural development. Report to Agro-ecological regions of Bangladesh. UNDP-FAO, BGD/81/ 035 Technical Report, vol. 2, pp. 570.
- Willey, R.W. 1990. Resource use in intercropping systems. *Agric. Water Manag.*, vol. 17, pp. 215–231.
- Yu, Y., Stomph, T.J., Makowski, D., van der Werf, W. 2015. Temporal niche differentiation increases the land equivalent ratio of annual intercrops: a meta-analysis. *Field Crop Res.*, vol. 184, pp. 133–144.
- Zhang, C.C., Dong, Y., Tang, L., Zheng, Y., Makowski, D., Yu, Y., Zhang, F.S., van der Werf, W. 2019. Intercropping cereals with faba bean reduces plant disease incidence regardless of fertilizer input; a meta-analysis. *Eur. J. Plant Pathol.* Doi: <https://doi.org/10.1007/s10658-019-01711-4>.