RESEARCH ARTICLE

MANAGEMENT OPTIONS AGAINST POD BORER (*MARUCA TESTULALIS* GEYER) AND POD SUCKING BUGS (*RIPTORTUS DENTIPES*) OF COWPEA (*VIGNA UNGUICULATA* (L.) WALP.)

Sagar Dahal^{a*}, Bhola Gautam^b, Bala Sharma^a, Kamal Neupane^a, Santosh Kandel^a, Samikshya Sedhai^a, Grace Tiwari^a, Laxmi Narayan Ojha^a^a Agriculture and Forestry University, Rampur, Chitwan, Nepal^b Asst. Professor, Agriculture and Forestry University, Rampur, Chitwan, Nepal* Corresponding Author e-mail: Sagardahal413@gmail.com

This is an open access article distributed under the Creative Commons Attribution License CC BY 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ARTICLE DETAILS

Article History:

Received 28 May 2020

Accepted 15 June 2020

Available online 18 June 2020

ABSTRACT

The efficacy of different management practices comprising cowpea/sorghum intercropping, microbial insecticide: Spinosad, neem-based botanical pesticide: neemix and synthetic insecticide: Chlorpyrifos 50 + Cypermethrin 5 were evaluated against pod borer and pod sucking bugs of cowpea. The experiment was laid out in completely randomized block design at Agriculture and Forestry University, Rampur, Nepal. All the treatments except cowpea/sorghum intercropping significantly reduced the flower infestation of pod borer larvae. The lowest number of infected flower and number of larvae per flower was observed in Spinosad treated plot. Also, the highest percentage of reduction in flower infestation was observed in Spinosad treated plots. Similarly, Chlorpyrifos 50 + Cypermethrin 5 treated plots showed the least number of pod bug infestation followed by Spinosad. The highest yield and lowest weight of damaged pod was recorded in Spinosad treated plots along with the lowest percentage infestation of pods by weight. The highest net income per hectare of land was recorded in Spinosad followed by Chlorpyrifos 50 + Cypermethrin 5 treatment. The marginal benefit cost ratio showed that the highest profit per unit rupee spent over control was obtained in Chlorpyrifos 50 + Cypermethrin 5 treatment followed by Spinosad. Considering the results obtained, Spinosad an eco-friendly, safe and less hazardous bio-pesticide has been recommended as a good approach for the management of cowpea pod borer and pod sucking bug.

KEYWORDS

botanical pesticide; microbial insecticide; eco-friendly; bio-pesticide

1. INTRODUCTION

Cowpea (*Vigna unguiculata* (L.) Walp.) is an annual herbaceous legume and is one of the most important grain legumes of Nepal. It is believed to have been originated from central Africa and is mainly grown for grain, green pods and leaves. The grain contains significant amount of protein with high biological value and thus it is also regarded as vegetable meat (Oyewale and Bamaïyi, 2013).

In Nepal, it is chiefly used as grain crop, as animal fodder, green manure crop, or as a vegetable. It is grown as vegetable in 4620.7 ha with productivity of 12.3 t/ha in Nepal. The released variety of cowpea are Malepatan-1, Surya, Prakash and Akash and the registered variety is Double Harvest (Joshi, et al., 2017).

About 21 different insect species are recorded to damage cowpea throughout its stages from seedling to maturity (Sardana & Verma, 1986; Choudhary, Hussain, Samota, & Nehra, 2017). Among them, pod borer (*Maruca testulalis*) and pod sucking bugs complex including *Riptortus dentipes* are considered destructive pest of cowpea in Africa and Asia and

cause serious loss in cowpea and may cause substantial economic loss if left uncontrolled (Jackai and Daoust, 2003; Baoua, et al., 2011).

Cowpea pod borer (*Maruca testulalis* Geyer) lays pale cream, translucent eggs singly or in batch on stems, young leaves, flowers and pods. The egg hatch in about 3-4 days and give rise to pale cream larvae with two rows of dark dots on their back which feeds inside the flower leading poor pod setting and pod formation. They also damage stem, peduncle, buds and leaves by eating and weaving them. The larvae in later stage of crop growth behaves as a pod borer and complete its larval and pupal development inside the pod leading poor pod formation, reduction in yield and decrease in the market value of the green pods (Shinde et al., 2017; Okeyo-Owuor et al., 1983).

The larvae feeds on leaves, stems, flower and pods and damage about 50-60% of flowers preventing them from setting pods. It has also been found to cause 80% yield loss (Nyiira, 1971; Nampala, et al., 2002). Pod borer is ubiquitous and very devastating which cause yield loss of about 80% in infested field and sometimes may lead to total crop failure if no control measures are adopted (Bett et al., 2017).

Quick Response Code

Access this article online

Website:

www.trab.org.my

DOI:

[10.26480/trab.01.2020.24.30](https://doi.org/10.26480/trab.01.2020.24.30)

Pod sucking bugs (*Riptortus dentipes* Fab.) damage the pods and seed of cowpea. The damage is caused by adult and nymph of the pod bug which rest on foliage and feeds on young shoots and green pods. They suck the sap from developing young shoots and green pods and results in seed deformation, premature seed drying, seed abortion and pod shriveling (Jackai et al., 2009). Their feeding punctures are undetectable, but they cause premature shriveling in young pods and half-filled seeds in older pods. The damage to cowpea increases with the increase in population of bugs and the yield reduction was more pronounced when the bug infest the cowpea during flowering stage compared to infestation during podding (Jackai et al., 1989).

Farmers are trying to reduce the damage of insect pests in cowpea by regularly spraying organophosphates and pyrethroids insecticides (Sreelakshmi, 2014). Use of insecticides and biopesticides may keep the population of insect pests below economically damaging levels but its use is uneconomical; due to low purchasing power of farmers and unsustainable; due to its serious effect on environment and human health (Bett et al., 2017; Kamara et al., 2007). It is commonly observed that the survivor individual become resistant to insecticides and thus more difficult to control (Sreelakshmi, 2014; Ekesi, 1999). So, for environment friendly and sustainable management of insect pests of cowpea, a holistic approach integrating all possible management strategies is essential. The objectives of this research are:

- To evaluate the different control methods of insect pest in cowpea.
- To recommend the environment friendly and effective method for controlling the insects under study.

2. MATERIAL AND METHODS

The study was carried out at Horticulture Farm of Agriculture and Forestry University, Rampur, Chitwan, Nepal from April, 2019 to June, 2019. The experiment was laid out in Randomized Complete Block Design (RCBD) with 5 treatments, each replicated four times. The treatments comprised synthetic chemical insecticide (Chlorpyrifos 50 + Cypermethrin 5), botanical neem-based pesticide (Neemix), microbial insecticide (Spinosad), intercropping with sorghum and control. The spray of chemical, botanical and microbial insecticide was done at an interval of 9 days and the sorghum was sown at the time of cowpea sowing and managed as additive intercrop throughout the crop period.

Table 1: Details of the insecticides experimented.

Common Name	Trade Name	Formulation Type	Dose
Chlorpyrifos 50 + Cypermethrin 5		EC	2 ml / l
Spinosad	Tracer	45% SC	0.3 ml / l
Azadirachtin	Neemix	EC	2 ml / l

Karma Stickless, a registered variety of cowpea with maturity 45 days and productivity 15t/ha, was planted in a plot of size 2.4 m x 2.4 m. The plots and blocks were spaced at 1 m distance each. Cowpea was sown at depth of 2-5 cm with 2 seeds per hole which was thinned after emergence to 1 seedling/hole (Joshi, et al., 2017; Ogunwolu, 1990). Gap filling to maintain optimum plant population was done 1 week after emergence (Maina et al., 2013). The spacing was 60 cm x 30 cm (Jakusko et al., 2013). Fertilizers were applied at the recommended dose and the field was kept weed free for 25 days by manual weeding (Kumar and Singh, 2004). Field was irrigated immediately after sowing and at every 3 days.

Ten flowers were randomly selected, excluding selection from the 5 randomly sampled plants, from each plot at 3 days interval and the number of infected flowers were counted. Also, the number of larvae in each infected flower were recorded. The flowers which contained the live

pod borer larvae or feeding damage to flower were considered the infected flower. Five plants from each treatment were randomly sampled and the total yield from these sampled plants were converted to Kg/ha. The damage to the pod by Pod borer larvae was quantified by taking the weight of infected pod i.e. the pod containing either live larvae or borer holes, from sampled plants of each treatment. The number of bugs were counted directly from the sampled plants of each treatments.

All the data collected were tabulated and drawn in Microsoft Excel. Before analysis, all the count data i.e. number of infected flowers, number of larvae per flower and number of bugs per sample plant were square root transformed. The data were analyzed using R, with analysis of variance (ANOVA). And wherever significance difference occurred, Duncan's Multiple Range test (DMRT) was done for multiple comparison and the separation and grouping of means. Also, percentage reduction in number of infected flowers for different treatments over control was calculated using modified Abbott's formula, which is as:

$$PROC = \left(1 - \left(\frac{Ta \times Cb}{Tb \times Ca} \right) \right) \times 100$$

Where,

PROC = Percentage reduction over control,

Ta = Population in treatment after spray,

Ca = Population in control after spray,

Tb = Population in treatment before spray,

Cb = Population in control before spray (Fleming & Ratnakaran, 1985)

Also, the marginal benefit cost ratio was calculated based on market price of cowpea, and the cost treatments (insecticide cost and spray cost). This ratio was calculated based on the formula used by Rahman et al. (2014), which is given by:

$$\text{Marginal BCR} = \frac{\text{Benefit over Control}}{\text{Cost of Treatment}}$$

3. RESULTS AND DISCUSSION

3.1 Effect of different treatments on flower incidence of *Maruca pod borer*

Three days after the first treatment, the mean number of infected flowers among different treatments varied significantly. Chlorpyrifos 50 + Cypermethrin 5 and Spinosad performed well and were statistically different ($p < 0.05$) from all other treatments. The lowest number of infected flower (0.5) was observed in Chlorpyrifos 50 + Cypermethrin 5 treated plot which was statistically identical ($p > 0.05$) to Spinosad treated plot. Whereas Neemix and Cowpea/Sorghum intercrop remained statistically similar ($p > 0.05$) compared to the control. Three days following first treatment, about 83 % reduction in number of infected flowers over control was observed in Chlorpyrifos 50 + Cypermethrin 5 treated plot and 60 % reduction over control was observed in Spinosad treated plot (Table 3).

Similarly, the mean number of borer infected flowers were highly significant between treatments both at three days and six days following the second treatment. The lowest number of infected flower (0.85) was observed in Spinosad which was statistically significant over control ($p < 0.05$). Neemix was statistically similar ($p > 0.05$) to Spinosad and also showed statistically significant ($p < 0.05$) effect compared to control. However, Chlorpyrifos 50 + Cypermethrin 5 and Cowpea/Sorghum intercrop were statistically similar to control (Table 2). Also, we observed about 86 % reduction in number of infected flowers over control in Spinosad treated plot and 72 % reduction over control in Neemix treated plot (Table 3).

Even after six days of second treatment, the most effective treatment was found as Spinosad which was statistically significant compared to control and cowpea/sorghum intercrop ($p < 0.05$) (Table 2). With Spinosad there was highest i.e. about 85 % reduction in number of infected flowers over control followed by Neemix (58 %) (Table 3).

Similar results were obtained by Randhawa & Saini (2015), as the author reported that Spinosad proved the most effective in reducing flower incidence of *Maruca vitrata*. Ameta *et al.* (2011) also reported that Spinosad brought significantly higher reduction in larval population and minimum flower and pod damage due to pod borers i.e. *Helicoverpa armigera* (Hubner) and *Maruca testulalis* (L.).

Table 2: Mean number of infected flowers on indicated days of treatments.

Treatments	DB1T	3DA1T	6DA1T	9DA1T	3DA2T	6DA2T
Control	1.83	1.85a	2.49	1.93	2.19a	2.21a
Chlorpyrifos 50 + Cypermethrin 5	1.62	0.5b	1.84	1.86	1.83abc	1.25ab
Spinosad	1.54	0.71b	2.38	2.27	0.85c	0.85b
Neemix	1.98	1.91a	2.17	2.13	1.14bc	1.54ab
Cowpea/Sorghum Intercrop	2.1	2.39a	2.33	2.44	2.09ab	2.04a
F Value	0.911 ns	11.22	1.559 ns	2.061 ns	3.75	3.49
C.V	27.61	33.39	17.92	15.76	37.86	38.59
P	0.488	0.0005	0.248	0.149	0.0333	0.0412

Means within a column with no letters in common are significantly different (Duncan's Multiple Range Test (DMRT), $p < 0.05$)

DB1T = Day before first treatment, 3DA1T = 3 Days after first treatment

6DA1T = 6 Days after first treatment, 9DA1T = 9 Days after first treatment

3DA2T = 3 Days after second treatment, 6DA2T = 6 Days after second treatment

ns = Non-Significant at $\alpha = 5\%$ ($p > 0.05$)

Table 3: Percentage reduction in number of infected flowers over control for different treatments.

Treatments	Percentage Reduction Over Control (PROC)		
	3DA1T	3DA2T	6DA2T
Chlorpyrifos 50 + Cypermethrin 5	83 %	25 %	49 %
Spinosad	60 %	86 %	85 %
Neemix	6 %	72 %	58 %
Cowpea/Sorghum intercrop	-28 %	44 %	44 %

3.2 Effect of different treatments on number of pod borer larvae per flower

Number of pod borer (*Maruca testulalis*, Geyer) larvae per flower varied significantly three days following the first spray. The lowest number of larvae per flower (0.16) was observed for Chlorpyrifos 50 + Cypermethrin 5 which is statistically at par with Spinosad (Table 4). Whereas Neemix and cowpea/sorghum intercrop showed very less effect on number of larvae per flower which were statistically similar to control. Three days after first treatment, Chlorpyrifos 50 + Cypermethrin 5 was found to reduce the number of larvae per flower by 88 % over control and Spinosad reduced the number of larvae per flower by 60 % over control (Table 5).

However, three days following the second treatment Spinosad showed the most promising effect with the lowest number of pod borer larvae per flower (0.26) and was statistically different from other treatments ($p < 0.05$). Neemix which was statistically similar to Spinosad also showed remarkable effect (Table 4). We observed about 89 % reduction in number of larvae per flower over control with Spinosad and 79 % reduction over control with Neemix (Table 5).

The number of larvae per flower varied significantly after six days of second treatment. Spinosad remained the most effective treatment in

controlling the pod borer larvae with the least number of larvae per flower (0.26) followed by Chlorpyrifos 50 + Cypermethrin 5 which is statistical at par with Neemix. However, cowpea/sorghum intercrop, as always, seemed ineffective against the pod borer and remained statistically insignificant with control ($p > 0.05$).

The above finding is coherent with the finding of Randhawa & Saini (2015) who reported Spinosad was responsible for maximum decrease in larval population of *Maruca vitrata*.

Table 4: Number of Larvae per flower on indicated days of treatments.

Treatments	DB1T	3DA1T	6DA1T	9DA1T	3DA2T	6DA2T
Control	0.66	0.63 a	1.04	0.63	0.71 a	0.69 a
Chlorpyrifos 50 + Cypermethrin 5	0.59	0.16 b	0.65	0.67	0.60 ab	0.39 ab
Spinosad	0.58	0.24 b	0.85	0.84	0.26 c	0.26 b
Neemix	0.74	0.65 a	0.86	0.78	0.36 bc	0.51 ab
Cowpea/Sorghum Intercrop	0.71	0.85 a	1.04	0.88	0.69 a	0.70 a
F Value	0.251 ns	9.129	2.582 ns	2.529 ns	4.554	3.416
C.V	39.63	38.25	22.94	17.71	35.83	39.56
P	0.903	0.0012	0.091	0.095	0.018	0.043

Means within a column with no letters in common are significantly different (Duncan's Multiple Range Test (DMRT), $p < 0.05$)

ns = Non-Significant at $\alpha = 5\%$ ($p > 0.05$)

Table 5: Percentage reduction in number of larvae per flowers over control for different treatments.

Treatments	Percentage Reduction Over Control (PROC)		
	3DA1T	3DA2T	6DA2T
Chlorpyrifos 50 + Cypermethrin 5	88%	37%	60%
Spinosad	60%	89%	89%
Neemix	7%	79%	65%
Cowpea/Sorghum intercrop	-78%	52%	50%

From the above results, Spinosad was the most effective management option against the pod borer (*Maruca testulalis* Geyer). We also observed that neemix also performed well and at many times it performed statistically at par with the chemical insecticide (Chlorpyrifos 50 + Cypermethrin 5). The percentage reduction of both flower infection and larval population per flower over control remained higher in neemix. We also observed that the Neemix performed more effective after second spray compared to first spray. Schmutterer suggests that botanical pesticides like neem have limited persistence in environment and repeated application and higher concentration is necessary for desired result (Schmutterer, 1990). Also, neem derivatives shows delayed effect compared to many other synthetic chemical insecticides.

3.3 Effect of different treatments on number of Riptortus pod bug per plant

Three days following the first treatment, all the treatments were significantly effective against the pod bug compared to the control. Chlorpyrifos 50 + Cypermethrin 5 was the most effective with no pod bug infestation in the treated plot and was statistically significant with all other treatments ($p < 0.05$). It was followed by Spinosad in its effectiveness with mean number of pod bugs per sampled plant being 0.67 and was statistically at par with neemix and cowpea/sorghum intercrop ($p > 0.05$) (Table 6). In terms of percentage reduction over control, we found that

cowpea/sorghum intercrop reduced the number of pod bugs by 92 % and Spinosad reduced it by 75 % over control (Table 7).

Similarly, the mean number of pod bugs (*Riptortus dentipes*) per sampled plant varied significantly at three days and six days after the second treatment. The lowest number of pod bugs were observed in Chlorpyrifos 50 + Cypermethrin 5 treated plots in both 3 days and 6 days after second treatment. Spinosad being statistically different to control ($p < 0.05$) also performed statistically at par with Chlorpyrifos 50 + Cypermethrin 5 at both 3 days and 6 days following the second treatment ($p > 0.05$). However, Neemix and cowpea/sorghum intercrop remained ineffective and were statistically similar to control ($p > 0.05$) both at 3 days and 6 days following the second treatment.

Chlorpyrifos 50 + Cypermethrin 5 reduced the number of pod bugs per sampled plant compared to control by 66 % at 3 days after and by 74 % at 6 days after the second treatment. It was followed by Spinosad which reduced the number of pod bugs per sampled plant compared to control by 47 % and 56 % at 3 days and 6 days following the second treatment (Table 7).

It showed that Chlorpyrifos 50 + Cypermethrin 5 was the most effective in management of pod bug (*Riptortus dentipes*). Spinosad was also effective against the pod bug being statistically similar to Chlorpyrifos 50 + Cypermethrin 5 at many times. Both of these showed statistically significant effect on pod bugs over control most of the time. A group researchers concluded that broad spectrum insecticides like chlorpyrifos were highly effective in suppressing the pod bug population and can be used for successful management of pod bugs in pigeon pea (Gopali et al., 2013). Similarly, some researchers reported that cypermethrin was more effective in decreasing pest densities in cowpea (Opolot et al., 2006). In other studies also demonstrated that cypermethrin was effective against the insect pests cowpea as the population of many targeted insects including pod sucking bug complex was found to be decreased (Mukendi et al., 2019). Narasimhamurthy and Keval revealed that spinosad recorded the highest percent population reduction of pod bug in pigeon pea (Narasimhamurthy and Keval, 2013).

Table 6: Number of pod bugs per sampled plant on indicated days of treatments.

TREATMENTS	DB1T	3Da1T	6Da1t	9DA1T	3DA2t	6DA2T
CONTROL	0.16	1.31 a	1.23	0.86	1.05 a	1.26 a
CHLORPYRIFOS 50 + CYPERMETHRIN 5	0	0 c	0.44	0.67	0.46 b	0.49 c
SPINOSAD	0.27	0.67 b	0.74	0.61	0.61 b	0.67 bc
NEEMIX	0.11	0.83 b	1.01	0.82	1.07 a	1.05 ab
COWPEA/SORGHUM INTERCROP	0.54	0.76 b	0.66	0.38	0.99 a	1.34 a
F VALUE	1.65 ns	11.07	2.519 ns	0.842 ns	5.972	6.045
C.V	148.87	39.56	47.49	62.23	27.64	31.01
P	0.227	0.0005	0.097	0.525	0.00698	0.00667

Means within a column with no letters in common are significantly different (Duncan's Multiple Range Test (DMRT), $p < 0.05$)

ns = Non-Significant at $\alpha = 5\%$ ($p > 0.05$)

Table 7: Percentage reduction in number of pod bugs per sampled plant over control for different treatments.

Treatments	Percentage reduction over control (PROC)		
	3DA1T	3DA2T	6DA2T
Chlorpyrifos 50 + Cypermethrin 5	-	66%	74%
Spinosad	75%	47%	56%
Neemix	24%	-16%	22%
Cowpea/Sorghum intercrop	92%	-264%	-336%

Cowpea/sorghum intercropping showed negative effect and was found to attract insect (Table 7). Similar finding was reported revealed that the infestation of pod borer and pod sucking bugs and there damage were significantly higher in cowpea/sorghum intercrop (Nampala et al., 2002). Ezueh also reported that intercropping cowpea with cereals like maize and sorghum has been found to increase Maruca borer and pod sucking bugs infestation (Ezueh, 1991). Gethi and Khaemba also recorded that cowpea combined with cereals such as maize and sorghum was more preferred and fed by pod bugs than that of pure cowpea stand and concluded that cowpea/cereal intercrop is undesirable for pod bug control (Gethi and Khaemba, 1991). A group researchers found that there was significantly higher pod bug and pod borer infestation and damage in cowpea/sorghum intercrop (Adipala et al., 2000).

3.4 Yield

It is evident from Table 8 that the mean total weight (Kg/ha), mean weight of infected pod (Kg/ha), mean weight of uninfected pod (Kg/ha) and percentage fruit infestation by weight are all highly significant. The highest total yield was recorded in Spinosad (16,943.2 Kg/ha) which is statistically at par with other treatments except Cowpea/sorghum intercrop which showed the lowest yield (5,249.7 Kg/ha) being statistically significant to other treatments ($p < 0.05$). Similarly, the highest weight of uninfected pod was also recorded in Spinosad (14,216 Kg/ha) which was statistically different to all other treatments ($p < 0.05$). The lowest weight of uninfected pod was recorded in Cowpea/sorghum intercrop (2,695.4 Kg/ha) which was statistically similar to control ($p > 0.05$).

Spinosad treated plot showed the lowest weight of infected pod (2,726.5 Kg/ha) and the percentage fruit infestation by weight was also lowest (16.2 %) and statistically significant to other treatment in spinosad treated plot ($p < 0.05$). However, though the weight of infected pod for Cowpea/sorghum intercrop is low and statistically similar to Spinosad, the percentage fruit infestation by weight is significantly higher than spinosad and is similar to control. Low weight of infected pod in cowpea/sorghum intercrop was only the result of its lowest total yield and not due to its effectiveness. Also, we can observe that the effectiveness of neemix and Chlorpyrifos 50 + Cypermethrin 5 was statistically similar to each other (Table 8).

The above findings corroborate with that of Randhawa and Saini, who reported that spinosad accounted to the minimum damage to the pod and highest grain yield in pigeon pea (Randhawa and Saini, 2015). Similar findings were reported recorded the lowest pod damage and highest grain yield in spinosad treated plots against cowpea spotted pod borer (Kaushik et al., 2016). In other study also recorded similar results who concluded that spinosad treated plants showed the lowest pod damage, lowest seed damage and highest grain yield (Rao et al., 2007). Sahoo also revealed that spinosad applied against insect pests of pigeon pea including pod borer and pod sucking bugs resulted lowest pod damage, lowest seed damage, lowest seed loss and higher seed yield (Sahoo, 2014).

Table 8: Effect of different treatments in yield of cowpea.

TREATMENTS	TOTAL YIELD (KG/HA)	WEIGHT OF INFECTED POD (KG/HA)	WEIGHT OF UNINFECTED POD (KG/HA)	% FRUIT INFESTATION BY WEIGHT
CONTROL	11,507.7 a	5,332.5 ab	6,175.2 bc	47.8 % a
CHLORPYRIPHOS 50 + CYPERMETHRIN 5	16,744.5 a	7,230.75 a	9,513.7 b	43.6 % a
SPINOSAD	16,943.2 a	2,726.50 c	14,216.7 a	16.2 % b
NEEMIX	12,636.2 a	4,791.25 bc	7,845.0 b	38.8 % a
COWPEA/ SORGHUM INTERCROP	5,249.7 b	2,554.35 c	2,695.4 c	47.5 % a
F VALUE	7.76	7.86	8.576	10.6
P	0.0043	0.0023	0.0016	0.00063

Means within a column with no letters in common are significantly different (Duncan's Multiple Range Test (DMRT), $p < 0.05$)

Figure 1: Effect of different treatments on Percentage of infestation of pod by weight.

3.5 Marginal Benefit : Cost Ratio

The highest net income was recorded in Spinosad (Rs. 3,00,750 per ha.) followed by Chlorpyrifos 50 + Cypermethrin 5 (Rs. 1,27,750 per ha.). The marginal benefit cost ratio showed that the highest profit per unit rupee spent over control (22.06) was obtained in Chlorpyrifos 50 + Cypermethrin 5 treatment followed by Spinosad (14.38). The net income of Rs. 59,750 per ha. was obtained from Neemix treated plot with marginal benefit cost ratio of 8.47. Cowpea/sorghum intercrop's performance was even lower than that of control (Table 9).

Table 9: Net income and marginal benefit cost ratio of different treatments.

TREATMENTS	UNINFECTED POD WEIGHT (KG/HA)	ADDITIONAL YIELD OVER CONTROL (KG/HA)	ADDITIONAL INCOME OVER CONTROL (RS./HA)	COST OF TREATMENT	NET INCOME (RS./HA)	MBCR
CONTROL	6,175.2	—	—	—	—	—
CHLORPYRIPHOS 50 + CYPERMETHRIN 5	9,513.7	3,338.5	1,33,540	5,790	1,27,750	22.06
SPINOSAD	14,216.7	8,041.5	3,21,660	20,910	3,00,750	14.38
NEEMIX	7,845	1,669.7	66,790	7,050	59,740	8.47
COWPEA / SORGHUM INTERCROP	458	-5,717.2	—	—	—	—

Note: Farm gate price of cowpea = Rs. 40/Kg. Cost of chlorpyrifos 50 + Cypermethrin 5 = Rs. 1200/liter, required amount (for 3 spray @ 2ml/l) = 4.2 liter/ha, Labor cost = Rs. 750, Total cost for Chlorpyrifos 50 + Cypermethrin 5 = Rs.5,790 (Rs. 1200 * 4.2 + Rs. 750). Cost of Spinosad = Rs. 32/ ml, required amount (3 spray @ 0.3 ml/l) = 630 ml/ha, Labor cost = Rs. 750, Total cost for Spinosad = Rs. 20,910 (Rs. 32 * 630 + Rs. 750) Cost of Neemix = Rs. 1500 / liter, required amount (3 spray @ 2 ml/liter) = 4.2 liter/ha, Labour Cost = Rs. 750, Total cost for neemix = Rs. 7,050 (Rs. 1500 * 4.2 + Rs. 750)

4. CONCLUSION

The result obtained from this study demonstrates that Spinosad was the most effective in managing pod borer (*Maruca testulalis* Geyer) in cowpea through reducing both the number of infected flower (flower incidence) and the larval population i.e. number of larvae per flower. Spinosad was also effective in reducing pod damage and accounts highest yield, lowest weight of infected pods, thus providing highest net return per hectare. Neemix, a botanical pesticide also performed impressively against *Maruca* pod borer. Chlorpyrifos 50 + Cypermethrin 5 proved the most effective in controlling the pod sucking bug (*Riptortus dentipes*) but Spinosad was also equally effective against it. However, the cowpea/sorghum intercrop was found ineffective, rather was found to aggravate the situation of insects incidence.

Spinosad has been reported as safe, environmental-friendly and less hazardous to human in many reports. And the results from this study showed Spinosad very effective against both cowpea pod bug and pod borer. So, the study has concluded that Spinosad (0.3 ml / liter of water) is effective against the test insects. Also, it has been recommended that the use of Spinosad is a good approach over other synthetic insecticides for the control of cowpea pod borer and pod sucking bugs. However, it has been suggested to conduct large scale, multilocation and multiyear studies and field trials before wider application of this technique. This study also

shows that Neemix, a neem-based pesticide can be a potential, environment friendly and sustainable option for insect pest management and suggests intensive studies.

ACKNOWLEDGEMENTS

We would like to express our very great appreciation to Assistant Professor Mr. Rajendra Regmi for his valuable and constructive suggestions during the planning and development of this research work.

REFERENCES

- Adipala, E., Nampala, P., Karungi, J., Isubikslu, P., 2000. A review on options for management of cowpea pests: experience from uganda. Integrated pest management review, 5 (3), Pp. 185-196.
- Ameta, O., Sharma, U.S., Jeengar, K., 2011. Efficacy of flubendiamide 480 SC against pod borers, *Helicoverpa armigera* (Hubner) and *Maruca testulalis* (L.) in pigeon pea. Indian Journal of Entomology, Pp. 191-195.
- Baoua, I., Ba, N., Agunbiade, T., Margam, V., Binso-Dabire, C., Antoine, S., Pittendrigh, B., 2011. Potential Use of *Sesbania pachycarpa* (Fabaceae: Papilionoidae) as a refugia for legume pod borer *Maruca Vitrata* (Lepidoptera: Carambidae). International Journal of Tropical Insect Science, 31 (4), Pp. 212-218.
- Bett, B., Gollasch, S., Moore, A., James, W., Armstrong, J., Walsh, T., Higgins, T.J., 2017, November. Transgenic cowpeas (*Vigna unguiculata* L. Walp) expressing *Bacillus thuringiensis* Vip3Ba protein are protected against the *Maruca* pod borer (*Maruca vitrata*). Plant Cell Tissue and Organ Culture (PCTOC), 131 (2), Pp. 335-345.
- Choudhary, A.L., Hussain, A., Samota, R., Nehra, S., 2017. Effect of biotic and abiotic factors on the incidence of aphid, *Aphis cracivora* Koch on Cowpea. Journal of Pharmacognosy and Phytochemistry, 6 (4), Pp. 1587-1590.
- Ekesi, S., 1999. Insecticide resistance in field populations of the legume pod-borer, *Maruca vitrata* Fabricius (Lepidoptera: Pyralidae), on cowpea, *Vigna unguiculata* (L.), Walp in Nigeria. International Journal of Pest Management, 45 (1), Pp. 57-59. doi:10.1080/096708799228058
- Ezueh, M., 1991. Prospects for Cultural and Biological Control of Cowpea Pests. International Journal of Tropical Insect Science, 12 (5-6), Pp. 585-592.
- Fleming, R., Ratnakaran, A., 1985. Evaluating Single Treatment Data Using Abbott's Formula With Reference to Insecticides. Journal of Economic Entomology, Pp. 1179-1181.
- Gethi, M., Khaemba, B., 1991. Damage by pod-sucking bugs on cowpea when intercropped with maize. Tropical Pest Management, 37 (3), Pp. 236-239. doi:10.1080/09670879109371591
- Gopali, J., Sharma, O., Yelshetty, S., Haveri, R.V., 2013. Effect of insecticides and biorationals against pod bug (*Clavigralla gibbosa*) in pigeonpea. Indian Journal of Agricultural Science, Pp. 582-585.
- Jackai, L., Atropo, P.K., Odebiyi, J.A., 1989. Use of the response of two growth stages of cowpea to different population densities of the coreid bug, *Clavigralla tomentosicollis* (Stål.) to determine action threshold levels. Crop Protection, 8 (6), Pp. 422-428.
- Jackai, L., Daoust, R., 2003. Insect Pests of Cowpeas. Annual Review of Entomology, 31, Pp. 95-119.
- Jackai, L.E., 2009. Studies on the feeding behaviour of *Cfavigralla tomentosicollis* (Stal.) (Hemiptera, Coreidae) and their potential use in bioassays for host plant resistance. Journal of Applied Entomology, Pp. 344-350. doi:10.1111/j.1439-0418.1984.tb02721.x
- Jakusko, B.B., Anasunda, U.I., Mustapha, A.B., 2013. Effect of inter-row spacing on some selected cowpea (*Vigna unguiculata* (L.) varieties in Yola, Adamawa State, Nigeria. IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS), 2 (3), Pp. 30-35.
- Joshi, B.K., Bhatta, M.R., Ghimire, K.H., Khanal, M., Gurung, B.S., Dhakal, R., Shapit, B., 2017. Released and Promising Crop Varieties of Mountain Agriculture in Nepal (1959-2016). LI-BIRD, Pokhara; NARC, Kathmandu and Bioversity International, Pokhara, Nepal.
- Kamara, A.Y., Chikoye, D., Omoigui, L., Dugje, I.Y., 2007. Influence of insecticide spraying regimes and cultivar on insect pests and yield of cowpea in the dry savannas of north-eastern Nigeria. Journal of Food, Agriculture & Environment, 5 (1), Pp. 154-158. doi:DOI: 10.13140/2.1.2385.9848
- Kaushik, A.K., Yadav, S.K., Srivastava, P., 2016. Field efficacy of insecticides and mixture against spotted pod borer, *Maruca Vitrata* Fabricius on cowpea. Ann. Pl. Protec. Sci., 24 (1), Pp. 89-92.
- Kumar, D., Singh, N.B., 2004. Cowpea in India. India: Scientific Publishers, India.
- Maina, Y.T., Degri, M.M., Sharah, H.A., 2013. Control of Pod-sucking bug *Riptortus dentipes* (Hemiptera: Alydidae) of Cowpea with aqueous Plant Extracts and Cymbush Super EC in Maiduguri, Nigeira. Journal of Health, Medicine and Nursing, 2, Pp. 8-12.
- Mukendi, R.T., Kanyenga, A.L., Baboy, L.L., Bugeme, D.M., Kalonji, A.M., Munyuli, T.M., 2019. Insecticide Spraying Regime to Control Insect Pest of Cowpea: Management and Monitored Application of Cypermethrin In Lomami Province, Dr Congo. Current Research in Agricultural Sciences, 6 (1), Pp. 37-53.
- Nampala, P., Ogenga-Latigo, M., Kyamanywa, S., Adipala, E., Oyobo, N., Jackai, L., 2002. Potential Impact of Intercropping on Major Cowpea Field Pests in Uganda. African Crop Science Journal, 10, Pp. 335-344.
- Narasimhamurthy, G., Keval, R., 2013. Field evaluation of some insecticides and bio-pesticides against pod bug, *Clavigralla gibbosa* (Spinola) in long duration pigeon pea. African Journal of Agricultural Research, 8 (38), Pp. 4876-4881.
- Nyiira, Z.M., 1971. The Status of Insect Pests of Cowpea (*vigna unguiculata* (L) Walp. in Uganda and their control. PANS Pest Articles & News Summaries, 17 (2), Pp. 194-197.
- Ogunwolu, E.O., 1990. Damage to cowpea by the legume pod borer, *Maruca testulalis* Geyer, as influenced by infestation density in Nigeria. Tropical Pest Management, 36 (2), Pp. 138-140. doi:10.1080/09670879009371457
- Okeyo-Owuor, J.B., Agwaro, P.O., Simbi, C.O., 1983. Studies on The Legume Pod-Borer *Maruca Testulalis* (Geyer)—V. Larval Population. International Journal of Tropical Insect Science, 4 (1-2), Pp. 75-81. doi: 10.1017/S1742758400004057
- Opolot, H., Agona, A., Kyamanywa, S., Mbata, G., Adipala, E., 2006. Integrated field management of cowpea pests using selected synthetic and botanical pesticides. Crop Protection - Crop Prot, 25, Pp. 1145-1152. doi: 10.1016/j.cropro.2005.03.019
- Oyewale, R.O., Bamaayi, L.J., 2013. Management of Cowpea Insect Pests. Scholars Academic Journal of Biosciences, 1 (5), Pp. 217-226.
- Rahman, A., Haquer, M., Alam, S., Mahmudunnabi, M., Dutta, N., 2014. Efficacy of Microbials as Insecticides for the management of Tomato (*Lycopersicon esculentum*) Fruitworm, *Helicoverpa armigera* (Hubner). The Agriculturists, 12 (1), Pp. 68-74.

- Randhawa, H., Saini, M., 2015. Efficacy of different insecticides against pod borer, (*Maruca vitrata* Geyer) in pigeon pea. Legume Research - An International Journal.
- Rao, G., Kumari, P., Rao, V., Reddy, Y., 2007. Evaluation of Spinosad and Indoxacarb for the management of legume pod borer, *Maruca vitrata* (Geyer) in pigeonpea. Journal of food legumes, Pp.126-127.
- Sahoo, G., 2014. Bio-Efficacy of ecofriendly pesticide in the suppression of insect pest in pigeon pea. College of Agriculture, Orissa University of Agriculture and Technology, Department of Entomology. Bhubaneswar: Orissa University of Agriculture and Technology. Retrieved May 15, 2020, from <https://krishikosh.egranth.ac.in/displaybitstream?handle=1/5810084082>
- Sardana, H.R., Verma, S., 1986. Preliminary studies on the prevalence of insect pests and their natural enemies on cowpea crop in relation to weather factors at Delhi. Indian Journal of Entomology, 48 (4), Pp.448-458.
- Schmutterer, H., 1990. Properties and potential of natural pesticides from the neem tree, *Azadirachta indica*. Annu. Rev. Entomol., Pp. 271-297.
- Shinde, K.G., Kumud, V.N., Raut, P.P., Desai, V.S., Mehendale, S.K., 2017. Biology of Pod Borer, *Maruca vitrata* (Geyer) Infesting Lablab Bean. . Int. J. Curr. Microbiol. App. Sci., 6 (9), Pp. 67-74. doi: <https://doi.org/10.20546/ijcmas.2017.609.007>
- Sreelakshmi, P., 2014. Insecticide Resistance In Field Populations Of The Cowpea Spotted Pod Borer, *Maruca Vitrata* Fabricius (Lepidoptera-Pyralidae)